

Technical University of Denmark



## Silent clusters

speak up!

**Gram, Lone**

*Published in:*  
Microbial Biotechnology

*Link to article, DOI:*  
[10.1111/1751-7915.12181](https://doi.org/10.1111/1751-7915.12181)

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Gram, L. (2014). Silent clusters: speak up! Microbial Biotechnology, 8(1), 13-14. DOI: 10.1111/1751-7915.12181

## DTU Library

Technical Information Center of Denmark

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## Silent clusters – speak up!

Lone Gram\*

Department of Systems Biology, Technical University of Denmark, Matematiktorvet bldg. 301, DK-2800 Kgs. Lyngby, Denmark.

Microorganisms have provided mankind with a multitude of useful compounds ranging from industrial enzymes to anti-cancer compounds and antibiotics. With the aging population, our need for treatment principles against cancer, Alzheimer's disease and metabolic disorders is increasing. Also, the occurrence and spread of antibiotic resistance in pathogenic microorganisms has become one of our main challenges. For a number of years, it was hoped that our advances in chemical synthesis would provide the desired novelty in terms of scaffolds and targets, however, while there has been success, it has also been realized that the complexity of natural (microbial) products far surpasses anything possible by chemical synthesis. Hence, drug discovery has returned to nature, especially microorganisms, as a source of novel chemistry. Recently, genome sequencing and mining has demonstrated that bioactive compounds hitherto believed to be produced by a eukaryotic organism (a marine sponge) were indeed synthesized by one of the dominant (uncultured) prokaryotes, '*Entothoonella*' further underlining the vast bioactive underestimated potential in microorganisms (Wilson *et al.*, 2014).

Classical bioassay guided fractionation has been used for decades as the major bioprospecting path, however, this is often a laborious process, and rediscovery of known compounds is a major challenge despite developments in de-replication strategies. The rapidly developing sequencing techniques are introducing sequencing of microbial genomes in almost any microbial research environment, and hence so-called genome mining has become a useful tool in discovery of novel microbial products (Jensen *et al.*, 2014). Despite the complexity of the bioactive secondary metabolites produced by microorganisms, several pathways are conserved at the modular level. This has allowed the development of software, such as ANTISMASH or NAPDoS, which can identify

characteristic structures of polyketide and non-ribosomal peptide synthetases (Weber 2014). Such genome mining processes have revealed that the bioactive potential in numerous microorganisms is likely much higher than previously discovered by bioassay-guided fractionation. This has been demonstrated for well-known producers of antibiotics, in *Actinobacteria* and fungi. However, it is also true for several groups of Gram-negative prokaryotes (Still *et al.*, 2014) that hitherto have not been thought of as useful producers of bioactive compounds. For instance in *Pseudoalteromonas luteoviolaceae*, 3–4 bioactive compounds have been isolated (violacein, indolmycin, pentabromopseudilin), however, genome mining reveals approximately 15 gene clusters likely encoding for pathways likely producing bioactives (H. Machado, unpublished).

It is believed that the reason for the 'non-discovery' of these potential novel compounds is largely because the genes are not expressed during normal laboratory cultivation; so-called silent gene clusters. Hence, the activation or use of such silent clusters is becoming a hot research topic. How to unlock this potential? How to make the silent clusters 'speak up'? Several paths are being pursued. One is based on synthetic biology with the approach of cloning the clusters and expressing in heterologous hosts. There are examples of success from this approach (Ross *et al.*, 2014), and more are likely to come. However, there are also several challenges. The gene clusters encoding putative bioactive compounds are typically large (10–100 kb) and often consists of many genes. Hence, the cloning and expression is significantly more challenging than cloning of a single gene and expressing its product. Also, the clusters/genes may fail to express in a heterologous host due to lack of signals naturally present in the native producer.

Another path involves the modification of gene expression in the natural producer organism in a general sense. In fungi, inhibitors of histone deacetylation can induce secondary metabolism by increasing promoter accessibility due to changes in chromatin structure. Analogously, expression of otherwise silent gene clusters in *Streptomyces coelicolor* has been demonstrated using histone deacetylation inhibitors as elicitors (Moore *et al.*, 2012). Japanese researchers have successfully used genetic modification of the producer microorganism to elicit bioactive compound production. Approaches include 'ribosome engineering' as well as overexpression of regulator proteins (Ochi and Hosaka, 2013).

Received 2 September, 2014; accepted 8 September, 2014.

\*For correspondence. E-mail gram@bio.dtu.dk; Tel. +45 45252586; Fax +45 45252501.

*Microbial Biotechnology* (2015) 8(1), 13–14

doi:10.1111/1751-7915.12181

**Funding Information** No funding information provided.

© 2014 The Author. *Microbial Biotechnology* published by John Wiley & Sons Ltd and Society for Applied Microbiology.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

A broad range of approaches all fall under the same overall strategy, involving variation of growth conditions of the potential producing organism with subsequent chemical analyses (LC-MS, MS/MS) and bioassays. A vast amount of studies have demonstrated that the production of secondary metabolites in microorganisms is dependent on growth substrate and environment. Stress signals, such as antibiotics (Seyedsayamdost, 2014), ppGpp, nutrient depletion or physical stress can induce silent clusters and production of bioactives. Also, several groups are working on culture approaches where the conditions are mimicking the natural chemical and/or biological environment of the potential producing microorganisms (Schroeckh *et al.*, 2009; Lincke *et al.*, 2010). This also covers switching from test tubes and flasks to using biofilm mode of growth, co-cultivation systems (König *et al.*, 2013) and natural co-occurring nutrients. Especially, the production of novel compounds as a result of co-culturing has been beautifully demonstrated by use of direct chemical probing (nanospray desorption electrospray ionization and microbial matrix-assisted laser desorption ionization-time of flight imaging mass spectrometry) (Traxler *et al.*, 2013). We demonstrated (Wietz *et al.*, 2011) that culturing marine bacteria on chitin, the dominant organic polymer in marine systems, as opposed to glucose dramatically altered the secondary metabolism, and the bacterium focused its production on bioactive compounds. In a simple agar screen for antibiotic activity, we recently found that if the producer organisms were grown on chitin, the number of isolates with antibacterial activity tripled (S. Giubergia, unpublished).

Likely, the future will see a combination of several approaches and technologies. Aligning manipulations of culture conditions (natural substrates, associate microbiota) with chemical (metabolomic molecular networking), genome and metagenome-based analyses is indeed likely to enhance our bio-discovery hit rate and provide novel understanding of the ecology and natural interactions of microorganisms.

### Conflict of interest

Authors have no conflict of interest to declare.

### References

- Jensen, P.R., Chavarria, K.L., Fenical, W., Moore, B.S., and Ziemert, N. (2014) Challenges and triumphs to genomics-based natural product discovery. *J Ind Microbiol Biotechnol* **41**: 203–209.
- König, C.C., Scherlach, K., Schroeckh, V., Horn, F., Nietzche, S., Brakhage, A.A., and Hertweg, C. (2013) Bacterium induces cryptic meroterpenoid pathway in the pathogenic fungus *Aspergillus fumigatus*. *Chem Bio Chem Comm* **14**: 938–942.
- Lincke, T., Behnken, S., Ishida, K., Roth, M., and Hertweg, C. (2010) Clostioamide: an unprecedented polythioamide antibiotic from the strictly anaerobic bacterium *Clostridium cellulolyticum*. *Angew Chem Int Ed Engl* **49**: 2011–2013.
- Moore, J.M., Bradshaw, E., Seipke, R.F., Hutchings, M.I., and McArthur, M. (2012) Use and discovery of chemical elicitors that stimulate biosynthetic gene clusters in *Streptomyces* bacteria. *Methods Enzymol* **517**: 367–385.
- Ochi, K., and Hosaka, T. (2013) New strategies for drug discovery: activation of silent or weakly expressed microbial gene clusters. *Appl Microbiol Biotechnol* **97**: 87–98.
- Ross, A.C., Gulland, L.E.S., Dorrestein, P.C., and Moore, B.S. (2014) Targeted capture and heterologous expression of the pseudoalteromonas alterochromide gene cluster in *Escherichia coli* represents a promising natural product exploratory platform. *ACS Synth Biol* (in press).
- Schroeckh, V., Scherlach, K., Nützmann, H.-W., Shelest, E., Schmidt-Heck, W., Schuemann, J., *et al.* (2009) Intimate bacterial-fungal interaction triggers biosynthesis of archetypal polyketides in *Aspergillus nidulans*. *Proc Natl Acad Sci USA* **106**: 14558–14563.
- Seyedsayamdost, M.R. (2014) High-throughput platform for the discovery of elicitors of silent bacterial gene clusters. *PNAS* **111**: 7266–7271.
- Still, P.C., Johnson, T.A., Theodore, C.M., Loveridge, S.T., and Crewes, P. (2014) Scrutinizing the scaffolds of marine biosynthetics from different source organisms: gram-negative cultured bacterial products enter centre stage. *J Nat Prod* **77**: 690–702.
- Traxler, M.F., Watrous, J.D., Alexandrov, T., Dorrestein, P.C., and Kolter, R. (2013) Interspecies interactions stimulate diversification of the *Streptomyces coelicolor* secreted metabolome. *MBio* **4**: e00459–13.
- Weber, T. (2014) *In silico* tools for the analysis of antibiotic biosynthesis pathways. *Int J Med Microbiol* **304**: 230–235.
- Wietz, M., Månsson, M., and Gram, L. (2011) Chitin stimulates production of the antibiotic andrimid in a *Vibrio corallilyticus* strain. *Environ Microbiol Rep* **3**: 559–564.
- Wilson, M.C., Mori, T., Rückert, C., Uria, A.R., Helf, M.J., Takada, K., *et al.* (2014) An environmental bacterial taxon with a large and distinct metabolic repertoire. *Nature* **506**: 58–62.