brought to you by



Conserved structure and expression of hsp70 paralogs in teleost fishes - DTU Orbit (08/11/2017)

Conserved structure and expression of hsp70 paralogs in teleost fishes

The cytosolic 70 KDa heat shock proteins (Hsp70s) are widely used as biomarkers of environmental stress in ecological and toxicological studies in fish. Here we analyze teleost genome sequences to show that two genes encoding inducible hsp70s (hsp70-1 and hsp70-2) are likely present in all teleost fish. Phylogenetic and synteny analyses indicate that hsp70-1 and hsp70-2 are distinct paralogs that originated prior to the diversification of the teleosts. The promoters of both genes contain a TATA box and conserved heat shock elements (HSEs), but unlike mammalian HSP70s, both genes contain an intron in the 5' UTR. The hsp70-2 gene has undergone tandem duplication in several species. In addition, many other teleost genome assemblies have multiple copies of hsp70-2 present on separate, small, genomic scaffolds. To verify that these represent poorly assembled tandem duplicates, we cloned the genomic region surrounding hsp70-2 in Fundulus heteroclitus and showed that the hsp70-2 gene copies that are on separate scaffolds in the genome assembly are arranged as tandem duplicates. Real-time quantitative PCR of F. heteroclitus genomic DNA indicates that four copies of the hsp70-2 gene are likely present in the F. heteroclitus genome. Comparison of expression patterns in F. heteroclitus and Gasterosteus aculeatus demonstrates that hsp70-2 has a higher fold increase than hsp70-1 following heat shock in gill but not in muscle tissue, revealing a conserved difference in expression patterns between isoforms and tissues. These data indicate that ecological and toxicological studies using hsp70 as a biomarker in teleosts should take this complexity into account

General information

State: Published

Organisations: National Institute of Aquatic Resources, Section for Marine Living Resources, University of British

Columbia

Authors: Metzger, D. C. (Ekstern), Hansen, J. H. (Intern), Schulte, P. M. (Ekstern)

Pages: 10-20

Publication date: 2016

Main Research Area: Technical/natural sciences

Publication information

Journal: Comparative Biochemistry and Physiology. Part D: Genomics and Proteomics

Volume: 18

ISSN (Print): 1744-117X

Ratings:

BFI (2017): BFI-level 1

Web of Science (2017): Indexed Yes

BFI (2016): BFI-level 1

Scopus rating (2016): SJR 0.921 SNIP 0.834 CiteScore 2.73

Web of Science (2016): Indexed yes

BFI (2015): BFI-level 1

Scopus rating (2015): SJR 1.023 SNIP 0.83 CiteScore 2.44

BFI (2014): BFI-level 1

Scopus rating (2014): SJR 0.931 SNIP 0.841 CiteScore 2.4

BFI (2013): BFI-level 1

Scopus rating (2013): SJR 0.814 SNIP 0.737 CiteScore 2.32

ISI indexed (2013): ISI indexed yes

BFI (2012): BFI-level 1

Scopus rating (2012): SJR 0.756 SNIP 0.88 CiteScore 2.52

ISI indexed (2012): ISI indexed yes Web of Science (2012): Indexed yes

BFI (2011): BFI-level 1

Scopus rating (2011): SJR 0.577 SNIP 0.536 CiteScore 1.79

ISI indexed (2011): ISI indexed yes

BFI (2010): BFI-level 1

Scopus rating (2010): SJR 0.673 SNIP 0.599

BFI (2009): BFI-level 1

Scopus rating (2009): SJR 0.606 SNIP 0.485

BFI (2008): BFI-level 1

Scopus rating (2008): SJR 0.432 SNIP 0.305 Scopus rating (2007): SJR 0.321 SNIP 0.28

Scopus rating (2006): SJR 0.108 SNIP 0.015 Original language: English DOIs:

10.1016/j.cbd.2016.01.007 Source: FindIt

Source-ID: 2291668697

Publication: Research - peer-review > Journal article - Annual report year: 2016