

Original Research

Unrestricted Paleolithic Diet is Associated with Unfavorable Changes to Blood Lipids in Healthy Subjects

MICHAEL M. SMITH^{#1}, ERIC T. TREXLER^{*2}, ALLAN J. SOMMER^{†4}, BROOKE E. STARKOFF^{‡3}, STEVEN T. DEVOR^{‡4}

¹ Department of Physical Education and Kinesiology, California State University, Bakersfield, Bakersfield, CA, USA; ² Department of Exercise and Sport Science, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, USA; ³ Department of Kinesiology, Sport Studies, and Physical Education, The College at Brockport, State University of New York, Brockport, New York, USA; ⁴ Department of Human Sciences, The Ohio State University, Columbus, Ohio, USA.

*Denotes undergraduate author, †Denotes graduate student author, ‡Denotes professional author

ABSTRACT

International Journal of Exercise Science 7(2) : 128-139, 2014. The Paleolithic (Paleo) diet is one modeled after the perceived food consumption of early human ancestors of the Paleolithic Era, consisting of mainly meat, fish, fruit, vegetables, eggs, and nuts. The purpose of this study was to examine the effects of a Paleo diet on blood lipids, including high-density lipoprotein (HDL), low-density lipoprotein (LDL), non-HDL cholesterol, triglycerides (TG), total cholesterol (TC), and the ratio between TC and HDL (TC/HDL) in a healthy population. Healthy subjects of both genders (24 males, 20 females) were asked to eat an *ad libitum* Paleo diet for 10 weeks. Prior to the intervention, body weight, body fat percentage (BF%), maximal oxygen consumption (VO₂max), TC, TG, HDL, and LDL were measured. These measurements were repeated following 10 weeks of a Paleo diet. As a whole, there was a significant increase in non-HDL (107.1±6.0 mg/dL to 120.2±6.5 mg/dL; *P*<0.01), LDL (93.1±5.4 mg/dL to 105.6±6.1 mg/dL; *P*<0.01), TC/HDL (3.0±0.2 to 3.3±0.2; *P*<0.05), and TC (168.8±5.4 mg/dL to 178.9±6.6 mg/dL; *P*<0.05) in healthy subjects following a Paleo diet. When stratified into groups based on initial blood lipid levels, deleterious changes were found in those with optimal HDL (82.1±3.2 mg/dL to 68.6±4.8 mg/dL; *P*<0.05), non-HDL (86.6±3.9 mg/dL to 101.4±4.8 mg/dL; *P*<0.01), TC (157.2±0.7 to 168.2±0.9 mg/dL; *P*<0.05), TC/HDL (2.5±0.1 to 2.7±0.1; *P*<0.05), and LDL (69.1±3.1 mg/dL to 83.5±4.1 mg/dL; *P*<0.01), whereas those within sub-optimal stratifications showed no significant changes. Subjects also decreased body weight (80.7±2.6 kg to 77.5±2.4 kg; *P*<0.001) and BF% (24.3±1.2% to 20.7±1.2%; *P* < 0.05). Our results demonstrate that an *ad libitum* unrestricted Paleo diet intervention is associated with deleterious changes to blood lipids in healthy subjects, despite concurrent improvements in body composition and cardiorespiratory fitness. Future research should focus on determining recommendations that embrace the positive aspects of the Paleo diet, while minimizing any deleterious impact on blood lipids in a healthy population.

INTRODUCTION

The Paleolithic (Paleo) diet is modeled after the perceived food consumption of early

human ancestors of the Paleolithic Era. Since the origin of this dietary concept, which dates back to the late 1970s (46), numerous books and articles have claimed

that increasing rates of “diseases of civilization”, including cardiovascular disease, metabolic syndrome, and type 2 diabetes, among others (12, 16), can be largely attributed to the deviation of the modern diet from the diet consumed during the majority of the evolutionary history of humans (10, 12, 46). Many prominent CrossFit athletes advocate adherence to the Paleo diet, which has contributed to a recent boost in the popularity of the diet. The common belief is that the Paleolithic Era pre-dated the advent of the cultivation and processing of plants and domestication of animals (16), though this idea has recently been challenged (35, 40). Researchers have attempted to reconstruct the diet of the early human based on archaeological evidence and studies involving modern hunter-gatherer societies, leading to a proposed early-human diet comprised of wild animal-source foods and uncultivated plant-source foods, devoid of dairy, legumes, and cereal grains (16).

The body of existing literature pertaining to the Paleo diet is characterized by inconsistency. Because the diet is defined by the avoidance of particular food sources rather than a specific macronutrient distribution, there is a large degree of variation in the macronutrient composition of various Paleo diet interventions. One of the earliest peer-reviewed papers on the Paleolithic diet suggests that the diet of the late Paleolithic Era was roughly 33% protein, 46% carbohydrate, and 21% fat (12). Recent interventions, which place more emphasis on food source than macronutrient distribution, have not replicated these suggested intakes. In two studies in which food rations were controlled, one diet derived 31.9% of its

caloric content from fat and 36.9% from carbohydrate (16), while the other derived 16% of calories from fat and 57% from carbohydrate (25).

Some degree of variation in macronutrient intake is to be expected in Paleo studies featuring free-eating subjects, but even the dietary guidelines enforced in such studies have been inconsistent. One study imposed limits on intakes of eggs, nuts, potatoes, and oils (33), one limited dried fruit intake along with the aforementioned foods (26), and another added fatty meats, honey, cured meats, and salted seafood to the list of limited foods, without any mention of egg intake (38). While previous interventions have utilized varying macronutrient distributions and somewhat inconsistent guidelines, the studies uniformly reinforce the basic characteristics of a Paleolithic diet: A diet based on lean meat, fish, eggs, nuts, fruit, and vegetables, and devoid of cereal grains, dairy, legumes, and processed sugars.

Authors of previous Paleo interventions have embraced the potential of the *ad libitum* Paleo diet. Authors of a 2006 study wrote that contemporary groups of hunter-gatherers “Stay lean and apparently reap health benefits similar to those induced by food restriction despite *ad libitum* availability of food.”(25) Another group of researchers found that a Paleo diet intervention yielded an improvement in glucose tolerance that appeared to be independent of energy intake and macronutrient distribution, prompting them to conclude that “...avoiding Western foods is more important than counting calories, fat, carbohydrate or protein.”(33) Although previous trials with free-eating subjects have recognized food sources like

meat, eggs, nuts, potatoes, and certain oils as part of the Paleolithic diet, such trials have imposed guidelines that restrict the consumption of one or more of the aforementioned foods (26, 33, 38).

A truly *ad libitum* application of the Paleo diet, without restrictions on intakes of fatty meat, eggs, nuts, and oils, could potentially be conducive to high intakes of total fat, saturated fat, and cholesterol. This style of dieting, without regard for daily intakes of total fat, saturated fat, or cholesterol, neglects guidelines set by the American Heart Association (AHA) regarding a heart-healthy diet aimed at lowering disease risk and supporting a healthy blood lipid profile (2). While much research has been done on the Paleo diet and the effects it has on overweight/obese and disease populations, there is a lack of research pertaining to the application of the Paleo diet in healthy, active populations. For these reasons, the current study aims to investigate the effects of a practical, *ad libitum* application of the Paleo diet on blood lipids in a sample of healthy, active subjects. To accomplish this objective, we recruited healthy, active adults and measured body composition and blood lipid markers before and after a 10-week Paleo diet. We hypothesized that a Paleo dietary intervention made without specific recommendations regarding macronutrient intake would be associated with an increase in low density lipoprotein (LDL), non-high density lipoprotein (n-HDL), and total cholesterol (TC), but no change in high density lipoprotein (HDL). Furthermore, we hypothesized that this diet would be associated with a decrease in body weight and body fat percentage.

METHODS

Participants

A total of 44 (24 males, 20 females) subjects fully completed the dietary intervention and returned for follow-up testing. All of the subjects provided written informed consent and all study methods and protocols were approved in advance by the Institutional Review Board at The Ohio State University.

Table 1. Subject characteristics

	Males (n=24)	Females (n=20)	Range
Age (years)	33.5 ± 0.3	31.2 ± 0.3	21.0 - 48.0
Height (in)	70.6 ± 0.1	64.8 ± 0.1	60.0 - 77.0
Weight (kg)	90.71 ± 0.25	68.02 ± 0.31	44.54 - 118.18
BMI (kg/m ²)	28.1 ± 0.1	25.1 ± 0.2	19.1 - 37.4
Body fat (%)	22.2 ± 0.3	26.6 ± 0.4	10.7 - 46.1
Lean mass (kg)	70.25 ± 0.37	49.00 ± 0.25	36.35 - 82.17
TC (mg/dL)	171.7 ± 8.7	164.7 ± 6.3	110.0 - 258.0
HDL (mg/dL)	50.6 ± 2.3	72.8 ± 3.8	35.0 - 100.0
LDL (mg/dL)	105.6 ± 7.8	76.8 ± 6.3	33.0 - 191.0
n-HDL (mg/dL)	118.0 ± 9.1	91.9 ± 6.7	40.0 - 204.0
TG (mg/dL)	72.6 ± 9.0	71.8 ± 6.9	45.0 - 226.0
TC/HDL	3.5 ± 0.2	2.3 ± 0.1	1.4 - 5.0

BMI = body mass index; in = inches; kg = kilograms; VO₂max = maximal oxygen consumption. All data are resting values and is presented as mean ± SEM.

Protocol

This study investigated the effect of a 10-week Paleo diet on body composition and blood lipids in healthy, active adults. Body composition using air displacement plethysmography and fasted state blood draws were performed in all subjects in the morning (7:30 a.m. to 11:30 a.m.) over a five-day period preceding the onset of the

intervention. Measurements were obtained following an overnight fast, and subjects refrained from exercise, alcohol, and caffeine for the 24 hours prior to testing. A total of 44 subjects completed the dietary intervention and returned for the post-training assessment of changes in the dependent variables of body composition and blood lipids. All returning subjects were assessed at the same time of day as the pre-training measures over a five-day period following the completion of the intervention. During the dietary intervention, all subjects regularly participated in a CrossFit-based, high-intensity circuit training program.

Dietary Intervention

A Paleolithic diet, as first described by Eaton and Konner, was implemented for all study participants (11). Subjects were advised to increase their consumption of lean meat, fish, eggs, nuts, fruit, and vegetables and were instructed to strictly avoid all grains, dairy products, and legumes. All modern, processed foods including any form of processed sugar, soft drinks, and coffees were also excluded from the diets of the subjects. No specific macronutrient recommendations were made, as the study design wanted to closely mimic a real world model that would incorporate food choices made by the average consumer. Intake of specific proportion of food categories (e.g. animal vs. plant foods) was also not given.

Blood Lipid Analysis

For all subjects, a finger-stick blood sample from a sterilized site was collected into a lithium heparin coated capillary tube by one trained investigator. Samples were immediately transferred into a point of care desktop lipid analyzer (Cholestech LDX,

San Diego, CA). This method has previously been shown to be within standards of accuracy and precision set forth by the National Cholesterol Education Program (NCEP) (41). High-density lipoprotein (HDL), total cholesterol (TC), and triglycerides (TG) were measured directly, while low density lipoprotein (LDL) was estimated using the Friedewald equation (17).

Table 2. Blood lipid risk stratification

	HDL (mg/dL)	LDL (mg/dL)	n-HDL (mg/dL)	TC/HDL	
Low	<40	Optimal	<100	Low <130	Optimal <3.4
Normal	40-60	Near optimal	100-129	Normal 130-189	Above optimal ≥3.5
High	≥60	Borderline High	130-159	High ≥190	

HDL = high density lipoprotein cholesterol; LDL = low density lipoprotein cholesterol; n-HDL = non-high density lipoprotein cholesterol; TC/HDL = ratio of total cholesterol to high density lipoprotein.

Body Composition

Percentage body fat was calculated using the Bod Pod air-displacement plethysmography device (Life Measurements Instruments, Concord, CA), which is shown to be an accurate method for assessing body composition in adults (3). Prior to measurement, the system was calibrated for volume using a cylinder of a known volume (50.312 L) and for mass using two 10 kg weights. Fasting-state body weight was measured to the nearest 0.1 kg and subjects entered the Bod Pod chamber wearing only a tight fitting swimsuit and swim cap. Body volume measurements were taken in duplicate and repeated if measures were not within 150 mL of each other (9). Body density was calculated as mass/body volume and body fat percentage was calculated by using Siri's formula (43). Body mass index (BMI) was

calculated as kg body mass divided by height in meters squared.

Graded Exercise Testing

All subjects performed a maximal treadmill exercise test before and after the training program using the Bruce protocol(5) to determine $\text{VO}_{2\text{max}}$. Subjects wore nose clips and breathed into a one-way mouthpiece, which allowed expired gases to be collected in a mixing chamber. Volume of expired air, oxygen consumption, and carbon dioxide production were determined by gas analyzers and a pneumotachometer attached to a calibrated, computerized metabolic cart (Parvomedics, Sandy, UT), which provides accurate and reliable results compared to the Douglas bag method(7). Oxygen consumption values were calculated every 15 s and the two highest consecutive values were averaged to determine absolute maximal oxygen consumption in L/min. Body weight was divided into absolute oxygen consumption to yield a value relative to body mass and was reported as relative $\text{VO}_{2\text{max}}$ in units of ml of O_2 /kg of body mass/min. The test was terminated and considered maximal when subjects reached self-determined exhaustion, and was verified by two of the following criteria: (1) plateau in oxygen consumption despite an increase in workload, (2) respiratory exchange ratio greater than 1.1, and (3) rating of perceived exertion of 18-20. Using these parameters have previously shown to be a reliable method of verifying $\text{VO}_{2\text{max}}$ has been attained, and provides statistically indistinguishable measurements compared to supramaximal testing(21).

Statistical Analysis

Changes in body composition and blood lipids were tested using a two-tailed, paired t-test. These values were tested as an entire group, and also in subsets that were stratified by initial values of blood lipids for each subject. Participants were stratified into subsets based on cutoffs (Table 2) published by the NCEP Adult Treatment Panel III (ATP III) (1). Two-tailed, paired t-tests were then used to test differences between pre- and post-intervention values of stratified blood lipids. Data are reported as mean \pm SEM. Statistical analysis was performed using STATA (version 11.1, College Station, TX). Statistical significance was defined *a priori* as the critical α -level of $P < 0.05$. Statistical power was calculated to be >99% for the variables collected from 44 subjects.

RESULTS

Characteristics of all the subjects who volunteered for the study are presented in Table 1. A total of 44 subjects participated in the study (24 males, 20 females). The average age of study participants was 33.5 ± 0.3 yr and 31.2 ± 0.3 yr for men and women, respectively. Following the dietary intervention, n-HDL (107.1 ± 6.0 mg/dL to 120.2 ± 6.5 mg/dL; $P < 0.01$), LDL (93.1 ± 5.4 mg/dL to 105.6 ± 6.1 mg/dL; $P < 0.01$), and TC (168.8 ± 5.4 mg/dL to 178.9 ± 6.6 mg/dL; $P < 0.05$) increased significantly from baseline in all participants (Figure 1).

Further, TC/HDL increased to a statistically significant degree (3.0 ± 0.2 to 3.3 ± 0.2 ; $P < 0.05$). Body fat percentage decreased significantly ($24.3 \pm 1.2\%$ to $20.7 \pm 1.2\%$; $P < 0.05$) compared with baseline values (Figure 1), as did body weight (80.7 ± 2.6 kg to 77.5 ± 2.4 kg; $P < 0.01$). When stratified by initial blood lipid levels based

on the ATP III, there was a significant decrease of HDL among subjects with the highest initial HDL (82.1 ± 3.2 mg/dL to 68.6 ± 4.8 mg/dL; $P < 0.05$; Figure 2) and significant increases of n-HDL among subjects with the lowest levels of n-HDL (86.6 ± 3.9 mg/dL to 101.4 ± 4.8 mg/dL; $P < 0.01$; Figure 3).




Figure 1. Blood lipids in healthy volunteer prior to and following a 10-week Paleolithic diet intervention. Non- high density lipoprotein (n-HDL), low density lipoprotein (LDL) and total cholesterol (TC) increased significantly from baseline, whereas no changes were observed with regard to high density lipoprotein (HDL) and triglycerides (TG). A significant decrease in body composition was observed compared to baseline. * $P < 0.05$; ** $P < 0.01$.

Furthermore, significant increases of LDL (69.1 ± 3.1 mg/dL to 83.5 ± 4.1 mg/dL; $P < 0.01$; Figure 4), TC (157.2 ± 0.7 mg/dL to 168.2 ± 0.9 mg/dL; $P < 0.05$), and TC/HDL (2.5 ± 0.1 to 2.7 ± 0.1 ; $P < 0.05$; Figure 5) were observed among subjects with the most optimal initial levels of each respective outcome variable. Relative maximal oxygen consumption also improved during the study (39.8 ± 1.2 ml/kg/min to 44.9 ± 8.2 ml/kg/min; $P < 0.001$; Table 3), presumably resulting from the concurrent regular participation in a CrossFit-based, high-intensity circuit training program, along

with a decrease in body weight. Despite the weight loss observed in study participants, absolute oxygen consumption also increased (3.18 ± 0.14 L/min to 3.46 ± 0.15 L/min; $P < 0.001$), which indicates an overall concurrent improvement of aerobic fitness.




Figure 2. High density lipoprotein (HDL) levels in healthy volunteers before and following a 10 week Paleolithic dietary intervention. When stratified by initial HDL levels, only subjects who presented with “High-HDL” were observed to have a significant decrease in HDL following a Paleolithic diet. * $P < 0.05$.




Figure 3. Non- high density lipoprotein (n-HDL) levels in healthy volunteers prior to and following a Paleolithic dietary intervention. When stratified by initial levels of n-HDL, only subjects with n-HDL considered to be “low” were measured to have a significant increase of n-HDL following 10 weeks of a Paleolithic diet. * $P < 0.05$.

Table 3. Changes observed after 10-week Paleo intervention.

	VO _{2max} (ml/kg/min)	VO _{2max} (L/min)	Body Fat (%)	TC (mg/dL)	LDL (mg/dL)	n-HDL (mg/dL)
Pre-test	39.82±7.72	3.18±0.14	24.32±7.63	168.8±5.4	93.1±5.4	107.1±6.0
Post-test	44.90±8.20**	3.46±0.15**	20.65±7.99**	178.9±6.6*	105.6±6.1**	120.2±6.5**

VO_{2max} = maximal oxygen consumption; TC = total cholesterol; LDL = low density lipoprotein, n-HDL = non-high density lipoprotein. *P<0.05, **P<0.01.




Figure 4. Low density lipoprotein (LDL) levels in healthy volunteers prior to and following a Paleolithic dietary intervention. When stratified by initial levels of LDL, only subjects with optimal LDL were measured to have a significant increase of LDL following 10 weeks of a Paleolithic diet. * $P < 0.05$.




Figure 5. Total cholesterol to high density lipoprotein ratio (TC/HDL) in healthy volunteers prior to and following a Paleolithic dietary intervention. When stratified by initial levels of TC/HDL, only subjects with optimal TC/HDL were measured to have a significant increase of TC/HDL following 10 weeks of a Paleolithic diet. * $P < 0.05$.

DISCUSSION

Our study featured a large number (n = 44) of healthy, active subjects that followed an *ad libitum* Paleo diet while participating in a CrossFit-based circuit training program. As hypothesized, we observed negative impacts on non-HDL, LDL, TC, and TC/HDL, no significant change in HDL, and a decrease in body weight and body fat percentage. Furthermore, when stratified by pre-test blood lipid values, the statistically significant effects primarily came from deleterious changes in the most

healthy subjects; subjects within optimal pre-test blood lipid stratifications saw statistically significant negative changes in HDL, non-HDL, LDL, TC, and TC/HDL. The Law of Initial Values might explain why these deleterious effects were so pronounced in subjects with the healthiest blood lipid profiles. Subjects with optimal initial blood lipids were unable to maintain their ideal blood lipid values after adhering to the Paleo diet for 10 weeks, whereas a similar effect would not likely be seen in subjects with poor initial blood lipid values. Simply limiting food sources to the list of

Paleo-approved foods could effectively improve the diet of very unhealthy populations, but the results of our study indicate that it may be difficult to maintain an ideal blood lipid profile while adhering to the commonly followed Paleo guidelines used in this intervention.

While some approaches to the Paleo diet provide guidelines that serve to limit the intake of total fat and saturated fat, such guidelines may not accurately reflect a truly *ad libitum* approach. Indeed, it is likely that many people who adhere to the Paleo diet merely limit themselves to an approved list of food sources and consume those accepted foods *ad libitum*. While previously conducted Paleo studies have utilized restrictive guidelines that help to limit total fat and saturated fat intakes (26, 33, 38), there is little data pertaining to the application of the less restrictive, more practical Paleo guidelines that do not impose limits on the intake of any approved food sources. Our study, which utilized the less restrictive guidelines that appear to be more commonly followed by many Paleo dieters, found deleterious effects on blood lipids following a 10-week Paleo intervention.

Previous studies have shown positive effects of Paleo interventions on a number of health markers (16, 25, 26, 33, 38), and in some cases, blood lipids (16, 26). However, these studies used more restrictive Paleo guidelines, and many used subjects that were initially overweight (16, 26, 33) and were either sedentary (16), had coronary artery disease (33), or had type 2 diabetes (26). There is a general lack of literature pertaining to Paleo interventions in healthy, highly active populations, and there is reason to question whether previous Paleo

studies accurately replicate the actual dietary intake of the free-living Paleo dieter. It is also not clear that the current Paleo diet embraces lifestyle habits that a true Paleolithic man would have faced, such as long periods of fasting performed concurrent to long periods of low intensity aerobic exercise(4).

Throughout the intervention, subjects completed a CrossFit-based exercise program while adhering to the Paleo diet, resulting in significant decreases in body weight and body fat percentage while significantly increasing maximal oxygen consumption, a common measure of cardiorespiratory fitness. In previous interventions utilizing diet and/or exercise, improvements in body composition and cardiorespiratory fitness have been accompanied by improvements in one or more blood lipid measurements, including TG (34, 42), TC (34, 42), LDL (34, 39, 42), HDL (31, 34), and TC/HDL (34), although it should be noted that some studies have shown HDL levels to drop during active weight loss, before increasing after weight stabilization is achieved (8). Additionally, cardiorespiratory fitness (19, 20, 24, 32) and weight loss (8) have both been independently associated with improved blood lipids. Previous literature would indicate that a combination of Paleo and vigorous exercise that improves cardiorespiratory fitness and body composition should also improve blood lipid measurements. However, while our subjects improved cardiorespiratory fitness and body composition, all blood lipid measurements were either unchanged or negatively impacted. Though the lack of HDL improvement could potentially be attributed to the possibility that many subjects had yet to sufficiently stabilize at

their post-test body weight, this does not explain the increases in n-HDL, LDL, TC/HDL, or TC experienced by the group, nor does it explain the absence of TG improvement. Our data indicate that the Paleo diet's deleterious impact on blood lipids was not only significant, but substantial enough to counteract the blood lipid improvements commonly seen with improved fitness and body composition.

It should be acknowledged that the Paleo diet has many positive characteristics. The guidelines allow dieters to consume plenty of high-protein foods and fruit, which have been shown to be highly satiating food sources (23). Direct comparison of the Mediterranean and Paleo diets has shown that subjects report higher levels of satiety per calorie from the Paleo diet (27). Furthermore, studies have shown that the low carbohydrate to protein ratio commonly associated with the Paleo diet improves glycemic control and body composition (14, 16, 28-30, 33).

The Paleo diet encourages the consumption of foods that are satiating, full of micronutrients, and often, great sources of fiber, essential fatty acids, and essential amino acids. Likewise, this diet restricts the consumption of a number of foods that are calorically dense and low in micronutrient content. Replacing highly processed, energy-dense foods with Paleo-approved foods is highly beneficial and achievable for many individuals. However, the diet unnecessarily restricts the consumption of foods that can contribute to a healthy diet, including dairy, whole grains, and legumes. Numerous studies have shown high dairy intake to be associated with decreased risk of all-cause mortality, ischemic heart disease, stroke, and diabetes (13), and

studies have shown an inverse relationship between high dairy or calcium intake and bodyweight, body fat, and BMI (45). In clinical trials, dairy has been shown to be more effective in terms of attenuating fat deposition (47) and increasing tibial cortical thickness (6) than equal amounts of supplementary calcium, prompting researchers to suggest that it is difficult to sufficiently replace the nutrients provided by dairy in the diet with supplements and non-dairy food sources (18, 22). Whole grains are associated with improved blood lipids and reduced risk of coronary artery disease, cardiovascular disease, type II diabetes, obesity, and some cancers (15, 36, 37, 44). Whole grains also contain a number of phytochemicals and bioactive compounds that may contribute to chronic disease prevention (37, 44). Some research has shown legumes to be inversely associated with coronary heart disease and cardiovascular disease, and soy protein is known to improve blood lipids by a small but significant degree (15). While those with Celiac Disease, gluten sensitivity, or lactose intolerance might wisely avoid gluten or lactose, a large body of literature indicates that dairy, legumes, and whole grains can contribute to a healthy, well-rounded diet in healthy individuals.

Our study does have some limitations. Our study did not impose a high level of control over our subjects— meals were not prepared for subjects, nor did the subjects stay in a metabolic ward for the duration of the study. Furthermore, our study did not include a control group and a low number of diet logs were returned ($n = 8$), which calls into question how accurately these logs portray the true dietary intake of the group and were therefore excluded from analysis. Despite these limitations, the

study used a large sample (n = 44), utilized a sample containing healthier, fitter subjects than previous Paleo interventions, featured a practical application of the Paleo diet that may represent the normal dietary habits of Paleo dieters more accurately than previous studies, and stratified the subjects to examine the diet's effects on the blood lipid profiles of subjects of varying pre-test blood lipid categories. Future studies with more controlled conditions will help to identify the exact dietary adjustments subjects must make to ensure the maintenance of a healthy blood lipid profile while adhering to this popular, practical application of the Paleo diet.

In conclusion, our study shows that the Paleo diet was significantly deleterious to blood lipid profiles in healthy subjects concurrently participating in a CrossFit-based, high-intensity circuit training program. Specifically, subjects with optimal initial blood lipid values demonstrated the greatest increase in LDL, TC/HDL, TC, and n-HDL values, along with the greatest decline in HDL values, following the 10-week Paleo diet intervention. Despite concurrent improvements in aerobic capacity and body composition noted in these subjects, the Paleo diet may have negated the positive effects of exercise on blood lipids.

REFERENCES

1. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation* 106(25):3143-421, 2002.
2. American Heart Association Nutrition C, Lichtenstein AH, Appel LJ, Brands M, Carnethon M, Daniels S, Franch HA, Franklin B, Kris-Etherton P, Harris WS, Howard B, Karanja N, Lefevre M, Rudel L, Sacks F, Van Horn L, Winston M, Wylie-Rosett J. Diet and lifestyle recommendations revision 2006: a scientific statement from the American Heart Association Nutrition Committee. *Circulation* 114(1):82-96, 2006.
3. Biaggi RR, Vollman MW, Nies MA, Brener CE, Flakoll PJ, Levenhagen DK, Sun M, Karabulut Z, Chen KY. Comparison of air-displacement plethysmography with hydrostatic weighing and bioelectrical impedance analysis for the assessment of body composition in healthy adults. *The American journal of clinical nutrition* 69(5):898-903, 1999.
4. Boullousa DA, Abreu L, Varela-Sanz A, Mujika I. Do olympic athletes train as in the Paleolithic era? *Sports medicine* 43(10):909-17, 2013.
5. Bruce RA, Hornsten TR. Exercise stress testing in evaluation of patients with ischemic heart disease. *Progress in cardiovascular diseases* 11(5):371-90, 1969.
6. Cheng S, Lyytikainen A, Kroger H, Lamberg-Allardt C, Alen M, Koistinen A, Wang QJ, Suuriniemi M, Suominen H, Mahonen A, Nicholson PH, Ivaska KK, Korpela R, Ohlsson C, Vaananen KH, Tyllavsky F. Effects of calcium, dairy product, and vitamin D supplementation on bone mass accrual and body composition in 10-12-y-old girls: a 2-y randomized trial. *Am J Clin Nutr* 82(5):1115-26; quiz 47-8, 2005.
7. Crouter SE, Antczak A, Hudak JR, DellaValle DM, Haas JD. Accuracy and reliability of the ParvoMedics TrueOne 2400 and MedGraphics VO2000 metabolic systems. *European Journal of Applied Physiology* 98(2):139-51, 2006.
8. Dattilo AM, Kris-Etherton PM. Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. *Am J Clin Nutr* 56(2):320-8, 1992.
9. Dempster P, Aitkens S. A new air displacement method for the determination of human body composition. *Medicine and Science in Sports and Exercise* 27(12):1692-7, 1995.
10. Eaton SB, Konner M. Paleolithic nutrition. A consideration of its nature and current implications. *N Engl J Med* 312(5):283-9, 1985.

11. Eaton SB, Konner M. Paleolithic nutrition. A consideration of its nature and current implications. *The New England journal of medicine* 312(5):283-9, 1985.
12. Eaton SB, Konner M, Shostak M. Stone agers in the fast lane: chronic degenerative diseases in evolutionary perspective. *Am J Med* 84(4):739-49, 1988.
13. Elwood PC, Pickering JE, Givens DI, Gallacher JE. The consumption of milk and dairy foods and the incidence of vascular disease and diabetes: an overview of the evidence. *Lipids* 45(10):925-39, 2010.
14. Farnsworth E, Luscombe ND, Noakes M, Wittert G, Argyiou E, Clifton PM. Effect of a high-protein, energy-restricted diet on body composition, glycemic control, and lipid concentrations in overweight and obese hyperinsulinemic men and women. *Am J Clin Nutr* 78(1):31-9, 2003.
15. Flight I, Clifton P. Cereal grains and legumes in the prevention of coronary heart disease and stroke: a review of the literature. *Eur J Clin Nutr* 60(10):1145-59, 2006.
16. Frassetto LA, Schloetter M, Mietus-Synder M, Morris RC, Jr., Sebastian A. Metabolic and physiologic improvements from consuming a paleolithic, hunter-gatherer type diet. *Eur J Clin Nutr* 63(8):947-55, 2009.
17. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clinical Chemistry* 18(6):499-502, 1972.
18. Fulgoni VL, 3rd, Keast DR, Auestad N, Quann EE. Nutrients from dairy foods are difficult to replace in diets of Americans: food pattern modeling and an analyses of the National Health and Nutrition Examination Survey 2003-2006. *Nutr Res* 31(10):759-65, 2011.
19. Gibbs BB, Brancati FL, Chen H, Coday M, Jakicic JM, Lewis CE, Stewart KJ, Clark JM. Effect of improved fitness beyond weight loss on cardiovascular risk factors in individuals with type 2 diabetes in the Look AHEAD study. *Eur J Prev Cardiol* 2012.
20. Grundy SM, Barlow CE, Farrell SW, Vega GL, Haskell WL. Cardiorespiratory fitness and metabolic risk. *Am J Cardiol* 109(7):988-93, 2012.
21. Hawkins MN, Raven PB, Snell PG, Stray-Gundersen J, Levine BD. Maximal oxygen uptake as a parametric measure of cardiorespiratory capacity. *Medicine and Science in Sports and Exercise* 39(1):103-7, 2007.
22. Heaney RP. Dairy and bone health. *J Am Coll Nutr* 28 Suppl 1:82S-90S, 2009.
23. Holt SH, Miller JC, Petocz P, Farmakalidis E. A satiety index of common foods. *Eur J Clin Nutr* 49(9):675-90, 1995.
24. Jette M, Sidney K, Quenneville J, Landry F. Relation between cardiorespiratory fitness and selected risk factors for coronary heart disease in a population of Canadian men and women. *CMAJ* 146(8):1353-60, 1992.
25. Jonsson T, Ahren B, Pacini G, Sundler F, Wierup N, Steen S, Sjoberg T, Ugander M, Frostegard J, Goransson L, Lindeberg S. A Paleolithic diet confers higher insulin sensitivity, lower C-reactive protein and lower blood pressure than a cereal-based diet in domestic pigs. *Nutr Metab (Lond)* 3:39, 2006.
26. Jonsson T, Granfeldt Y, Ahren B, Branell UC, Palsson G, Hansson A, Soderstrom M, Lindeberg S. Beneficial effects of a Paleolithic diet on cardiovascular risk factors in type 2 diabetes: a randomized cross-over pilot study. *Cardiovasc Diabetol* 8:35, 2009.
27. Jonsson T, Granfeldt Y, Erlanson-Albertsson C, Ahren B, Lindeberg S. A paleolithic diet is more satiating per calorie than a mediterranean-like diet in individuals with ischemic heart disease. *Nutr Metab (Lond)* 7:85, 2010.
28. Layman DK, Baum JI. Dietary protein impact on glycemic control during weight loss. *J Nutr* 134(4):968S-73S, 2004.
29. Layman DK, Boileau RA, Erickson DJ, Painter JE, Shiue H, Sather C, Christou DD. A reduced ratio of dietary carbohydrate to protein improves body composition and blood lipid profiles during weight loss in adult women. *J Nutr* 133(2):411-7, 2003.

30. Layman DK, Shiue H, Sather C, Erickson DJ, Baum J. Increased dietary protein modifies glucose and insulin homeostasis in adult women during weight loss. *J Nutr* 133(2):405-10, 2003.
31. Lee MG, Park KS, Kim DU, Choi SM, Kim HJ. Effects of high-intensity exercise training on body composition, abdominal fat loss, and cardiorespiratory fitness in middle-aged Korean females. *Appl Physiol Nutr Metab* 2012.
32. Lee S, Kuk JL, Katzmarzyk PT, Blair SN, Church TS, Ross R. Cardiorespiratory fitness attenuates metabolic risk independent of abdominal subcutaneous and visceral fat in men. *Diabetes Care* 28(4):895-901, 2005.
33. Lindeberg S, Jonsson T, Granfeldt Y, Borgstrand E, Soffman J, Sjostrom K, Ahren B. A Palaeolithic diet improves glucose tolerance more than a Mediterranean-like diet in individuals with ischaemic heart disease. *Diabetologia* 50(9):1795-807, 2007.
34. Martins RA, Verissimo MT, Coelho e Silva MJ, Cumming SP, Teixeira AM. Effects of aerobic and strength-based training on metabolic health indicators in older adults. *Lipids Health Dis* 9:76, 2010.
35. Mercader J. Mozambican grass seed consumption during the Middle Stone Age. *Science* 326(5960):1680-3, 2009.
36. Mozaffarian D, Appel LJ, Van Horn L. Components of a cardioprotective diet: new insights. *Circulation* 123(24):2870-91, 2011.
37. Okarter N, Liu RH. Health benefits of whole grain phytochemicals. *Crit Rev Food Sci Nutr* 50(3):193-208, 2010.
38. Osterdahl M, Kocturk T, Koochek A, Wandell PE. Effects of a short-term intervention with a paleolithic diet in healthy volunteers. *Eur J Clin Nutr* 62(5):682-5, 2008.
39. Paillard T, Lafont C, Costes-Salon MC, Dupui P, Riviere D, Vellas B. Cholesterol reduction and increased cardiovascular fitness following a 12 weeks brisk walking. *J Nutr Health Aging* 6(2):138-40, 2002.
40. Revedin A, Aranguren B, Becattini R, Longo L, Marconi E, Lippi MM, Skakun N, Sinitsyn A, Spiridonova E, Svoboda J. Thirty thousand-year-old evidence of plant food processing. *Proc Natl Acad Sci U S A* 107(44):18815-9, 2010.
41. Santee J. Accuracy and precision of the Cholestech LDX System in monitoring blood lipid levels. *Am J Health Syst Pharm* 59(18):1774-9, 2002.
42. Saremi A, Asghari M, Ghorbani A. Effects of aerobic training on serum omentin-1 and cardiometabolic risk factors in overweight and obese men. *J Sports Sci* 28(9):993-8, 2010.
43. Siri WE. Body composition from fluid spaces and density: analysis of methods. 1961. *Nutrition* 9(5):480-91; discussion , 92, 1993.
44. Slavin JL, Martini MC, Jacobs DR, Jr., Marquart L. Plausible mechanisms for the protectiveness of whole grains. *Am J Clin Nutr* 70(3 Suppl):459S-63S, 1999.
45. Van Loan M. The role of dairy foods and dietary calcium in weight management. *J Am Coll Nutr* 28 Suppl 1:120S-9S, 2009.
46. Voegtlin WL. *The stone age diet : based on in-depth studies of human ecology and the diet of man*. 1st ed. New York: Vantage Press; 1975, xvii, 277 p. p.
47. Zemel MB, Shi H, Greer B, Dirienzo D, Zemel PC. Regulation of adiposity by dietary calcium. *FASEB J* 14(9):1132-8, 2000.