

Post Activation Potentiation of Back Squat and Trap Bar Deadlift on Acute Sprint Performance

Chmiel, J., Carillo, J., Cerone, D., Phillips, J., Swensen, T., Kaye, M. Ithaca College, Ithaca, NY

PURPOSE: We investigated the ability of back squat (BS) and trap bar deadlift (TBD) to elicit post activation potentiation (PAP) and hence improve 40m sprint performance in college age female rowers. **METHODS:** Twenty division III collegiate female rowers, whose $\bar{X} \pm SD$ for age, height, weight, and VO_2 Max were 19.2 ± 1.1 y, 1.7 ± 0.06 m, 67.4 ± 6.8 kg and 42.5 ± 3.9 $ml \cdot kg^{-1} \cdot min^{-1}$, completed a 40m sprint timed at 10m, 20m, and 40m with a timing system. Based on sprint times, subjects were divided into two groups: BS and TBD. Subsequently, we determined their one repetition maximum (1 RM) for the BS and TBD. One week later, subjects completed a 40m sprint, and then three repetitions at 90% of their 1RM for BS or TBD; after 7 min of active rest, they completed another 40m sprint. A dynamic warm-up and active cool down preceded and followed all testing sessions. Data were analyzed with three 2x2 repeated measures ANOVA. **RESULTS:** We found both conditions significantly increased 20m and 40m sprint times as shown below:

Sprint Times (s) by Group

	Trap Bar Dead Lift (n=10)			Back Squat (n = 10)		
	10m	20m	40m	10m	20m	
40m						
Pre	2.06 ± 0.11	3.53 ± 0.19	6.48 ± 0.44	2.10 ± 0.09	3.62 ± 0.14	6.61 ± 0.32
Post	2.07 ± 0.11	3.60 ± 0.19*	6.57 ± 0.42*	2.10 ± 0.09	3.66 ± 0.17*	6.67 ± 0.36*

Note. * = $p < 0.05$

CONCLUSION: The data show that three repetitions at 90% of 1RM for BS or TBD did not elicit PAP in female college rowers. These findings may be related to sex, load, training incompatibility, and sprinting proficiency in this subject population.