

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/109066/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Caballero, Javier, Peralta, Cecilia, Molla, Antonella, Del Valle, Eleodoro E, Caballero, Primitivo, Berry, Colin ORCID: <https://orcid.org/0000-0002-9943-548X>, Felipe, Veronica, Yaryura, Pablo and Palma, Leopolda 2018. Draft genome sequence of *Bacillus cereus* CITVM-11.1, a strain exhibiting interesting antifungal activities. *Journal of Molecular Microbiology and Biotechnology* 28 (1) , pp. 47-51. 10.1159/000487597 file

Publishers page: <http://dx.doi.org/10.1159/000487597>
<<http://dx.doi.org/10.1159/000487597>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



1 **Draft genome sequence of *Bacillus cereus* CITVM-11.1, a strain exhibiting**
2 **interesting antifungal activities.**

3

4 Javier Caballero^{a,b}, Cecilia Peralta^{c,d}, Antonella Molla^d, Eleodoro E. Del Valle^e,
5 Primitivo Caballero^{a,b}, Colin Berry^f, Verónica Felipe^d, Pablo Yaryura^{c,d}, and Leopoldo
6 Palma^{c,d*}

7

8 ^aInstituto de Agrobiotecnología, CSIC-Gobierno de Navarra, 31192 Mutilva Baja,
9 Navarra, Spain.

10 ^bDepartamento de Producción Agraria, Universidad Pública de Navarra, Campus
11 Arrosadía s/n, 31006, Pamplona, Navarra, Spain.

12 ^cCentro de Investigaciones y Transferencia de Villa María (CIT-VM), Consejo Nacional
13 de Investigaciones Científicas y Técnicas (CONICET), Universidad Nacional de Villa
14 María, Argentina.

15 ^dInstituto Académico Pedagógico de Ciencias Básicas y Aplicadas (IAPCByA),
16 Universidad Nacional de Villa María (UNVM), Villa María, Argentina.

17 ^eFacultad de Ciencias Agrarias, Universidad Nacional del Litoral, Esperanza, Santa Fe,
18 Argentina.

19 ^fCardiff School of Biosciences, Cardiff University, Park Place, Cardiff, CF10 3AT, UK

20

21 Javier Caballero and Cecilia Peralta contributed equally to this work

22 *Corresponding author: Leopoldo Palma (palma.leopoldo@gmail.com)

23 Instituto Académico Pedagógico de Ciencias Básicas y Aplicadas, Universidad
24 Nacional de Villa María, Avenida Arturo Jauretche 1555, Villa María (5900), Córdoba,
25 Argentina. Tel.: +54-(0353)-4539100 int. 3069; Fax: +54-(0353)-4539117

26 **Abstract**

27 *Bacillus cereus* is a Gram-positive spore-forming bacterium possessing an important
28 and historical record as human-pathogenic bacterium. However, several strains of this
29 species exhibit an interesting potential to be used as plant-growth promoting
30 rhizobacteria. Here, we report the draft genome sequence of *Bacillus cereus* strain
31 CITVM-11.1, which consists of 37 contig sequences accounting for 5,746,486 bp, with
32 a GC content of 34.8% and 5,752 predicted protein-coding sequences. Several of them
33 could potentially be involved in plant-bacterium interactions and may contribute to the
34 strong antagonistic activity shown by this strain against the charcoal rot fungus
35 *Macrophomina phaseolina*. This genomic sequence also showed a number of genes that
36 may confer to this strain resistance against several polluting heavy metals and for the
37 bioconversion of mycotoxins.

38

39 *Bacillus cereus* is a Gram-positive and ubiquitous spore-forming bacterium that
40 has been isolated from a wide range of ecosystems including water, dead insects, soil
41 samples, the rhizosphere, the gut of several animals but is also associated with food
42 poisoning by consumption of rice-based dishes [[Krawczyk et al., 2015](#)]. This bacterium
43 is causing, after *Salmonella* and *Staphylococcus aureus*, the highest number of
44 collective food poisoning outbreaks in Europe [[Ramarao and Sanchis, 2013](#)]. *B. cereus*
45 food poisoning causes gastroenteritis which can be manifested in two different types of
46 illness, one vomiting (emetic) form that resembles *S. aureus* infections and the
47 diarrhoeal form, with a similar symptomatology to the infections caused by *Clostridium*
48 *perfringens* [[Ramarao and Sanchis, 2013](#)].

49 Nevertheless, several strains of this species have demonstrated potential to be
50 used as plant growth promoting rhizobacteria (PGPR) since they are capable of

51 exhibiting antagonistic activities against several phytopathogenic microorganisms
52 [[Kumar et al., 2014b](#)] and inducing plant-systemic resistance against phytopathogenic
53 bacteria such as *Pseudomonas syringae* [[Niu et al., 2011](#)].

54 In this work, we report the draft genome sequence of *Bacillus cereus* strain
55 CITVM-11.1, which was isolated from a soil sample obtained in a field of alfalfa plants
56 (*Medicago sativa* L.) in the province of Córdoba, Argentina [[Felipe et al., 2016](#)]. This
57 strain exhibited strong antagonistic activity *in vitro*, against the charcoal rot fungus
58 *Macrophomina phaseolina* by causing inhibition of hyphal development and impaired
59 formation of sclerotia [[Felipe et al., 2016](#)]. This finding was consistent with other *B.*
60 *cereus* strains that have demonstrated their potential for the biocontrol of some
61 phytopathogenic fungi, bacteria, and plant-parasitic nematodes both in *in vitro* assays
62 and through *in vivo* trials [[Kumar et al., 2014b](#); [Martinez-Alvarez et al., 2016](#)].

63 Purified total DNA from *B. cereus* CITVM-11.1 was obtained using the Wizard
64 genomic DNA purification kit (Promega), following the instructions for the isolation of
65 DNA from Gram-positive bacteria. Total DNA, which in some strains may be
66 composed of the bacterial chromosome and a variable number of plasmids, was
67 electrophoresed in 1% agarose gels stained with SYBR Safe (Thermo Fisher Scientific).

68 Genome sequencing was performed at Stabvida (Portugal) by using high-
69 throughput Illumina sequencing technology with a genomic coverage of 1000×.
70 Genome assembly was performed by assembling (*de novo*) the Illumina reads with
71 Geneious R10 (Biomatters) into 37 contigs totalling 5,746,486 bp, with a maximum
72 contig size of 695,448 bp and a G+C content of 34.8 %. Genome annotation was
73 performed with the NCBI Prokaryotic Genome Annotation Pipeline (2017 release),
74 although it was also analysed with RAST [[Aziz et al., 2008](#)], which produced a total of
75 5,752 protein-coding sequences (CDs) plus 71 RNA genes (rRNAs and tRNAs) and 5

76 non-coding RNAs.

77 Phylogenetic analysis using *gyrB* gene sequence and following the methodology
78 described by Bavykin et al. (2004), showed that *B. cereus* strain CITVM-11.1 belongs
79 to Cereus B subgroup located at Cluster I inside the *Bacillus cereus* group [[Bavykin et](#)
80 [al., 2004](#)] (Figure S1, supplementary material).

81 From the 5,752 predicted protein-coding sequences, several of them could be
82 potentially involved in plant-bacterium interactions (e.g. auxin biosynthesis) and the
83 previously reported antagonistic activity against *M. phaseolina* (Figure 1).

84 Genes potentially involved in the biosynthesis of thiopeptides or thiazolyl
85 peptides have been found in the genome. The thiopeptide cyclothiazomycin B1 (CTB1)
86 is an antifungal cyclic thiopeptide isolated from a *Streptomyces* sp. that produces
87 growth inhibition and morphological changes of hyphae and induces fragility of the
88 fungal cell wall by binding chitin [[Mizuhara et al., 2011](#)] and capable of producing
89 growth inhibition of fungal species such as *Fusarium*, *Aspergillus* and *Penicillium* spp
90 [[Mizuhara et al., 2011](#)]. A similar impaired growth has been produced in the charcoal-
91 root fungus *M. phaseolina* on exposure to *B. cereus* strain CITVM-11.1, as previously
92 reported (Felipe et al., 2016). Wang et. al (2010) analysed the biosynthetic gene cluster
93 responsible of the production of cyclothiazomycin thiopeptide in *Streptomyces*
94 *hygroscopicus* 10-22 [[Wang et al., 2010](#)] and described a gene cluster model for the
95 biosynthesis of cyclothiazomycin that involves several genes encoding putative
96 functional enzymes, namely: Ser and Thr dehydratases, enzymes producing the thertiary
97 thioether and an epoxide hydrolase [[Wang et al., 2010](#)]. Homologous genes to those
98 described by Wang wet al. (2010) have been found at the genome of *B. cereus* CITVM-
99 11.1 at contig No.12 (Thr-dehydratase, L-serine dehydratase, thioestearase) and contig
100 No. 23 (epoxide hydrolase, thioestearase and a thioazol kinase), even though they are

101 not organized in a biosynthetic gene cluster. Some *B. cereus* strains have been also
102 described as thiopeptide producing strains showing growth inhibition of *Aspergillus*
103 *flavus* and *Fusarium oxysporum*, although genes responsible of this thiopeptide
104 production have not been yet described [[Kumar et al., 2014a](#); [Kumar et al., 2014b](#)].
105 Other genes showing significant similarity with chitinase enzymes and surfactins were
106 also found in the genome and may be contributing to the antifungal activity exhibited by
107 this *B. cereus* strain.

108 Gene cluster analysis using anti SMASH (antibiotics & secondary metabolites
109 analysis shell) [[Weber et al., 2015](#)] showed that this strain potentially harbours 50
110 biosynthetic gene clusters. From them, best predicted gene clusters may be potentially
111 involved in i) the synthesis and accumulation of polyhydroxyalkanoates with 100 % of
112 the genes exhibiting similarity, ii) the production of the non-ribosomal peptide
113 bacillobactin (siderophore) with 46 % of the genes exhibiting similarity, iii) synthesis of
114 the non-ribosomal peptide bacitracin (antibiotic) with 100 % of the genes exhibiting
115 similarity, iv) the synthesis of the bacteriocin Thuricin H with 60 % of the genes
116 exhibiting similarity and v) the production of the siderophore petrobactin with 100 % of
117 the genes exhibiting similarity (Figure S2, supplementary material).

118 Contig 27 was automatically circularized by Geneious R10 as a putative plasmid
119 of 10,741 bp in size. Circularization of contigs occurs when running Geneious R10 *de*
120 *novo* assembly tool and a pair of reads of each end of the contig match and also such
121 reads must not intersect with each other in any other part of the contig. Accordingly,
122 agarose gel electrophoresis of total DNA showed an additional band consistent with the
123 presence of a plasmid (Fig 2A)". We have named this plasmid pBC11.1. Two RAST
124 annotated genes on the plasmid might be related to the mobilization (horizontal transfer
125 by conjugation) of the plasmid whereas two others, have been annotated by RAST as a

126 macrolide-efflux protein and a putative mercury resistance protein (Figure 2B).
127 Acquisition of antimicrobial-resistance genes in bacteria can occur by means of self-
128 transmissible plasmids (conjugative plasmids). These plasmids usually harbour all the
129 genes involved in mating-pore formation as well as the essential *mob* gene (encoding
130 DNA relaxase) and the recognition sequence commonly known as origin of transfer
131 (*oriT*) [[Ramsay et al., 2016](#)]. Despite the *mob* gene was found in pBC11.1 plasmid, we
132 could not effectively predict any known putative *oriT* sequence in this plasmid.

133 In addition, the genomic sequence also exhibits other RAST annotated genes
134 that could be related to the metabolism of several heavy metals that pollute the
135 environment, namely: i) for arsenic (As), three arsenic efflux pump proteins, one
136 arsenical resistance operon repressor and two arsenical-resistance proteins; ii) for
137 copper (Cu), a membrane protein for copper uptake and a copper resistance protein D;
138 ii) for cobalt (Co), zinc (Zn) and cadmium (Cd); three cobalt-zinc-cadmium resistance
139 proteins; iv) for mercury (Hg), one predicted gene, located at the plasmid pBC11.1,
140 potentially encode for a mercury resistance protein; v) for aluminium (Al), an
141 aluminium resistance protein and vi) for tellurium (Te), one tellurite-resistance protein
142 and three tellurium-resistance proteins. Some of the heavy metals mentioned above, e.g.
143 Zn, Cu, Ni, Co with chromium (Cr), are necessary as micronutrients, playing vital roles
144 in metabolic and physiological processes of microorganisms, plants and animals.
145 However, non-essential heavy metals such as silver (Ag), As, Cd, Pb and Hg are not
146 necessary for living organisms and their presence in soil and water sources pollute
147 ecosystems [[Fashola et al., 2016](#)].

148 The genomic sequence of *B. cereus* strain CITVM-11.1 also exhibits several
149 enzyme-coding genes that might be involved in the biotransformation of mycotoxins
150 [[Loi et al., 2017](#)]. Such genes, harboured at CITVM-11.1 strain, encode the following

151 enzymes: i) oxidases, peroxidases, reductases and manganese peroxidases (potential
152 aflatoxin-degrading enzymes); ii) carboxylesterases, aminotransferases and esterase
153 (potential fumonisin-degrading enzymes) and iii) cytochrome P450 and
154 glycosyltransferases (potential trichothecenes-degrading enzymes) [Loi et al., 2017].
155 Thus, *B. cereus* CITVM-11.1 could be a good source of enzymes for reducing
156 mycotoxin accumulation of staple food commodities [Loi et al., 2017], although it has
157 been shown to be a β -hemolytic strain (data not shown) that contains genes coding
158 known enterotoxins and their elimination may be necessary.

159 In this work, we report the draft genome sequence of *B. cereus* CITVM-11.1,
160 which showed a strong antagonistic activity against the charcoal rot fungus *M.*
161 *phaseolina*. This draft genome sequence provides an overview of the genes that could
162 be involved in plant-microbe interactions and the development of antagonistic activities
163 against phytopathogenic fungi, as well as indicating the potential of this strain to
164 tolerate the toxic activity of a number of heavy metals. The preliminary results
165 presented in this work encourage us to perform deeper studies, in order to elucidate both
166 the biocontrol and bioremediation potential of this strain, which deserves to be further
167 investigated.

168 This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank
169 under the accession number [MVFX00000000](#). The version described in this paper is the
170 first version, [MVFX01000000](#).

171

172 **Acknowledgements**

173 This work was supported by a research grant awarded by the Universidad Nacional de
174 Villa María (UNVM) and the Instituto de Investigación-UNVM (Disp. N° 03/2016,
175 Anexo I). We also thank the Centro de Investigación y Transferencia Villa María (CIT-

176 VM) for their support provided on this research project.

177

178 **Disclosure Statement**

179 The authors declare no conflict of interests.

180 **References**

- 181 Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass
182 EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK,
183 Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V,
184 Wilke A, Zagnitko O. The RAST server: rapid annotations using subsystems technology. BMC
185 Genomics 2008; 9:1-15.
- 186 Bavykin SG, Lysov YP, Zakhariev V, Kelly JJ, Jackman J, Stahl DA, Cherni A. Use of 16S
187 rRNA, 23S rRNA, and *gyrB* gene sequence analysis to determine phylogenetic relationships of
188 *Bacillus cereus* group microorganisms. J Clin Microbiol 2004; 42:3711-3730.
- 189 Fashola MO, Ngole-Jeme VM, Babalola OO. Heavy Metal Pollution from Gold Mines:
190 Environmental Effects and Bacterial Strategies for Resistance. Int J Environ Res Public Health
191 2016; 13.
- 192 Felipe V, Palma L, Yaryura P. Antagonistic activity of *Bacillus* sp. strain isolated in Córdoba,
193 Argentina against *Macrophomina phaseolina* (Tassi) Goid. . Rev Argent Microbiol 2016;
194 <http://dx.doi.org/10.1016/j.ram.2016.08.008>.
- 195 Krawczyk AO, de Jong A, Eijlander RT, Berendsen EM, Holsappel S, Wells-Bennik MH,
196 Kuipers OP. Next-Generation Whole-Genome Sequencing of Eight Strains of *Bacillus cereus*,
197 Isolated from Food. Genome Announc 2015; 3.
- 198 Kumar SN, Nambisan B, Mohandas C. Purification and identification of two antifungal cyclic
199 dipeptides from *Bacillus cereus* subsp. *thuringiensis* associated with a rhabditid
200 entomopathogenic nematode especially against *Fusarium oxysporum*. J Enzyme Inhib Med
201 Chem 2014a; 29:190-197.
- 202 Kumar SN, Sreekala SR, Chandrasekaran D, Nambisan B, Anto RJ. Biocontrol of *Aspergillus*
203 species on peanut kernels by antifungal diketopiperazine producing *Bacillus cereus* associated
204 with entomopathogenic nematode. PLoS One 2014b; 9:e106041.
- 205 Loi M, Fanelli F, Liuzzi VC, Logrieco AF, Mule G. Mycotoxin Biotransformation by Native
206 and Commercial Enzymes: Present and Future Perspectives. Toxins (Basel) 2017; 9.

207 Martinez-Alvarez JC, Castro-Martinez C, Sanchez-Pena P, Gutierrez-Dorado R, Maldonado-
208 Mendoza IE. Development of a powder formulation based on *Bacillus cereus* sensu lato strain
209 B25 spores for biological control of *Fusarium verticillioides* in maize plants. World J Microbiol
210 Biotechnol 2016; 32:75.

211 Mizuhara N, Kuroda M, Ogita A, Tanaka T, Usuki Y, Fujita K. Antifungal thiopeptide
212 cyclothiazomycin B1 exhibits growth inhibition accompanying morphological changes via
213 binding to fungal cell wall chitin. Bioorg Med Chem 2011; 19:5300-5310.

214 Niu DD, Liu HX, Jiang CH, Wang YP, Wang QY, Jin HL, Guo JH. The plant growth-
215 promoting rhizobacterium *Bacillus cereus* AR156 induces systemic resistance in *Arabidopsis*
216 *thaliana* by simultaneously activating salicylate- and jasmonate/ethylene-dependent signaling
217 pathways. Mol Plant Microbe Interact 2011; 24:533-542.

218 Ramarao N, Sanchis V. The pore-forming haemolysins of bacillus cereus: a review. Toxins
219 (Basel) 2013; 5:1119-1139.

220 Ramsay JP, Kwong SM, Murphy RJ, Yui Eto K, Price KJ, Nguyen QT, O'Brien FG, Grubb
221 WB, Coombs GW, Firth N. An updated view of plasmid conjugation and mobilization in
222 *Staphylococcus*. Mob Genet Elements 2016; 6:e1208317.

223 Wang J, Yu Y, Tang K, Liu W, He X, Huang X, Deng Z. Identification and analysis of the
224 biosynthetic gene cluster encoding the thiopeptide antibiotic cyclothiazomycin in *Streptomyces*
225 *hygroscopicus* 10-22. Appl Environ Microbiol 2010; 76:2335-2344.

226 Weber T, Blin K, Duddela S, Krug D, Kim HU, Bruccoleri R, Lee SY, Fischbach MA, Muller
227 R, Wohlleben W, Breitling R, Takano E, Medema MH. antiSMASH 3.0-a comprehensive
228 resource for the genome mining of biosynthetic gene clusters. Nucleic Acids Res 2015;
229 43:W237-243.

230

231


232

233 **Figure captions**



234 **Figure 1:** Potential plant-bacterium interaction and PGPR related features predicted and
235 annotated by RAST are highlighted in red.

236 **Figure 2:** A) Agarose gel electrophoresis of total DNA showing the genomic and
237 plasmid DNA (MM: molecular marker). B) Map of the circularized contig sequence
238 (contig 27).

239 **Figure 1**



241 **Figure 2**



242 A)

243


244

245

246


247

B)



249

250 **Figure S1:** Genetic relationship of *B. cereus* CITVM-11.1 and other strains based on *gyrB* gene sequences [Bavykin et al., 2004].



257 **Figure S2:** Potential biosynthetic gene clusters associated to secondary metabolites production identified with antiSMASH [[Weber et al., 2015](#)].

258

259