

UNIVERSITY OF WESTMINSTER



## WestminsterResearch

<http://www.wmin.ac.uk/westminsterresearch>

### **Phage display antibody against gill proteins of the tropical clam.**

**J-P. Gourdine<sup>1</sup>**  
**E.J. Smith-Ravin<sup>1</sup>**  
**Pamela Greenwell<sup>2</sup>**

<sup>1</sup> Biology, Universite des Antilles et de la Guyane, Pointe-a-Pitre, Guadeloupe, France

<sup>2</sup> School of Biosciences, University of Westminster

This is an electronic version of an article published in *Molecular and Cellular Proteomics*, 2 (9). 759, September 2003. *Molecular and Cellular Proteomics* is available online at:

<http://www.mcponline.org/cgi/reprint/2/9/702>

---

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners. Users are permitted to download and/or print one copy for non-commercial private study or research. Further distribution and any use of material from within this archive for profit-making enterprises or for commercial gain is strictly forbidden.

---

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch.  
(<http://www.wmin.ac.uk/westminsterresearch>).

In case of abuse or copyright appearing without permission e-mail [wattsn@wmin.ac.uk](mailto:wattsn@wmin.ac.uk).

## 39.4

## Phage Display Antibody Against Gill Proteins of the Tropical Clam

**J.-P. Gourdine<sup>1</sup>, E. J. Smith-Ravin<sup>1</sup>, and P. Greenwell<sup>2</sup>**

<sup>1</sup>Biology, Université des Antilles et de la Guyane, Pointe-à-Pitre, Guadeloupe, France; and <sup>2</sup>University of Westminster, London, United Kingdom

Introduction: *Codakia orbicularis* (Bivalvia, Lucinidae) is a tropical clam, which lives in areas of high sulfur concentrations and which contains endosymbiotic chemoautotrophic bacteria in its gills. Gill proteins are thought to be involved in sulfur metabolism and in symbiont/host interactions. We used the Recombinant Phage display Antibody System (RPAS) to raise Single chain Fragment variable (ScFv) antibodies against these proteins. Material and Methods: Griffin.1 library of phage display antibodies was used. Three rounds of selection of phage panning were performed on immunotubes coated with total extract of native gill proteins. The bound M13 phages were isolated and tested by ELISA and the positive clones were selected for ScFv production. The presence of ScFv DNA inserts was assessed by PCR. These chimeric antibodies were purified by ammonium sulfate precipitation or IMAC, and used in Dot-blot and Western-blot assays as well as in plate antigen capture combined with SDS-PAGE. Results and Discussion: Five out of 60 isolated clones were used for ScFv production. Purification on IMAC gave ScFv antibodies of about 30 000 Daltons at a higher yield compared to ammonium sulfate precipitation. Antigen capture showed a strong positive reaction to a 15 000 Dalton protein, a major band in the electrophoretic profile of gill proteins. These recombinant antibodies are being tested by immunohistochemistry on gill sections. Total gill extract has been used for the first time to raise phage antibodies against clam proteins, which antibodies might be useful in the isolation and study of the properties of the said proteins.