

Original citation:

Perry, Motty and Reny, Philip J.. (2016) How to count citations if you must. American Economic Review, 106 (9). pp. 2722-2741.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/82103>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

How To Count Citations If You Must*

Motty Perry
Department of Economics
University of Warwick
Coventry CV4 7AL

Philip J. Reny[†]
Department of Economics
University of Chicago
Chicago, IL, 60637 USA.

April 2016

Abstract

Citation indices are regularly used to inform critical decisions about promotion, tenure, and the allocation of billions of research dollars. Nevertheless, most indices (e.g., the *h*-index) are motivated by intuition and rules of thumb, resulting in undesirable conclusions. In contrast, five natural properties lead us to a unique new index, the *Euclidean index*, that avoids several shortcomings of the *h*-index and its successors. The Euclidean index is simply the Euclidean length of an individual's citation list. Two empirical tests suggest that the Euclidean index outperforms the *h*-index in practice.

Keywords: citation indices, axiomatic, scale invariance, Euclidean length

1. Introduction

Citation indices attempt to provide useful information about a researcher's publication record by summarizing it with a single numerical score. They provide government agencies, departmental and university committees, administrators, faculty, and students with a simple and potentially informative tool for comparing one researcher to another, and are regularly used to inform critical decisions about funding, promotion, and tenure. With decisions of this magnitude on the line, one should approach the problem of developing a good index as systematically as possible. Doing so here, we are led to a unique new index.¹ This new index, called the *Euclidean index*, is simply the Euclidean length of an individual's citation list.

*We wish to thank Pablo Beker, Faruk Gul, Glenn Ellison, Sergiu Hart, Thierry Marchant, Andrew Oswald, Herakles Polemarchakis, and the editor Debraj Ray and three anonymous referees for helpful comments, Dan Feldman for superb research assistance, and we gratefully acknowledge financial support as follows: Perry from the Israel Science Foundation, and Reny from the National Science Foundation (SES-0922535, SES-1227506). The authors declare that they have no relevant or material financial interests that relate to the research described in this paper.

[†]To whom correspondence should be addressed. E-mail: preny@uchicago.edu; Phone: 773-702-8192.

¹Clearly, reducing a research record to a single index number entails a loss of information. Consequently, no single index number is intended to be sufficient for making decisions about funding, promotion etc. It is but one tool among many for such purposes. But it is enough to hold the view that no tool should move one toward unsound or inconsistent decisions and it is this principle that forms the basis of our analysis.

Perhaps the best-known citation index beyond a total citation count is the h -index (Hirsch 2005). A scholar's h -index is the number, h , of his/her papers that each have at least h citations. By design, the h -index limits the effect of a small number of highly cited papers, a feature which, though well intentioned, can produce intuitively implausible rankings. For example, consider two researchers, one with 10 papers, each with 10 citations (h -index = 10), and another with 8 papers, each with 100 (or even 1000) citations (h -index = 8).

Improvements to the h -index have been suggested. Consider, for example, the g -index (Egghe 2006a, 2006b), which is the largest number g such that the total citation count of the g most cited papers is at least g^2 . The g -index is intended to correct for the insensitivity of the h -index to the number of citations received by the h most cited papers. Many other variations have been suggested since. Yet, like the h -index, they are ad hoc measures based almost entirely on intuition and rules of thumb with insufficient justification given for choosing them over the infinitely many other unchosen possibilities. But for a novel empirical approach to selecting among a new class of h -indices, see Ellison (2012, 2013).²

To illustrate the difficulties that can arise when a systematic approach to choosing an index is not followed, let us consider the very practical and well-recognized problem of comparing the records of individuals in different fields or subfields.³ There is compelling evidence to suggest that for such comparisons to be meaningful, one must rescale each individual's citation list by dividing each entry in it by the average number of citations in that individual's field. Indeed, Radicchi et. al. (2008) observe that while the distributions of citations vary widely across a variety of fields (from agricultural economics to nuclear physics), after rescaling by the average number of citations within a field, the distributions all become virtually identical (see Section 4).

With this in mind, suppose that two macroeconomists, A and B , are being considered for a single position and it is noted that A has the higher h -index and so should be the preferred candidate. But before a final decision is made, a second position as well as a new candidate, C , become available. Candidate C , however, is an industrial organization economist and so a rescaling of the citation lists is necessary to compare the three records across the two fields. But a serious difficulty arises. The rescaling has reversed the ranking of the two macroeconomists. That is, applying the h -index to the rescaled lists produces the ranking: B preferred to C preferred to A , and it is now entirely unclear which one of the two macroeconomists should be hired!

²Ellison (2013) introduces the following class of generalized Hirsch indices. For any $a, b > 0$ and for any citation list, $h_{(a,b)}$ is the number of papers, h , that each have at least ah^b citations. Ellison estimates a and b so that $h_{(a,b)}$ gives the best fit in terms of ranking economists at the top 50 U.S. universities in a manner that is consistent with the observed labor market outcome. Ellison (2012) is a follow-up paper that instead uses data on computer scientists.

³See, e.g. Radicchi et. al. (2008) and the references therein.

But do such reversals occur in practice? According to data collected by Ellison (2013),⁴ the average number of citations per paper among macroeconomists at the top 50 U.S. universities is 98, while among IO economists it is 55.⁵ That is, macroeconomists are cited 1.8 ($= 98/55$) times more often per paper, on average, than IO economists. Applying the h -index to Ellison’s (2013) data, both before and after reducing the macroeconomists’ citations by a factor of 1.8, we find that 95% of them experience at least one pairwise ranking change with another macroeconomist and 60% of them experience at least one strict ranking reversal (see Figure 3 in Section 5).^{6,7} So, for the h -index, the problem of ranking reversals is quite real.

To avoid this and other difficulties, we take an axiomatic approach in our search for an index.⁸ The methodology of this approach is to select, with care, a number of basic properties that an index should have. The advantage of this approach is that it focuses attention on the properties that an index should possess rather than the functional form that it should take. After all, it is the properties we desire an index to possess that should determine its functional form, not the other way around. The five properties that we identify are quite natural for the problem at hand and lead uniquely to the Euclidean index.

Still, one might wonder how well the Euclidean index performs in practice relative to the h -index. To get a sense of this with Ellison’s (2013) data, we assigned a score to each index by considering how well its ranking of economists at the top 50 U.S. universities matched the NRC’s ranking of the departments in which they are employed. More specifically, we increased an index’s score by 1 whenever its ranking of two economists agreed with the NRC’s ranking of their departments and we decreased an index’s score by 1 otherwise, ignoring all ties.⁹ The Euclidean index outscored the h -index in this test and it also outscored a whole family of indices that satisfy a strict subset of our axioms (see Figure 4 in Section 5).¹⁰ Thus,

⁴See Section 5 for more details.

⁵These figures pertain to individuals who are classified by Ellison (2013) as being in a unique field, in this case either IO or macroeconomics. There are 88 out of 226 such macroeconomists and 64 out of 120 such IO economists.

⁶The changes in rank that occur in the data often arise between individuals with similar initial h -indices. Such reversals are then less likely to occur between two randomly chosen individuals. Of course recruitment targets are not chosen at random. To the extent that departments tend to consider individuals of similar stature, and to the extent that the h -index is positively correlated with stature, the frequency of reversals reported here is potentially quite relevant in practice.

⁷One might instead suggest rescaling the h -index itself since this will not affect the within-field ranking. But unless one can justify why the h -index is preferable to an ordinally equivalent index, say $h + \sqrt{h}$, this adjustment too will lead to difficulties. See Section 4.

⁸We are certainly not the first to do so. See e.g., Queseda, 2001; Woeginger, 2008a and 2008b; Marchant, 2009a, 2009b; and Chambers and Miller (2014). But in each of these cases, except Marchant 2009b, the stated goal is to provide axioms for a pre-existing citation index. As far as we are aware, the functional form that our axioms identify has not been proposed for use as a citation index until now.

⁹That is, we assign scores according to Kendall’s (1938) rank correlation coefficient.

¹⁰All of the indices in the family outscore the h -index.

at least with respect to this test, the Euclidean index performs rather well.

The Euclidean index is intended to judge an individual’s record *as it stands*. It is not intended as a means to predict an individual’s record at some future date. If a prediction is important to the decision at hand (as in tenure decisions, for example) then separate methods must first be used to obtain a predicted citation list to which our index can then be applied.¹¹

Finally, it should be noted that our index is ordinal. Indeed, we would argue that virtually all citation indices are ordinal because they do not provide decision makers (administrators, tenure committees, funding committees, etc.) with information such as individual A is “worth twice as much” as individual B . On the other hand, citation indices can and do provide usefully interpretable ordinal information such as “ A is equivalent to an individual who receives twice as many citations as B on every paper” (see Section 3.1). It is up to decision-makers to apply this interpretable information to decide how much A should be paid relative to B , or how much teaching A should be assigned relative to B , or how much funding A should receive relative to B , etc. Given our ordinal perspective, we will have little to say about assigning scores to scholarly journals (e.g., Palacios-Huerta and Volij, 2004 and 2014) since such scores are typically used to translate publications from distinct journals into a common cardinal currency.

The rest of the paper proceeds as follows. In Section 2, we list the five properties that characterize the Euclidean index. Section 3 contains the statement of our main result as well as a brief discussion of the roles of each property. Section 4 discusses the important issue of how to eliminate the well-recognized biases that arise when comparing citation lists of individuals from different fields or subfields. Understanding this issue is central to understanding why we introduce the property that we call “scale invariance.” Section 5 provides the details of the two empirical exercises described above.

2. Five Properties for Citation Indices

In this section, we present five properties that lead to a unique index. Unless otherwise stated, these properties should be thought of as pertaining to citation lists whose cited papers differ only in their number of citations but otherwise have common characteristics, e.g., year of publication, field, number of authors, etc. Taking into account differences in such characteristics is an important issue that is touched upon in Section 4.

Citation indices work as follows. First, an individual’s record is summarized by a finite list of numbers, typically ordered from highest to lowest, where the i -th number in the list

¹¹We thank Glenn Ellison for helpful discussions on this point.

is the number of citations received by the individual's i -th most highly cited paper. The list of citations is then operated upon by some index function to produce the individual's index number. As already mentioned in the introduction (see also Section 4), one must sometimes rescale one individual's citation list to adequately compare it to another individual's list. Consequently, one must be prepared to consider noninteger lists. We therefore take as our domain the set \mathbb{L} consisting of all finite nonincreasing sequences of nonnegative real numbers. Any element of \mathbb{L} is called a *citation list*.¹²

A *citation index* is any continuous function, $\iota : \mathbb{L} \rightarrow \mathbb{R}$, that assigns to each citation list a real number.¹³ Consider first the following four basic properties. A discussion of them follows.

1. **Monotonicity.** The index value of a citation list does not fall when a new paper with sufficiently many citations is added to the list.
2. **Independence.** The index's ranking of two citation lists does not change when a new paper is added to each list and each of the two new papers receives the same number of citations.
3. **Depth Relevance.** It is not the case that, for every citation list, the index weakly increases when any paper in the list is split into two and its citations are divided in any way between them.
4. **Scale Invariance.** The index's ranking of two citation lists does not change when each entry of each list is multiplied by any common positive scaling factor.

The first property, monotonicity, says that for each citation list there is a number (which can depend on the list) such that the index does not fall if a new paper with at least that number of citations is added to the list. All indices that we are aware of satisfy this property. In fact, all indices that we are aware of satisfy the stronger property that they weakly increase when any new paper is added, regardless of how few citations it receives. In contrast, our less restrictive monotonicity property allows (but does not require) the index to fall if a new but infrequently cited paper is added. Note that monotonicity says nothing about the behavior of the index when an existing paper receives additional citations.

¹²Implicit in the convention to list the numbers in decreasing order is that the index value would be the same were the numbers listed in any other order.

¹³Continuity means that for any citation list x , if a sequence of citation lists x^1, x^2, \dots each of whose length is the same as the finite length of x , converges to x in the Euclidean sense, then $\iota(x^1), \iota(x^2), \dots$ converges to $\iota(x)$. The use of the Euclidean metric here is not important. Any equivalent metric, e.g., the taxicab metric, would do.

The second property, independence, is natural given that our index is intended to compare the records – as they currently stand – of any two individuals.¹⁴ It says, in particular, that a tie between two records cannot be broken by adding identical papers to each record. The h -index fails to satisfy independence. For example, if individual A has 10 papers with 10 citations each, and B has 5 papers with 15 citations each, then A 's h -index is 10 and B 's is 5. But if they each produce 10 identical new papers that receive 15 citations each, then A 's h -index is still 10, but B 's h -index increases from 5 to 15. Note that the independence property is agnostic about the effect on the index of adding new papers. Independence allows that adding identically cited papers to two records could separately increase or decrease the indices, so long as the rankings are not reversed.

It is often suggested that a good index should encourage “quality over quantity,” i.e., encourage a smaller number of highly cited papers over a larger number of infrequently cited papers. The third property, depth relevance, takes a very weak stand on this important trade-off. It says that it should *not* be the case that for any fixed number of citations, the index is maximized by spreading them as thinly as possible across as many publications as possible. Depth relevance is satisfied by all indices that we are aware of with the exception of the total citation count.

When distinct fields have significantly different average numbers of citations per paper, citation lists must be rescaled in order to make meaningful cross-field comparisons (see Section 4). The fourth property, scale invariance, ensures that the final ranking of the population is independent of whether the lists are scaled down relative to the disadvantaged field or scaled up relative to the advantaged field. Because the h -index is not scale invariant it can, as already observed in the introduction, reverse the order of individuals in the same field when their lists are adjusted to account for differences across fields (Section 5 illustrates this with empirical data).

In addition to the four basic properties above, let us now introduce a fifth property that we call “directional consistency.” It is motivated as follows.

Consider two individuals with equally ranked citation lists who, over the next year, each receive the same number of additional citations on their most cited paper, and the same number but fewer additional citations on their second most cited paper, etc. Suppose that at the end of the year their lists remain equally ranked. Directional consistency requires that if their papers continue to accrue citations at those same yearly rates, then their lists will continue to remain equally ranked year after year. The formal statement is as follows.

¹⁴Marchant (2009b) appears to be the first to apply this independence property to the bibliometric index problem.

5. Directional Consistency $\forall x, y, d \in \mathbb{L}$, if $\iota(x) = \iota(y)$ and $\iota(x + d) = \iota(y + d)$, then $\iota(x + \lambda d) = \iota(y + \lambda d)$ for all $\lambda > 1$.¹⁵

Observe that each of properties 1-5 is ordinal in the sense that if an index satisfies any one of the properties, then any monotone transformation of that index also satisfies that property. The independence of properties 1-5 is established in the next section following the statement of Theorem 3.1.

3. The Euclidean Index

Call two citation indices *equivalent* if one of them is a monotone transformation of the other. We now state our main result, whose proof is in the appendix.

Theorem 3.1. A citation index satisfies properties 1-5 *if and only if* it is equivalent to the index that assigns to any citation list, (x_1, \dots, x_n) , the number

$$\sqrt{\sum_{i=1}^n x_i^2}.$$

Since this number is the Euclidean length of (x_1, \dots, x_n) , we may call this index the *Euclidean index*, and we denote it by ι_E .

It may be helpful to briefly discuss the roles of properties 1-5. Even though we make no such assumption directly, properties 1-4, together with continuity, imply that the index strictly increases when any existing paper receives additional citations. Independence then implies that the (continuous) index must be equivalent to one that is additively separable (a direct application of Debreu 1960). The symmetry built into the index (see footnote 12) implies that the summand in the additively separable representation consists of a single common function, which must be strictly increasing. Scale invariance then implies that the index must be equivalent to an index of the form $(\sum_{i=1}^n x_i^\sigma)^{1/\sigma}$ for some $\sigma > 0$, and depth relevance implies that $\sigma > 1$. Thus, as Part I of the proof of Theorem 3.1 shows (see the appendix), properties 1-4 determine the index up to the single parameter $\sigma > 1$. Property 5 pins σ down to the value $\sigma = 2$.¹⁶

¹⁵If, for example, $x \in \mathbb{R}_+^n \cap \mathbb{L}$ and $d \in \mathbb{R}_+^m \cap \mathbb{L}$ and $n < m$, then $x + d = (x_1 + d_1, \dots, x_n + d_n, d_{n+1}, \dots, d_m)$.

¹⁶It is interesting to note that properties 1-4 alone imply that for any two lists $x, y \in \mathbb{L} \cap \mathbb{R}^n$, if $x_1 \geq y_1$ and $x_1 + x_2 \geq y_1 + y_2$ and ... $x_1 + \dots + x_n \geq y_1 + \dots + y_n$, with at least one inequality strict, then the index must rank x strictly above y . This follows directly from Hardy et. al.'s (1934) well-known result on second-order stochastic dominance since $\sum_{i=1}^n x_i^\sigma$ is an increasing and strictly convex function for any $\sigma > 1$. Thus, some lists can be compared without appealing to property 5.

While the functional form $(\sum_{i=1}^n x_i^\sigma)^{1/\sigma}$ is novel for citation indices, it has a long tradition in other areas. It is an l_σ norm from classical mathematics, a CES utility function as derived in Burk (1936), an Atkinson (1970) measure of income inequality, and a Foster et. al. (1984) poverty index. In addition, there are several well-known axiomatic characterizations already in the literature, including Burk (1936), Roberts (1980), and Blackorby and Donaldson (1982). The main ingredients of each of these characterizations include assumptions of continuity, symmetry, independence, coordinatewise monotonicity, and scale invariance. In contrast, we do not impose any monotonicity conditions on the coordinates of the index function. The monotonicity property specified here is quite distinct from those in the literature because it refers to the effect of increasing the dimension of the vector that is being evaluated. For example, the index function $\sum_i (x_i - 10)^3$ is not an increasing function of any of its coordinates, yet it satisfies our monotonicity property because adding a paper with more than 10 citations increases the index. Relatedly, the literature cited above focuses on domains whose dimension is fixed at a particular value of n (and it is often assumed that $n \geq 3$) whereas our analysis must include and patch together all possible dimensions $n \in \{1, 2, \dots\}$. Finally, we provide a novel condition that isolates the value $\sigma = 2$.

Properties 1-5 are independent. That is, that no single property is implied by the other four. This can be demonstrated by displaying, for each property, a citation index that satisfies the other four properties but is not equivalent to the Euclidean index identified in Theorem 3.1. Finding five such indices is not difficult.¹⁷

3.1. Linear Homogeneity

Of all the equivalent indices satisfying properties 1-5, only the Euclidean index $\iota_E(x) = (\sum_i x_i^2)^{1/2}$ and its positive multiples are homogeneous of degree 1. This turns out to be convenient.

For example, suppose that the Euclidean index of x is twice that of y , i.e., $\iota_E(x) = 2\iota_E(y)$. How might one describe how the two lists compare to one another in terms that would be helpful to an administrator or someone outside the field in question? The answer is that one could say that “list x is as good as the list that has as many papers as in y , but receives twice as many citations on each of them.”

The reason that this statement is correct is that, by linear homogeneity, $2\iota_E(y) = \iota_E(2y)$. Therefore, $\iota_E(x) = 2\iota_E(y)$ if and only if $\iota_E(x) = \iota_E(2y)$, and this last equality means precisely

¹⁷For example, $\iota = 1/n$ satisfies all properties but monotonicity; $\iota = \max(x_1, \dots, x_n)$ satisfies all properties but independence; $\iota = \sum_{i=1}^n x_i$ satisfies all properties but depth relevance; $\iota = \sum_{i=1}^n x_i^2 + n$ satisfies all properties but scale invariance; and $\iota = \sum_{i=1}^n x_i^3$ satisfies all properties but directional consistency. In addition, the lexicographic order satisfies all properties but continuity. None of these indices is equivalent to the Euclidean index.

what is stated in the quotation marks in the previous paragraph.

Thus, the Euclidean index, ι_E , is particularly convenient because the ratio, λ , of the Euclidean index of x to that of y tells us that “list x is as good as the list that has as many papers as list y , but receives λ times as many citations on each of them.” In particular, if an individual’s Euclidean index is twice that of the median Euclidean index in his field, then the individual is equivalent to one who produces twice as many citations per paper as the median researcher in his field.

4. Why Rescale the Lists?

Our justification for the scale invariance property rests on the claim that the appropriate way to adjust for differences in fields is to divide each entry in an individual’s citation list by the average number of citations in that individual’s field. We now support this claim by reviewing the work in this direction due to Radicchi et. al. (2008).¹⁸




Fig. 1: Histograms of raw citations across fields in 1999, where c_0 is the average number of citations per paper published in 1999 in that field. (From Radicchi et. al., 2008).

In Figure 1, we reproduce Figure 1 in Radicchi et. al. (2008), showing normalized (relative to the total number of citations) histograms of unadjusted citations received by papers published in 1999 within each of 5 fields out of 20 considered. The main point of this figure is to show that the distributions of unadjusted citations differ widely across fields. For example, in Developmental Biology, a publication with 100 citations is 50 times more frequent than in Aerospace Engineering. However, after adjusting each citation received by a year-1999 paper by dividing it by the average number of citations of all the year-1999

¹⁸It should be noted that, for the most part, Radicchi et. al. hold fixed the year of publication so that the age of the papers considered is the same.

papers in the same field, the distinct histograms in Figure 1, remarkably, align to form single histogram, shown in Figure 2 (in both Figures, the axes are on a logarithmic scale).¹⁹

We encourage the reader to consult the Radicchi et. al. paper for details, but let us point out that several statistical methods are used there to verify what seems perfectly clear from the figures, namely, that dividing by the average number of citations per paper in one’s field corrects for differences in citations across fields in the very strong sense that, after the adjustment, the distributions of citations are virtually identical across fields.²⁰




Fig. 2: Histograms of adjusted citations across fields in 1999. (From Radicchi et. al., 2008).

4.1. Why not rescale the index itself?

Another way to adjust for differences in fields might be to adjust the index itself. For example, a popular method for comparing h -indices across fields is to divide each individual’s h -index by the average h -index in his/her field.²¹ This produces, for each individual, an adjusted h -index, called the h_s -index in the literature, that is then used to compare any two individuals in the population, regardless of field. This adjustment has the obvious but important property

¹⁹A consequence of this alignment is that the suggested adjustment is unique in the following sense. Restricting to papers published in 1999, let \tilde{c}_j denote the random variable describing the distribution of citations in field j , and let \bar{c}_j denote its expectation. The suggested adjustment is to divide any field j paper’s number of citations, c , by \bar{c}_j , giving the adjusted entry c/\bar{c}_j . If any increasing functions ϕ_1, ϕ_2, \dots align the distributions across fields – i.e., are such that the adjusted random variables $\phi_1(\tilde{c}_1), \phi_2(\tilde{c}_2), \dots$ all have the same distribution – then there is a common increasing function f such that $\phi_j(c) = f\left(\frac{c}{\bar{c}_j}\right)$ for every j .

²⁰In Radicchi et al. (2008), the common histogram in Figure 2 is estimated as being log normal with variance 1.3 (the mean must be 1 because of the rescaling).

²¹This h -index adjustment is suggested by Kaur et. al. (2013) and it is used by the online citation analysis tool “Scholarometer” (see <http://scholarometer.indiana.edu>), whose popularity for computing and comparing the h -indices of scholars across disciplines seems to be growing.

that it does not affect the ranking of individuals within the same field. The following example shows, however, that the final ranking of the population depends on the particular numerical representation of the h -index that is employed.

Example 4.1. *There are two mathematicians, M_1 and M_2 , and two biologists, B_1 and B_2 , with each individual representing half of his field. Their h -indices are 2, 8, 7 and 27, respectively so that their ranking according to the raw h -index is*

$$M_1 \prec B_1 \prec M_2 \prec B_2, \quad (4.1)$$

where \prec means “is ranked worse than.” To compare mathematicians to biologists, let us divide each of the mathematician’s indices by their field’s average $((2+8)/2 = 5)$ and divide each of the biologist’s h -indices by their field’s average $((7+27)/2 = 17)$. The resulting adjusted indices, h_s , are respectively, 0.4, 1.6, 0.41, and 1.58, producing the final adjusted ranking,

$$M_1 \prec B_1 \prec B_2 \prec M_2. \quad (4.2)$$

But suppose that instead of starting the exercise with the h -index, we start with the monotonic transformation $h^* = h + \sqrt{h}$. The h^* -indices of M_1, M_2, B_1 , and B_2 are, respectively, $2 + \sqrt{2}$, $8 + \sqrt{8}$, $7 + \sqrt{7}$, and $27 + \sqrt{27}$, i.e., 3.4, 10.8, 9.6, and 32.2, yielding the same ranking (4.1) as the ordinally equivalent h -index.

However, if for h^* we follow the same procedure and divide each individual’s h^* -index by the average h^* in his/her field (i.e., divide the mathematicians’ h^* -indices by 7.1 and divide the biologists’ by 20.9), then the final adjusted ranking becomes,

$$B_1 \prec M_1 \prec M_2 \prec B_2. \quad (4.3)$$

Comparing (4.2) with (4.3), we see that the ranking of M_1 and B_1 and of M_2 and B_2 depend on which of the ordinally equivalent indices h or h^* that one begins with.²² Absent a compelling argument for choosing one over the other, one must conclude that this adjustment procedure is not well-founded.

²²Indeed, the ranking in (4.2) is impossible to obtain through any rescaling of the mathematicians’ and the biologists’ h^* indices because there is no $\lambda > 0$ such that $\lambda(2 + \sqrt{2}) < (7 + \sqrt{7}) < (27 + \sqrt{27}) < \lambda(8 + \sqrt{8})$.

5. Two Empirical Exercises

We conclude by providing the details of the two empirical exercises described in the Introduction.²³ In both exercises we rely on a dataset constructed by Glen Ellison who graciously made it available to us. Ellison (2013) describes his data as follows: “The dataset includes citation records of almost all faculty at the top 50 economics departments in the 1995 NRC ranking. Faculty lists were taken from the departmental websites in the fall of 2011. Citation data for each economist were collected from Google Scholar. Economists were classified into one or more of 15 subfields primarily by mapping keywords appearing in descriptions of research interests on departmental or individual websites. Slightly less than half of economists are classified as working in a single field.”

When one looks, even casually, at Ellison’s citation data, one cannot help but notice the large differences in the average numbers of citations across fields, from a high of 108 per paper in international trade to a low of 30 per paper in economic history. To correct for these differences when comparing scholars from different fields, one must first rescale the individual citation lists, as explained in section 4, by dividing each scholars’ citations, paper by paper, by the average number of citations in that paper’s field. For scholars whose papers are in a single field, all citations should be divided by the average number of citations in that field.

Our first exercise illustrates that use of the h -index can lead to serious practical problems when efforts are made to correct for differences in fields of study because rescaling has a nontrivial effect on the h -index’s within-field rankings. In this exercise we restricted attention to individuals in Ellison’s (2013) data who are classified as working in a single field. For the 88 individuals who are classified as macroeconomists, the average number of citations per paper is 97.9, while the average number of citations per paper for the 64 IO economists is 55.17. Therefore, to adequately compare macroeconomists to IO economists one must scale down the macroeconomists’ citation lists by a factor of $(97.9)/(55.17) = 1.77$.

Figure 3 shows that the h -index’s within-field ranking of macroeconomists is nontrivially affected by this rescaling. Each numbered space on the horizontal axis corresponds to a macroeconomist, with space k denoting the macroeconomist with the k -th highest h -index before rescaling. The heights of the bars indicate either the total number of ranking changes (lighter colored bars) or the number of strict ranking reversals (darker colored bars) experienced by each macroeconomist as a result of rescaling.

For example, after rescaling all of the macroeconomists’ lists, macroeconomist number 47 experiences 8 within-field strict ranking reversals and a total of 13 within-field ranking

²³We are very grateful to the editor and referees for encouraging us to include some empirical facts along with our theoretical analysis.

changes. That is, there are 8 macroeconomists (the height of the darker-colored bar) who either have a strictly higher h -index than macroeconomist 47 before rescaling but a strictly lower h -index after or vice versa; and there are 5 additional macroeconomists (for a total of 13, the height of the lighter-colored bar) who either have the same h -index as macroeconomist 47 before rescaling but a different h -index after or vice versa.

While a number of macroeconomists experienced no ranking changes at all (e.g., macroeconomist number 1 was top-ranked both before and after rescaling), 95% of the 88 macroeconomists experienced at least one pairwise change in their h -index ranking as a result of rescaling, and 60% experienced at least one strict pairwise ranking reversal. In contrast, the Euclidean index, being scale invariant, generates the same within-field ranking of macroeconomists before and after rescaling.




Fig. 3: The h -index is susceptible to ranking changes after rescaling for differences in fields.

Our second exercise is intended to provide an empirically relevant and reasonably objective comparison between the Euclidean index and the h -index. Inspired by Ellison (2013),

and again using his data, we tested each index to see how well its ranking of economists at the top 50 U.S. universities matched the NRC’s ranking of the departments in which they are employed. To do this, we first rescaled the citation lists of each economist by the average number of citations in his/her field (with economic history as the numeraire). We then assigned a score to each index as follows. Starting from a score of zero, we increased an index’s score by 1 whenever its strict ranking of two economists agreed with the NRC’s strict ranking of their departments and we decreased an index’s score by 1 otherwise, ignoring all ties. Summing all the +1’s and –1’s and then dividing by the total number of distinct pairs of economists yields the index’s score, a number between –1 and +1. That is, we computed Kendall’s (1938) rank correlation coefficient (or “tau coefficient”) between the ranking of economists according to each of the two indices’ (*h*-index and Euclidean index) and the NRC’s ranking of their departments.

The Euclidean index outscored the *h*-index in this test, producing a score of 0.1846, over 14% higher than the *h*-index’s score of 0.1614. And both indices’ rankings of economists reject (with a p-value of 0.4% for the Euclidean index and 2.6% for the *h*-index) the null hypothesis of mutual independence with the NRC’s ranking.²⁴

In addition to testing the Euclidean index against the *h*-index, we tested the Euclidean index against a whole family of indices that nest the Euclidean index. Specifically, we computed Kendall’s rank correlation coefficient between the ranking of economists given by the “ σ -index” $(\sum_{i=1}^n x_i^\sigma)^{1/\sigma}$ with $\sigma > 1$, and the NRC’s ranking of their departments. We did this for all values of σ between 1 and 4. The family of σ -indices with $\sigma > 1$ is precisely the family of indices that satisfy properties 1-4 (see the discussion following Theorem 3.1 in Section 3).

²⁴Under the null hypothesis of mutual independence between an index’s ranking and the NRC’s ranking, the scores are (see Kendall 1938) well approximated by a normal distribution with mean zero and standard deviation $2/(3\sqrt{n}) = 0.071$, where $n = 88$ is the number of economists who are classified as working in the single field of macroeconomics.




Fig. 4: The Euclidean index outperforms the h-index in matching labor market data.

Figure 4 (solid line) graphs the value of Kendall's correlation coefficient as a function of σ . For comparison, we have also included in Figure 4 the value of Kendall's coefficient achieved by the Euclidean index (solid dashed line) and the lower value achieved by the h -index (dotted line). As is evident from the figure, for all values of σ between 1 and 4, Kendall's coefficient for the σ -index is larger than that for the h -index. Moreover, Kendall's coefficient achieves its maximum value of 0.1849 when $\sigma = 1.85$, a value of σ that is remarkably close to the value of $\sigma = 2$ that defines the Euclidean index. Thus, the Euclidean index, in addition to outscoring the h -index, outscores a whole family of indices satisfying properties 1-4, namely the family of σ -indices for values of $\sigma > 1$ that are not too close to 1.85.

A. Appendix

Proof of Theorem 3.1. Since sufficiency is clear, we focus only on necessity. We separate the proof of necessity into two parts. Part I imposes only properties 1-4, and shows that the index must be equivalent to $\sum_i x_i^\sigma$ for some $\sigma > 1$. Part II shows that for any $\sigma > 1$, the index $\sum_i x_i^\sigma$ satisfies property 5 only if $\sigma = 2$.

Proof of Part I. Suppose that the (continuous) citation index $\iota : \mathbb{L} \rightarrow \mathbb{R}$ satisfies properties 1-4. For every n , it will be convenient to extend ι from the set of nonincreasing and nonnegative n -vectors to all of \mathbb{R}_+^n by defining, for any $(x_1, \dots, x_n) \in \mathbb{R}_+^n$, $\iota(x_1, \dots, x_n) := \iota(X_1, X_2, \dots, X_n)$, where X_i is i -th order statistic of x_1, \dots, x_n . Thus ι is now defined on the extended domain $\mathbb{L}^* := \cup_n \mathbb{R}_+^n$ in such a way that it is symmetric; i.e., if $x \in \mathbb{L}^*$ is a permutation of $x' \in \mathbb{L}^*$, then $\iota(x) = \iota(x')$. Clearly, the extended function ι satisfies the extensions to \mathbb{L}^* of properties 1-4, and for any $n \geq 1$ its restriction to \mathbb{R}_+^n is continuous. We work with the extended index ι from now on.

Let us first show that adding a paper with zero citations leaves the index unchanged. Let z be any element of \mathbb{L}^* . We wish to show that $\iota(z) = \iota(z, 0)$. By depth relevance, there is a list $(x_1, w) \in \mathbb{L}^*$, where w may be the empty list, and there are numbers $y_1, y_2 \geq 0$

with $y_1 + y_2 = x_1$ such that $\iota(x_1, w) > \iota(y_1, y_2, w)$. By independence, $\iota(x_1) > \iota(y_1, y_2)$ and by scale invariance, $\iota(\lambda x_1) > \iota(\lambda y_1, \lambda y_2)$ for all $\lambda > 0$. Taking the limit as $\lambda \rightarrow 0^+$ gives, by continuity, that $\iota(0) \geq \iota(0, 0)$. Applying independence once gives $\iota(z, 0) \geq \iota(z, 0, 0)$ and applying it again gives $\iota(z) \geq \iota(z, 0)$. For the reverse inequality note that by monotonicity, there is $m > 0$ such that $\iota(m, 0) \geq \iota(0)$. By scale invariance, $\iota(\lambda m, 0) \geq \iota(0)$ for all $\lambda > 0$. Taking the limit as $\lambda \rightarrow 0^+$ gives, by continuity, that $\iota(0, 0) \geq \iota(0)$. Then, by independence, $\iota(z, 0, 0) \geq \iota(z, 0)$, and applying independence once more gives $\iota(z, 0) \geq \iota(z)$, as desired.

We next show that $\iota(1) > \iota(0)$. By monotonicity, there is $m > 0$ such that $\iota(m, 0) \geq \iota(0)$. Since by the previous paragraph $\iota(m, 0) = \iota(m)$, we have $\iota(m) \geq \iota(0)$. Since $m > 0$, scale invariance implies that $\iota(1) \geq \iota(0)$. It remains to show that $\iota(1) \neq \iota(0)$. Suppose, by way of contradiction, that $\iota(1) = \iota(0)$. Then, by scale invariance, $\iota(x_1) = \iota(0)$ for all $x_1 \geq 0$. Then for all $x_2 \geq 0$, by independence, $\iota(x_1, x_2) = \iota(x_2, 0) = \iota(x_2) = \iota(0)$, where the second equality follows from the previous paragraph. Continuing in this manner shows that $\iota(x_1, \dots, x_n) = \iota(0)$ for all n and all $x_1, \dots, x_n \geq 0$, contradicting depth relevance. We conclude that $\iota(1) > \iota(0)$.

We next show that, for every $n \geq 1$, ι is *strongly increasing* on \mathbb{R}_+^n i.e.,

$$\iota(x'_i, x_{-i}) > \iota(x_i, x_{-i}), \quad \forall x \in \mathbb{R}_+^n, \quad \forall i, \quad \text{and} \quad \forall x'_i > x_i. \quad (\text{A.1})$$

To establish (A.1) it suffices to show, by independence, that $x'_i > x_i$ implies $\iota(x'_i) > \iota(x_i)$; note that x_i and x'_i here are real numbers (e.g., x_i is a list consisting of 1 paper with x_i citations). By scale invariance, it then suffices to show that $\iota(1) > \iota\left(\frac{x_i}{x'_i}\right)$. So, let $\lambda = x_i/x'_i < 1$ and suppose, by way of contradiction, that $\iota(\lambda) \geq \iota(1)$. Then $\lambda > 0$ since $\iota(1) > \iota(0)$ as shown above. Hence, we may apply scale invariance to $\iota(\lambda) \geq \iota(1)$ using the scaling factor λ . Doing so k times gives $\iota(\lambda^{k+1}) \geq \iota(\lambda^k) \geq \dots \geq \iota(\lambda) \geq \iota(1)$. Taking the limit as $k \rightarrow \infty$ of $\iota(\lambda^{k+1}) \geq \iota(1)$ implies, because ι is continuous and $\lambda < 1$, that $\iota(0) \geq \iota(1)$. But this contradicts $\iota(1) > \iota(0)$ and so establishes (A.1).

For any $n \geq 1$, let $\iota_n : \mathbb{R}_+^n \rightarrow \mathbb{R}$ be the restriction of ι to \mathbb{R}_+^n . For any subset I of $\{1, \dots, n\}$ and for any $x \in \mathbb{R}_+^n$ let $x_I = (x_i)_{i \in I}$ and let $I^c = \{1, \dots, n\} \setminus I$.

We next show that ι_n satisfies Blackorby and Donaldson's (1982) "complete strict separability" condition.²⁵ That is, $\forall x, x', y \in \mathbb{R}_+^n$ and $\forall I \subseteq \{1, \dots, n\}$,

$$\iota_n(x_I, x_{I^c}) \geq \iota_n(x_I, y_{I^c}) \text{ iff } \iota_n(x'_I, x_{I^c}) \geq \iota_n(x'_I, y_{I^c}).^{26} \quad (\text{A.2})$$

To see that (A.2) holds, note that $\iota_n(x_I, x_{I^c}) \geq \iota_n(x_I, y_{I^c})$ iff $\iota(x_I, x_{I^c}) \geq \iota(x_I, y_{I^c})$ iff $\iota(x_{I^c}) \geq \iota(y_{I^c})$ iff $\iota(x'_I, x_{I^c}) \geq \iota(x'_I, y_{I^c})$ iff $\iota_n(x'_I, x_{I^c}) \geq \iota_n(x'_I, y_{I^c})$ as desired, where the second and third equivalences follow from independence.

Because, for each n , the continuous and symmetric function $\iota_n : \mathbb{R}_+^n \rightarrow \mathbb{R}$ satisfies (A.1), (A.2), and scale invariance, we may apply Theorem 2, part (i), in Blackorby and Donaldson (1982) (see also Theorem 6 in Roberts 1980) to conclude that for all $n \geq 3$, there exist

²⁵Or, equivalently, Debreu's (1960) "factor-independence" condition.

²⁶Where, e.g., $x_I = (x_i)_{i \in I}$.

$a_n \in \mathbb{R}$ and a strictly increasing $\phi_n : \mathbb{R}_{++} \rightarrow \mathbb{R}$ such that,

$$\iota_n(x) = \phi_n \left(\left(\sum_i x_i^{a_n} \right)^{\frac{1}{a_n}} \right), \text{ for all } x \in \mathbb{R}_{++}^n,$$

where we adopt the convention that $(\sum_i x_i^{a_n})^{\frac{1}{a_n}} := \prod_i x_i$ if $a_n = 0$. From this we may conclude that ι_n is equivalent to $(\sum_i x_i^{a_n})^{\frac{1}{a_n}}$ on the *strictly positive* orthant \mathbb{R}_{++}^n .

We claim that $a_n > 0$. Indeed, let $w(\alpha) = (1, \dots, 1, \alpha) \in \mathbb{R}_+^n$ and suppose first that $a_n < 0$. Then $\lim_{\alpha \rightarrow 0^+} (\sum_i (w_i(\alpha))^{a_n})^{\frac{1}{a_n}} = 0$. Consequently, because the functional form of ι_n above implies that $\iota_n(1, 0, \dots, 0) \leq \iota_n(w(\alpha))$, we have $\iota_n(1, 0, \dots, 0) \leq \lim_{\alpha \rightarrow 0^+} \iota_n(w(\alpha)) = \lim_{\alpha \rightarrow 0^+} \phi_n \left((\sum_i (w_i(\alpha))^{a_n})^{\frac{1}{a_n}} \right) < \phi_n(\lambda n^{\frac{1}{a_n}}) = \iota_n(\lambda \mathbf{1}_n)$ holds for any $\lambda > 0$, where the strict inequality follows because ϕ_n is strictly increasing and $\lim_{\alpha \rightarrow 0^+} (\sum_i (w_i(\alpha))^{a_n})^{\frac{1}{a_n}} = 0 < \lambda n^{\frac{1}{a_n}}$. Hence, $\iota_n(1, 0, \dots, 0) \leq \lim_{\lambda \rightarrow 0^+} \iota_n(\lambda \mathbf{1}_n) = \iota_n(0, \dots, 0)$, which contradicts (A.1). A similar argument holds for $a_n = 0$.

Hence, for every $n \geq 3$, a_n is strictly positive and ι_n is equivalent to $\sum_{i=1}^n x_i^{a_n}$ on \mathbb{R}_{++}^n . But since both ι_n and $\sum_{i=1}^n x_i^{a_n}$ are continuous on \mathbb{R}_+^n we may conclude that ι_n is equivalent to $\sum_{i=1}^n x_i^{a_n}$ on \mathbb{R}_+^n . And because adding a paper with zero citations does not change the value of the index, we may further conclude that $a_n = a_{n+1}$ for all $n \geq 3$.

Thus, we have so far shown that there exists $\sigma > 0$ such that, for any $n \geq 3$ and for any lists $x, y \in \mathbb{R}_+^n$ of the same length n , $\iota_n(x) \geq \iota_n(y)$ iff $\sum_{i=1}^n x_i^\sigma \geq \sum_{i=1}^n y_i^\sigma$. It remains only to characterize the index when comparing lists whose common length is less than 3, and when comparing lists of different lengths. But such comparisons can be made by first adding zeros to the two lists to be compared so that the lengths of the resulting lists are equal and at least 3. Since adding zeros does not affect the value of the index the correct comparison can now be obtained by applying to the lengthened lists the function $\sum_i x_i^\sigma$. Hence, we may conclude that ι is equivalent to the index that assigns to any citation list $(x_1, \dots, x_n) \in \mathbb{L}^*$ of any length n the number $\sum_{i=1}^n x_i^\sigma$.

It remains only to show that $\sigma > 1$. But this follows from depth relevance because $0 < \sigma \leq 1$ implies $(x_1^\sigma + x_2^\sigma) \geq (x_1 + x_2)^\sigma \forall x_1, x_2 \geq 0$, which implies $(x_1^\sigma + x_2^\sigma) + \sum_i z_i^\sigma \geq (x_1 + x_2)^\sigma + \sum_i z_i^\sigma$ for all nonnegative $z = (z_1, \dots, z_n)$, which implies $\iota(x_1, x_2, z) \geq \iota(x_1 + x_2, z)$, and so depth relevance fails.²⁷ This completes the proof of Part I.

Proof of Part II. We must show that if $\sigma > 1$ and $\iota(x) = \sum_i x_i^\sigma$ satisfies property 5, then $\sigma = 2$. In fact, we will show the even stronger result that if $\sigma > 1$ and $\sum_i x_i^\sigma$ satisfies property 5 for lists of length 2 then $\sigma = 2$. So, from now on we restrict attention to lists of length 2, i.e., to vectors in $\mathbb{L} \cap \mathbb{R}_+^2$.

Let $x = ((1/2)^{1/\sigma}, (1/2)^{1/\sigma})$ and let $y = (1, 0)$. Then,

$$\iota(x) = \iota(y) = 1$$

By Lemma A.1 below, there exists $d_1 > 1$ such that

$$((1/2)^{1/\sigma} + d_1)^\sigma + ((1/2)^{1/\sigma} + 1)^\sigma = (1 + d_1)^\sigma + 1. \quad (\text{A.3})$$

²⁷For $\sigma \in (0, 1]$, the function $(x_1^\sigma + x_2^\sigma) - (x_1 + x_2)^\sigma$ has nonnegative partial derivatives when $x_1, x_2 > 0$ and so it achieves a minimum value of 0 on \mathbb{R}_+^2 at, for example, $x_1 = x_2 = 0$.

Since $d_1 > 1$, the vector $(d_1, 1)$ is in $\mathbb{L} \cap \mathbb{R}_+^2$. Letting $d = (d_1, 1)$, (A.3) says that,

$$\iota(x + d) = \iota(y + d).$$

Therefore, by property 5, we must have,

$$\iota(x + \lambda d) = \iota(y + \lambda d), \quad \forall \lambda > 1.$$

That is,

$$((1/2)^{1/\sigma} + \lambda d_1)^\sigma + ((1/2)^{1/\sigma} + \lambda)^\sigma = (1 + \lambda d_1)^\sigma + \lambda^\sigma, \quad \forall \lambda > 1.$$

Dividing this equality by λ^σ , letting $\alpha = (1/2)^{1/\sigma}$, and letting $\beta = 1/\lambda$ gives

$$(\beta\alpha + d_1)^\sigma + (\beta\alpha + 1)^\sigma = (\beta + d_1)^\sigma + 1, \quad \forall \beta \in (0, 1). \quad (\text{A.4})$$

Assume, by way of contradiction, that σ is not a positive integer. Then, for any integer $k = 1, 2, 3, \dots$ we may differentiate (A.4) k times with respect to β and take the limit of the resulting equality as $\beta \rightarrow 0$ giving,

$$\alpha^k (d_1)^{\sigma-k} + \alpha^k = (d_1)^{\sigma-k}. \quad (\text{A.5})$$

Solving for d_1 gives,

$$d_1 = \left(\frac{\alpha^k}{1 - \alpha^k} \right)^{\frac{1}{\sigma-k}} = (2^{k/\sigma} - 1)^{\frac{1}{k-\sigma}}, \quad (\text{A.6})$$

where the second equality uses the definition $\alpha = (1/2)^{1/\sigma}$.

Since (A.6) holds for every $k = 1, 2, 3, \dots$ it must also hold in the limit as $k \rightarrow \infty$. We claim that

$$\lim_{k \rightarrow \infty} (2^{k/\sigma} - 1)^{\frac{1}{k-\sigma}} = 2^{1/\sigma}. \quad (\text{A.7})$$

To see this, replace k with the continuous variable $z \in (1, \infty)$ and take the limit of the log of the expression of interest. This gives,

$$\begin{aligned} \lim_{z \rightarrow \infty} \log (2^{z/\sigma} - 1)^{\frac{1}{z-\sigma}} &= \lim_{z \rightarrow \infty} \frac{\log (2^{z/\sigma} - 1)}{z - \sigma} \\ &= \lim_{z \rightarrow \infty} \frac{(2^{z/\sigma} - 1)^{-1} 2^{z/\sigma} \log(2^{1/\sigma})}{1} \\ &= \log(2^{1/\sigma}), \end{aligned}$$

where the second equality follows by l'Hopital's rule This proves (A.7).

Since, (A.6) must hold, in particular, for $k = 1$, and also in the limit, we must have

$$2^{1/\sigma} = (2^{1/\sigma} - 1)^{\frac{1}{1-\sigma}}.$$

Raising both sides to the power $1 - \sigma$ gives,

$$2^{(1-\sigma)/\sigma} = 2^{1/\sigma} - 1.$$

Since the left-hand side is $(1/2)(2^{1/\sigma})$, we obtain

$$(1/2)(2^{1/\sigma}) = 1,$$

or, equivalently,

$$2^{1/\sigma} = 2.$$

But this last equality implies that $\sigma = 1$, which is a contradiction since $\sigma > 1$. Hence, σ must be an integer greater than 1.

Next, assume by way of contradiction, that $\sigma \neq 2$, i.e., that σ is an integer greater than 2. Since (A.6) is valid for all $k = 1, \dots, \sigma - 1$, it is valid, in particular for $k = \sigma - 1$ and $k = \sigma - 2 \geq 1$. Hence,

$$(2^{(\sigma-1)/\sigma} - 1)^{\frac{1}{(\sigma-1)-\sigma}} = (2^{(\sigma-2)/\sigma} - 1)^{\frac{1}{(\sigma-2)-\sigma}},$$

which is equivalent to,

$$(2(2^{-1/\sigma}) - 1)^{-1} = (2(2^{-2/\sigma}) - 1)^{-1/2},$$

which, after cross-multiplying and squaring, is equivalent to

$$\begin{aligned} 2(2^{-2/\sigma}) - 1 &= (2(2^{-1/\sigma}) - 1)^2 \\ &= 4(2^{-2/\sigma}) - 4(2^{-1/\sigma}) + 1. \end{aligned}$$

Gathering terms, this is equivalent to,

$$\begin{aligned} 0 &= 2(2^{-2/\sigma}) - 4(2^{-1/\sigma}) + 2 \\ &= 2(2^{-1/\sigma} - 1)^2, \end{aligned}$$

which is a contradiction, because the last expression on the right-hand side is positive for all finite values of $\sigma > 1$. We conclude that $\sigma = 2$. ■

Lemma A.1. *There exists $d_1 > 1$ that solves (A.3).*

Proof. Since the functions in (A.3) are continuous, it suffices to show: (a) when $d_1 = 1$ in (A.3), the left-hand side is larger than the right-hand side, and (b) the right-hand side of (A.3) is larger than the left-hand side for all d_1 sufficiently large.

Starting with (a), set $d_1 = 1$ in (A.3). We wish to show that

$$2((1/2)^{1/\sigma} + 1)^\sigma > 2^\sigma + 1. \tag{A.8}$$

The left-hand side of (A.8) is equal to $(1 + 2^{1/\sigma})^\sigma$. Taking the σ -th root of both sides, we see that (A.8) holds iff,

$$1 + 2^{1/\sigma} > (2^\sigma + 1)^{1/\sigma},$$

which can be re-written as,

$$(1^\sigma + 0^\sigma)^{1/\sigma} + (1^\sigma + 1^\sigma)^{1/\sigma} > (2^\sigma + 1^\sigma)^{1/\sigma}.$$

But this last strict inequality holds by the Minkowski inequality for the vectors $(1, 0)$, $(1, 1)$ and $(2, 1)$ under the l_p norm $\|(a_1, a_2)\|_p = (a_1^p + a_2^p)^{1/p}$ with $p = \sigma > 1$, because $(1, 0)$ and

(1, 1) are not proportional.²⁸ This establishes (A.8) and therefore (a). We now turn to (b).
 Let $\alpha = (1/2)^{1/\sigma}$. It suffices to show that,

$$\lim_{w \rightarrow \infty} [(1+w)^\sigma - (\alpha+w)^\sigma] = +\infty.$$

We can establish this limit as follows.

$$\begin{aligned} \lim_{w \rightarrow \infty} [(1+w)^\sigma - (\alpha+w)^\sigma] &= \lim_{r \rightarrow 0^+} [(1 + \frac{1}{r})^\sigma - (\alpha + \frac{1}{r})^\sigma] \\ &= \lim_{r \rightarrow 0^+} \frac{[(r+1)^\sigma - (\alpha r + 1)^\sigma]}{r^\sigma} \\ &= \lim_{r \rightarrow 0^+} \frac{[(r+1)^{\sigma-1} - \alpha(\alpha r + 1)^{\sigma-1}]}{r^{\sigma-1}} \\ &= +\infty, \end{aligned}$$

where the third equality follows by l'Hopital's rule because $\sigma > 0$, and the fourth since the numerator of the third line tends to $1 - \alpha > 0$ and the denominator tends to 0 from the right because $\sigma > 1$. This proves (b). ■

References.

- Atkinson, A. B. (1970): "On the measurement of inequality," *Journal of Economic Theory*, 2, 244–263.
- Blackorby, C. and D. Donaldson (1982): "Ratio-Scale and Translation-Scale Fair Interpersonal Comparability without Domain Restrictions: Admissible Social-Evaluation Functions," *International Economic Review*, 23, 249-268.
- Burk (Bergson), A. (1936): "Real income, expenditure proportionality, and Frisch's 'New Methods of Measuring Marginal Utility'," *The Review of Economic Studies*, 4, 33-52.
- Chambers, C. P., and A. D. Miller (2014): "Scholarly Influence," *Journal of Economic Theory*, 151 571-583.
- Debreu, G. (1960): "Topological Methods in Cardinal Utility," In Kenneth J. Arrow, Samuel Karlin, and Patrick Suppes, eds., *Mathematical methods in the social sciences*, Stanford University Press, Stanford, California, 16-26.
- Egghe, L. (2006a): "An improvement of the h-index: The g-index," *ISSI Newsletter*, 2, 8–9.
- Egghe, L. (2006b): "Theory and practice of the g-index," *Scientometrics*, 69 (1):131–152.
- Ellison, G. (2012): "Assessing Computer Scientists Using Citation Data," Department of Economics, MIT, <http://economics.mit.edu/files/7784>, and NBER.
- Ellison, G. (2013): "How Does the Market Use Citation Data? The Hirsch Index in Economics," *American Economic Journal: Applied Economics*, 5, 63-90.
- Foster, J., J. Greer and E. Thorbecke (1984): "A class of decomposable poverty measures," *Econometrica*, 3, 761–766.

²⁸We thank Sergiu Hart for this Minkowski inequality argument.

- Hardy G. H., J. E. Littlewood, and G. Polya (1934): *Inequalities*. Cambridge University Press.
- Hirsch, J. E. (2005): “An index to quantify an individual’s scientific research output,” *Proceedings of the National Academy of Sciences*, 102, 16569–16572.
- Kendall, M. G. (1938): “A New Measure of Rank Correlation,” *Biometrika*, 30, 81–89.
- Kaur J., F. Radicchi, and F. Menczer (2013), “Universality of scholarly impact metrics,” *Journal of Informetrics*, 7, 924– 932.
- Marchant, T. (2009a): “An axiomatic characterization of the ranking based on the h-index and some other bibliometric rankings of authors,” *Scientometrics*, 80, 325–342.
- Marchant, T. (2009b): “Score-based bibliometric rankings of authors,” *Journal of the American Society for Information Science and Technology*, 60, 1132-1137.
- Palacios-Huerta, I., and O. Volij (2004): “The Measurement of Intellectual Influence,” *Econometrica*, 72, 963-977.
- Palacios-Huerta, I., and O. Volij (2014), “Axiomatic Measures of Intellectual Influence,” *International Journal of Industrial Organization*, 34, 85-90.
- Perry, M., and P. J. Reny (2014): “How to Count Citations if You Must,” working paper version, <https://sites.google.com/site/philipjreny/home/research>.
- Quesada, A. (2011): “Axiomatics for the Hirsch index and the Egghe index,” *Journal of Infometrics*, 5, 476-480.
- Radicchi, F., S. Fortunato, and C. Castellano (2008): “Universality of citation distributions: Toward an objective measure of scientific impact,” *Proceedings of the National Academy of Sciences*, 105 (45):17268–17272.
- Roberts, K. W. S. (1980): “Interpersonal Comparability and Social Choice Theory,” *Review of Economic Studies*, 47,421-446.
- Woeginger, G. L. (2008a): “An axiomatic analysis of Egghe’s g-index,” *Journal of Infometrics*, 2, 364-368.
- Woeginger, G. L. (2008b): “An axiomatic characterization of the Hirsch-index,” *Mathematical Social Sciences*, 56, 224-232