

Original citation:

Hoffman, Karen P., Playford, E. Diane, Grill, Eva, Soberg, Helene L. and Brohi, Karim. (2016) Minimum data set to measure rehabilitation needs and health outcome after major trauma : application of an international framework. *European journal of physical and rehabilitation medicine*, 52 (3). pp. 331-46.

Permanent WRAP URL:

<http://wrap.warwick.ac.uk/82142>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Published version: <http://www.minervamedica.it/en/journals/europa-medicophysica/article.php?cod=R33Y2016N03A0331>

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP URL' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

ORIGINAL ARTICLE

**Minimum data set to measure rehabilitation needs and health outcome after major
trauma: Application of an international framework**

SHORT TITLE: REHABILITATION NEEDS AND HEALTH OUTCOME AFTER MAJOR
TRAUMA

Karen P. HOFFMAN ¹*, Diane E. PLAYFORD ², Eva GRILL ³, Helene L. SOBERG ⁴,
Karim BROHI ¹

¹Centre for Trauma Sciences, Blizard Institute, Queen Mary University of London, London, UK; ²Queen Square National Hospital for Neurology and Neurosurgery, UCL Institute of Neurology, London, UK; ³Biometry and Epidemiology (IBE), Institute for Medical Informatics, Ludwig-Maximilians-Universität Munich, Munich, Germany; ⁴Department of Physical Medicine and Rehabilitation, Oslo University Hospital, Oslo, Norway

*Corresponding author: Karen Hoffman, Centre for Trauma Sciences, Blizard Institute, Queen Mary University of London, 4 Newark Street, London, E1 2AT. E-mail: K.hoffman@qmul.ac.uk

ABSTRACT

BACKGROUND: Measurement of long term health outcome after trauma remains non-standardized and ambiguous which limits national and international comparison of burden of injuries. The World Health Organization (WHO) has recommended the application of the International Classification of Function, Disability and Health (ICF) to measure rehabilitation and health outcome worldwide. No previous poly-trauma studies have applied the ICF comprehensively to evaluate outcome after injury.

AIM: To apply the ICF categorization in patients with traumatic injuries to identify a minimum data set of important rehabilitation and health outcomes to enable national and international comparison of outcome data.

DESIGN: A mixed methods design of patient interviews and an on-line survey.

SETTING: An ethnically diverse urban major trauma center in London.

POPULATION: Adult patients with major traumatic injuries (poly-trauma) and international health care professionals (HCPs) working in acute and post-acute major trauma settings.

METHODS: Mixed methods investigated patients and health care professionals (HCPs) perspectives of important rehabilitation and health outcomes. Qualitative patient data and quantitative HCP data were linked to ICF categories. Combined data were refined to identify a minimum data set of important rehabilitation and health outcome categories.

RESULTS: Transcribed patient interview data (N.=32) were linked to 234 (64%) second level ICF categories. Two hundred and fourteen HCPs identified 121 from a possible 140 second level ICF categories (86%) as relevant and important. Patients and HCPs strongly agreed on ICF body structures and body functions categories which include temperament, energy and drive, memory, emotions, pain and repair function of the skin. Conversely, patients prioritised domestic tasks, recreation and work compared to HCP priorities of self-care and mobility.

Twenty six environmental factors were identified. Patient and HCP data were refined to recommend a 109 possible ICF categories for a minimum data set.

CONCLUSIONS: The comprehensive measurement of health outcomes after trauma is important for patients, health professionals and trauma systems. An internationally applied ICF minimum data set will standardize the language used and concepts measured after major trauma to enable national and international comparison of outcome data.

CLINICAL REHABILITATION IMPACT: A minimum ICF data set for trauma will standardize rehabilitation language and provide a minimum dataset to capture outcome in trauma systems to enable comparison and service improvement.

(Cite this article as: Hoffman KP, Playford DE, Grill E, Soberg HL, Brohi K. Minimum data set to measure rehabilitation needs and health outcome after major trauma: Application of an international framework. Eur J Phys Rehabil Med 2016;)

Key words: Rehabilitation - Multiple trauma - Outcomes assessment (Health Care).

Major trauma involving multiple injuries is a leading contributor to the global burden of disease and mortality for working aged adults.¹ However, as survival from injury improves with modern trauma care, so does the importance of the rehabilitation and health outcomes for these patients.^{2, 3} Many severely injured trauma patient will experience prolonged periods in critical care which often leads to additional non-injury related morbidity.^{4, 5} The accurate and comprehensive assessment of health outcomes, which includes physical, psychological and social wellbeing,^{6, 7} as well as the impact on the family and society is fundamental to inform future efforts to improve lives after injury.⁸ However existing generic outcome measures only assess a small proportion of the overall burden experienced by trauma patients.^{9, 10} Moreover,

the lack of validated trauma specific outcome measures has caused uncertainty amongst experienced trauma clinicians regarding which key domains should be measured as part of a minimum data set.¹¹⁻¹³ As a consequence, little is known about the multidimensional effects that major traumatic injury has on outcomes important to patients.^{14, 15} Many measures fail to evaluate the patients' perception of their health outcome and recovery, and consequently there may be little agreement between patient priorities and goals set by rehabilitation professionals.¹⁶ Moreover, the patients' performance within their social context needs to be considered as part of the rehabilitation process, as support and systems can either hinder or help recovery.¹⁷ This improved knowledge would inform rehabilitation priorities to support patient recovery and facilitate cost effective trauma service delivery and resource utilization.¹¹ Outcomes derived from a minimum data set would enable assessment of trauma system performance while patients return to health after injury.^{18, 19}

Recent developments of trauma systems in the United Kingdom and Europe necessitate a robust, patient centered data set to measure health outcome after trauma. Not only are prehospital and in-hospital outcomes important, factors such as the impact of critical care, secondary complications and population burden after discharge need to be considered.⁸ In line with current methodology, new health measures should be developed within an internationally recognized framework,²⁰ involving patients and health care professionals to ensure wide acceptance and validity. The International Classification of Function, Disability and Health (ICF)²¹ is a hierarchically organized framework consisting of approximately 1400 health categories. The World Health Organisation (WHO) has recommended its application in all health outcome studies.²² ICF Core Sets have been applied in many rehabilitation settings and health conditions including traumatic brain injury²³ and spinal cord injury.²⁴ However, patients with multiple traumatic injuries have additional, potentially more complex needs to those with isolated brain or spinal injury. Recent publications have called for the development

of core sets specifically for use in trauma outcome evaluation and rehabilitation.²⁵⁻²⁷ The ICF health categories most applicable for multiply injured patients are yet to be reported.

The primary aim of this study was to identify important ICF health and rehabilitation categories from the perspective of patients and health care professionals (HCPs). We aimed to investigate patient and HCP perspectives of important health outcomes following trauma with the ICF framework as a reference to identify a minimum data set of important rehabilitation and health outcomes to enable national and international comparison of outcome data. Secondly we wished to evaluate to what degree components of body functions, body structures and activities and participation are affected by trauma in relation to contextual issues such as the environmental and personal factors.

Materials and methods

ICF definitions

The ICF is a comprehensive and international framework that considers human functioning as a whole. Part one captures health and disability in terms of how body functions (b), body structures (s) and activities and participation (d) are affected. Part two comprises of contextual factors and includes environmental (e) and personal factors (Figure 1). All aforementioned components, except personal factors, consist of several chapters which contain hierarchically organized categories including second and third level sub-categories. Each category contains specific health issues and becomes more detailed in second and third level sub-categories, similar to ICD-10 categories. For example: b1 Mental functions' (first level category), b114 Orientation functions' (second level category), b1142 Orientation to person (third level category).

Study design

Mixed methods were used to investigate important health outcomes after trauma. We conducted qualitative semi-structured, face-to-face interviews with patients. Health care professionals (HCPs) participated in an international on-line questionnaire to identify important health issues after trauma. Ethical approval was gained from an NHS Research Ethics Committee.

Qualitative patient interviews

STUDY POPULATION

Participants were recruited from an urban major trauma centre in London that admits on average 1800 patients with multiple traumatic injuries each year. Discharged adult patients (≥ 18 years of age) were invited to participate via a postal invitation, which was followed up by a telephone call. A convenience sample of patients with varying ages, genders, time since injury, injury mechanisms, and injury severity were recruited. The severity of injury was categorised using the Injury Severity Score (ISS).²⁸ We aimed to capture all potential problems rather than which problems occur at specific timeframes, therefore patients were approached at a variety of time frames since injury as health problems and the prevalence of symptoms may change over time. This form of purposive sampling was used to obtain a wide-ranging sample of this heterogeneous patient population and has been used successfully in previous studies.²⁹ A sample size of 40-45 participants was chosen whilst aiming for a 50% recruitment rate to ensure sufficient interview data. Previous qualitative studies suggest that data saturation can be reached with 20-30 interviews.³⁰ Data saturation is the point at which no new data is generated by participant interviews in an effort to reduce bias and increase trustworthiness.

DATA COLLECTION

Where possible, interviews were scheduled to coincide with other hospital appointments to reduce patient burden. Written consent was obtained prior to all interviews. Semi-structured interviews were conducted by one researcher (Karen Hoffman, KH) over a three month period using an interview guide containing six questions based on the components of the ICF and rehabilitation priorities. These questions related to body functions, body structures, carrying out usual activities, participating in meaningful tasks and issues related to environmental barriers or facilitators and personal factors. A final question allowed for any other comments. Interviews were digitally recorded and transcribed verbatim.

On-line questionnaire for health care professionals

INCLUSION CRITERIA

Any HCP registered with a professional body with at least five years' experience of working in trauma and able to read and write English were able to participate. A health care professional was defined as a doctor, nurse, physiotherapist, occupational therapist, psychologist, social worker, dietician and speech and language therapist (non-exhaustive list). It was not the intention to get the view of specific HCPs, but rather to obtain an overview of important rehabilitation categories as identified by trauma experts. This would facilitate application of a minimum data set for trauma rehabilitation in multi-disciplinary team settings.

SAMPLE SIZE AND RECRUITMENT

Previous ICF studies recruited between 21 and 126 HCPs to develop condition specific ICF Core Sets.³¹⁻³³ Their recruitment rate varied between 50%-70% and completion rate between 36-50%.³¹⁻³³ In order to ensure sufficient representation from an international trauma

community we aimed to recruit 70% of invited participants with a completion rate of 50%. We invited two groups of HCPs to participate. Group one was a purposively selected cohort known to be experts in trauma care and were invited to participate via email. The trauma experts were from international settings with representation from five of the World Health regions and were identified via peer review publications, trauma organizations and societies. The second group of trauma HCPs were recruited through web based invitations posted on profession specific or special interest group websites.

DATA COLLECTION

Consent to participate was obtained at the start of the online questionnaire, using Survey Monkey® online web platform.³⁴ Rather than presenting all 1400 detailed ICF categories for the online questionnaire, researchers identified 140 ICF categories from the Acute and Post-acute Comprehensive ICF Core Sets,³⁵⁻³⁷ previously developed by experts working in orthopedic, neurological and cardio-vascular settings using a Delphi process. These specific core sets were selected for their similarities to the early stages and post-acute injury rehabilitation, scientific rigor used to develop them and to reduce respondent burden. During the online questionnaire HCPs were asked to rate the prevalence and importance of ICF categories for trauma patients. Some health issues may not be very prevalent (common) but could have a significant impact on rehabilitation and outcome and should be given greater importance.

Data analysis for patient interviews and on-line questionnaire

Transcribed interviews were analyzed using the meaning condensation procedure, typically used during ICF core set development.³⁸ Transcribed text was reviewed and grouped into meaningful units or themes. In a second step, health and rehabilitation concepts contained in

the meaningful units were identified. A meaningful unit can contain more than one concept, *e.g.* “I get tired when walking”. This contains concepts related to mobility (walking) and energy levels (fatigue). In a final step, each identified concept was systematically linked to ICF categories according to published ICF linking rules.^{39, 40} Detailed third level categories were converted into less detailed second level categories to enable comparison with HCP data which included only second level ICF categories. Absolute and relative frequencies were calculated. Relative frequency aims to establish prevalence, and refers to the number of patients who mentioned a specific category, thus the category was recorded only once per interview despite possibly being mentioned several times. Absolute frequency was used to capture the magnitude or total number of times a category was mentioned, where some patients may mention the same category or concept more than once. For example, many patients mentioned *pain* several times during the interview. Once ICF categories were identified, a process of member checking, sometimes referred to as respondent validation,²⁹ was completed in an effort to ensure all aspects important for patients were captured. Member checking helps to validate data accuracy and reduce bias and patients returned summaries of transcribed data, confirming important categories. In line with previous ICF patient perspective studies, categories with a relative frequency between $\geq 5\%$ and $< 10\%$ were reported and regarded as low to moderate frequency categories.^{41, 42} Categories with a frequency $\geq 15\%$ were considered as high frequency categories⁴² and relevant for comparison with categories identified by HCPs.

HCP questionnaire data were analyzed using descriptive statistics. Each of the 140 ICF categories contained two response options. One captured how prevalent or common a category was (not common, common or very common) and the other how important a category was (not important, important, very important). The average for both options are

presented as a percentage, representing how many HCPs regarded a specific ICF category as common (prevalent) or important.

Previous consensus conference cut off points of $\geq 50\%$ were used to indicate low to moderate HCP agreement.⁴³ Categories with a combined frequency (prevalence and importance) of $\geq 70\%$ were regarded as high frequency categories⁴² and relevant for comparison with categories identified by HCPs.

Finally, moderate to high frequency ICF categories identified by patients and HCPs were compared and condensed. All categories which had a patient identified relative frequency of $\geq 15\%$ and all HCP categories with a combined frequency (prevalence and importance) of $\geq 70\%$ were included in the final minimum data set representing moderate to high frequencies of relevant ICF categories. Consensus between HCP reported data was set at 70% to ensure adequate level of agreement.^{33, 44}

Results

Patients

Forty four patients were invited to participate and 35 (80%) consented to be interviewed (Table I). Participants that did not provide written consent (N.=9) were younger males (30 years), more recently injured (3-9 month post injury) and more severely injured (ISS: 25). Of these, six did not attend their out-patient clinic appointment after providing verbal telephone consent, two opted not to participate as they were too busy and one patient could not be reached after he was included in the sample. Health concepts captured from three of the interviews were excluded as they related to non-trauma chronic medical issues. The majority of patient participants were male (87%, N.=27) with an average age of 37 years (range 18-75). Participants with blunt injuries were older (42 years vs. 25 years) compared to participants

with penetrating trauma, but similar severity of injury (ISS: 20 vs. ISS 22) (Table I). Finally, those who had suffered blunt trauma had longer timeframes since injury (9 months vs. 4 months) in comparison to penetrating injuries.

A total of 2742 health concepts were extracted from the interviews. These were linked to 388 second and third level ICF categories. Third level categories were converted to a total of 234 second level ICF categories. Of these, 146 (62%) had a relative frequency of $\geq 5\%$ and were identified by two or more patients (Tables II-V). Due to the high level of categories identified by less than two patients (38%), data saturation was inferred. Fifty five concepts were too general to be linked or related to personal factors which currently do not have ICF categories. An average of 86 ICF categories was identified per patient (range: 15 to 182 categories).

Health care professionals

Three hundred and twenty nine HCPs consented to participate and 214 completed the questionnaire (65%), (Table VI). From the 210 invited trauma experts in group one, 179 agreed to participate and 128 (72%) completed the questionnaire. Group two, had 179 participants with a completion rate of 48% (N.=86).

The majority of HCPs who commenced the questionnaire (N.=256; 78%) had more than eight years' trauma experience (Table VI), and 66% were based in an acute care setting (N.=216). Nineteen percent (N.=61) of participants worked in both the acute and post-acute setting. The majority of respondents (N.=224; 68%) were allied health professionals (AHPs) including therapists and nurses. Participating doctors (N.=105; 32%) were from emergency medicine, general surgery, traumatology, critical care, anesthesia, vascular and orthopedic surgery and rehabilitation settings. Of the 140 ICF categories presented to the HCPs 86% were regarded as relevant for trauma patients ($\geq 50\%$ common and important).

Comparison of patient and HCP responses

Low to moderate frequency categories of patient ($\geq 5\%$) and HCP ($\geq 50\%$) data were combined resulting in 192 categories with a frequency above 5% for patient data and 50% for HCP. Patients identified some ICF categories which were not included (NI) in the HCP questionnaire and vice versa (Tables II-V). The greatest overlap of categories between patients and HCP was seen in the body functions (b) and body structures (s) component. There was a high level of agreement for 7 body function categories which were temperament and personality functions (b126), energy and drive functions (b130), memory functions (b144), emotional functions (b152), experience of self and time functions (b180), sensation of pain (b280) and repair function of the skin (b820) (Table II). Seven body functions were considered important by HCPs, but not by patients. These were level of consciousness (b110), psychomotor problems (b147), perceptual problems (b156), blood pressure (b420), respiratory functions (b440), muscle endurance (b740) and gait pattern (b770). In terms of *body structures*, patient identified an additional eight structures (Table III). Very few patients discussed issues related to body structures during the interviews and lower extremity (s750) was mentioned the most frequently (63%). The biggest discrepancy was seen for structures of the brain (s110), spinal cord (s120), the shoulder region (s720) and areas of skin (s810) (Table III).

There was less agreement between patients and HCPs in areas of activities and participation (d) (Table IV). Fifteen categories identified by patients were not included in the HCP questionnaire. In contrast, four categories within the *Learning and applying Knowledge* (d1) chapter were identified as important by clinicians but were rarely reported by patients. Patients prioritized tasks such as housework, shopping and helping others as part of *domestic life* (chapter 6) compared to HCPs that prioritised items of self-care (chapter 5). Both patients and HCPs agreed on the importance of walking (d450), looking after one's health (d570) and

handling stress and other psychological demands (d240). They also agreed on family relationships (d760). Patients identified additional relationships such as social (d750) and intimate relationships (d770). Remunerative employment (d850) does not occur in the comprehensive ICF core set and was not included in the HCP questionnaire, although economic self-sufficiency (d870) is, and this was regarded as very important by HCPs. Similarly recreation and leisure (d920) was very important for patients, whereas problems with engaging in community life (d910) were identified in the HCP questionnaire.

There were also discrepancies in the environmental factors (e) component (Table V). Environmental categories, which refer to contextual factors, are coded as barriers or facilitators to functioning. The most important environmental facilitators identified by both patients and HCPs were support and attitudes of immediate family members (e310, e410), support from and attitudes of health care professionals (e355, e450) and health services, systems and policies (e580). Many patients (41%) described social security, systems and policies (e570) as a barrier to recovery where HCPs reported these as facilitators. The third level category *drugs* (e1101) or medication was aggregated to the second level category substances for personal consumption (e110) which was seen as both a barrier and a facilitator for patients. HCPs identified 742 personal factors which could either help or hinder recovery such as age, education and attitude. Personal factors in patient interviews related to dealing with death of a relative involved in the incident, the impact of injuries on retirement, attitudes towards recovery and the content of nightmares.

Table VII presents the final list combined of ICF categories important for rehabilitation and health outcome as frequently identified by patients and HCPs. These are the combined categories with a relative frequency of $\geq 15\%$ for patient data and/or a frequency of $\geq 70\%$ for HCPs categories. The final list consists of 109 categories comprising of 29 body functions, 11 body structures, 41 activity and participation categories and 28 environmental factors. There

was 100% agreement for body function and body structure categories. Patients identified 39 (93%) of the activity and participation categories compared to 27 (62%) identified by HCPs. Environmental factors had an 88% overlap between HCPs and patients.

Discussion

This mixed methods study describes the breadth and importance of rehabilitation and health outcomes after severe injury, using an internationally accepted framework of function and disability. We demonstrate that it is possible to engage both trauma patients and HCPs in rehabilitation research using the ICF as a reference with acceptable levels of recruitment and participation, well above our expected threshold of 50%. Both patients and health care practitioners identify a broad range of categories, the majority of which are not captured by individual outcome measures.⁹ There was substantial agreement between patients and HCPs in areas related to body structure and body function. However within the activity, participation and environmental components there were categories which patients deemed important but were not considered significant or prevailing by HCPs. Conversely some areas considered important by HCPs were not frequently recognized by patients. There was disagreement in some contextual factors of the environment as to whether particular factors were more likely to be barriers or facilitators to recovery. One previous study used the ICF to as a conceptual framework to develop a list of all deficits (LOAD) to capture the overall burden of traumatic injuries.⁸ Population burden was calculated using a combination of 20 domains consisting of individual, family and societal deficits. The purpose of the Injury LOAD framework was to list all deficits and highlight the importance of each factor to prompt the scientific community to recognize the wider impact of injury and to develop population metrics.⁸ Whilst this work is complementary to our study it has a strong emphasis

on cost burden, presented as population burden which is not the focus of our work. Rather, the application of a minimum data set will improve standardized data collection to enable comparison and description of rehabilitations needs, input and outcomes of trauma populations, where after burden can be accurately evaluated.

Measuring health outcome of major trauma is complex, particularly due to varied injury patterns and heterogeneous patient populations. By dividing patient and HCP data into high and low frequency categories in our study we defined common and important problems faced by trauma patients. This methodology, frequently used in ICF studies, appears to provide useful insights into trauma patients' recovery priorities. For example, trauma outcomes research has historically used clinician rated dependency measures such as the Barthel Index⁴⁵ or disability measures such as the Functional Independence Measure.⁴⁶ These are most often applied in sub-acute and primary rehabilitation settings, whereas other outcome studies measured longer term function using health related quality of life (HRQL) measures thus not capturing all aspects of health outcome. Safety and independence in self-care is often a discharge requirement⁴⁷ and HCPs may spend some time getting patients independent or provide equipment to reduce risk prior to discharge.³ This focus may not be important for patients' during or after acute hospitalization. Our data demonstrate that work, finance and domestic tasks were of greater importance to patients than self-care activities. Others have reported a discrepancy between rehabilitation goals described by trauma patients' and their rehabilitation professionals with only a 40% agreement related to work or education and 19% agreement for recreation and leisure.¹⁶ The combined data, and proposed data set provides a comprehensive set of key domains which should be considered for trauma outcomes assessment. This may focus rehabilitation priorities on patients-driven outcomes and increase engagement between patients and HCPs.¹⁶ It will also direct and facilitate health service delivery and research.⁴⁸

Existing outcome measures individually capture only a fraction of the health outcomes identified in this study. In a recent systematic review,⁹ we found that commonly used measures collectively assessed a maximum of 29 ICF categories (8% of the total 2nd level categories and 2% of overall ICF). The majority of these measures were developed with a narrow scope based on chronic disease models,⁴⁹ and not specifically for trauma.⁵⁰ Only one measure, the Trauma Outcomes Profile (TOP)⁵¹ used in one study, captured 61 ICF categories (17% second level categories). However, 14 of these related to pain in different body structures, thus only 47 (13%) unique ICF categories. Furthermore, less than 2% of all environmental factors were captured with these outcome measures⁹ despite evidence of the impact of education, access to medical insurance, trauma systems and support services has on outcome after injury.⁵² Thus, there is a real concern that studies using existing measures of health outcomes after trauma do not comprehensively reflect the greater impact of injury on patients' lives.

As an example, post-traumatic stress disorder (PTSD) is not evaluated by existing generic outcome measures⁵³ although PTSD scales are used intermittently in specific patient populations such as the military⁵⁴ and critical care.⁵ In our study patients did not diagnose themselves with PTSD but rather described factors that impact on their ability to manage stress after trauma. These included stress management (d240) which was the third most limiting factor in activity and participation. Patients also reported difficulty in managing temperament (b126); struggling to look after their own health (d570); and loss of productivity in terms of remunerative employment (d850). Similarly, environmental factors such as limited access to health care services, health care professional attitudes and lack of support systems or service are known to affect the recovery burden.⁵² Assessment of these factors are of importance and should be considered during rehabilitation and outcome assessment, especially given the discrepancy between patient and HCP in relation to which environmental

factors are barriers or facilitators to recovery. The importance of family support and relationships (d760 and e310) has been highlighted in traumatic brain injury rehabilitation.⁵⁵

It was identified by more than half of the patients and nearly 90% of HCPs despite a lack of research on the impact of family support and relationships in major trauma.

Finally, work (d850) and leisure (d970) were the most important activity and participation categories identified by patients but are not routinely measured in trauma outcome studies despite the evidence that emphasizes the importance of these aspects.⁵⁶ Productivity losses, due to traumatic injuries, are estimated to be billions of pounds each year due to the demographic make-up of trauma patients, such as age, education and compensation status,⁵⁷

⁵⁸ The benefit of sport and leisure activities was demonstrated in survivors of spinal cord injury.⁵⁹ Our data demonstrate that outcomes that are not captured by existing measures are non-trivial despite widespread recognition of their important impact on recovery.

The ICF framework shows clear potential in its ability to capture rehabilitation and health outcomes of trauma patients. Our methodology has produced a potential minimum data set of 109 ICF categories that may be used to measure rehabilitation priorities and health outcome of injured patients in the future. It also prompts clinicians to consider contextual issues such as environmental factors as these will impact recovery and access to services. The coded categories derived from this study will enable national and international comparison of rehabilitation and health outcome after injury using a standardized language.

There are a number of limitations to this study. Patient interviews were conducted on a small sample from only one trauma center albeit in an ethnically diverse major city. Secondly, patients were at home and this possibly reflects the prioritization of domestic tasks over self-care or other tasks. Moreover, people of different age groups may have different perceptions and attitudes to functional limitations and older people may report no difficulty in a task despite changes in the way they used to perform the task.⁶⁰ These differences and perceptions

were not analyzed or captured during the interviews but will add important depth to understanding the degree of difficulty experienced by some. HPCs identified health issues related to body functions which were not identified by patients and were excluded from the final data set due to low frequencies. This could be attributed to the fact that the majority of HCPs were based in acute services and had knowledge of acute problems, rather than community services. HPCs are also knowledgeable about specific body functions such as consciousness (b110) and psychomotor problems (b147) which is not necessarily problematic for patients in the community.

Other issues to consider are the impact of missing data in terms of patients consented to participate (N.=32 from 44 patients invited; 73%) and completion rate of the on-line questionnaire (N.=214 from 329 HCP's invited; 65%). Although a sample size of 32 participants are acceptable to ensure data saturation in qualitative interviews,⁶¹ it need to be recognized that the patients that declined to participate were younger and more severely injured. They have similar characteristics to those described as "lost to follow-up" in other trauma studies,^{62, 63} and may have different problems and priorities from the sample interviewed. We therefor recognize that more work needs to be done to ensure their needs are captured in ICF categories to enable generalization to all trauma patients. In addition, even though we achieved an above average response- and completion rate for the on-line questionnaire,⁶⁴⁻⁶⁶ we acknowledge that there are a substantial amount of missing data for non-respondents (N.=115; 35%). A variety of factors contributed to this for which we received feedback. These include the length of the questionnaire, invitations to clinicians that are too busy or HCPs regarding the topic as irrelevant to their area of practice. We did not analyze the characteristic of non-respondents or in-complete questionnaires. As a consequence this may limit generalization and application of results due to an under representation of the views of certain HCPs or the international community. Whilst

recognizing this, we aim to address some of these issues at an international consensus conference with appropriate international representation from a variety of health care professionals.

We propose an international consensus conference, which includes patients, patient representatives and professionals, to further debate issues and reach agreement of the principal categories to include in a minimum dataset for trauma. The final categories can also be grouped in terms of setting. Acute and post-acute settings may have different priorities, resulting in some body structure categories being made redundant. This will warrant wide application and international acceptance to ensure implementation and adherence to allow comparison of rehabilitation and outcome after trauma.

Conclusions

We have used the internationally recognized ICF framework to describe the range and complexity of health outcomes after injury. The comprehensive assessment of on-going health issues of injured patients is important for individuals, institutions, regional trauma systems, science and society. The strong consensus between an international group of trauma experts and patients presents an opportune prospect for the application of a trauma framework to collect international trauma outcome data. We present the candidate categories for an ICF-based minimum data set for the development of modern health outcome assessment in trauma care.

References

1. Murray CJ, Vos T, Lozano R, Naghavi M, Flaxman AD, Michaud C, *et al.* Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2197-223.
2. Cameron PA, Gabbe BJ, McNeil JJ. The importance of quality of survival as an outcome measure for an integrated trauma system. *Injury* 2006;37:1178-84.

3. Gabbe BJ, Simpson PM, Sutherland AM, Wolfe R, Fitzgerald MC, Judson R, *et al.* Improved functional outcomes for major trauma patients in a regionalized, inclusive trauma system. *Ann Surg* 2012;255:1009-15.
4. Wieske L, Dettling-Ihnenfeldt DS, Verhamme C, Nollet F, van Schaik IN, Schultz MJ, *et al.* Impact of ICU-acquired weakness on post-ICU physical functioning: a follow-up study. *Crit Care* 2015;19:196.
5. Peris A, Bonizzoli M, Iozzelli D, Migliaccio ML, Zagli G, Bacchereti A, *et al.* Early intra-intensive care unit psychological intervention promotes recovery from post traumatic stress disorders, anxiety and depression symptoms in critically ill patients. *Crit Care* 2011;15:R41.
6. Constitution of the World Health Organisation W. Constitution of the World Health Organisation. New York: World Health Organisation; 1946; [Internet], Available from: http://www.who.int/governance/eb/who_constitution_en.pdf [cited 2016, May 2].
7. NICE NIOHaCE. Rehabilitation after critical illness [CG83]. London: National Institute of Health and Care Excellence; 2009.
8. Lyons RA, Finch CF, McClure R, van Beeck E, Macey S. The injury List of All Deficits (LOAD) Framework--conceptualizing the full range of deficits and adverse outcomes following injury and violence. *Int J Inj Contr Saf Promot* 2010;17:145-59.
9. Hoffman K, Cole E, Playford ED, Grill E, Soberg HL, Brohi K. Health Outcome after Major Trauma: What Are We Measuring? *PLoS One* 2014;9:e103082.
10. Grill E, Mittrach R, Müller M, Mutschler W, Schwarzkopf SR. [Systematic review of measurement instruments and concepts used for functioning outcome in multiple trauma]. *Unfallchirurg* 2010;113:448-55.
11. Khan F, Amatya B, Hoffman K. Systematic review of multidisciplinary rehabilitation in patients with multiple trauma. *Br J Surg.* 2012 Jan;99 Suppl 1:88-96. PubMed PMID: 22441861. Epub 03/28. eng.
12. Sleat GK, Ardolino AM, Willett KM. Outcome measures in major trauma care: a review of current international trauma registry practice. *Emerg Med J* 2011;28:1008-12.
13. Brasel KJ, Deroon-Cassini T, Bradley CT. Injury severity and quality of life: whose perspective is important? *J Trauma* 2010;68:263-8.
14. Van Horn ER. Recovery from traumatic injury: Trauma patients' perceptions of facilitators and barriers. *J Burn Care Res* 2010;31:540-50.
15. Anke AG, Fugl-Meyer AR. Life satisfaction several years after severe multiple trauma--a retrospective investigation. *Clin Rehabil* 2003;17:431-42.
16. Soberg HL, Finset A, Roise O, Bautz-Holter E. Identification and comparison of rehabilitation goals after multiple injuries: an ICF analysis of the patients', physiotherapists' and other allied professionals' reported goals. *J Rehabil Med* 2008;40:340-6.
17. Brown M. Perspectives on outcome: what disability insiders and outsiders each bring to the assessment table. *Arch Phys Med Rehabil* 2009;90(11 Suppl):S36-40.
18. Santana MJ, Straus S, Gruen R, Stelfox HT. A qualitative study to identify opportunities for improving trauma quality improvement. *J Crit Care* 2012;27:738.e1-7.
19. Stelfox HT, Boyd JM, Straus SE, Gagliardi AR. Developing a patient and family-centred approach for measuring the quality of injury care: a study protocol. *BMC Health Serv Res* 2013;13:31.
20. Rauch A, Cieza A, Stucki G. How to apply the International Classification of Functioning, Disability and Health (ICF) for rehabilitation management in clinical practice. *Eur J Phys Rehabil Med* 2008;44:329-42.
21. Who WHO. International Classification of Functioning, Disability and Health: ICF. Geneva: World Health Organisation, 2001.

22. Stucki G, Cieza A, Melvin J. The International Classification of Functioning, Disability and Health (ICF): a unifying model for the conceptual description of the rehabilitation strategy. *J Rehabil Med* 2007;39:279-85.
23. Laxe S, Zasler N, Tschiesner U, López-Blazquez R, Tormos JM, Bernabeu M. ICF use to identify common problems on a TBI neurorehabilitation unit in Spain. *NeuroRehabilitation* 2011;29:99-110.
24. Cieza A, Kirchberger I, Biering-Sorensen F, Baumberger M, Charlifue S, Post MW, *et al.* ICF Core Sets for individuals with spinal cord injury in the long-term context. *Spinal Cord* 2010;48:305-12.
25. Rimmer JH. Use of the ICF in identifying factors that impact participation in physical activity/rehabilitation among people with disabilities. *Disabil Rehabil* 2006;28:1087-95.
26. Schwarzkopf SR, Grill E, Dreinhöfer KE. [Using the international classification of functioning, disability and health (ICF) in trauma care]. *Unfallchirurg* 2010;113:441-7.
27. Stucki G, Stier-Jarmer M, Grill E, Melvin J. Rationale and principles of early rehabilitation care after an acute injury or illness. *Disabil Rehabil* 2005;27:353-9.
28. Baker SP, O'Neill B, Haddon W, Jr., Long WB. The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *J Trauma* 1974;14:187-96.
29. Barbour RS. Checklists for improving rigour in qualitative research: a case of the tail wagging the dog? *BMJ* 2001;322:1115-7.
30. Mason M. Sample Size and Saturation in PhD Studies Using Qualitative Interviews. [Single Contributions]: *Forum Qualitative Sozialforschung/Forum: Qualitative Social Research*; 2010 [updated 2010-08-24; cited 2014 18th August]; 11/3,art. 8:[*Forum Qualitative Sozialforschung/Forum: Qualitative Social Research*, 11/3, art. 8]; [Internet]. Available from: <http://www.qualitative-research.net/index.php/fqs/article/view/1428/3027> [cited 2016 May 2].
31. Weigl M, Cieza A, Andersen C, Kollerits B, Amann E, Stucki G. Identification of relevant ICF categories in patients with chronic health conditions: a Delphi exercise. *J Rehabil Med* 2004:12-21.
32. Lemberg I, Kirchberger I, Stucki G, Cieza A. The ICF Core Set for stroke from the perspective of physicians: a worldwide validation study using the Delphi technique. *Eur J Phys Rehabil Med* 2010;46:377-88.
33. Boonen A, van Berkel M, Kirchberger I, Cieza A, Stucki G, van der Heijde D. Aspects relevant for functioning in patients with ankylosing spondylitis according to the health professionals: a Delphi study with the ICF as reference. *Rheumatology (Oxford)* 2009;48:997-1002.
34. SurveyMonkey. SurveyMonkey ® online web platform. Palo Alto, California, USA: SurveyMonkey Inc; [Internet]. Available from: www.surveymonkey.com [cited 2016, MAY 2].
35. Grill E, Strobl R, Muller M, Quittan M, Kostanjsek N, Stucki G. ICF Core Sets for early post-acute rehabilitation facilities. *J Rehabil Med* 2011;43:131-8.
36. Grill E, Ewert T, Chatterji S, Kostanjsek N, Stucki G. ICF Core Sets development for the acute hospital and early post-acute rehabilitation facilities. *Disabil Rehabil* 2005;27:361-6.
37. Müller M, Stier-Jarmer M, Quittan M, Strobl R, Stucki G, Grill E. Validation of the comprehensive ICF Core Sets for patients in early post-acute rehabilitation facilities. *J Rehabil Med* 2011;43:102-12.
38. Malterud K. Systematic text condensation: a strategy for qualitative analysis. *Scand J Public Health* 2012;40:795-805.
39. Cieza A, Brockow T, Ewert T, Amman E, Kollerits B, Chatterji S, *et al.* Linking health-status measurements to the international classification of functioning, disability and health. *J Rehabil Med* 2002;34:205-10.

40. Cieza A, Geyh S, Chatterji S, Kostanjsek N, Ustun B, Stucki G. ICF linking rules: an update based on lessons learned. *J Rehabil Med* 2005;37:212-8.
41. Mueller M, Lohmann S, Thul P, Weimann A, Grill E. Functioning and health in patients with cancer on home-parenteral nutrition: a qualitative study. *Health Qual Life Outcomes* 2010;8:41.
42. Sveen U, Ostensjo S, Laxe S, Soberg HL. Problems in functioning after a mild traumatic brain injury within the ICF framework: the patient perspective using focus groups. *Disabil Rehabil* 2012;35:749-57.
43. Grill E, Quittan M, Huber EO, Boldt C, Stucki G. Identification of relevant ICF categories by health professionals in the acute hospital. *Disabil Rehabil* 2005;27:437-45.
44. Sumsion T. The Delphi Technique: An Adaptive Research Tool. *Br J Occup Ther* 1998;61:153-6.
45. Wade DT, Collin C. The Barthel ADL Index: a standard measure of physical disability? *Int Disabil Stud* 1988;10:64-7.
46. Hetherington H, Earlam RJ, Kirk CJ. The disability status of injured patients measured by the functional independence measure (FIM) and their use of rehabilitation services. *Injury* 1995;26:97-101.
47. Kripalani S, Jackson AT, Schnipper JL, Coleman EA. Promoting effective transitions of care at hospital discharge: A review of key issues for hospitalists. *J Hosp Med* 2007;2:314-23.
48. Jordan J, Dowswell T, Harrison S, Lilford RJ, Mort M. Health needs assessment. Whose priorities? Listening to users and the public. *BMJ* 1998;316:1668-70.
49. Hobart JC, Cano SJ, Zajicek JP, Thompson AJ. Rating scales as outcome measures for clinical trials in neurology: problems, solutions, and recommendations. *Lancet Neurol* 2007;6:1094-105.
50. Gabbe BJ, Williamson OD, Cameron PA, Dowrick AS. Choosing outcome assessment instruments for trauma registries. *Acad Emerg Med* 2005;12:751-8.
51. Attenberger C, Amsler F, Gross T. Clinical evaluation of the Trauma Outcome Profile (TOP) in the longer-term follow-up of polytrauma patients. *Injury*. Netherlands: Elsevier Ltd; 2012. p. 1566-74.
52. Resnik LJ, Allen SM. Using International Classification of Functioning, Disability and Health to understand challenges in community reintegration of injured veterans. *J Rehabil Res Dev* 2007;44:991-1006.
53. Soberg HL, Bautz-Holter E, Roise O, Finset A. Mental health and posttraumatic stress symptoms 2 years after severe multiple trauma: self-reported disability and psychosocial functioning. *Arch Phys Med Rehabil* 2010;91:481-8.
54. Schwartz I, Tsenter J, Shochina M, Shiri S, Kedary M, Katz-Leurer M, *et al.* Rehabilitation outcomes of terror victims with multiple traumas. *Arch Phys Med Rehabil* 2007;88:440-9.
55. Griffin JM, Friedemann-Sánchez G, Jensen AC, Taylor BC, Gravely A, Clothier B, *et al.* The invisible side of war: families caring for US service members with traumatic brain injuries and polytrauma. *J Head Trauma Rehabil* 2012;27:3-13.
56. Holtslag HR, Post MW, van der Werken C, Lindeman E. Return to work after major trauma. *Clin Rehabil* 2007;21:373-83.
57. Finkelstein EA, Corso PS, Miller TR. Incidence and economic burden of injuries in the United States. Oxford: Oxford University Press; 2007.
58. Gabbe BJ, Harris IA, Collie A, Cameron PA. Does access to compensation have an impact on recovery outcomes after injury? Comment. *Med J Aust* 2010;193:188; author reply 90.

59. McVeigh SA, Hitzig SL, Craven BC. Influence of sport participation on community integration and quality of life: a comparison between sport participants and non-sport participants with spinal cord injury. *J Spinal Cord Med* 2009;32:115-24.
60. Ramos-Pichardo JD, Cabrero-Garcia J, Gonzalez-Llopis L, Cabanero-Martinez MJ, Munoz-Mendoza CL, Sanjuan-Quiles A, *et al.* What do older people understand by mobility-related difficulties? *Arch Gerontol Geriatr* 2014;59:122-30.
61. Baker SE, Edwards R. How many qualitative interviews is enough; [Internet]. Available from: http://eprints.ncrm.ac.uk/2273/4/how_many_interviews.pdf [cited 2016, May 2].
62. Langley J, Johnson S, Slatyer M, Skilbeck CE, Thomas M. Issues of loss to follow-up in a population study of traumatic brain injury (TBI) followed to 3 years post-trauma. *Brain Inj* 2010;24:939-47.
63. Krellman JW, Kolakowsky-Hayner SA, Spielman L, Dijkers M, Hammond FM, Bogner J, *et al.* Predictors of follow-up completeness in longitudinal research on traumatic brain injury: findings from the National Institute on Disability and Rehabilitation Research traumatic brain injury model systems program. *Arch Phys Med Rehabil* 2014;95:633-41.
64. Nulty DD. The adequacy of response rates to online and paper surveys: What can be done? *Assessment & Evaluation in Higher Education* 2008;33:301-14.
65. McColl E, Jacoby A, Thomas L, Soutter J, Bamford C, Steen N, *et al.* Design and use of questionnaires: a review of best practice applicable to surveys of health service staff and patients. *Health Technol Assess* 2001;5:1-256.
66. Cook JV, Dickinson HO, Eccles MP. Response rates in postal surveys of healthcare professionals between 1996 and 2005: an observational study. *BMC Health Serv Res* 2009;9:160.

Funding.— Diane Playford was supported by the National Institute for Health Research University College London Hospitals Biomedical Research Centre.

Conflicts of interest.—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript.

Acknowledgements.—We would like to thank all the patients' and HCPs' that participated in the interviews and the online questionnaire. A special thanks to Melissa Selb for her support with this project.

Article first published online: January 8, 2016. - Manuscript accepted: January 4, 2016. -

Manuscript revised: November 27, 2015. - Manuscript received: June 15, 2015.

Figure 1.—ICF framework illustrating four different components body function (b), body structure (s), activity and participation (d) and environmental factors (e). Personal factors currently do not have categories and codes. Numbers indicate total number of second, third and fourth level categories in each component.