

Circulation Research Compendium on Atrial Fibrillation:

Atrial Fibrillation Compendium: Historical Context and Detailed Translational Perspective on an Important Clinical Problem
The Clinical Profile and Pathophysiology of Atrial Fibrillation: Relationships Among Clinical Features, Epidemiology, and Mechanisms

Emerging Directions in the Genetics of Atrial Fibrillation

Cellular and Molecular Electrophysiology of Atrial Fibrillation Initiation, Maintenance and Progression

Role of the Autonomic Nervous System in Atrial Fibrillation: Pathophysiology and Therapy

Mathematical Approaches to Understanding and Imaging Atrial Fibrillation: Significance for Mechanisms and Management

Atrial Fibrillation Therapy Now and in the Future: Drugs, Biologicals, and Ablation

Stanley Nattel, Guest Editor

Role of the Autonomic Nervous System in Atrial Fibrillation Pathophysiology and Therapy

Peng-Sheng Chen, Lan S. Chen, Michael C. Fishbein, Shien-Fong Lin, Stanley Nattel

Abstract: Autonomic nervous system activation can induce significant and heterogeneous changes of atrial electrophysiology and induce atrial tachyarrhythmias, including atrial tachycardia and atrial fibrillation (AF). The importance of the autonomic nervous system in atrial arrhythmogenesis is also supported by circadian variation in the incidence of symptomatic AF in humans. Methods that reduce autonomic innervation or outflow have been shown to reduce the incidence of spontaneous or induced atrial arrhythmias, suggesting that neuromodulation may be helpful in controlling AF. In this review, we focus on the relationship between the autonomic nervous system and the pathophysiology of AF and the potential benefit and limitations of neuromodulation in the management of this arrhythmia. We conclude that autonomic nerve activity plays an important role in the initiation and maintenance of AF, and modulating autonomic nerve function may contribute to AF control. Potential therapeutic applications include ganglionated plexus ablation, renal sympathetic denervation, cervical vagal nerve stimulation, baroreflex stimulation, cutaneous stimulation, novel drug approaches, and biological therapies. Although the role of the autonomic nervous system has long been recognized, new science and new technologies promise exciting prospects for the future. (*Circ Res.* 2014;114:1500-1515.)

Key Words: heart failure ■ myocardial infarction

Autonomic nervous system activation can induce significant and heterogeneous changes of atrial electrophysiology and induce atrial tachyarrhythmias, including atrial tachycardia and atrial fibrillation (AF). The importance of the autonomic nervous system in atrial arrhythmogenesis is also supported by circadian variation in the incidence of symptomatic AF in humans.¹ Methods that reduce autonomic innervation or outflow have been shown to reduce the incidence of spontaneous or induced atrial arrhythmias.²⁻⁶ The latter studies suggest that neuromodulation may be helpful in controlling AF. In this review, we focus on the relationship between

the autonomic nervous system and the pathophysiology of AF and the potential benefit and limitations of neuromodulation in the management of this arrhythmia.

Cardiac Autonomic Innervation

The heart is richly innervated by the autonomic nerves. The ganglion cells of the autonomic nerves are located either outside the heart (extrinsic) or inside the heart (intrinsic). Both extrinsic and intrinsic nervous systems are important for cardiac function and arrhythmogenesis.⁷⁻¹⁰ The vagal nerves include axons that come from various nuclei in the medulla. The

Original received February 19, 2014; revision received March 17, 2014; accepted March 27, 2014.

From the Department of Medicine, Division of Cardiology, The Krannert Institute of Cardiology (P.-S.C., S.-F.L.) and Department of Neurology (L.S.C.), Indiana University School of Medicine, Indianapolis; Department of Pathology and Laboratory Medicine, University of California, Los Angeles (M.C.F.); Institute of Biomedical Engineering, National Chiao Tung University, Hsinchu, Taiwan (S.-F.L.); and Department of Medicine, Montreal Heart Institute and Université de Montréal, Montreal, Quebec, Canada (S.N.).

Correspondence to Peng-Sheng Chen, MD, Indiana University School of Medicine, 1800 N Capitol Ave, E475, Indianapolis, IN 46202. E-mail chenpp@iu.edu
© 2014 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.114.303772

Nonstandard Abbreviations and Acronyms

AF	atrial fibrillation
APD	action potential duration
CaMKII	Ca ²⁺ /calmodulin-dependent protein kinase type II
EAD	early afterdepolarization
LA	left atrium
PAT	paroxysmal atrial tachycardia
PKA	protein kinase A
PV	pulmonary vein
SGNA	stellate ganglion nerve activity
SR	sarcoplasmic reticulum
VNA	vagal nerve activity
VNS	vagal nerve stimulation

extrinsic sympathetic nerves come from the paravertebral ganglia, including the superior cervical ganglion, middle cervical ganglion, the cervicothoracic (stellate) ganglion, and the thoracic ganglia.¹¹ The intrinsic cardiac nerves are found mostly in the atria and are intimately involved in atrial arrhythmogenesis. Figure 1 is a highly simplified illustration of the cardiac autonomic innervation and sites reported to be relevant in neuromodulation to control atrial arrhythmia. Among them,




Figure 1. Autonomic innervation and neuromodulation. Neiguan P6 is an acupoint used in a clinical trial of atrial fibrillation (AF).¹⁹ The black dots indicate sites used by various investigators for neuromodulation to control AF. See Neuromodulation section for details. VLCCN indicates ventral lateral cervical cardiac nerve; and VMCCN, ventromedial cervical cardiac nerve. Illustration Credit: Ben Smith.

the stellate ganglion is a major source of cardiac sympathetic innervation. The stellate ganglion connects with multiple intrathoracic nerves and structures, as well as skin.^{12–15} Figure 2 shows immunohistochemical staining of the major autonomic structures that innervate the heart. The ganglion cells within the stellate ganglion mostly (>90%) stain positive for tyrosine hydroxylase, the rate-limiting enzyme responsible for the synthesis of catecholamines (Figure 2A). However, there are also ganglion cells that are negative for that enzyme (Figure 2B). The negatively stained cells (Figure 2C) stain positively for choline acetyltransferase (Figure 2D),¹⁶ an enzyme responsible for the synthesis of the neurotransmitter acetylcholine. Tyrosine hydroxylase–positive ganglion cells are also found in the cervical vagal nerve of dogs (Figure 2E) and humans.^{17,18} These findings suggest that the sympathetic components in the vagal nerve may serve as a source of sympathetic tone. Because cells that stain positive for tyrosine hydroxylase may also stain positive for choline acetyltransferase (Figure 2E), ganglion cells in the autonomic nerve structures are not only dedicated to produce catecholamines.

Like the stellate ganglion, the vagal nerves also have a complex structure containing mixed nerve types. A large portion of the vagus trunk contains sensory and motor nerves.²⁰ In addition to the parasympathetic structure that sends fibers to various parts of the body,²¹ a sympathetic component is known to be present in the vagal nerves based on physiological observations.^{22–24} These findings were subsequently confirmed with immunohistochemical staining that documented the presence of tyrosine hydroxylase–positive nerve fibers in human and canine vagal nerves.^{17,18,25–27} As shown in Figure 3, the tyrosine hydroxylase–positive nerves are distributed mostly in the periphery of the vagal nerve (Figure 3A–3E), but occasionally tyrosine hydroxylase–positive nerves can extend into the center of the vagal nerve (Figure 3F). Similar findings are found in the thoracic vagal nerves.²⁶ Vagal nerve recordings in ambulatory dogs showed that in 3 dogs isolated vagal nerve activation induces tachycardia (Figure 3G), consistent with activation of the sympathetic component of the vagal nerves.

In addition to these extrinsic cardiac nerves, the heart is also well innervated by the intrinsic cardiac nerves.^{9,28} Histological study of human pulmonary vein (PV)–left atrium (LA) junction²⁹ showed that numerous autonomic nerves are present. The nerve densities are the greatest in the LA within 5 mm of the PV–LA junction and are higher in the epicardium than endocardium. Adrenergic and cholinergic nerves are strongly colocalized at tissue and cellular levels. A significant proportion (30%) of ganglion cells expresses dual adrenergic-cholinergic phenotypes (ie, stain positive for both tyrosine hydroxylase and cholineacetyltransferase). Because these nerve structures are highly colocalized, it is difficult to perform radiofrequency catheter ablation that selectively eliminates purely sympathetic or parasympathetic arms of the autonomic nervous system.

Neuroplasticity

In addition to the complex anatomic and physiological interactions between various nerve structures, cardiac autonomic innervation is also constantly remodeling, especially during disease states. Pathological examinations of diseased hearts by Vracko et al^{30,31} showed findings consistent with cardiac




Figure 2. Presence of both adrenergic and cholinergic nerves structures in the extrinsic cardiac nervous system. **A**, Low-power view of the left stellate ganglion, showing numerous ganglion cells and nerve fibers stained positively for tyrosine hydroxylase (TH). Although most of the ganglion cells are TH positive, some ganglion cells were negative (**B**). **C**, TH staining of a different stellate ganglion, showing TH-negative cells (arrows). These same cells stained positive for cholineacetyltransferase (ChAT, arrows in **D**). Some cells stain positive for both markers. Reprinted from Shen et al¹⁶ with permission of the publisher. **E** and **F**, TH and cholineacetyltransferase stains, respectively, of the canine left cervical vagal nerve. Arrows point to cells that stained positive for both markers. Reprinted from Onkka et al¹⁷ with permission of the publisher.

neural remodeling. Cao et al³² injected nerve growth factor into the left stellate ganglion and induced robust cardiac nerve sprouting in normal canine hearts. The same effects are

observed with low-amplitude electric stimulation of the left stellate ganglion.³³ Zhou et al³⁴ performed a study of the mechanisms of nerve sprouting using a canine model of myocardial




Figure 3. Tyrosine hydroxylase and cholineacetyltransferase staining of the cervical vagal nerves. **A**, A low-power view of the right cervical vagal nerve stained with tyrosine hydroxylase. There are 2 distinct nerve bundles in this nerve. The tyrosine hydroxylase stain of the smaller (**B**) and the larger (**D**) bundles in **A**. The brown color identifies the positively stained nerves. Note that tyrosine hydroxylase-positive nerves are located in the periphery of the nerve bundle. **C** and **E**, Cholineacetyltransferase staining of the same structures as in **B** and **D**, respectively. Note that cholineacetyltransferase-positive components are widely distributed in the cervical vagal nerve. **F**, The tyrosine hydroxylase-positive nerve structure (red arrow) in the middle of the cervical vagal nerve. The objective lens used in **A** was $\times 4$, with a calibration bar of 0.2 mm in length. The objective lens used in **B–F** was $\times 20$, with a calibration bar of 0.2 mm in length. **G**, Activation of vagal nerve alone is associated increased heart rate, a finding consistent with the activation of the sympathetic component of the vagal nerve. SGNA indicates stellate ganglion nerve activity; and VNA, vagal nerve activity. Reprinted from Onkka et al¹⁷ with permission of the publisher.

infarction. The results show a persistent elevation of nerve growth factor levels in aorta and coronary sinus within 1 month after myocardial infarction. Nerve growth factor and growth-associated protein 43 are transported retrogradely to the left stellate ganglion through retrograde axonal transport. The increased nerve growth factor then triggers nerve sprouting at the noninfarcted ventricles and atria.³⁵ Increased atrial sympathetic innervation is associated with increased incidence and duration of AF in those animals. These studies show that, although cardiac injury is limited to the ventricle, neural remodeling may occur throughout the heart. Cardiac diseases, such as myocardial infarction, can potentially increase nerve activities and promote the development of both atrial and ventricular arrhythmias.

Autonomic Remodeling and AF

There is an association between abnormal autonomic innervation and AF in both animal models and in humans. The abnormal autonomic innervation may be important in the mechanisms of AF.^{36–40} Jayachandran et al⁴¹ used [C-11] hydroxyephedrine to label sympathetic nerve terminals in dogs with pacing-induced AF and documented heterogeneously increased atrial sympathetic innervation. The increased sympathetic nerve densities were later confirmed by immunohistochemical staining using antibody against tyrosine hydroxylase in dogs with pacing-induced AF.⁴² Atrial nerve sprouting and sympathetic hyperinnervation also occur after ventricular myocardial infarction and are associated with increased incidence and duration of AF.³⁵ Consistent with these results, atrial sympathetic nerve densities are also significantly increased in patients with chronic AF.⁴³ Multiple other studies have also documented the pathophysiological importance of autonomic remodeling in various animal models and in humans.^{44–47} In addition to atrial sympathetic hyperinnervation, diseases also cause remodeling of extracardiac nerve structures in both experimental animals and in humans.^{48–50}

Cellular Mechanisms of Cardiac Autonomic Neurotransmission and Signaling

Sympathetic neurotransmission results from the excitation of sympathetic nerve terminals via electric impulses traveling down the efferent postsynaptic sympathetic nerves, which originate in sympathetic ganglia like the stellates. The production, release, reuptake, and degradation of sympathetic neurotransmitters are an extremely complex and highly regulated process.⁵¹ This regulation is essential to ensure that the critically important function of adrenergic control is well tuned to physiological needs under a wide range of conditions. In brief, the principal neurotransmitter norepinephrine is synthesized in neural cell bodies and transported and concentrated in vesicles in nerve varicosities adjacent to adrenergic receptors, where it is released by nerve depolarization through a Ca^{2+} -dependent process. In addition to norepinephrine, these vesicles contain smaller amounts of a variety of other biologically active substances such as opioids, chromogranin, and other neuropeptides.⁵¹ Rapid uptake mechanisms limit the amount of norepinephrine that can access adrenergic receptors, and norepinephrine is also rapidly degraded by a variety of enzymes such as monoamine oxidase. In addition to reuptake

and enzymatic degradation, norepinephrine action is controlled by negative feedback through presynaptic receptors, particularly α_2 -adrenergic, dopamine, and muscarinic receptors.⁵¹ Systemically circulating epinephrine released from the adrenal medulla also contributes to cardiac sympathetic activation, especially in conditions of generalized sympathetic activation.

Norepinephrine interacts with a variety of adrenergic receptors on cardiomyocytes to execute adrenergic actions. The detailed biochemistry of adrenergic receptor pharmacology is complex, and the interested reader is referred to an excellent recent review.⁵² Here, we will focus primarily on the β -adrenergic receptor and its downstream signaling relevant to AF (Figure 4). The β -adrenergic receptor is a member of the enormous family of 7-transmembrane domain G-protein-coupled receptors and includes 3 subtypes, β_{1-3} , of which β_1 -receptors are most relevant to atrial arrhythmias. The G-protein system includes 3 subunits: α , β , and γ . The $G\beta$ and $G\gamma$ subunits bind to each other and are often referred to together as the $G\beta\gamma$ subunit. A variety of $G\alpha$ subunits exist, but the principal adrenergic $G\alpha$ subunit is the $G\alpha_s$ or stimulatory subunit. When the β -receptor is unoccupied, most $G\alpha_s$ is bound to $G\beta\gamma$. Norepinephrine binding to the β -receptor leads to GTP binding of the $G\alpha_s$ subunit, lowering its affinity to $G\beta\gamma$, which dissociates and allows the free $G\alpha_s$ subunit to activate adenylate cyclase, which converts ATP to cAMP, the primary β -adrenergic second messenger. cAMP activates protein kinase A (PKA), which exerts a wide range of effects by phosphorylating membrane proteins, including Ca^{2+} -handling proteins and ion channels.

Acetylcholine is synthesized from choline and acetylcoenzyme-A via choline acetyltransferase, primarily in cholinergic nerve terminals where it is concentrated in synaptic vesicles. Like sympathetic neurotransmitter production and release, acetylcholine biology is highly regulated and subject to feedback inhibition via presynaptic muscarinic receptors.⁵³ Released acetylcholine is rapidly broken down by acetylcholinesterase. Acetylcholinesterase is remarkably efficient at breaking down acetylcholine and greatly limits the spread of acetylcholine from its release site. Consequently, the effects of acetylcholine are localized, allowing for spatial heterogeneity of acetylcholine effects under vagal activation, a property that is important in AF.

The cardiac cholinergic receptor is an M_2 type-2 muscarinic subtype. M_2 -acetylcholine receptors are also G-coupled, with the inhibitory G-protein $G\alpha_i$ being the principal subtype bound to $G\beta\gamma$. When acetylcholine interacts with the M_2 -receptor, $G\alpha_i$ -GTP interaction occurs, and as for adrenergic receptors, this causes dissociation of $G\beta\gamma$ subunits from $G\alpha_i$ (Figure 4). However, unlike adrenergic activation, which uses $G\alpha_s$ as the main signaling G-protein, cholinergic effects result predominantly from $G\beta\gamma$ activation of the ligand-gated K^+ channel I_{KACH} , composed of Kir3.1 and Kir3.4 subunits.⁵⁴ I_{KACH} activation produces an outward K^+ current that flows throughout the depolarized phases of the cardiac action potential, resulting in substantial reduction in action potential duration (APD).

Autonomic Regulation of Atrial Cardiomyocyte Electrophysiology

The principal molecular mechanisms by which autonomic influences affect AF likelihood are illustrated in Figure 4. Please

note that another article in this compendium deals in detail with the cellular machinery underlying AF.⁵⁵ In this article, we will limit ourselves to the specific mechanisms underlying autonomic AF promotion. The principal arrhythmogenic targets of β -adrenergic stimulation relate to cardiomyocyte Ca^{2+} handling. The main business of β -adrenergic activation in the heart is to enhance cardiac output during fight-or-flight reactions. Accordingly, β -adrenergic stimulation enhances virtually all process controlling Ca^{2+} entry, storage, and release in the heart. These effects are initiated by PKA and amplified by Ca^{2+} /calmodulin-dependent protein kinase type II (CaMKII). PKA and CaMKII phosphorylate many of the same proteins (albeit at different sites): the L-type Ca^{2+} channel (I_{CaL}), the sarcoplasmic reticulum (SR) Ca^{2+} release channel ryanodine receptor 2, and phospholamban.⁵⁶ I_{CaL} phosphorylation

increases voltage-dependent Ca^{2+} entry through the plasma membrane. Ryanodine receptor 2 phosphorylation amplifies Ca^{2+} -dependent Ca^{2+} release from the SR. Together, these actions greatly augment the systolic Ca^{2+} transient and thereby contraction strength. Phospholamban binds to and inhibits the SR Ca^{2+} transporter, SR Ca^{2+} ATPase, the principal mechanism responsible for maintaining SR Ca^{2+} stores and restoring low diastolic Ca^{2+} levels after the systolic Ca^{2+} transient to allow diastolic relaxation/filling. Adrenergically induced phospholamban phosphorylation by PKA and CaMKII dissociates phospholamban from SR Ca^{2+} ATPase, disinhibiting SR Ca^{2+} ATPase Ca^{2+} pumping into the SR. Under acute stress conditions, adrenergic activation provides an essential boost to Ca^{2+} -dependent cardiac function. However, under conditions predisposing to Ca^{2+} -dependent triggered activity,^{57,58}




Figure 4. Molecular basis for autonomic contributions to atrial fibrillation substrate. β -adrenergic receptor (β -AR) activation causes GTP binding to the G_{α_s} subunit, allowing it to dissociate from $G\beta$ and γ subunits and activate adenylyl cyclase (AC), which converts ATP to cAMP. cAMP activates protein kinase A (PKA), which phosphorylates a range of Ca^{2+} -handling proteins including the L-type Ca^{2+} channel (LTCC), ryanodine receptor (RyR2), and phospholamban (PLB). PLB phosphorylation causes it to dissociate from the sarcoplasmic reticulum (SR) Ca^{2+} ATPase (SERCA2a), removing SERCA2a from PLB inhibition and activating SR Ca^{2+} uptake. RyR2 phosphorylation increases RyR2 open probability, enhancing the systolic Ca^{2+} transient but also enhancing diastolic Ca^{2+} leak. Adrenergic stimulation also increases Ca^{2+} binding to calmodulin (CaM), activating Ca^{2+} /CaM-dependent kinase type II (CaMKII), which phosphorylates many of the same proteins as PKA. Ca^{2+} /CaM also activates calcineurin (Cn), which dephosphorylates nuclear factor of activated T cells (NFAT), allowing it to translocate to the nucleus and activate hypertrophic and profibrotic gene programs. LTCC phosphorylation increases I_{CaL} and shifts its voltage dependence to cause larger window currents. Adrenergic stimulation also inhibits inward rectifier K^+ current (I_{K1}) and enhances slow delayed rectifier K^+ current (I_{Ks}). Cholinergic activation of muscarinic type-2 (M_2) acetylcholine receptors (AChRs) causing GTP binding to G_{α_i} , releasing $G\beta\gamma$, and allowing it to activate the acetylcholine-dependent K^+ current (I_{KACh}). CICR indicates calcium-induced calcium release; HDAC, histone deacetylases; MEF2, myocyte enhancer factor-2; and NCX1, sodium calcium exchanger.

the enhanced Ca^{2+} -loading/release conditions produced by adrenergic stimulation strongly promote arrhythmogenesis. In a canine model of chronic atrial ischemia, aberrant Ca^{2+} release responsible for ectopic activity requires adrenergic drive to manifest.⁵⁹

Autonomic modulation has significant effects on cardiac ion channels. In addition to the acetylcholine-induced activation of I_{KACh} , a host of ion channels are affected by adrenergic tone.⁶⁰ The most important of these are I_{CaL} , already discussed, the slow delayed rectifier K^+ current I_{Ks} , and the inward rectifier I_{K1} . I_{Ks} is strongly enhanced by adrenergically induced PKA phosphorylation,⁶¹ allowing it to offset the increased inward current resulting from adrenergic enhancement of I_{CaL} and prevent early afterdepolarizations (EADs).⁶² I_{K1} is important in setting the resting potential, contributing to repolarization reserve⁶³ and governing AF dynamics.⁶⁴ I_{K1} is typically inhibited via α -adrenergic receptor stimulation.⁶⁵

Autonomic Effects on Mechanisms Governing AF Occurrence

The potential basis for autonomic nervous system promotion of AF is summarized in Figure 5. AF can result from focal or re-entrant mechanisms.^{66,67} Focal mechanisms are important in 2 ways: they may act as a trigger on a susceptible substrate or by firing rapidly provide an AF-maintaining driver. Adrenergic activation may promote focal activity via each of the principal cellular mechanisms: enhanced automaticity (Figure 5A), EADs (dashed tracings; Figure 5B), or delayed afterdepolarization-associated triggered activity (red dashed tracings; Figure 5C). I_{K1} provides a diastolic outward current that prevents spontaneous phase-4 depolarization to the threshold potential by the pacemaker funny current that underlies spontaneous automaticity. Automaticity is enhanced by reduced I_{K1} , which can result from α -adrenergic stimulation, or increased funny current, produced by β -adrenergic activation.⁶⁸ Phase-2 EAD-induced ectopic activity (red dashed tracings; Figure 5B) likely underlies the increased risk of AF in patients with congenital long-QT syndrome.⁶⁹ β -adrenergic activation enhances plateau I_{CaL} (via PKA/CaMKII phosphorylation), increasing EAD likelihood, particularly when adrenergic augmentation of I_{Ks} is deficient (eg, in long-QT syndrome type 1). Phase-3 EADs can be associated with APD prolongation (blue tracing; Figure 5B). It may occur as the result of electrotonic current across steep repolarization gradients between phase-2 EAD and the adjacent repolarized tissues or occur as the result of low I_{K1} .⁷⁰ In comparison, a late phase-3 EAD (green tracings; Figure 5B) is associated with shortened rather than prolonged APD.⁷¹ If there is simultaneous activation of the sympathetic nervous system that increases the intracellular Ca^{2+} transient and parasympathetic nervous system that activates I_{KACh} , then APD is shortened while the Ca^{2+} transient is large and long. A short APD and a large Ca^{2+} transient create a condition for late phase-3 EADs, which can induce triggered activity and AF (solid green tracing; Figure 5B).^{71,72} Because PVs naturally have short APDs, they are particularly prone to develop these Ca^{2+} transient-triggered arrhythmias.^{38,40,73} Delayed afterdepolarizations (Figure 5C) result from diastolic ryanodine receptor 2 Ca^{2+} leak, favored by β -adrenergic enhancement of cell

Ca^{2+} loading and increased ryanodine receptor 2 open probability because of PKA/CaMKII phosphorylation.

The precise details of mechanisms maintaining re-entry (Figures 5D), such as the structure and number of circuits, role of rotors, remain controversial.⁷⁴ However, shortened refractoriness promotes functional re-entry in all conceptual models. Vagal stimulation strongly abbreviates atrial refractoriness by augmenting I_{KACh} . Furthermore, the refractoriness-abbreviating effects of vagal activation show strong regional variation, much more so than adrenergic effects; this regional variability underlies particularly strong AF-promoting effects of vagal tone.⁷⁵

Finally, structural remodeling is known to be an important contributor to AF persistence.⁶⁶ Increased Ca^{2+} /calmodulin binding caused by β -adrenergic stimulation activates the protein phosphatase calcineurin (Figure 4). Calcineurin dephosphorylates the transcription factor nuclear factor of activated T cells, allowing it to translocate into the nucleus and alter gene transcription, inducing hypertrophic and profibrotic gene expression programs. Adrenergic stimulation also promotes structural remodeling via other actions, including actions mediated by CaMKII, oxidative stress, and signaling via an alternate $\text{G}\alpha$ subunit, $\text{G}\alpha_q$.⁵²

Autonomic Nerve Activity and Atrial Arrhythmias

Direct recording of autonomic nerve activity can provide insight into its role in atrial arrhythmogenesis in animal models. Long-term recording of nerve activity in ambulatory animals was first successfully performed by Barrett et al.⁷⁶ Stable cardiac nerve activity was then recorded in the heart, allowing for the relationships between neural activity and arrhythmogenesis.⁷⁷

Stellate ganglion nerve activity (SGNA) and vagal nerve activity (VNA) increase after the induction of heart failure by ventricular tachypacing.²⁶ Increased nerve activity was directly associated with paroxysmal atrial tachycardias (PAT) in these dogs. A canine model of intermittent atrial tachypacing was then developed, with rapid atrial pacing for 6 to 7 days, followed by 1 nonpaced day to observe PAT and paroxysmal atrial fibrillation (PAF) without pacing artifacts. Intermittent LA tachypacing causes sympathetic hyperinnervation, PAF, and PAT.⁴ Simultaneous sympathovagal discharges commonly precede arrhythmias, implicating them as triggers. Figure 6A shows a typical example of PAF, with sinus arrhythmia in the first 20 seconds, followed by an abrupt increase in SGNA and VNA and PAF. Figure 6B shows an example of PAT to PAF transition that occurs frequently both in this animal model and in humans. Figure 6C is a 6-second close-up of the same episode shown in Figure 6B, straddling the initiation of PAF. An initial increase in VNA (1) followed by increased SGNA (2) is followed by an acceleration of PAT from 521 bpm to 562 bpm. A second increase in VNA (3) followed closely by a massive burst of SGNA (4) precedes the onset of PAF by ≈ 3 seconds. About 73% of PAT and PAF episodes were preceded by simultaneous sympathovagal discharges. Optical mapping data implicate Ca^{2+} -initiated triggered activity in atrial arrhythmogenesis resulting from parasympathetic activation in transgenic mice that develop a fibrotic AF substrate because of overexpression of constitutively activated transforming growth factor- $\beta 1$. These findings are consistent with a previous study⁷⁸

Adrenergic and Cholinergic Contributions to AF Mechanisms




Figure 5. Mechanisms by which autonomic tone can promote atrial fibrillation (AF). Top, Action potential changes showing cellular mechanisms by which adrenergic activation can lead to focal ectopic firing. Black dotted tracings represent normal reference action potentials in each panel. **A**, Enhanced automaticity. **B**, Early afterdepolarizations (EADs). **C**, Delayed afterdepolarization (DADs). Contributions from adrenergic activation alone are shown by red tracings, whereas that from cholinergic activation (combined with adrenergic activation) by green tracings. Adrenergic stimulation in the setting of impaired repolarization reserve can cause phase-2 EADs (red dashed tracings in **B**). Most phase-3 EADs are also associated with prolonged action potential duration (APD; blue dashed tracings in **B**). Combined adrenergic/vagal discharge can produce late phase-3 EADs (green dashed tracings in **B**) because of a prolonged and enhanced Ca^{2+} transient that outlasts I_{KACH} -induced accelerated repolarization. **D**, Tissue-level arrhythmia mechanisms, with focal ectopic activity maintaining AF as a driver or acting on vulnerable re-entrant substrates. Parasympathetic firing discharges acetylcholine, producing spatially heterogeneous action potential and refractory period abbreviation that promotes the occurrence and maintenance of re-entrant activity. CaT indicates calcium transient; LA, left atrium; NCX, sodium calcium exchanger; RA, right atrium; RyR, ryanodine receptor; and SR, sarcoplasmic reticulum.

that showed AF induction by simultaneous acetylcholine and isoproterenol infusion into the sinus node artery of anesthetized dogs.

Direct recordings from both the extrinsic nervous system (left stellate ganglion and left thoracic vagal nerve) and the intrinsic cardiac nervous system (including superior left ganglionated plexi and ligament of Marshall) were performed to distinguish their relative role in AF development.⁷⁵ After intermittent rapid atrial pacing, ambulatory dogs displayed spontaneous PATs before the development of persistent AF. Atrial tachyarrhythmias were invariably preceded by intrinsic cardiac nerve activity. These findings further support the importance of autonomic ganglia in the pathogenesis of AF associated with atrial tachycardia remodeling.⁷⁹ Because histological studies show extensive colocalization of adrenergic and cholinergic nerve structures in the intrinsic cardiac nerves,²⁹ it is possible that the simultaneous activation of these 2 arms of autonomic nervous system may be involved in arrhythmia initiation.

Autonomic Nerve Activity and Persistent AF

In most patients with AF, rate control is not inferior to rhythm control as a management strategy.⁸⁰ It is known that the inferior vena cava–inferior atrial ganglionated plexus (also known as the inferior right or right inferior ganglionated plexi)⁸¹ is important in modulating atrioventricular node conduction. Direct electric stimulation of these ganglionated plexi may slow ventricular rate during AF in human patients.⁸² Ambulatory recordings of bilateral cervical VNA and inferior vena cava–inferior atrial ganglionated plexus nerve activity during persistent AF show that in most but not all dogs, the left vagal nerve controls the atrioventricular node, whereas the right vagal nerve controls the sinus node.¹⁸ The only nerve structure that consistently controls atrioventricular nodal conduction is the inferior vena cava–inferior atrial ganglionated plexus. Figure 7 shows an example in which inferior vena cava–inferior atrial ganglionated plexus nerve activity is associated with abrupt reduction of ventricular rate during persistent atrial fibrillation. VNA may sometimes be associated with acceleration of heart rate,




Figure 6. Two examples of paroxysmal atrial fibrillation (PAF). **A**, Sinus rhythm to AF conversion. **B**, Atrial tachycardia to AF conversion. **C**, Magnified from the center of **B** (line segment above ECG), showing that the elevated vagal nerve activity (VNA) accelerated atrial rate, leading to paroxysmal reduction of ventricular rate (prolonged RR interval) before conversion from paroxysmal atrial tachycardia (PAT) to PAF. LA indicates left atrium; and SGNA, stellate ganglion nerve activity. Reprinted from Tan et al⁴ with permission of the publisher. AVN indicates atrioventricular node.

probably because of activation of the sympathetic component within the vagal nerves.^{17,27} Thus, the ventricular rate during sustained AF is controlled by collaboration among different nerve structures.

Coordination Among Nerve Structures and the Development of AF

Detailed analysis and integration of nerve activity over time have revealed several previously unappreciated patterns of nerve activation.^{18,83–85} First, the correlation between SGNA and VNA was found to fall into 2 different basic patterns. In a minority of dogs, the 2 nerve structures would fire simultaneously (group 1). In the remaining dogs, the SGNA and VNA fired separately (ie, one would activate, whereas the other was quiescent; group 2). The group 1 dogs, which tend to have simultaneous sympathovagal discharges, have more PAT episodes at baseline and faster induction of sustained AF by rapid pacing than the remaining (group 2) dogs that had an L-shaped correlation, indicating temporally separate sympathetic and vagal activity. Perhaps because these dogs were followed for relatively short periods of time (weeks), each dog continued to show a consistent pattern of nerve firing. However, in a subsequent study when 1 dog was followed for ≈6 months, a switch from group 1 to group 2 was observed.⁸⁵ If sympathovagal correlation is important in the development of atrial tachyarrhythmias and AF, the changing patterns of sympathovagal correlation suggest the possibility of dynamically varying arrhythmia susceptibility.

In addition to SGNA and VNA, both linear and L-shaped correlations have been observed between cervical VNA and the inferior vena cava–inferior atrial ganglionated plexus.¹⁸ In 5 of the 6 dogs studied, an L-shaped relationship was present between right VNA and left VNA during AF. In the remaining 1 dog, a linear correlation was noted between right and left VNA. These findings indicate that right and left cervical vagal nerves do not randomly activate relative to each other. Rather, most typically one would activate when the other is quiescent. In a small minority of dogs, they almost always activate together. Coactivation of these 2 nerves may be associated with rapid ventricular rate, suggesting that there might be coactivation of the sympathetic nervous system. Another important finding is that the intrinsic nerves (inferior vena cava–inferior atrial ganglionated plexus nerve activity) show a linear correlation with left VNA in a dog with L-shaped correlation between left and right VNA. This indicates that the left VNA almost always fires together with inferior vena cava–inferior atrial ganglionated plexus, whereas the right VNA fires at a different time and does not control the inferior vena cava–inferior atrial ganglionated plexus. Observations such as these clearly indicate that extrinsic and intrinsic nervous systems do not activate randomly in ambulatory dogs. Rather, a high degree of coordination is present among these nerve structures.

Neuromodulation as a Therapeutic Approach

Because different autonomic nerve structures coordinate their activation with each other, interruption or modification of the




Figure 7. Local control of atrioventricular node conduction during persistent atrial fibrillation. Slowing of ventricular rate (VR) was associated with inferior vena cava–inferior atrial ganglionated plexus nerve activity (IVC–IAGPNA) without either right vagal nerve activity (RVNA) or left vagal nerve activity (LVNA). Subsequent simultaneous activation of right vagal nerve activity and left vagal nerve activity resulted in a rapid ventricular rate. Because of the presence of abundant sympathetic nerves within the vagus,¹⁷ these observations suggest that sympathetic component within the vagal nerves have accelerated the ventricular rate. LEGM is the local bipolar electrogram. Reprinted from Park et al¹⁸ with permission of the publisher.

activity in one structure may change the pattern of activation of another. These changes may convey therapeutic effects, including arrhythmia control. Some methods of neuromodulation are already in place in clinical use. Others are still being tested in the animal laboratory or clinical trials. Common sites for neuromodulation are labeled by black dots in Figure 1.

Sympathetic and Vagal Denervation

Because autonomic nerve activity can act as a direct trigger of PAF,^{4,86} it is logical to test the hypothesis that stellate ganglion ablation can reduce the incidence of AF. Accordingly, cryoablation of the lower portion of both left and right stellate ganglia, sparing the upper portion of the stellate ganglia to prevent Horner syndrome,⁸⁷ along with the T2 to T4 thoracic sympathetic ganglia, was performed in dogs. The vagi were denervated by ablating the superior cardiac branch of the left thoracic vagal nerve. The locations of these structures are shown in Figure 1. One major consequence of cryoablation was a lack of heart rate response to SGNA and VNA. A second major consequence was a delay in the development of sustained AF in response to atrial tachypacing. Whereas control dogs developed sustained AF in 2 to 4 weeks, the group subjected to cryoablation required 3 to 12 weeks of atrial pacing to sustain AF.⁴ A third effect of cryoablation was a suppression of premature atrial contractions and elimination of episodes of PAT and PAF typically associated with intermittent rapid atrial pacing. These findings support the notion that simultaneous sympathovagal discharges contribute importantly to atrial arrhythmogenesis. Because cryoablation only delayed but did not prevent sustained AF, autonomic nerve activity is not the only factor determining AF maintenance. Dogs with pacing-induced heart failure develop both prolonged sinus pauses and PAT.²⁶ Cryoablation of bilateral stellate and T2 to T4 thoracic ganglia significantly reduces PAT and prolonged sinus pause episodes induced by sympathetic discharges in dogs with pacing-induced heart failure.⁸⁸

The above studies suggest that cardiac sympathetic denervation might be useful in controlling PAT and PAF by reducing sympathetic outflow to the heart. However, these studies have multiple limitations. One limitation is that in the canine model PAT and PAF were induced by rapid pacing of

either the atria or the ventricles. In contrast, the established risk factors for AF in humans include age, male sex, systolic and diastolic heart failure, valvular heart disease, myocardial infarction, hypertension, diabetes mellitus, obesity, and cigarette smoking.⁸⁹ The canine model of PAT and PAF may not be applicable to humans. A second limitation is that the stellate ganglion and T2 to T4 sympathetic ganglia are not easily accessible in humans. However, the invention of videoscopic left cardiac denervation⁹⁰ may reduce the procedural complexity of this approach. A third limitation is that the nervous system is highly plastic. It is possible that reinnervation can occur after the denervation procedures and negate the effects of denervation. A fourth limitation is that surgical removal of the stellate ganglion causes irreversible changes of the sympathetic nervous system. The long-term effects of sympathetic denervation in patients with AF are unknown.

Vagal Nerve Stimulation

Because of the above limitations, it is highly desirable to develop a neuromodulation method that can be easily terminated, without causing permanent damage to the autonomic structures. Transvenous parasympathetic nerve stimulation can be used as a method of ventricular rate control during AF.⁹¹ However, vagal nerve stimulation (VNS) can also be used in the animal laboratory as a method to induce or maintain sustained AF.^{92,93} Many studies have documented the effects of neural stimulation or ablation in inducing or controlling cardiac arrhythmias.^{94–97} The effects of neural stimulation may not be limited to the area directly innervated by the modified nerve structures. For example, stimulating the afferent cervical vagal nerve in cats suppresses sympathetic discharges.⁹⁸ Because cervical vagal nerves are accessible through surgical approaches, they are the prime target for neural modulation with the hope that their stimulation will achieve therapeutic effects distant beyond the nerves stimulated. A documented success is the use of left cervical VNS to suppress epilepsy in humans.⁹⁹ Vanoli et al¹⁰⁰ showed that chronic VNS can prevent ventricular fibrillation and sudden cardiac death in conscious dogs with a healed myocardial infarction. Others showed that VNS might be used to attenuate heart failure development in dogs,¹⁰¹ rats¹⁰² and humans.^{103–105} Although most of these studies used

stimulus strength sufficient to reduce heart rate, low-level VNS, defined by a stimulus strength 1 V below the threshold needed to reduce heart rate, is effective in suppressing AF induction in open-chest-anesthetized dogs.^{106,107} Because VNS opposes sympathetic actions at both pre- and postjunctional levels,^{108,109} VNS may achieve the therapeutic effects by suppressing sympathetic outflow to the heart. To test this hypothesis, Shen et al⁵ performed continuous low-level VNS in a canine model of PAF while continuing to record SGNA and VNA. Consistent with the observations of Schwartz et al,⁹⁸ VNS may immediately suppress SGNA when the stimulator is turned on. However, chronic VNS is associated with further reduction of SGNA. The effects of VNS are most apparent in the morning when the SGNA is most active. The VNS reduced the number of sympathetic discharge episodes and shortened the average duration of discharges. Because of the reduced duration of sympathetic discharges, the SGNA caused less heart rate acceleration during VNS than at baseline. The effects of VNS are not permanent. Rather, SGNA normalizes at the cessation of low-level VNS. In addition to its effects on SGNA, low-level VNS also significantly reduces the number of PAT episodes.

Because VNS has chronic effects on SGNA, VNS might have caused the remodeling of the stellate ganglion. Immunostaining of the left stellate ganglion in dogs with and without VNS showed that low-level VNS decreased the density of nerve structures (presumably sympathetic) staining positive for tyrosine hydroxylase. Although a majority (>90%) of the ganglion cells normally stain positive for tyrosine hydroxylase, a small minority of cells show no tyrosine hydroxylase staining (Figure 2). There was a 3-fold increase in the prevalence of tyrosine hydroxylase-negative cells in VNS group compared with controls. In a different group of dogs, small-conductance calcium-activated K channel subtype 2 protein expression in the VNS group was found to be ≈50% higher than in the control group.¹⁶ Immunostaining also showed that the density of nerve structures stained with small-conductance calcium-activated K channel subtype 2 antibody was higher in the VNS group than in the control group. There was significantly increased small-conductance calcium-activated K channel subtype 2 protein staining in the periphery of ganglion cells compared with the cell center. This was not observed in normal control dogs. In addition, there were significantly more ganglion cells without immunoreactivity to tyrosine hydroxylase in dogs with VNS (average, 11.4%) than in control (4.9%), again showing an ≈2- to 3-fold increase of the tyrosine hydroxylase-negative cells in the VNS group. Furthermore, a high percentage of tyrosine hydroxylase-negative cells stained positive for choline acetyltransferase. The increased percentage of these cells suggests that VNS might cause phenotypic switching between adrenergic and cholinergic nerves. Figure 8 shows a summary of the stellate ganglion remodeling induced by VNS. The chronic effects of VNS can be partially explained by stellate ganglion remodeling, including increased small-conductance calcium-activated K channel subtype 2 proteins and the reduction of tyrosine hydroxylase-positive ganglion cells.

Baroreflex Stimulation and Exercise

Exercise training results in functional modulation of autonomic balance. Exercise may activate parasympathetic nervous system

through changes of plasma volume (baroreflex stimulation)¹¹⁰ or via augmented baroreflex responsiveness and increased cardiomyocyte sensitivity to cholinergic stimulation.¹¹¹ In the case of exercise training, enhanced sensitivity to acetylcholine seems to be because of reduced expression of a family of proteins called regulators of G-protein signaling,¹¹⁰ which have GTPase activity and terminate acetylcholine-induced I_{KACH} activation by breaking down $G\alpha_s$ -associated GTP. Endurance exercise training increases AF susceptibility in rats via increased parasympathetic tone accompanied by atrial dilation and mild fibrosis.¹¹¹ These observations parallel clinical observations of an importantly increased prevalence of AF in endurance athletes.¹¹² However, chronic exercise training may be beneficial for the management of AF by improving rate control.¹¹³ It is possible to use implantable devices to stimulate the carotid sinus directly and activate the baroreflex.^{114,115} Similar to VNS, baroreflex stimulators can sharply decrease sympathetic nerve activity and lower blood pressure among responders.¹¹⁶ The reduced sympathetic nerve activity may be, in part, responsible for the improved rate control during AF. Although strong baroreflex stimulation may reduce atrial effective refractory period and promote AF, low-level baroreflex stimulation only causes moderate shortening of atrial effective refractory period.^{117,118} Additional studies are needed to determine whether low-level baroreflex stimulation can be used to control cardiac arrhythmias by reducing sympathetic tone without massively shortening the atrial effective refractory period.

Ganglionated Plexus Ablation

Intrinsic cardiac nerve activity invariably precedes the onset of AF in ambulatory dogs.⁸⁶ If these findings are applicable to humans, then ablation of the ganglionated plexi of the intrinsic cardiac nervous system with surgical or catheter ablation techniques may be effective in controlling AF. Earlier non-randomized observational studies showed that PV denervation may enhance the long-term outcome of circumferential ablation of PAF.¹¹⁹ These findings enhanced the theory that hyperactivity of local cardiac ganglionated plexi plays a role in the generation and maintenance of AF.³⁹ One approach to ganglionated plexus ablation is to use high-frequency stimulation to identify ganglionated plexi before ablation.¹²⁰ Others used an anatomically based approach without high-frequency stimulation.^{121,122} Because ganglionated plexus ablation is a new procedure, it is possible that there is a bias in favor of reporting positive results. Katritsis et al¹²³ performed a prospective randomized clinical trial, exposing 242 patients with PAF to PV isolation alone, ganglionated plexus ablation alone (anatomic approach), and PV isolation plus ganglionated plexus ablation. After 2 years of follow-up, freedom from AF or atrial tachycardia was achieved in 56%, 48%, and 74% of patients in the PV isolation, ganglionated plexus ablation, and PV isolation+ganglionated plexus ablation groups, respectively ($P=0.0036$). The authors concluded that the addition of ganglionated plexus ablation to PV isolation confers a significantly higher success rate compared with either PV isolation or ganglionated plexus ablation alone in patients with PAF. In addition to catheter ablation, minimally invasive surgical procedures have been used for PV isolation and ganglionated plexus ablation, with significant improvement in the




Figure 8. Changes of type 2 small-conductance calcium-activated K (SK2) protein in the left stellate ganglion (LSG) with low-level vagal nerve stimulation. **A**, Representative Western blots show that the signal ratio of SK2 protein to GAPDH of vagal nerve stimulation dogs (group 1) was significantly higher than that of control (group 2). **B**, There is an upregulation of SK2 protein level in the LSG in group 1 dogs after being normalized to GAPDH. **C** and **D**, Representative immunostaining of SK2 protein in the left stellate ganglion. The density of SK2-positive nerve structures (as pointed by a black arrowhead) is significantly higher in group 1 dogs (**C**) compared with group 2 dogs (**D**). **E** and **F**, Representative low-power view of immunostaining of SK2 protein in the LSG that clearly demonstrates higher SK2 density in group 1 dogs (**E**) compared with group 2 dogs (**F**). **G** and **H**, Immunofluorescence confocal microscope images of the LSG from group 1 and group 2 dogs, respectively. Blue colored dots show the nuclei stained with 4',6-diamidino-2-phenylindole. Red color marks the SK2 protein. Note a significantly increased SK2 staining in the periphery of ganglion cells but decrease in the cytosol of group 1 (**G**). In contrast, in group 2 LSG, the SK2 staining was homogeneous (**H**). AU indicates arbitrary units. Calibration bar, 50 μm for **C–F** and 20 μm for **G** and **H**.

outcome.¹²⁴ The clinical evidence to date seems to support the use of ganglionated plexus ablation as an adjunctive procedure in AF ablation.

Renal Sympathetic Denervation

Preliminary clinical trials conducted by various investigators suggest that renal sympathetic denervation through an endovascular approach is effective in controlling drug-resistant hypertension.^{125,126} Other work showed that renal sympathetic denervation can reduce sympathetic nerve activity.¹²⁷ Because sympathetic nerve activity is important in blood pressure control,^{76,85} reduction of sympathetic outflow may, in part, explain the reduction of blood pressure in some patients. The same effects may also be useful in controlling AF. There are ongoing clinical studies testing the hypothesis that concomitant renal denervation may improve the outcomes from catheter ablation of AF.^{128,129} Renal sympathetic denervation has also been used for ventricular rate control in AF and for reduction of AF episodes in patients with sleep apnea.^{128–130} Preclinical studies suggest that long-term renal denervation may be beneficial in treating rats with heart failure induced by myocardial

infarction.¹³¹ It is possible that renal sympathetic denervation may benefit cardiac arrhythmic control by improving myocardial function in heart failure. The latter hypothesis is being tested by several studies listed in clinicaltrials.org. The results of those studies should advance the field by defining the benefits and risks of renal sympathetic denervation. It remains to be seen if successful treatment of heart failure can also result in reduced incidence of AF in those trials. Recently, the first large-scale randomized sham controlled clinical trial¹³² failed to document the efficacy of renal denervation in patients with resistant hypertension.¹³³ The implications of this outcome for the concept and application of renal sympathetic denervation are certainly major and will undoubtedly motivate careful reflection and additional investigation.¹³⁴

Somatic Sensory Stimulation for Neuromodulation

Various forms of somatic sensory stimulation can produce autonomic reflex responses, depending on the visceral organs and somatic afferents that are stimulated.¹³⁵ Yu et al¹³⁶ developed a noninvasive transcutaneous approach to deliver low-level VNS to the tragus of the ear to treat cardiac arrhythmias such

as AF. The authors found that low-level tragus stimulation can reverse pacing-induced atrial remodeling and suppress AF inducibility, suggesting possible value in treatment of AF. An alternative approach to neuromodulation is acupuncture, which is widely practiced for pain control, although the clinical efficacy remains unproven.^{137,138} Lomuscio et al¹⁹ showed that acupuncture using Neiguan, Shenmen, and Xinshu spots might prevent arrhythmia recurrences in patients with persistent AF after electric cardioversion. These 2 studies applying cutaneous stimulation raise the possibility of using somatic sensory stimulation to achieve neuromodulation. A possible mechanistic rationale is that the somata of the skin sympathetic nerves originate from the middle cervical and stellate ganglion, the same ganglia that innervate the heart.¹³ However, the limitations of these studies are considerable, and extensive further investigations and clinical trials will be needed to optimize and test the efficacy of cutaneous neuromodulation in the management of AF.

Effects of Neuromodulation on the Structure and Function of the Heart

In addition to changes in the structure and function of the nervous systems, neuromodulation may also exert direct effects on the structure and function of the heart. Chronic norepinephrine infusion in dogs can reduce cardiac sympathetic nerve density, decrease myocardial norepinephrine uptake activity, and downregulate cardiac β adrenoceptors, reproducing that which occurs in heart failure.^{139,140} Successful treatment of heart failure may result in the improvement of cardiac norepinephrine uptake and attenuate sympathetic nerve terminal abnormalities.^{141,142} Because neuromodulation methods may reduce sympathetic outflow, it may help normalize the cardiac sympathetic innervation and improve receptor function in diseased hearts. In addition to suppressing sympathetic outflow, vagal nerve and epicardial ganglionated plexi stimulations may be anti-inflammatory^{101,143,144} and may improve LA function and suppress the development of LA fibrosis.¹⁴⁵ Renal sympathetic denervation may control AF through modification of the atrial substrates.⁶ These findings suggest that neuromodulation may achieve its therapeutic effects, in part, by causing beneficial structural and functional remodeling in the heart.

Autonomic Nervous System Targets for Antiarrhythmic Drug Therapy

Given the apparent importance of the autonomic nervous system in AF, it should be possible to identify autonomic targets for drug therapy. β -Blockade has moderate but statistically significant effects to prevent AF recurrence after electric cardioversion.¹⁴⁶ With more research, it may be possible to identify patients to target based on particularly important autonomic contributions to their AF. One such group is patients undergoing cardiac surgery, for which there is evidence of an important role of Ca^{2+} homeostasis abnormalities in postoperative AF.¹⁴⁷ Prophylactic β -blockers are particularly effective in preventing postoperative AF,¹⁴⁸ illustrating the applicability of the concept. Based on the importance of I_{KACH} in AF, selective blockers are being developed, with some success in preclinical studies.¹⁴⁹ Biological therapies targeting G-proteins have been applied to modulate atrioventricular nodal function and

control the ventricular response in AF,¹⁵⁰ as well as to prevent AF induction in a vagal model.¹⁵¹ These studies offer a proof of principle for biological therapies targeting specific components of G-protein autonomic effectors, with possible greater specificity and efficacy in the future.

Conclusions

Autonomic nerve activity plays an important role in the initiation and maintenance of AF, and modulating autonomic nerve function may contribute to AF control. Potential therapeutic applications include ganglionated plexus ablation, renal sympathetic denervation, cervical VNS, baroreflex stimulation, cutaneous stimulation, novel drug approaches, and biological therapies. Although the role of the autonomic nervous system has long been recognized, new science and new technologies promise exciting prospects for the future.

Acknowledgment

We thank Medtronic, St Jude, Boston Scientific, and Cyberonics Inc for donating equipment to our research laboratories.

Sources of Funding

Research reported in this article was supported by the National Heart, Lung and Blood Institute of the National Institutes of Health under award number P01HL78931, R01HL71140, and R21HL106554, a Piansky Endowment (M.C. Fishbein), a Medtronic-Zipes Endowment (P.-S. Chen), the Canadian Institutes of Health Research (6957, 43565, S. Nattel), and the Indiana University Health-Indiana University School of Medicine Strategic Research Initiative.

Disclosures

P.-S. Chen has the following patents relevant to the materials described in this review: US Patents 6351668, 6353757, 6398800, 6487450, 6824538, and 7266410. The other authors report no conflicts.

References

1. Viskin S, Golovner M, Malov N, Fish R, Alroy I, Vila Y, Laniado S, Kaplinsky E, Roth A. Circadian variation of symptomatic paroxysmal atrial fibrillation. Data from almost 10 000 episodes. *Eur Heart J*. 1999;20:1429–1434.
2. Leiria TL, Glavinovic T, Armour JA, Cardinal R, de Lima GG, Kus T. Longterm effects of cardiac mediastinal nerve cryoablation on neural inducibility of atrial fibrillation in canines. *Auton Neurosci*. 2011;161:68–74.
3. Richer LP, Vinet A, Kus T, Cardinal R, Ardell JL, Armour JA. Alpha-adrenoceptor blockade modifies neurally induced atrial arrhythmias. *Am J Physiol Regul Integr Comp Physiol*. 2008;295:R1175–R1180.
4. Tan AY, Zhou S, Ogawa M, Song J, Chu M, Li H, Fishbein MC, Lin SF, Chen LS, Chen PS. Neural mechanisms of paroxysmal atrial fibrillation and paroxysmal atrial tachycardia in ambulatory canines. *Circulation*. 2008;118:916–925.
5. Shen MJ, Shinohara T, Park HW, et al. Continuous low-level vagus nerve stimulation reduces stellate ganglion nerve activity and paroxysmal atrial tachyarrhythmias in ambulatory canines. *Circulation*. 2011;123:2204–2212.
6. Wang X, Zhao Q, Huang H, Tang Y, Xiao J, Dai Z, Yu S, Huang C. Effect of renal sympathetic denervation on atrial substrate remodeling in ambulatory canines with prolonged atrial pacing. *PLoS One*. 2013;8:e64611.
7. Janes RD, Johnstone DE, Brandys JC, Armour JA. Functional and anatomical variability of canine cardiac sympathetic efferent pathways: implications for regional denervation of the left ventricle. *Can J Physiol Pharmacol*. 1986;64:958–969.
8. Janes RD, Brandys JC, Hopkins DA, Johnstone DE, Murphy DA, Armour JA. Anatomy of human extrinsic cardiac nerves and ganglia. *Am J Cardiol*. 1986;57:299–309.
9. Armour JA, Murphy DA, Yuan BX, Macdonald S, Hopkins DA. Gross and microscopic anatomy of the human intrinsic cardiac nervous system. *Anat Rec*. 1997;247:289–298.

10. Ardell JL. The cardiac neuronal hierarchy and susceptibility to arrhythmias. *Heart Rhythm*. 2011;8:590–591.
11. Kawashima T. The autonomic nervous system of the human heart with special reference to its origin, course, and peripheral distribution. *Anat Embryol (Berl)*. 2005;209:425–438.
12. Ellison JP, Williams TH. Sympathetic nerve pathways to the human heart, and their variations. *Am J Anat*. 1969;124:149–162.
13. Baron R, Jänig W, With H. Sympathetic and afferent neurones projecting into forelimb and trunk nerves and the anatomical organization of the thoracic sympathetic outflow of the rat. *J Auton Nerv Syst*. 1995;53:205–214.
14. Taniguchi T, Morimoto M, Taniguchi Y, Takasaki M, Totoki T. Cutaneous distribution of sympathetic postganglionic fibers from stellate ganglion: a retrograde axonal tracing study using wheat germ agglutinin conjugated with horseradish peroxidase. *J Anesth*. 1994;8:441–449.
15. Pagé PL, Dandan N, Savard P, Nadeau R, Armour JA, Cardinal R. Regional distribution of atrial electrical changes induced by stimulation of extracardiac and intracardiac neural elements. *J Thorac Cardiovasc Surg*. 1995;109:377–388.
16. Shen MJ, Hao-Che Chang, Park HW, George Akingba A, Chang PC, Zheng Zhang, Lin SF, Shen C, Chen LS, Chen Z, Fishbein MC, Chiamvimonvat N, Chen PS. Low-level vagus nerve stimulation upregulates small conductance calcium-activated potassium channels in the stellate ganglion. *Heart Rhythm*. 2013;10:910–915.
17. Onkka P, Maskoun W, Rhee KS, Hellyer J, Patel J, Tan J, Chen LS, Vinters HV, Fishbein MC, Chen PS. Sympathetic nerve fibers and ganglia in canine cervical vagus nerves: localization and quantitation. *Heart Rhythm*. 2013;10:585–591.
18. Park HW, Shen MJ, Han S, Shinohara T, Maruyama M, Lee YS, Shen C, Hwang C, Chen LS, Fishbein MC, Lin SF, Chen PS. Neural control of ventricular rate in ambulatory dogs with pacing-induced sustained atrial fibrillation. *Circ Arrhythm Electrophysiol*. 2012;5:571–580.
19. Lomuscio A, Belletti S, Battezzati PM, Lombardi F. Efficacy of acupuncture in preventing atrial fibrillation recurrences after electrical cardioversion. *J Cardiovasc Electrophysiol*. 2011;22:241–247.
20. Foly JO, DuBois FS. Quantitative studies of the vagus nerve in the cat. I. The ratio of sensory and motor fibers. *J Comp Neurol*. 1937;67:49–67.
21. Henry TR. Therapeutic mechanisms of vagus nerve stimulation. *Neurology*. 2002;59:S3–S14.
22. Randall WC, Priola DV, Pace JB. Responses of individual cardiac chambers to stimulation of the cervical vagosympathetic trunk in atropinized dogs. *Circ Res*. 1967;20:534–544.
23. Armour JA, Randall WC. Functional anatomy of canine cardiac nerves. *Acta Anat (Basel)*. 1975;91:510–528.
24. Armour JA, Hopkins DA. Anatomy of the extrinsic efferent autonomic nerves and ganglia innervating the mammalian heart. In: Randall WC, ed. *Nervous Control of Cardiovascular Function*. New York, NY: Oxford University Press; 1984:21–45.
25. Kawagishi K, Fukushima N, Yokouchi K, Sumitomo N, Kakegawa A, Moriizumi T. Tyrosine hydroxylase-immunoreactive fibers in the human vagus nerve. *J Clin Neurosci*. 2008;15:1023–1026.
26. Ogawa M, Zhou S, Tan AY, Song J, Gholmieh G, Fishbein MC, Luo H, Siegel RJ, Karagueuzian HS, Chen LS, Lin SF, Chen PS. Left stellate ganglion and vagal nerve activity and cardiac arrhythmias in ambulatory dogs with pacing-induced congestive heart failure. *J Am Coll Cardiol*. 2007;50:335–343.
27. Seki A, Tan J, Chen PS, Fishbein MC. Are there sympathetic nerve fibers in human cervical and thoracic vagus nerves? *Circulation*. 2013;128:A12365 (abstract).
28. Armour JA. Potential clinical relevance of the 'little brain' on the mammalian heart. *Exp Physiol*. 2008;93:165–176.
29. Tan AY, Li H, Wachsmann-Hogiu S, Chen LS, Chen PS, Fishbein MC. Autonomic innervation and segmental muscular disconnections at the human pulmonary vein-atrial junction: implications for catheter ablation of atrial-pulmonary vein junction. *J Am Coll Cardiol*. 2006;48:132–143.
30. Vracko R, Thorning D, Frederickson RG. Nerve fibers in human myocardial scars. *Hum Pathol*. 1991;22:138–146.
31. Vracko R, Thorning D, Frederickson RG. Fate of nerve fibers in necrotic, healing, and healed rat myocardium. *Lab Invest*. 1990;63:490–501.
32. Cao JM, Chen LS, KenKnight BH, Ohara T, Lee MH, Tsai J, Lai WW, Karagueuzian HS, Wolf PL, Fishbein MC, Chen PS. Nerve sprouting and sudden cardiac death. *Circ Res*. 2000;86:816–821.
33. Swissa M, Zhou S, Gonzalez-Gomez I, Chang CM, Lai AC, Cates AW, Fishbein MC, Karagueuzian HS, Chen PS, Chen LS. Long-term sub-threshold electrical stimulation of the left stellate ganglion and a canine model of sudden cardiac death. *J Am Coll Cardiol*. 2004;43:858–864.
34. Zhou S, Chen LS, Miyauchi Y, Miyauchi M, Kar S, Kangavari S, Fishbein MC, Sharifi B, Chen PS. Mechanisms of cardiac nerve sprouting after myocardial infarction in dogs. *Circ Res*. 2004;95:76–83.
35. Miyauchi Y, Zhou S, Miyauchi M, Omichi C, Okuyama Y, Hamabe A, Hayashi H, Mandel WJ, Fishbein MC, Chen LS, Chen PS, Karagueuzian HS. Induction of atrial sympathetic nerve sprouting and increased vulnerability to atrial fibrillation by chronic left ventricular myocardial infarction. *Circulation*. 2001;104:II-77.
36. Arora R. Recent insights into the role of the autonomic nervous system in the creation of substrate for atrial fibrillation: implications for therapies targeting the atrial autonomic nervous system. *Circ Arrhythm Electrophysiol*. 2012;5:850–859.
37. Volders PG. Novel insights into the role of the sympathetic nervous system in cardiac arrhythmogenesis. *Heart Rhythm*. 2010;7:1900–1906.
38. Patterson E, Jackman WM, Beckman KJ, Lazzara R, Lockwood D, Scherlag BJ, Wu R, Po S. Spontaneous pulmonary vein firing in man: relationship to tachycardia-pause early afterdepolarizations and triggered arrhythmia in canine pulmonary veins in vitro. *J Cardiovasc Electrophysiol*. 2007;18:1067–1075.
39. Scherlag BJ, Patterson E, Po SS. The neural basis of atrial fibrillation. *J Electrocardiol*. 2006;39:S180–S183.
40. Patterson E, Po SS, Scherlag BJ, Lazzara R. Triggered firing in pulmonary veins initiated by in vitro autonomic nerve stimulation. *Heart Rhythm*. 2005;2:624–631.
41. Jayachandran JV, Sih HJ, Winkle W, Zipes DP, Hutchins GD, Olgin JE. Atrial fibrillation produced by prolonged rapid atrial pacing is associated with heterogeneous changes in atrial sympathetic innervation. *Circulation*. 2000;101:1185–1191.
42. Chang CM, Wu TJ, Zhou S, Doshi RN, Lee MH, Ohara T, Fishbein MC, Karagueuzian HS, Chen PS, Chen LS. Nerve sprouting and sympathetic hyperinnervation in a canine model of atrial fibrillation produced by prolonged right atrial pacing. *Circulation*. 2001;103:22–25.
43. Nguyen BL, Fishbein MC, Chen LS, Chen PS, Masroor S. Histopathological substrate for chronic atrial fibrillation in humans. *Heart Rhythm*. 2009;6:454–460.
44. Ng J, Villuendas R, Cokic I, Schliamsner JE, Gordon D, Koduri H, Benefield B, Simon J, Murthy SN, Lomasney JW, Wasserstrom JA, Goldberger JJ, Aistrup GL, Arora R. Autonomic remodeling in the left atrium and pulmonary veins in heart failure: creation of a dynamic substrate for atrial fibrillation. *Circ Arrhythm Electrophysiol*. 2011;4:388–396.
45. Arora R, Ulphani JS, Villuendas R, et al. Neural substrate for atrial fibrillation: implications for targeted parasympathetic blockade in the posterior left atrium. *Am J Physiol Heart Circ Physiol*. 2008;294:H134–H144.
46. Ajijola OA, Yagishita D, Patel KJ, Vaseghi M, Zhou W, Yamakawa K, So E, Lux RL, Mahajan A, Shivkumar K. Focal myocardial infarction induces global remodeling of cardiac sympathetic innervation: neural remodeling in a spatial context. *Am J Physiol Heart Circ Physiol*. 2013;305:H1031–H1040.
47. Vaseghi M, Lux RL, Mahajan A, Shivkumar K. Sympathetic stimulation increases dispersion of repolarization in humans with myocardial infarction. *Am J Physiol Heart Circ Physiol*. 2012;302:H1838–H1846.
48. Ajijola OA, Wisco JJ, Lambert HW, Mahajan A, Stark E, Fishbein MC, Shivkumar K. Extracardiac neural remodeling in humans with cardiomyopathy. *Circ Arrhythm Electrophysiol*. 2012;5:1010–1116.
49. Nguyen BL, Li H, Fishbein MC, Lin SF, Gaudio C, Chen PS, Chen LS. Acute myocardial infarction induces bilateral stellate ganglia neural remodeling in rabbits. *Cardiovasc Pathol*. 2012;21:143–148.
50. Han S, Kobayashi K, Joung B, Piccirillo G, Maruyama M, Vinters HV, March K, Lin SF, Shen C, Fishbein MC, Chen PS, Chen LS. Electroanatomic remodeling of the left stellate ganglion after myocardial infarction. *J Am Coll Cardiol*. 2012;59:954–961.
51. Francis GS. Modulation of peripheral sympathetic nerve transmission. *J Am Coll Cardiol*. 1988;12:250–254.
52. Lympferopoulos A, Rengo G, Koch WJ. Adrenergic nervous system in heart failure: pathophysiology and therapy. *Circ Res*. 2013;113:739–753.
53. Oberhauser V, Schwertfeger E, Rutz T, Beyersdorf F, Rump LC. Acetylcholine release in human heart atrium: influence of muscarinic autoreceptors, diabetes, and age. *Circulation*. 2001;103:1638–1643.
54. Nerbonne JM, Kass RS. Molecular physiology of cardiac repolarization. *Physiol Rev*. 2005;85:1205–1253.
55. Heijman J, Voigt N, Nattel S, Dobrev D. Cellular and molecular electrophysiology of atrial fibrillation initiation, maintenance, and progression. *Circ Res*. 2014;114:1483–1499.
56. Nattel S, Dobrev D. The multidimensional role of calcium in atrial fibrillation pathophysiology: mechanistic insights and therapeutic opportunities. *Eur Heart J*. 2012;33:1870–1877.

57. Yeh YH, Wakili R, Qi XY, Chartier D, Boknik P, Kääh S, Ravens U, Coutu P, Dobrev D, Nattel S. Calcium-handling abnormalities underlying atrial arrhythmogenesis and contractile dysfunction in dogs with congestive heart failure. *Circ Arrhythm Electrophysiol.* 2008;1:93–102.
58. Voigt N, Heijman J, Wang Q, Chiang DY, Li N, Karck M, Wehrens XH, Nattel S, Dobrev D. Cellular and molecular mechanisms of atrial arrhythmogenesis in patients with paroxysmal atrial fibrillation. *Circulation.* 2014;129:145–156.
59. Nishida K, Qi XY, Wakili R, Comtois P, Chartier D, Harada M, Iwasaki YK, Romeo P, Maguy A, Dobrev D, Michael G, Talajic M, Nattel S. Mechanisms of atrial tachyarrhythmias associated with coronary artery occlusion in a chronic canine model. *Circulation.* 2011;123:137–146.
60. Zicha S, Tsuji Y, Shiroshita-Takeshita A, Nattel S. Beta-blockers as anti-arrhythmic agents. In: Kass RS, Clancy CC, eds. *Handbook of Experimental Pharmacology—Basis and Treatment of Cardiac Arrhythmias.* Heidelberg, Germany: Springer Publishing; 2005:235–266.
61. Bartos DC, Giudicessi JR, Tester DJ, Ackerman MJ, Ohno S, Horie M, Gollob MH, Burgess DE, Delisle BP. A KCNQ1 mutation contributes to the concealed type 1 long QT phenotype by limiting the Kv7.1 channel conformational changes associated with protein kinase A phosphorylation. *Heart Rhythm.* 2014;11:459–468.
62. Han W, Wang Z, Nattel S. Slow delayed rectifier current and repolarization in canine cardiac Purkinje cells. *Am J Physiol Heart Circ Physiol.* 2001;280:H1075–H1080.
63. Fink M, Giles WR, Noble D. Contributions of inwardly rectifying K⁺ currents to repolarization assessed using mathematical models of human ventricular myocytes. *Philos Trans A Math Phys Eng Sci.* 2006;364:1207–1222.
64. Pandit SV, Berenfeld O, Anumonwo JM, Zaritski RM, Kneller J, Nattel S, Jalife J. Ionic determinants of functional reentry in a 2-D model of human atrial cells during simulated chronic atrial fibrillation. *Biophys J.* 2005;88:3806–3821.
65. Sato R, Koumi S. Modulation of the inwardly rectifying K⁺ channel in isolated human atrial myocytes by alpha 1-adrenergic stimulation. *J Membr Biol.* 1995;148:185–191.
66. Andrade J, Khairy P, Dobrev D, Nattel S. The clinical profile and pathophysiology of atrial fibrillation: relationships among clinical features, epidemiology, and mechanisms. *Circ Res.* 2014;114:1453–1468.
67. Nattel S. Paroxysmal atrial fibrillation and pulmonary veins: relationships between clinical forms and automatic versus re-entrant mechanisms. *Can J Cardiol.* 2013;29:1147–1149.
68. DiFrancesco D. The role of the funny current in pacemaker activity. *Circ Res.* 2010;106:434–446.
69. Lemoine MD, Duverger JE, Naud P, Chartier D, Qi XY, Comtois P, Fabritz L, Kirchhoff P, Nattel S. Arrhythmogenic left atrial cellular electrophysiology in a murine genetic long QT syndrome model. *Cardiovasc Res.* 2011;92:67–74.
70. Maruyama M, Lin SF, Xie Y, Chua SK, Joung B, Han S, Shinohara T, Shen MJ, Qu Z, Weiss JN, Chen PS. Genesis of phase 3 early afterdepolarizations and triggered activity in acquired long-QT syndrome. *Circ Arrhythm Electrophysiol.* 2011;4:103–111.
71. Burashnikov A, Antzelevitch C. Reinduction of atrial fibrillation immediately after termination of the arrhythmia is mediated by late phase 3 early afterdepolarization-induced triggered activity. *Circulation.* 2003;107:2355–2360.
72. Choi EK, Chang PC, Lee YS, Lin SF, Zhu W, Maruyama M, Fishbein MC, Chen Z, Rubart-von der Lohe M, Field LJ, Chen PS. Triggered firing and atrial fibrillation in transgenic mice with selective atrial fibrosis induced by overexpression of TGF- β 1. *Circ J.* 2012;76:1354–1362.
73. Patterson E, Lazzara R, Szabo B, Liu H, Tang D, Li YH, Scherlag BJ, Po SS. Sodium-calcium exchange initiated by the Ca²⁺ transient: an arrhythmia trigger within pulmonary veins. *J Am Coll Cardiol.* 2006;47:1196–1206.
74. Pandit SV, Jalife J. Rotors and the dynamics of cardiac fibrillation. *Circ Res.* 2013;112:849–862.
75. Liu L, Nattel S. Differing sympathetic and vagal effects on atrial fibrillation in dogs: role of refractoriness heterogeneity. *Am J Physiol.* 1997;273:H805–H816.
76. Barrett CJ, Ramchandra R, Guild SJ, Lala A, Budgett DM, Malpas SC. What sets the long-term level of renal sympathetic nerve activity: a role for angiotensin II and baroreflexes? *Circ Res.* 2003;92:1330–1336.
77. Jung BC, Dave AS, Tan AY, Gholmieh G, Zhou S, Wang DC, Akingba AG, Fishbein GA, Montemagno C, Lin SF, Chen LS, Chen PS. Circadian variations of stellate ganglion nerve activity in ambulatory dogs. *Heart Rhythm.* 2006;3:78–85.
78. Sharifov OF, Fedorov VV, Beloshapko GG, Glukhov AV, Yushmanov AV, Rosenshtraukh LV. Roles of adrenergic and cholinergic stimulation in spontaneous atrial fibrillation in dogs. *J Am Coll Cardiol.* 2004;43:483–490.
79. Nishida K, Maguy A, Sakabe M, Comtois P, Inoue H, Nattel S. The role of pulmonary veins vs. autonomic ganglia in different experimental substrates of canine atrial fibrillation. *Cardiovasc Res.* 2011;89:825–833.
80. Wyse DG, Waldo AL, DiMarco JP, Domanski MJ, Rosenberg Y, Schron EB, Kellen JC, Greene HL, Mickel M, Dalquist JE, Corley SD. Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) Investigators. A comparison of rate control and rhythm control in patients with atrial fibrillation. *N Engl J Med.* 2002;347:1825–1833.
81. Hou Y, Scherlag BJ, Lin J, Zhang Y, Lu Z, Truong K, Patterson E, Lazzara R, Jackman WM, Po SS. Ganglionated plexi modulate extrinsic cardiac autonomic nerve input: effects on sinus rate, atrioventricular conduction, refractoriness, and inducibility of atrial fibrillation. *J Am Coll Cardiol.* 2007;50:61–68.
82. Rossi P, Bianchi S, Barretta A, Della Scala A, Kornet L, De Paulis R, Bellisario A, D'Addio V, Pavaci H, Miraldi F. Post-operative atrial fibrillation management by selective epicardial vagal fat pad stimulation. *J Interv Card Electrophysiol.* 2009;24:37–45.
83. Shen MJ, Choi EK, Tan AY, Han S, Shinohara T, Maruyama M, Chen LS, Shen C, Hwang C, Lin SF, Chen PS. Patterns of baseline autonomic nerve activity and the development of pacing-induced sustained atrial fibrillation. *Heart Rhythm.* 2011;8:583–589.
84. Shinohara T, Shen MJ, Han S, Maruyama M, Park HW, Fishbein MC, Shen C, Chen PS, Lin SF. Heart failure decreases nerve activity in the right atrial ganglionated plexus. *J Cardiovasc Electrophysiol.* 2012;23:404–412.
85. Hellyer J, George Akingba A, Rhee KS, Tan AY, Lane KA, Shen C, Patel J, Fishbein MC, Chen PS. Autonomic nerve activity and blood pressure in ambulatory dogs. *Heart Rhythm.* 2014;11:307–313.
86. Choi EK, Shen MJ, Han S, Kim D, Hwang S, Sayfo S, Piccirillo G, Frick K, Fishbein MC, Hwang C, Lin SF, Chen PS. Intrinsic cardiac nerve activity and paroxysmal atrial tachyarrhythmia in ambulatory dogs. *Circulation.* 2010;121:2615–2623.
87. Schwartz PJ, Locati EH, Moss AJ, Crampton RS, Trazzi R, Ruberti U. Left cardiac sympathetic denervation in the therapy of congenital long QT syndrome. A worldwide report. *Circulation.* 1991;84:503–511.
88. Ogawa M, Tan AY, Song J, Kobayashi K, Fishbein MC, Lin SF, Chen LS, Chen. Cryoablation of extrinsic cardiac sympathetic nerves markedly reduces atrial arrhythmias in ambulatory dogs with pacing-induced heart failure. *Heart Rhythm.* 2008;5:S54.
89. Benjamin EJ, Chen PS, Bild DE, et al. Prevention of atrial fibrillation: report from a national heart, lung, and blood institute workshop. *Circulation.* 2009;119:606–618.
90. Coleman MA, Bos JM, Johnson JN, Owen HJ, Deschamps C, Moir C, Ackerman MJ. Videoscopic left cardiac sympathetic denervation for patients with recurrent ventricular fibrillation/malignant ventricular arrhythmia syndromes besides congenital long-QT syndrome. *Circ Arrhythm Electrophysiol.* 2012;5:782–788.
91. Schauer P, Scherlag BJ, Scherlag MA, Goli S, Jackman WM, Lazzara R. Ventricular rate control during atrial fibrillation by cardiac parasympathetic nerve stimulation: a transvenous approach. *J Am Coll Cardiol.* 1999;34:2043–2050.
92. Goldberger AL, Pavelec RS. Vagally-mediated atrial fibrillation in dogs: conversion with bretylium tosylate. *Int J Cardiol.* 1986;13:47–55.
93. Wang Z, Pagé P, Nattel S. Mechanism of flecainide's antiarrhythmic action in experimental atrial fibrillation. *Circ Res.* 1992;71:271–287.
94. Schauer P, Scherlag BJ, Pitha J, Scherlag MA, Reynolds D, Lazzara R, Jackman WM. Catheter ablation of cardiac autonomic nerves for prevention of vagal atrial fibrillation. *Circulation.* 2000;102:2774–2780.
95. Scherlag BJ, Yamanashi WS, Schauer P, Scherlag M, Sun YX, Hou Y, Jackman WM, Lazzara R. Endovascular stimulation within the left pulmonary artery to induce slowing of heart rate and paroxysmal atrial fibrillation. *Cardiovasc Res.* 2002;54:470–475.
96. Elvan A, Huang XD, Pressler ML, Zipes DP. Radiofrequency catheter ablation of the atria eliminates pacing-induced sustained atrial fibrillation and reduces connexin 43 in dogs. *Circulation.* 1997;96:1675–1685.
97. Elvan A, Pride HP, Eble JN, Zipes DP. Radiofrequency catheter ablation of the atria reduces inducibility and duration of atrial fibrillation in dogs. *Circulation.* 1995;91:2235–2244.
98. Schwartz PJ, Pagani M, Lombardi F, Malliani A, Brown AM. A cardio-cardiac sympathovagal reflex in the cat. *Circ Res.* 1973;32:215–220.
99. Terry R. Vagus nerve stimulation: a proven therapy for treatment of epilepsy strives to improve efficacy and expand applications. *Conf Proc IEEE Eng Med Biol Soc.* 2009;2009:4631–4634.
100. Vanoli E, De Ferrari GM, Stramba-Badiale M, Hull SS Jr, Foreman RD, Schwartz PJ. Vagal stimulation and prevention of sudden death

- in conscious dogs with a healed myocardial infarction. *Circ Res*. 1991;68:1471–1481.
101. Zhang Y, Popovic ZB, Bibeovski S, Fakhry I, Sica DA, Van Wagoner DR, Mazgalev TN. Chronic vagus nerve stimulation improves autonomic control and attenuates systemic inflammation and heart failure progression in a canine high-rate pacing model. *Circ Heart Fail*. 2009;2:692–699.
 102. Li M, Zheng C, Sato T, Kawada T, Sugimachi M, Sunagawa K. Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats. *Circulation*. 2004;109:120–124.
 103. De Ferrari GM, Sanzo A, Schwartz PJ. Chronic vagal stimulation in patients with congestive heart failure. *Conf Proc IEEE Eng Med Biol Soc*. 2009;2009:2037–2039.
 104. De Ferrari GM, Crijns HJ, Borggreve M, Milasinovic G, Smid J, Zabel M, Gavazzi A, Sanzo A, Dennert R, Kuschyk J, Raspopovic S, Klein H, Swedberg K, Schwartz PJ; CardioFit Multicenter Trial Investigators. Chronic vagus nerve stimulation: a new and promising therapeutic approach for chronic heart failure. *Eur Heart J*. 2011;32:847–855.
 105. De Ferrari GM, Schwartz PJ. Vagus nerve stimulation: from pre-clinical to clinical application: challenges and future directions. *Heart Fail Rev*. 2011;16:195–203.
 106. Li S, Scherlag BJ, Yu L, Sheng X, Zhang Y, Ali R, Dong Y, Ghias M, Po SS. Low-level vagosympathetic stimulation: a paradox and potential new modality for the treatment of focal atrial fibrillation. *Circ Arrhythm Electrophysiol*. 2009;2:645–651.
 107. Yu L, Scherlag BJ, Li S, Sheng X, Lu Z, Nakagawa H, Zhang Y, Jackman WM, Lazzara R, Jiang H, Po SS. Low-level vagosympathetic nerve stimulation inhibits atrial fibrillation inducibility: direct evidence by neural recordings from intrinsic cardiac ganglia. *J Cardiovasc Electrophysiol*. 2011;22:455–463.
 108. Vanhoutte PM, Levy MN. Prejunctional cholinergic modulation of adrenergic neurotransmission in the cardiovascular system. *Am J Physiol*. 1980;238:H275–H281.
 109. Takahashi N, Zipes DP. Vagal modulation of adrenergic effects on canine sinus and atrioventricular nodes. *Am J Physiol*. 1983;244:H775–H781.
 110. Stanley J, Peake JM, Buchheit M. Cardiac parasympathetic reactivation following exercise: implications for training prescription. *Sports Med*. 2013;43:1259–1277.
 111. Guasch E, Benito B, Qi X, et al. Atrial fibrillation promotion by endurance exercise: demonstration and mechanistic exploration in an animal model. *J Am Coll Cardiol*. 2013;62:68–77.
 112. Sorokin AV, Araujo CG, Zweibel S, Thompson PD. Atrial fibrillation in endurance-trained athletes. *Br J Sports Med*. 2011;45:185–188.
 113. Reed JL, Mark AE, Reid RD, Pipe AL. The effects of chronic exercise training in individuals with permanent atrial fibrillation: a systematic review. *Can J Cardiol*. 2013;29:1721–1728.
 114. Scheffers JJ, Kroon AA, Schmidli J, et al. Novel baroreflex activation therapy in resistant hypertension: results of a European multi-center feasibility study. *J Am Coll Cardiol*. 2010;56:1254–1258.
 115. Tordoir JH, Scheffers I, Schmidli J, Savolainen H, Liebeskind U, Hansky B, Herold U, Irwin E, Kroon AA, de Leeuw P, Peters TK, Kieval R, Cody R. An implantable carotid sinus baroreflex activating system: surgical technique and short-term outcome from a multi-center feasibility trial for the treatment of resistant hypertension. *Eur J Vasc Endovasc Surg*. 2007;33:414–421.
 116. Heusser K, Tank J, Engeli S, Diedrich A, Menne J, Eckert S, Peters T, Sweep FC, Haller H, Pichlmaier AM, Luft FC, Jordan J. Carotid baroreceptor stimulation, sympathetic activity, baroreflex function, and blood pressure in hypertensive patients. *Hypertension*. 2010;55:619–626.
 117. Linz D, Mahfoud F, Schotten U, Ukena C, Neuberger HR, Wirth K, Böhm M. Effects of electrical stimulation of carotid baroreflex and renal denervation on atrial electrophysiology. *J Cardiovasc Electrophysiol*. 2013;24:1028–1033.
 118. Linz D, Ukena C, Mahfoud F, Neuberger HR, Böhm M. Atrial autonomic innervation: a target for interventional antiarrhythmic therapy? *J Am Coll Cardiol*. 2014;63:215–224.
 119. Pappone C, Santinelli V, Manguso F, Vicedomini G, Gugliotta F, Augello G, Mazzone P, Tortorello V, Landoni G, Zangrillo A, Lang C, Tomita T, Mesas C, Mastella E, Alfieri O. Pulmonary vein denervation enhances long-term benefit after circumferential ablation for paroxysmal atrial fibrillation. *Circulation*. 2004;109:327–334.
 120. Po SS, Nakagawa H, Jackman WM. Localization of left atrial ganglionated plexi in patients with atrial fibrillation. *J Cardiovasc Electrophysiol*. 2009;20:1186–1189.
 121. He B, Scherlag BJ, Nakagawa H, Lazzara R, Po SS. The intrinsic autonomic nervous system in atrial fibrillation: a review. *ISRN Cardiol*. 2012;2012:490674.
 122. Zhang Y, Wang Z, Zhang Y, Wang W, Wang J, Gao M, Hou Y. Efficacy of cardiac autonomic denervation for atrial fibrillation: a meta-analysis. *J Cardiovasc Electrophysiol*. 2012;23:592–600.
 123. Katritsis DG, Pokushalov E, Romanov A, Giazitzoglou E, Siontis GC, Po SS, Camm AJ, Ioannidis JP. Autonomic denervation added to pulmonary vein isolation for paroxysmal atrial fibrillation: a randomized clinical trial. *J Am Coll Cardiol*. 2013;62:2318–2325.
 124. Zheng S, Li Y, Han J, Zhang H, Zeng W, Xu C, Jia Y, Wang J, Guo K, Jiao Y, Meng X. Long-term results of a minimally invasive surgical pulmonary vein isolation and ganglionic plexi ablation for atrial fibrillation. *PLoS One*. 2013;8:e79755.
 125. Esler MD, Krum H, Schlaich M, Schmieder RE, Böhm M, Sobotka PA; Symplicity HTN-2 Investigators. Renal sympathetic denervation for treatment of drug-resistant hypertension: one-year results from the Symplicity HTN-2 randomized, controlled trial. *Circulation*. 2012;126:2976–2982.
 126. Esler MD, Krum H, Sobotka PA, Schlaich MP, Schmieder RE, Böhm M; Symplicity HTN-2 Investigators. Renal sympathetic denervation in patients with treatment-resistant hypertension (The Symplicity HTN-2 Trial): a randomised controlled trial. *Lancet*. 2010;376:1903–1909.
 127. Schlaich MP, Sobotka PA, Krum H, Lambert E, Esler MD. Renal sympathetic-nerve ablation for uncontrolled hypertension. *N Engl J Med*. 2009;361:932–934.
 128. Ahmed H, Miller MA, Dukkupati SR, Cammack S, Koruth JS, Gangireddy S, Ellsworth BA, D'Avila A, Domanski M, Gelijns AC, Moskowitz A, Reddy VY. Adjunctive renal sympathetic denervation to modify hypertension as upstream therapy in the treatment of atrial fibrillation (H-FIB) study: clinical background and study design. *J Cardiovasc Electrophysiol*. 2013;24:503–509.
 129. Hogarth AJ, Dobson LE, Tayebjee MH. During ablation for atrial fibrillation, is simultaneous renal artery ablation appropriate? *J Hum Hypertens*. 2013;27:707–714.
 130. Linz D, Hohl M, Nickel A, Mahfoud F, Wagner M, Ewen S, Schotten U, Maack C, Wirth K, Böhm M. Effect of renal denervation on neurohumoral activation triggering atrial fibrillation in obstructive sleep apnea. *Hypertension*. 2013;62:767–774.
 131. Nozawa T, Igawa A, Fujii N, Kato B, Yoshida N, Asanoi H, Inoue H. Effects of long-term renal sympathetic denervation on heart failure after myocardial infarction in rats. *Heart Vessels*. 2002;16:51–56.
 132. Kandzari DE, Bhatt DL, Sobotka PA, O'Neill WW, Esler M, Flack JM, Katzen BT, Leon MB, Massaro JM, Negoita M, Oparil S, Rocha-Singh K, Straley C, Townsend RR, Bakris G. Catheter-based renal denervation for resistant hypertension: rationale and design of the SYMPPLICITY HTN-3 Trial. *Clin Cardiol*. 2012;35:528–535.
 133. Bhatt DL, Kandzari DE, O'Neill WW, D'Agostino R, Flack JM, Katzen BT, Leon MB, Liu M, Mauri L, Negoita M, Cohen SA, Oparil S, Rocha-Singh K, Townsend RR, Bakris GL, Investigators SH-. A controlled trial of renal denervation for resistant hypertension. *N Engl J Med*. 2014;370:1393–1401.
 134. Floras JS. Renal denervation for drug-resistant hypertension: suffering its original sin, seeking redemption. *Can J Cardiol*. 2014;30:476–478.
 135. Kimura A, Sato A. Somatic regulation of autonomic functions in anesthetized animals—neural mechanisms of physical therapy including acupuncture. *Jpn J Vet Res*. 1997;45:137–145.
 136. Yu L, Scherlag BJ, Li S, Fan Y, Dyer J, Male S, Varma V, Sha Y, Stavrakis S, Po SS. Low-level transcutaneous electrical stimulation of the auricular branch of the vagus nerve: a noninvasive approach to treat the initial phase of atrial fibrillation. *Heart Rhythm*. 2013;10:428–435.
 137. White AR, Ernst E. A systematic review of randomized controlled trials of acupuncture for neck pain. *Rheumatology (Oxford)*. 1999;38:143–147.
 138. Ezzo J, Berman B, Hadhazy VA, Jadad AR, Lao L, Singh BB. Is acupuncture effective for the treatment of chronic pain? A systematic review. *Pain*. 2000;86:217–225.
 139. Himura Y, Felten SY, Kashiki M, Lewandowski TJ, Delehanty JM, Liang CS. Cardiac noradrenergic nerve terminal abnormalities in dogs with experimental congestive heart failure. *Circulation*. 1993;88:1299–1309.
 140. Dong E, Yatani A, Mohan A, Liang CS. Myocardial beta-adrenoceptor down-regulation by norepinephrine is linked to reduced norepinephrine uptake activity. *Eur J Pharmacol*. 1999;384:17–24.
 141. Kawai H, Fan TH, Dong E, Siddiqui RA, Yatani A, Stevens SY, Liang CS. ACE inhibition improves cardiac NE uptake and attenuates

- sympathetic nerve terminal abnormalities in heart failure. *Am J Physiol.* 1999;277:H1609–H1617.
142. Cha YM, Redfield MM, Shah S, Shen WK, Fishbein MC, Chen PS. Effects of omapatrilat on cardiac nerve sprouting and structural remodeling in experimental congestive heart failure. *Heart Rhythm.* 2005;2:984–990.
 143. Martelli D, McKinley MJ, McAllen RM. The cholinergic anti-inflammatory pathway: a critical review [published online ahead of print]. *Auton Neurosci.* 2013. pii: S1566-0702(13)00777-7. doi: 10.1016/j.autneu.2013.12.007.
 144. Rossi P, Ricci A, De Paulis R, et al. Epicardial ganglionated plexus stimulation decreases postoperative inflammatory response in humans. *Heart Rhythm.* 2012;9:943–950.
 145. Kusunose K, Zhang Y, Mazgalev TN, Van Wagoner DR, Thomas JD, Popović ZB. Impact of vagal nerve stimulation on left atrial structure and function in a canine high-rate pacing model. *Circ Heart Fail.* 2014;7:320–326.
 146. Nergårdh AK, Rosenqvist M, Nordlander R, Frick M. Maintenance of sinus rhythm with metoprolol CR initiated before cardioversion and repeated cardioversion of atrial fibrillation: a randomized double-blind placebo-controlled study. *Eur Heart J.* 2007;28:1351–1357.
 147. Nattel S. Ionic determinants of atrial fibrillation and Ca²⁺ channel abnormalities: cause, consequence, or innocent bystander? *Circ Res.* 1999;85:473–476.
 148. Arsenaault KA, Yusuf AM, Crystal E, Healey JS, Morillo CA, Nair GM, Whitlock RP. Interventions for preventing post-operative atrial fibrillation in patients undergoing heart surgery. *Cochrane Database Syst Rev.* 2013;1:CD003611.
 149. Machida T, Hashimoto N, Kuwahara I, Ogino Y, Matsuura J, Yamamoto W, Itano Y, Zamma A, Matsumoto R, Kamon J, Kobayashi T, Ishiwata N, Yamashita T, Ogura T, Nakaya H. Effects of a highly selective acetylcholine-activated K⁺ channel blocker on experimental atrial fibrillation. *Circ Arrhythm Electrophysiol.* 2011;4:94–102.
 150. Bauer A, McDonald AD, Nasir K, Peller L, Rade JJ, Miller JM, Heldman AW, Donahue JK. Inhibitory G protein overexpression provides physiologically relevant heart rate control in persistent atrial fibrillation. *Circulation.* 2004;110:3115–3120.
 151. Aistrup GL, Cokic I, Ng J, Gordon D, Koduri H, Browne S, Arapi D, Segon Y, Goldstein J, Angulo A, Wasserstrom JA, Goldberger JJ, Kadish AH, Arora R. Targeted nonviral gene-based inhibition of Gα(i/o)-mediated vagal signaling in the posterior left atrium decreases vagal-induced atrial fibrillation. *Heart Rhythm.* 2011;8:1722–1729.

Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Role of the Autonomic Nervous System in Atrial Fibrillation: Pathophysiology and Therapy

Peng-Sheng Chen, Lan S. Chen, Michael C. Fishbein, Shien-Fong Lin and Stanley Nattel

Circ Res. 2014;114:1500-1515

doi: 10.1161/CIRCRESAHA.114.303772

Circulation Research is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2014 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circres.ahajournals.org/content/114/9/1500>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Circulation Research* is online at:
<http://circres.ahajournals.org/subscriptions/>