

# Impact Of An Interactive Care Plan On Patient Activation In Heart Failure Inpatients

Dominique Comer, PharmD  
Jefferson School of Population Health

Abstract recently presented at the ISPOR International Annual Meeting: Washington, DC

Dominique Comer, PharmD<sup>1</sup>; David Wright, MPH<sup>2</sup>; Julia Southerton<sup>2</sup>; Angie Zanke<sup>2</sup>; Joseph Couto, PharmD, MBA<sup>1</sup>

<sup>1</sup>Jefferson School of Population Health, Philadelphia, PA; <sup>2</sup>Institute for Interactive Patient Care, Bethesda, MD

# Funding

- Research was completed by the Institute for Interactive Patient Care (IIPC) in conjunction with the Jefferson School of Population Health.
- Funding for this research was provided by GetWellNetwork, Inc.

# What exactly is Patient Activation?

- As defined by Hibbard et al:
- “An engaged patient who demonstrates an understanding that actions determine health outcomes and that they have the confidence to institute change.”


# Patient Activation

- Patient's playing a more active role in their care and advocating for themselves
- Seeks to improve the following areas:
  - Health Literacy
  - Clinical Decision Making
  - Self Care
  - Patient Safety

# Implications of Patient Activation

- Patients who are activated<sup>1,2</sup> :
  - Participate in self-management behaviors
  - Have higher medication adherence
  - Have higher quality of life scores
  - Improved outcomes in patients with chronic conditions

MAY 2012, VOLUME 1, ISSUE 3



# Journal of Comparative Effectiveness Research

## Senior Editors:

Eugene Rich & Sheldon Greenfield

## Editorials

Patient engagement: the critical catalyst to health reform in the USA

DE Couto & DM Comer

Lessons learned from the Practice-Based Opportunities for Weight Reduction (POWER) trial  
JJ Jerome, RR Rubin, JM Clark *et al.*

## News & Views

### News

#### Interview

Health technology assessment in Asia  
B Yang

### Review

Pharmacologic and nonpharmacologic thromboprophylactic strategies in atrial fibrillation  
T Alam, CA Clyne & CM White

### Research Article

Prioritizing comparative effectiveness research for cancer diagnostics using a regional stakeholder approach  
G Klein, LS Gold, SD Sullivan *et al.*

## Spotlight Article Series:

**A translation table for patient-centered comparative effectiveness research**

### Foreword

A translation table for patient-centered comparative effectiveness research  
SR Tunis, DA Messner, P Mohr, RE Glicklich & RW Dubois

### Reviews

Changing the clinical research paradigm to account for comparative effectiveness research  
S Greenfield & SH Kaplan

Which design for which question? An exploration toward a translation table for comparative effectiveness research  
VM Montori, SP Kim, GH Guyatt & ND Shah

### Research Article

Incorporating stakeholder perspectives in developing a translation table framework for comparative effectiveness research  
RE Glicklich, MB Leavy, P Velentgas *et al.*

### Commentary

Comparative effectiveness research: a view from the other side of the pond  
MD Rawlins

ISSN 2042-6305

Future  
Medicine part of



# Background

- Heart failure (HF) is a serious chronic condition that requires significant self-care to manage effectively and reduce associated avoidable readmissions.
- The inpatient acute care setting provides ample opportunities to educate and engage patients in their care to increase patient activation and improve patient outcomes<sup>2</sup>.


# Background

- Patient activation requires a demonstrated understanding of their condition, medications and lifestyle change requirements prior to being discharged from the hospital.
- Patient activation can be measured using the PAM survey

# PAM

- The Patient Activation Measure (PAM)
  - Validated and reliable
  - Original 22-item survey, abbreviated 13-item survey
  - Assesses self-reported knowledge, confidence and skill for self-management of chronic disease

# PAM Activation Levels



<http://www.insigniahealth.com/solutions/patient-activation-measure/>

# Heart Failure Interactive Care Plan


- 4-Phase Care Plan designed to better prepare patient to manage condition
- Nurse Activated
- Focus on understanding:
  - Condition
  - Signs/Symptoms
  - Medications
  - Lifestyle change

# Interactive Care Plan: Phase 1






# Interactive Care Plan: Phases 2-3



# Interactive Care Plan: Phase 4



# Objective

- To determine if the use of the Heart Failure Interactive Care Plan results in an improvement in PAM-13 scores during the course of hospitalization.

# Study Subjects

- Inclusion Criteria
  - Patients 18 years and over with a primary diagnosis of heart failure (ICD-9 code 428.X) in hospitals utilizing GetWellNetwork
- Exclusion Criteria
  - Unable to demonstrate proficiency in literary assessment

# Methods

- Pre-test post-test design
- Patient activation measured using the PAM-13
- Two-tailed paired t-test
  - Difference in pre-test v. post-test


# Results

- 275 patients in total
  - Six participating hospitals
  - Six month time span
- An increase of 0.38 in levels of activation ( $p < 0.001$ ) was seen after the implementation of the care plan.

# Patient Activation Scores

Pre-Intervention

Post-Intervention




# Results

- When data were stratified to remove all patients with an initial activation level of 4, the remaining 107 patients showed an average improvement of 1.05 levels of activation ( $p < 0.001$ ).

# Results

- All-cause readmission rates were promising for one hospital site:

The Indiana Heart Hospital All-Cause Readmissions Trend



# Further Analysis

- Gender and age were only available for a subset of the 275 patients that participated in the study (n=113).
- Males showed a generally higher level of activation at pre-test (3.34) compared to females (3.18), although the difference was not statistically significant.
- Both males and females demonstrated statistically significant increases in activation ( $p < 0.01$ ).

# Further Analysis

Description by Age Group

Age Group	Pre-intervention	Post-intervention
<50	3.5	3.8
50-59	3.28	3.74
60-69	3.28	3.67
>70	3.32	3.62

# Limitations

- Limited demographics were collected
  - Limitation of available technology and man-power
- Demographics were not complete
- Most patients in the study were already in the top activation level, pre-test

# Future Research

- Future research includes analysis on patient activation and length of stay
- Further investigation of readmission rates, specific to heart failure

# Conclusion

- Interactive patient-centered educational tools such as the Heart Failure Interactive Care Plan can be used to improve patient activation in heart failure inpatients.

# Resources

- 1. Mosen DM, Schmittiel J, Hibbard J, Sobel D, Remmers C, Bellows J. Is patient activation associated with outcomes of care for adults with chronic conditions? *J Ambul Care Manage*. 2007 Jan-Mar;30(1):21-9.
- 2. Chu ES, Hakkarinen D, Evig C, Page S, Keniston A, Dickinson M, Albert RK. Underutilized time for health education of hospitalized patients. *J of Hosp Med*. 2008; 3(3):238-246.

# QUESTIONS???