

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): J. Kaler, S.L.S. Daniels, J.L. Wright, L.E. Green

Article Title: A randomised factorial design clinical trial to investigate the impact of parenteral long acting oxytetracycline, foot trimming and flunixin meglumine on time to recovery from lameness and foot lesions in sheep lame with footrot

Year of publication: in press

Link to published version:

<http://www.wiley.com/bw/journal.asp?ref=0891-6640&site=1>

Publisher statement: This is the pre-peer-reviewed version of the following article: Kaler, J.. et al. A randomised factorial design clinical trial to investigate the impact of parenteral long acting oxytetracycline, foot trimming and flunixin meglumine on time to recovery from lameness and foot lesions in sheep lame with footrot. Journal of Veterinary Internal Medicine, which will be published in final form at <http://www.wiley.com/bw/journal.asp?ref=0891-6640&site=1>

1 **Randomised clinical trial of long acting oxytetracycline, foot trimming and flunixin**
2 **meglumine on time to recovery in sheep with footrot**

3 J. Kaler, S.L.S. Daniels, J.L. Wright, L.E. Green

4 Department of Biological Sciences, University of Warwick, Coventry, CV4 7AL, UK

5

6 Short title: [Treatment for footrot](#)

7

8 Keywords: Ovine, infectious disease, *Dichelobacter nodosus*, antibacterials

9 Abbreviations: NSAID, Non steroidal anti-inflammatory drug; OR, Odds Ratio

10

11 The trial was carried out on a commercial farm in England under supervision of researchers from

12 University of Warwick, UK.

13

14 The study was supported by funds from the Biotechnology and Biological Sciences Research

15 Council BBE01870X1.

16

17 Acknowledgements: We thank the farmer for help and Graham Medley for discussion.

18

19 Corresponding author: J. Kaler, Dept. of Biological Sciences, University of Warwick, Coventry,

20 CV4 7AL, UK; j.kaler@warwick.ac.uk

21

1 **Abstract**

2 **Background:** Internationally, foot trimming is used by most farmers and parenteral antibacterials
3 by some, to treat sheep with footrot. Non steroidal anti-inflammatory drugs (NSAID) are
4 sometimes used. No clinical trials have compared these treatments.

5 **Objectives:** To investigate the above treatments on time to recovery from lameness and foot
6 lesions in sheep with footrot.

7 **Animals:** 53 sheep with footrot on a commercial farm in England.

8 **Methods:** In a randomised factorial design, the sheep were allocated to 6 treatment groups. The
9 treatments were oxytetracycline spray to all sheep (positive control) and one or more of:
10 [parenteral administration of long acting oxytetracycline](#), flunixin meglumine and foot trimming
11 on day 1 or day 6 of diagnosis. Follow-up was for 15 days. Time to recovery from lameness and
12 lesions was investigated with discrete- time survival models.

13 **Results:** There was significant association ($p<0.05$) between recovery from lameness and lesions.
14 Sheep receiving antibacterials parenterally recovered faster from lameness (odds ratio (OR): 4.92
15 (1.20-20.10)) [and](#) lesions (OR: 5.11 (1.16-22.4)) than positive controls [whereas](#) sheep foot
16 trimmed on day 1 (lameness- OR: 0.05 (0.005-0.51); lesions- OR: 0.06 (0.008-0.45)) or day 6 of
17 diagnosis (lameness-OR: 0.07 (0.01-0.72); lesions- OR: 0.07 (0.01-0.56)) recovered slowly than
18 positive controls. NSAID had no significant effect on recovery.

19 **Conclusions and Clinical Importance:** If foot trimming on day 1 or 6 of diagnosis was stopped
20 and parenteral antibacterials were used, then over 1 million sheep/annum lame with footrot in the
21 UK, would recover more rapidly with benefits to productivity. Globally, this figure would be
22 much higher.

23

1 **Introduction**

2 Footrot causes lameness in sheep throughout the world and is an important health, welfare and
3 economic concern. Over 90% of flocks in the UK have lame sheep, with a within flock
4 prevalence of ~ 10%^{1,2}. Warm, wet environmental conditions favour the spread of footrot³ and
5 the severity of clinical presentation. Virulent footrot presents clinically³ with separation of hoof
6 horn from the sensitive dermis with a characteristic smell⁴. It causes approximately 30%, and
7 interdigital dermatitis (often benign footrot⁵) 50 – 60%, of [lameness in sheep in the UK](#)². Benign
8 and virulent footrot are caused by *Dichelobacter nodosus*. Over 95% of isolates of *D. nodosus*
9 from the UK are virulent by laboratory tests⁵, irrespective of serogroup and clinical presentation.
10 Laboratory diagnosis is of limited value to diagnose individuals because culture and PCR give
11 false positive and negative [results](#). It is used to determine flock infection and control in countries
12 where there are control policies [whereas](#) elsewhere clinical presentation is used.

13 Treatments for footrot include one or more of: trimming hoof horn, parenteral antibacterials and
14 topical bactericide^{6,7}. The most popular treatment for footrot is trimming and applying a topical
15 bactericide, with more than 90% farmers in the UK using this treatment in 2000⁸ and 2004⁹.
16 Trimming removes excess horn and exposes footrot lesions to air^{7,10}; ‘letting air in’ was
17 recommended to combat *D. nodosus* because of its anaerobic nature^{6,11}. Some practitioners
18 propose severe trimming of the hoof horn at diagnosis¹², however, many have moved away from
19 this and recommend that hoof horn is trimmed carefully 5 days after treatment, when the lesions
20 have start to heal^{7,13}, there is no evidence for or against this latter recommendation. All foot
21 trimming runs the risk of damaging the sensitive dermis of the foot, causing pain and
22 granulomatous proliferations^{3,7}.

23 In the UK, two epidemiological studies led to the hypothesis that parenteral and topical
24 antibacterials were the most effective treatment for footrot^{8,14}. From clinical trials in Australia

1 and the UK ~ 90% sheep were cured after treatment with penicillin and streptomycin¹⁵,
2 lincomycin and lincospectin¹⁶, erythromycin¹⁷ and long acting oxytetracycline^{18,19}. However, in
3 these trials no observations were made on lameness, only lesions, and all sheep were examined
4 for the first time 4-5 weeks after treatment so there is no estimate of time to recovery. Foot
5 trimming was combined with all treatments so the impact of trimming on recovery was not
6 elucidated. Some farmers treat lame sheep with NSAID which have anti-inflammatory, anti-
7 pyretic and analgesic properties that provide symptomatic relief from inflammatory conditions²⁰,
8 ²¹; their efficacy has not been evaluated.

9 There is no clinical trial that compares all treatments under similar conditions. The aim of the
10 current study was to investigate the effects of long acting parenteral oxytetracycline, foot
11 trimming on the same day or day 6 of diagnosis and flunixin meglumine (NSAID) on the time to
12 recovery from lameness and foot lesions in sheep lame with footrot.

13

1 **Materials and Methods**

2 *Study design*

3 A randomised factorial-design clinical trial was conducted on a farm in England, with a history of
4 sheep with footrot between October and December 2007. The flock comprised 250, 9-month-old
5 ewes that were out-wintered and not pregnant. The farmer put three replicates of 18, 32 and 10
6 lame sheep into a field on 18/10/2007, 14/11/2007 and 5/12/2007 respectively. The lame sheep
7 were identified with an ear tag and their locomotion²² was scored by researchers. The feet were
8 inspected and sheep with footrot (separation of the hoof horn and a characteristic smell) were
9 recruited into the trial. The severity of lesions on each foot was recorded²³. These sheep were
10 lame for up to 2 weeks before they were recruited in the trial. Sheep lame for any other reason
11 were treated using recommended treatments⁷ and returned to the main flock.

12

13 *Allocation to treatment groups*

14 The 53 sheep (replicate 1=14, replicate 2=29, and replicate 3=10) were matched by maximum
15 lesion severity and randomly allocated to one of six treatments (Table 1) by selecting a coloured
16 ball from a bag. The treatments were antibacterial aerosol spray to all sheep and then [parenteral
17 administration of antibacterials on day 1 of diagnosis \(n=34\), NSAID on day 1 of diagnosis
18 \(n=8\), foot trimming on day 1 \(n=15\) and foot trimming on day 6 \(n=21\) of diagnosis in an
19 incomplete factorial design^{24, 25}](#). The experimental protocol was approved by the Home Office as
20 a comparison of currently used methods to treat footrot and therefore ethically acceptable. The
21 researchers and farmer also discussed the treatments and the farmer approved them for his sheep.

22

23 The parenteral antibacterial was a long acting preparation of oxytetracycline^a (200mg/ml) at a
24 dose of 1ml/10kg bodyweight by deep intramuscular injection with a maximum dose of 5ml per

1 site. The aerosol spray was terramycin^b (oxytetracycline hydrochloride PhEur (3.93% w/w). The
2 NSAID used was flunixin meglumine BP^c at a dose of 2ml/45kg by intramuscular injection.
3 Sheep were diagnosed and treated by two researchers (SLSD and JLW) trained by JK. All foot
4 trimming was done by the farmer who was blinded to the treatment allocation. This farmer had
5 been sheep farming for 15 years and did not trim feet severely when he trimmed them
6 (observations of SLSD and JLW). All selected sheep on day 1 and only sheep still lame on day 6
7 of diagnosis allocated to be trimmed were foot trimmed.

8

9 *Follow up*

10 Sheep locomotion was scored²² and their feet inspected and lesions scored on three occasions
11 each week for up to 15 days. All sheep not returning to locomotion score zero by 15 days were
12 inspected and those with footrot or interdigital dermatitis were treated with parenteral
13 antibacterials and antibacterial spray.

14

15 *Statistical analysis*

16 *Outcome and model set up*

17 Data were stored in Microsoft Access. The outcomes of interest were time to recovery from
18 lameness and time to recovery from footrot lesions. A sheep was defined as recovered from
19 lameness when it had a locomotion score of zero for two consecutive observations. Recovery
20 from footrot lesions occurred when the lesion had healed and there was no foul smell or exudates.
21 Discrete time survival analysis²⁶ modelling hazard probability for recovery was used to analyse
22 the data^d. For both lameness and lesions the time to recovery was divided into three discrete-time
23 periods 1-5 days (T₁), 6-10 days (T₂) and 11-15 days (T₃) and sheep either recovered or not in
24 each time period. The hazard probability is the conditional probability that an event occurs in a

1 particular time period, given that it has not occurred in the previous time period (s) and is
2 described by an odds ratio.

3 Data were converted into sheep-period format with a separate record for each discrete time that a
4 sheep had a probability of recovering. For each of these time periods a binary variable was used
5 to indicate whether recovery occurred or not. Sheep were excluded from all subsequent time
6 periods once they had recovered and those not recovered by 15 days were right censored.

7

8 The model took the form²⁶:

$$9 \text{ logit } h(T_{it}) = [\alpha_1 T_{1it} + \alpha_2 T_{2it} + \alpha_3 T_{3it}] + [\beta_1 X_{1it} + \beta_2 X_{2it} + \dots + \beta_p X_{pit}]$$

10

11 ***Predictor variables***

12 The predictor variables were discrete time periods (T_1, T_2, T_3), did or did not receive [parenteral](#)
13 [administration of antibacterial or NSAID](#) on day 1 of diagnosis, the time dependent variable no
14 foot trimming, foot trim on day 1 or day 6 of diagnosis, with a positive control of receiving
15 topical antibacterials (Table 2), locomotion score at the start of the trial (locomotion score ≤ 2 or
16 3), number of feet affected with footrot (1 or 2), replicate (1, 2 or 3) and severity of footrot
17 lesions (score 1 or 2). Interactions between the variables were investigated. Observations that
18 clustered within sheep were included as a random effect.

19 The adjusted odds ratio (OR) (the hazard probability ratio) and associated 95% confidence
20 intervals were estimated. The model fit was evaluated by examining the deviance residuals at
21 sheep-period level and the sum of squared deviance residuals for each sheep's contribution to the
22 deviance statistic²⁶. The association between recovery from lameness and healing of footrot
23 lesions was assessed using a χ^2 test for each time period. The predicted probability for recovery
24 from lameness and footrot lesions in the discrete time period was plotted.

1 **Results**

2 On day 1 of diagnosis, 1 sheep had a locomotion score 1, 19 a score 2 and 33 had a locomotion
3 score 3. Sixty eight percent of sheep (36/53) had one foot affected with footrot and 32% (17) had
4 two feet affected and 60% (32) had a footrot severity score 1 and 40% (21) had a severity score 2.
5 The environmental conditions were fairly similar for each replicate; temperatures ranged from 7
6 to 11° C with a rainfall of 45 to 50 mm during the trial²⁷. Sheep were moved straight to pasture
7 after treatment, the weather was mostly dry for the 24 hours after treatment but moisture levels
8 due to dew are unknown. The proportion of sheep allocated to each treatment in each replicate
9 was similar (Table 2).

10
11 Approximately 51% (27/53) of sheep recovered from lameness during T₁, 55% (14/26) during T₂
12 and 58% (7/12) during T₃. The time to recovery from lameness and lesions was significantly
13 associated (p<0.05). Lameness and foot lesions resolved in the same time period except for 4 and
14 1 sheep that were still lame at the end of T₁ and T₂ respectively but had no lesions; these sheep
15 recovered from lameness in the following time period. Recovery rates were not significantly
16 different between the three discrete time periods (Table 3). Five sheep were still lame and had
17 lesions at the end of T₃; all had been foot trimmed, one had been treated with parenteral
18 antibacterials.

19
20 Sixty-five percent (22/34) of sheep that received antibacterials [administered parenterally](#)
21 recovered from lameness and 76% (26/34) from lesions in T₁. Nine of the remaining 12 sheep and
22 5 / 8 sheep with lesions, recovered from lameness and lesions respectively in T₂. Only 26%
23 (5/19) of sheep that did not receive [parenteral administration of antibacterials](#) recovered from
24 both lameness and lesions in T₁ and a further 36% (5/14) and 43% (6/14) recovered from

1 lameness and lesions respectively in T₂. Sheep that received [parenterally administered](#)
2 [antibacterials](#) were significantly more likely to have a faster recovery from both lameness (OR:
3 4.92 (1.20-20.10)) and lesions (OR: 5.11 (1.16-22.40)) than positive controls. There was no
4 significant effect of NSAID on the time to recovery from lameness (Table 3).

5
6 Four out of 15 trimmed sheep recovered from lameness and 5/15 from lesions in T₁. There were
7 12 sheep that were still lame by day 6 of diagnosis that were foot trimmed, 3/12 recovered from
8 both lameness and lesions in T₂. Sheep that were trimmed on day 1 (lameness - OR: 0.05 (0.005-
9 0.51); footrot lesions- OR: 0.06 (0.008-0.45)) or day 6 (lameness- OR: 0.07 (0.01-0.72); footrot
10 lesions- OR: 0.07 (0.01-0.56)) of diagnosis were significantly less likely to recover in the time
11 period than sheep that were not trimmed (Table 3).

12
13 The overall estimated hazard probability for recovery from both lameness and lesions within 5
14 days of treatment was highest in sheep that were treated with [parenterally administered](#)
15 [antibacterials](#) without foot trimming (Figure 1) followed by positive controls and sheep that
16 were foot trimmed and given [parenteral administration of antibacterials](#) and was lowest in sheep
17 that were foot trimmed and not given [parenteral administration of antibacterials](#).

18
19 Sheep with a locomotion score 3 took longer to recover from lameness (OR: 0.05 (0.01-0.24))
20 and lesions (OR: 0.04 (0.007-0.21)) than sheep with locomotion score ≤ 2 . There was no
21 significant effect of number of feet affected or severity of footrot lesions or replicate on the time
22 to recovery from lameness or lesions (Table 3).

23

1 There were no statistically significant interactions between treatments. The models provided a
2 reasonable fit to the data (Hosmer–Lemeshow statistic, lameness ($p=0.93$), lesions ($p=0.52$)).
3 There were no extreme observations with large residuals in the index plots of deviance residuals.

4 **Discussion**

5 In this trial sheep treated with [parenteral administration of long acting oxytetracycline](#) (together
6 with an oxytetracycline spray) were significantly more likely to recover from footrot lesions and
7 lameness within 5 days of treatment compared with sheep that were foot trimmed with or without
8 [parenteral administration of antibacterials](#) or positive controls (Table 3). The explanation for this
9 rapid response to treatment might be that *D. nodosus* is a bacterium susceptible to all antibacterial
10 classes and that the [antibacterials administered parenterally](#) penetrated deep into the dermis
11 where *D. nodosus* can be present ¹⁵. The results fit with the cure percents for [parenteral](#)
12 [administration of antibacterials](#) from other studies but we recorded time to recovery from both
13 lameness and lesions which was rapid. The results from the current study suggest that [parenteral](#)
14 [administration of antibacterials](#) is very effective (with 75% sheep recovering within 5 days) in
15 minimising the time that sheep are lame and thus treating all lame sheep with footrot with them
16 could not only improve the health and welfare of the sheep but could also minimise the effects of
17 [chronic lameness on loss of body condition and reduced productivity, such as lambs born and](#)
18 [lamb growth rate](#) ²⁸ .

19
20 Sheep in the current study were moved straight to pasture after treatment with parenteral
21 antibacterials. Previous evidence suggests that a dry environment for 24 hrs after parenteral
22 treatment is important ¹⁵ for improved efficacy of short acting parenteral antibacterials. Thus it is
23 possible that for long acting preparations with therapeutic levels present in serum for at least 72

1 hours ²⁹ , provision of a dry environment for up to 24 hrs is not be essential, especially
2 considering that the majority of sheep recovered within 5 days of treatment.

3 Foot trimming sheep lame with footrot without [administration of antibacterials parenterally](#) was
4 associated with the longest time to recovery from lameness and lesions both when sheep were
5 trimmed on day 1 or day 6 of diagnosis, suggesting that foot trimming lame sheep had a
6 detrimental effect on recovery whether the lesions were active or healing. For discrete time model
7 the odds of recovery from lameness or lesions within each 5 day period were similar for sheep
8 trimmed on day 1 or day 6 of diagnosis, suggesting that time of trimming did alter the time to
9 recovery differentially. In addition, lame sheep trimmed on day 6 of diagnosis were compared to
10 sheep still lame at day 6, avoiding selection bias. Trimmed sheep were lame and had lesions for
11 twice as long as the non-trimmed sheep, and presumably were infectious for twice as long, if we
12 assume that infectiousness is correlated with the presence of lesions. The delay in recovery
13 among trimmed sheep might be because trimming the hoof horn aggravated the foot and delayed
14 healing because of physical damage to the foot. The farmer did not trim feet severely and it is not
15 unreasonable to assume that his trimming practice is similar to other farmers who trim feet
16 without causing them to bleed. It is quite possible that had there been 'severe' trimming and
17 damage to the dermis that the time to recovery might have been longer.

18 The combination of antibacterials [administered parenterally](#), foot trimming and foot spraying had
19 a similar time to recovery from lameness and lesions as antibacterial spray alone, suggesting that
20 [parenteral administration of an antibacterial agent](#) negated the negative effect of foot trimming
21 but did not lead to as rapid a recovery from lameness as these antibacterials without foot
22 trimming. Quite remarkably, sheep that were not foot trimmed and did not receive parenteral
23 antibacterials, but only an oxytetracycline spray (as were all sheep in this study) recovered more
24 quickly than those foot trimmed and given topical spray (Figure 1), suggesting that foot trimming

1 cannot be recommended as a ‘second best’ treatment where farmers do not wish to use
2 antibacterials [administered parenterally](#) but rather it is detrimental to recovery from lameness and
3 lesions.

4 In the current study, sheep with a higher locomotion score at the start of the study took longer to
5 recover, suggesting an association between the severity of lameness and time to recovery from
6 footrot lesions and lameness. There was no significant association between the severity of the
7 initial lesion (even in univariate analysis) and the time to recovery from lesions or lameness, also
8 reported elsewhere ^{17, 19}. It is possible that the classification of under-run area (Table 3) was
9 insufficient to detect a difference or that the sheep rate of healing is such that severe lesions heal
10 as rapidly as mild lesions. A positive association between presence of lesions and lameness was
11 also observed in a longitudinal study⁹. We hypothesise that lesions cause lameness.
12 Consequently, sheep with old and chronic lesions and / or extensive damage to the wall horn
13 might take longer to recover from lameness if the lesions take longer to heal. However in an
14 intervention study²³ sheep lame for several months after treatment with parenteral antibacterials
15 recovered in median time of 4 days. This might suggest that given appropriate treatment the
16 propensity for the sheep foot to heal is great.

17 The lack of association between the number of feet with footrot and time to recovery from footrot
18 contrasts with a previous study ¹⁹ where sheep with 1 or 2 feet affected had higher cure rates than
19 sheep with 3 or 4 feet affected. However, because sheep had a maximum of two feet affected in
20 [the current study](#) it was not possible to investigate the relationship between more than this
21 number of affected feet and recovery.

22 NSAID did not change the time to recovery from lameness significantly, possibly because 8
23 sheep received NSAIDS or because they were given for only one day. A previous study ³⁰
24 suggested that after 3-days administration of flunixin meglumine noxious mechanical

1 stimulation to sheep with footrot responses were comparable to healthy sheep. NSAID do
2 produce analgesia for up to 24 hrs and so one injection might help ewes to continue to feed and
3 allow their lambs to feed, it certainly should not be discounted because of the results from the
4 current study.

5 No non-lame sheep were foot trimmed in this study and it is only possible to hypothesise on the
6 impact of routine foot trimming from other studies ^{8, 14, 31} where trimming was positively
7 correlated to lameness. Foot trimming might cause damage to the foot and spread or increase
8 susceptibility to *D. nodosus* or it might be indirectly causal if farmers trim rather than treat lame
9 sheep ³². Untreated sheep with footrot can remain lame for many months.

10 This study is an example of reduction and refinement [in use of animals in experiments](#). A
11 factorial design comparing the effects of several treatments individually and in combination used
12 a relatively small number of sheep (53) with sufficient power, possibly with the exception of
13 NSAID. In the absence of a gold standard laboratory test, sheep were diagnosed using
14 characteristic clinical presentation. The researchers were not blinded to treatment allocation
15 which could lead to bias, however, during follow up, [sheep number was used to identify](#)
16 [individual animals](#) and it seems unlikely that researchers remembered the identity of a sheep and
17 its treatment. The use of one flock and one environment gave consistency to compare the
18 effectiveness of the treatments and avoided confounding factors such as differences in breed, age,
19 micro and macro environment and other factors ¹⁹ that vary between farms. Further studies are
20 required to externally validate these findings and to investigate the effect of trimming feet of non-
21 lame sheep.

22

23

24

1 **Footnotes**

2 ^a Intervet/ Schering-Plough Animal Health, UK

3 ^b Pfizer Limited, UK

4 ^c Intervet/ Schering-Plough Animal Health, UK

5 ^d Stata 10.0, Statacorp, USA

6

7 **References**

8 1. Grogono-Thomas R, Johnston AM. A Study of Ovine Lameness. MAFF Final Report
9 MAFF Open Contract OC59 45K. DEFRA Publications, London; 1997.

10

11 2. Kaler J, Green LE. Naming and recognition of six foot lesions of sheep using written and
12 pictorial information: A study of 809 English sheep farmers. *Prev Vet Med* 2008; 83:52-64.

13

14 3. Egerton JR, Yong WK, Riffkin GG. Foot Rot and Foot Abscess of Ruminants, CRC
15 1989; 235-48.

16

17 4. Winter A. Lameness in sheep: 1, Diagnosis. *In Practice* 2004; 26:58–63.

18

19 5. Moore LJ, Wassink GJ, Green LE, Grogono-Thomas R. The detection and
20 characterisation of *Dichelobacter nodosus* from cases of ovine footrot in England and Wales.
21 *Veterinary Microbiology* 2005; 108:57–67.

22

23 6. Anonymous. Lameness in sheep, DEFRA publications, [ADMAIL 6000, London; 2003](#).

24

- 1 7. Winter A. Lameness in sheep 2. Treatment and control. In Practice 2004; 26:130-39.
- 2
- 3 8. Wassink GJ, Grogono-Thomas R, Moore LJ, Green LE. Risk factors associated with the
- 4 prevalence of footrot in sheep from 1999 to 2000. Veterinary Record 2003; 152:351-58.
- 5
- 6 9. Kaler J. Epidemiological investigations into lameness in sheep. PhD Thesis University of
- 7 Warwick 2008.
- 8
- 9 10. Abbott KA, Lewis CJ. Current approaches to the management of ovine footrot. The
- 10 Veterinary Journal 2005; 169: 28-41.
- 11
- 12 11. Bulgin MS, Lincoln SD, Lane VM, Matlock M. Comparison of treatment methods for the
- 13 control of contagious ovine foot rot. J Am Vet Med Assoc 1986; 189:194-6.
- 14
- 15 12. Duncanson G. Ovine Footrot: a closer look. Vet times 2009; VT37.09: 8-9.
- 16
- 17 13. Anonymous. Better returns from reducing flock lameness. EBLEX publications, [EBLEX](#),
- 18 [Graphic House, Ferrars Road, Huntingdon, Cambridgeshire; 2008.](#)
- 19
- 20 14. Green LE, Wassink GJ, Grogono-Thomas R, Moore LJ, Medley GF. Looking after the
- 21 individual to reduce disease in the flock: A binomial mixed effects model investigating the
- 22 impact of individual sheep management of footrot and interdigital dermatitis in a prospective
- 23 longitudinal study on one farm. Prev Vet Med 2007;78:172-78.
- 24

- 1 15. Egerton JR, Parsonson IM, Graham NP. Parenteral chemotherapy of ovine foot-rot.
2 Australian Veterinary Journal 1968; 44: 275-83.
3
- 4 16. Venning CM, Curtis MA, Egerton JR. Treatment of virulent footrot with lincomycin and
5 spectinomycin. Australian Veterinary Journal 1990; 67(7):258-60.
6
- 7 17. Ware JK, Scrivener CJ, Vizard AL. Efficacy of erythromycin compared with
8 penicillin/streptomycin for the treatment of virulent footrot in sheep. Australian Veterinary
9 Journal 1994; 71: 88-89.
10
- 11 18. Grogono-Thomas R, Wilsmore AJ, Simon AJ, Izzard KA. The use of long-acting
12 oxytetracycline for the treatment of ovine footrot. British Veterinary Journal 1994;150 :561-68.
13
- 14 19. Jordan D, Plant JW, Nicol HI, Jessep TM, Scrivener CJ. Factors associated with the
15 effectiveness of antibacterial treatment for ovine virulent footrot. Australian Veterinary Journal
16 1996; 73 :211-15.
17
- 18 20. Fitzpatrick J, Scott M, Nolan A. Assessment of pain and welfare in sheep. Small
19 Ruminant Research 2006; 62 (1-2 SPEC. ISS.): 55-61.
20
- 21 21. Flecknell P. Analgesia from a veterinary perspective. Br J Anaesth 2008; 101 :121-24.
22
- 23 22. Kaler J, Wassink GJ, Green LE. The inter- and intra-observer reliability of a locomotion
24 scoring scale for sheep. The Veterinary Journal 2009; 180(2):189-94.

- 1
- 2 23. Hawker EM. An intervention study to minimise footrot in sheep. MSc.University of
3 Warwick 2008.
- 4
- 5 24. Montgomery A, Peters T, Little P. Design, analysis and presentation of factorial
6 randomised controlled trials. *BMC Medical Research Methodology* 2003;3:26.
- 7 25. Chow SC, Liu JP. Design and analysis of clinical trials. New York: Wiley; 1998.
- 8
- 9 26. Singer JD, Willett JB. Applied Longitudinal Analysis: Modeling Change and Event
10 Occurrence. Oxford: Oxford University Press; 2003.
- 11
- 12 27. Anonymous. Pitsford Hall Weather Station, Northampton 2007. (Accessed 2008).
- 13
- 14 28. Marshall DJ, Walker RI, Cullis BR, Luff MF. The effect of footrot on body weight and
15 wool growth of sheep. *Australian Veterinary Journal* 1991; 68: 45–49.
- 16
- 17 29. Punch PI, Costa ND, Chambers ED, et al. Plasma and tear concentrations of antibacterials
18 administered parenterally to cattle. *Res Vet Sci* 1985; 39:179–187.
- 19
- 20 30. Welsh EM, Nolan AM. Effect of flunixin meglumine on the thresholds to mechanical
21 stimulation in healthy and lame sheep. *Res Vet Sci* 1995; 58: 61-66.
- 22

- 1 31. Kaler J, Green LE. Farmers' practices and factors associated with the prevalence of all
2 lameness and lameness attributed to interdigital dermatitis and footrot in sheep
3 flocks in England in 2004. *Prev Vet Med* 2009; 92: 52-59.
- 4 32. Kaler J, Green LE. Recognition of lameness and decisions to catch for inspection
5 among sheep farmers and specialists in GB. *BMC Veterinary Research* 2008; 4(1) :
6 41.
- 7
- 8
- 9

1 Table 1: Number of sheep and treatment given in the factorial design

	Parenteral antibacterial at day 1 of diagnosis	Foot trimming at day 1 of diagnosis	Foot trimming at day 6 of diagnosis	Non steroidal anti- inflammatory drug at day 1 of diagnosis
Treatment group 1 (n=9)	9	0	0	0
Treatment group 2 (n=8)	8	0	0	8
Treatment group 3 (n=7)	7	7	0	0
Treatment group 4 (n=10)	10	0	10*	0
Treatment group 5 (n=8)	0	8	0	0
Treatment group 6 (n=11)	0	0	11*	0
Total (n=53)	34	15	21	8

2 * 5 lame sheep from Treatment group 4 and 7 lame sheep from Treatment group 6 were trimmed
3 because 5 and 4 sheep from these groups recovered before day 6 respectively.

4

1 Table 2: Distribution of sheep in treatment groups by replicate and footrot severity

Treatment Group	Replicate 1		Replicate 2		Replicate 3		Footrot severity score 1		Footrot severity score 2	
	Number of sheep	Percent (%)	Number of sheep	Percent (%)	Number of sheep	Percent (%)	Number of sheep	Percent (%)	Number of sheep	Percent (%)
1	2	14.3	5	17.2	2	20.0	5	15.6	4	19.0
2	2	14.3	5	17.2	1	10.0	5	15.6	3	14.3
3	2	14.3	4	13.8	1	10.0	4	12.5	3	14.3
4	3	21.4	5	17.2	2	20.0	6	18.8	4	19.0
5	2	14.3	4	13.8	2	20.0	5	15.6	3	14.3
6	3	21.4	6	20.7	2	20.0	7	21.9	4	19.0
Total	14	100.0	29	100.0	10	100.0	32	100.0	21	100.0


2

1 Table 3: Multivariable discrete time survival analysis of time to recovery from lameness and
 2 footrot lesions of 53 sheep with footrot after treatment with combinations of parenteral long
 3 acting oxytetracycline at day 1 of diagnosis, NSAID at day 1 of diagnosis and foot trimming at
 4 day 1 or day 6 of diagnosis

Predictor Variable	Lameness				Footrot lesions			
	No. of sheep obs.	OR	95% C.I.	p-value	No. of sheep obs.	OR	95% C.I.	p-value
T ₃ (time period 3)	12	Ref.			11	Ref.		
T ₁ (time period 1)	53	0.28	0.04-1.98	0.20	53	0.15	0.02-1.31	0.09
T ₂ (time period 2)	26	1.71	0.26-11.04	0.57	22	0.71	0.08-6.11	0.75
Parenteral LAO at day 1 of diagnosis- No	42	Ref.			41	Ref.		
Yes	49	4.92	1.20-20.1	0.02	45	5.11	1.16-22.4	0.03
NSAID at day 1 of diagnosis-No	81	Ref.			78	^		
Yes	10	1.64	0.28-9.57	0.57	8			
Foot Trimming at day 1 of diagnosis . No	76	Ref.			71	Ref.		
Yes	15	0.05	0.005-0.51	0.003	15	0.06	0.008-0.45	0.006
Foot Trimming at day 6 of diagnosis- No	79	Ref.			74	Ref.		
Yes	12	0.07	0.01-0.72	0.009	12	0.07	0.01-0.56	0.01
Locomotion score at the start of the trial= ≤ 2#	26	Ref.			25	Ref.		
Locomotion score at the start of the trial= 3##	65	0.05	0.01-0.24	<0.001	61	0.04	0.007-0.21	<0.001
No. of feet affected =1	59	Ref.			56	Ref.		
No. of feet affected =2	32	0.84	0.24-2.93	0.79	30	0.84	0.26-2.65	0.77
Replicate 1	20	Ref.			20	Ref.		
Replicate 2	47	0.71	0.16-3.09	0.65	43	0.92	0.26-3.21	0.90
Replicate 3	24	0.43	0.08-2.05	0.28	23	0.67	0.18-2.51	0.56
Footrot severity score 1*	53	Ref.			50	Ref.		
Footrot severity score 2**	38	1.28	0.38-4.27	0.68	36	0.83	0.21-3.24	0.80

5 LAO, long acting oxytetracycline; OR, odds ratio; obs., observations; Ref., reference; # ,Visible nodding of head in
 6 time with short stride and/or uneven posture, shortened stride on one leg; ## ,Uneven posture, shortened stride on
 7 one leg, excessive flicking of head more than nodding in time with short stride; *,<50% of heel/sole/wall
 8 separation;** ≥50% and <100% heel/sole/wall separation; ^ coefficient could not be estimated because all sheep
 9 recovered in T₁

1 Figure 1: The fitted hazard probability of recovery from lameness and footrot lesions within 5
 2 days when treated with or without a parenteral antibacterial injection with no foot trimming or
 3 foot trimming at discrete times T1 and T2



4
 5
 6 Black bar: parenteral antibacterial and no foot trimming; **White bar with grid**: no parenteral
 7 antibacterial injection and no foot trimming; **White bar with cross hatches**: parenteral
 8 antibacterial injection and foot trimming; **Grey bar**: no parenteral antibacterial injection and foot
 9 trimming.
 10