



Seeking cyclonic activity records in speleothems from central Pacific: preliminary sample screening

Isabelle Couchoud, Samuel Etienne, Russell N. Drysdale, John C. Hellstrom,
Christoph Spötl, Yves Perrette

► To cite this version:

Isabelle Couchoud, Samuel Etienne, Russell N. Drysdale, John C. Hellstrom, Christoph Spötl, et al.. Seeking cyclonic activity records in speleothems from central Pacific: preliminary sample screening . Climate Change: The Karst Record VII (KR7), Sep 2014, Melbourne, Australia. 1 p., <10.1007/s00382-013-2035-y>. <hal-01261372>

HAL Id: hal-01261372

<https://hal.archives-ouvertes.fr/hal-01261372>

Submitted on 25 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Seeking cyclonic activity records in speleothems from central Pacific: preliminary sample screening


Isabelle Couchoud^{1,3}, Samuel Etienne², Russell Drysdale^{1,3}, John Hellstrom⁴, Christoph Spötl⁵, Yves Perrette¹

Figure 1. Atmospheric circulation in southern, central and western Pacific (ABOM-CSIRO, 2011).



Fig 2. Rurutu island and cave locations




Figure 3. Frequency of tropical cyclones between 1970 and 2009 in French Polynesia (Larrieu & Chiron 2010).




Figure 4. Records of daily min. (blue) and max. (red) temperatures on Rurutu island and in caves of Varirepo (green) and Tauapee (orange and pink). The hatched lines (light blue and orange) mark the min. and max. values reached in the caves during the year.




Figure 5. Rainfall (black) and drip rate under two stalactites (green and pink) in Tauapee cave, recorded every 20 min.

INTRODUCTION

The western half of Tropical Pacific is the planet's hot spot of cyclogenesis, with an average of 10 cyclones impacting the SW Pacific region per year. During El Niño years, cyclone activity migrates eastwards and affects the islands and populations of the central Pacific (Terry & Etienne, 2010). To evaluate the spatial and temporal evolution of this hazard in the context of global climate change, it is necessary to improve our knowledge about its natural evolution, on longer time-scale and at high resolution.

Speleothems can register, under given hydrological and geomorphological conditions, cyclonic events as abrupt variations of their oxygen isotope ratios or as mud-layers (e.g. Benoit Frappier et al., 2007, 2014; Nott et al., 2007).

Focusing on the Australes archipelago, a frequently hit region of southern Polynesia, this exploratory project aims at: 1) identifying speleothems capable of recording cyclones; 2) reconstructing a chronicle of cyclonic activity over the past few millennia.

SITE AND METHODS

We found caves hosting speleothems on the uplifted atoll of Rurutu (Fig.2), located in the eastern part of the South Pacific Convergence Zone (SPCZ; Fig.1). Activity and position of the SPCZ are essentially controlled by ENSO on annual scale, and by the IPO (Interdecadal Pacific Oscillation) on longer scales (Saltinger et al. 2014). Maximal cyclone occurrence is during El Niño periods (Wang et Fiedler, 2006). Rurutu has been hit every 7 to 10 years along the period 1970-2009 (Fig. 3).

In order to evaluate the capacity of the sampled speleothems to register cyclones hitting the island, we need to study the response of the hydrological system to meteorological variations, and the physico-chemical conditions of calcite precipitation. We will then identify among the sampled speleothems the most suitable specimens for detailed study according to their growth period and rate.

- Monitoring:
 - meteo station: Pluvimate + min-max thermometer + cumulative rain gauge
 - $\delta^{18}\text{O}$ and δD of daily rain water and monthly drip water + trace elements
 - buttons for cave temperature (3h interval) and Stalagmate for drip counting (20 min interval)
 - modern calcite for $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ comparison with drip water and cave temperature
- Speleothem analyses:
 - in caves, screening of candidates to provide useful proxy-records of recent cyclone activity
 - stratigraphy and petrography
 - $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ variations (increment 50-100 μm or 1mm; CF-IRMS Univ. of Innsbruck & Univ. de Savoie)
 - fluorescence imaging for layer counting (Univ. de Savoie)
 - dating by $^{230}\text{Th}/^{234}\text{U}$ (MC-ICP-MS Univ. of Melbourne) and ^{14}C (AMS ANSTO).

MONITORING DATA

Tauapee cave:
- temperature is stable over the year (-1 °C maximal amplitude);
- isotopic analyses of modern calcite and drip water provide calculated temperature consistent with the measured one.
=> The cave atmosphere is buffered and calcite precipitation occurs in near-equilibrium conditions.

Dripping:
- Dripping can get low but never stops, even during dry season.
- 2 drip counters show synchronous variations but differing in amplitude
- rainfall signal is buffered and modulated by the karst filter
- Two precipitation systems in the vadose zone, giving slow or fast responses to recharge events
=> The buffering effect of the vadose zone on Tauapee gives more regular drip rates, probably more mineralised waters leading to faster stalagmite growth rate, but more water mixing and smoother isotopic signals, inconvenient for targeting abrupt events.
- Drip water isotopic composition falls on the local meteoric water line => it still reflects rainwater composition.

Varirepo cave:
- Larger T_c amplitude (3.5 °C): the cave is more ventilated.
- Cooler T_c than in Tauapee due to its pit-shaped morphology (cold air trap)
- Modern calcite and dripwater provide a calculated T_c slightly inferior (-2 °C) to the mean temperature: maybe ventilation is not strong enough to generate strong kinetic fractionation and isotopic signal could still be exploitable.

SPELEOTHEM ANALYSES: PRELIMINARY RESULTS

Ideal cyclone-sensitive speleothems would have the following properties: active at the time of sampling and recording the past few millennia, with fast growth rate, made of clean primary calcite, precipitated without kinetic fractionation, and fed by water with a short residence time allowing for the isotopic signal of short events like cyclones to be transferred but long enough for the drip water to be supersaturated.

The screening for this type of speleothem is still in progress and the results presented here are very preliminary.


U-Th dating: low U concentration (9-40ng/g; host rock is reef limestone); significant detrital content, involving large corrections and large final age uncertainties.

Growth rate is highly variable, from ~0.12 mm/yr to ~0.03mm/yr.

Tau12A: $\delta^{18}\text{O}$ profile shows some quite large variations (~3‰), with several abrupt events that we will try to identify in other archives. $\delta^{13}\text{C}$ signal amplitude is very large (~4 to ~14‰) suggesting the effect of prior calcite precipitation. The $\delta^{18}\text{O}$ signal gets heavier when the $\delta^{13}\text{C}$ does, which could reflect the rainfall amount in this context.

Var12C: $\delta^{18}\text{O}$ profile shows values between -3.1 and -6.3‰, with decadal or centennial fluctuations, superimposed on a general trend which is consistent with the one of the $\delta^{13}\text{C}$ profile without being correlated, which implies that calcite precipitation could occur in conditions close to isotopic equilibrium. Once properly dated, this record should provide interesting information about past rainfall regimes on the island.

Var12-A: $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ profiles are very similar, signalling the effect of kinetic fractionation altering the record of the climate signal. Moreover, the $\delta^{13}\text{C}$ values are very high (+2 to -9‰) compared to the other stalagmites and to what would be expected under such soil. However, if kinetic processes are linked to the variations in drip rate or to the cave hydrogeology, they may inform us qualitatively about the effective rainfall through time. In such a case, we could focus on the longer term trends.



REFERENCES :

1. Terry J.P., Etienne S. (2010). Tempestuous times in the South Pacific islands. *Science*, 328, 5977, 428-429.
2. Benoit Frappier A. et al. (2007). Two millennia of tropical cyclone-induced mass loss in a northern Yucatan stalagmite: multiple overlapping climatic hazards during the Maya Terminal Classic "megadroughts". *Geophysical Research Letters*, 34, 5148-5157.
3. Benoit Frappier A. et al. (2007). Stalagmite stable isotope record of recent tropical cyclone events. *Geology*, 35(2), 111-114.
4. Nott J. et al. (2007). Growth frequency variability of landfalling tropical cyclones at centennial, century-long, seasonal and decadal scales. *Earth and Planetary Science Letters*, 255(3-4), 367-372.
5. Larrieu S. et Chiron T. (2010). Les îles de Polynésie française face à l'âge cyclonique. *Vertigo*, 10(3).
6. Samaniego M. et al. (2014). A new index for variations in the position of the South Pacific convergence zone 1910/11-2011/2012. *Climate Dynamics*, DOI 10.1007/s00338-013-2035-y.
7. Wang, C. & Fiedler, P. C. (2006). ENSO variability and the eastern tropical Pacific: a review. *Progress in Oceanography*, 69, p. 239-266.

PERSPECTIVES

Dating (in progress): more U-Th dates will be run on the oldest parts; ^{14}C dating will take place at ANSTO. It will focus on finding the "bomb pulse" and defining the dead carbon proportion for constraining calcite growth during recent centuries until it is viable to use U-Th dating. The potential for dating by laminae counting needs to be tested; petrographic changes are too faint so fluorescent laminae are being investigated.

Hydrologic monitoring indicates storage and mixing of the order of several weeks to months. Sampling is continuing in both caves to clarify this, as water transfer rate may change with the recharge, thus according to the season. This aspect needs consideration when looking for single abrupt events such as cyclones, as the signal of the ^{18}O -depleted rainwater may be masked unless the quantity is large enough to flush and replace the stored water.

Different drip sites in the same cave react differently to rainfall events. Water residence time and mineralization will change accordingly: some stalagmites grow faster than others, some stop seasonally while others grow steadily. Thus, some of these stalagmites will be more suitable to provide interannual climate variation records whilst others, despite a noisier signal, will be capable to inform us about seasonal evolution or even the occurrence of cyclones, if their growth rate is sufficient to allow the necessary sampling.

These stalagmites will also be useful to establish climate reconstructions of multi-annual resolution across the last millennium in the Australes archipelago, extending instrumental data about the hydrological variability in the Central Pacific linked to shifts in the SPCZ.