

University of Warwick institutional repository

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Suren Poghosyan and Daniel Ueltschi

Article Title: Abstract cluster expansion with applications to statistical mechanical systems

Year of publication: 2009

Link to published version: <http://dx.doi.org/10.1063/1.3124770>

Publisher statement: None

## Abstract cluster expansion with applications to statistical mechanical systems

Suren Poghosyan<sup>1,2,a)</sup> and Daniel Ueltschi<sup>1,2,b)</sup>

<sup>1</sup>*Institute of Mathematics, Armenian National Academy of Science, Marshal Bagramian 24-B, Yerevan 375019, Armenia*

<sup>2</sup>*Department of Mathematics, University of Warwick, Coventry CV4 7AL, England*

(Received 3 December 2008; accepted 2 April 2009; published online 8 May 2009)

We formulate a general setting for the cluster expansion method and we discuss sufficient criteria for its convergence. We apply the results to systems of classical and quantum particles with stable interactions. © 2009 American Institute of Physics. [DOI: [10.1063/1.3124770](https://doi.org/10.1063/1.3124770)]

### I. INTRODUCTION

The method of cluster expansions was introduced in the 1930s in statistical mechanics in order to study gases of classical interacting particles. Its main achievement, from the point of view of physics, may be the derivation of the van der Waals equation of state for weakly interacting systems. The method was made rigorous by mathematical physicists in the 1960s, see Ref. 23 and references therein.

The method split afterwards. One branch involves continuous systems, with applications to classical systems,<sup>19,17,5</sup> quantum systems,<sup>10,11,21</sup> or quantum field theory.<sup>12,16,2,6</sup> The other branch involves polymer systems, i.e., discrete systems with additional internal structure.<sup>13,7,3,18,25,9,14</sup> An important step forward was the article of Kotecký and Preiss with its simplified setting and its elegant condition for the convergence of the cluster expansion.<sup>15</sup>

The methods for proving the convergence are diverse. Let us mention the study of Kirkwood–Salsburg equations that involves correlation functions (see Ref. 23 and references therein), the algebraic approach of Ruelle,<sup>23</sup> combinatorial approaches using tree identities,<sup>19,5,2,6</sup> and inductions for discrete systems.<sup>7,3,18</sup>

Important and useful surveys were written by Brydges,<sup>4</sup> Pfister,<sup>20</sup> and Abdesselam and Rivasseau.<sup>1</sup> Recent articles have been devoted to combinatorial aspects<sup>25,8,14</sup> and to weakening the assumptions.<sup>9,8,22</sup>

The method of cluster expansions applies when the objects do not interact much; this is the case when they are far apart (low density) or when interactions are weak. An extension of the criterion of Ref. 15 that takes into account these two aspects was proposed in Ref. 26; it applies to both discrete and continuous systems.

All abstract (i.e., general) approaches involve restrictions that correspond to repulsive interactions. Yet the old results for classical and quantum systems only assume *stable* interactions, that may include an attractive part. The aim of the present article is to propose a general approach that applies to discrete and continuous systems with repulsive or stable interactions. Our proof is split into several independent steps and this helps clarify the situation.

The setting and the results are presented in Sec. II. We consider applications to classical systems of particles in Sec. III, to polymer systems in Sec. IV, and to the quantum gas in Sec. V. A fundamental tree estimate is derived in Sec. VI, and the theorems of Sec. II are proved in Sec. VII.

<sup>a)</sup>Electronic mail: [suren.poghosyan@unicam.it](mailto:suren.poghosyan@unicam.it).

<sup>b)</sup>Electronic mail: [daniel@ueltschi.org](mailto:daniel@ueltschi.org). URL: <http://www.ueltschi.org>.

## II. CLUSTER EXPANSIONS

We consider a set  $\mathbb{X}$  whose elements may represent widely different objects—in the three applications considered in this article, an element  $x \in \mathbb{X}$  represents (i) the position of a classical particle, (ii) a polymer, i.e., a connected set of  $\mathbb{Z}^d$ , and (iii) a closed Brownian bridge. For the general abstract theory, we assume the structure of a measure space,  $(\mathbb{X}, \mathcal{X}, \mu)$ , with  $\mu$  a complex measure. We denote  $|\mu|$  the total variation (absolute value) of  $\mu$ . Let  $u$  and  $\zeta$  be complex measurable symmetric functions on  $\mathbb{X} \times \mathbb{X}$ , that are related by the equation

$$\zeta(x, y) = e^{-u(x, y)} - 1. \quad (2.1)$$

We allow the real part of  $u$  to take the value  $+\infty$ , in which case  $\zeta(x, y) = -1$ . In typical applications  $u(x, y)$  represents the interactions between  $x$  and  $y$ , and the value  $+\infty$  corresponds to a hard core repulsion. We define the “partition function” by

$$Z = \sum_{n \geq 0} \frac{1}{n!} \int d\mu(x_1) \cdots \int d\mu(x_n) \exp \left\{ - \sum_{1 \leq i < j \leq n} u(x_i, x_j) \right\}, \quad (2.2)$$

or, equivalently,

$$Z = \sum_{n \geq 0} \frac{1}{n!} \int d\mu(x_1) \cdots \int d\mu(x_n) \prod_{1 \leq i < j \leq n} (1 + \zeta(x_i, x_j)). \quad (2.3)$$

The term  $n=0$  of the sums is understood to be 1.

The main goal of cluster expansions is to express the partition function as the exponential of a convergent series of “cluster terms.” The main difficulty is to prove the convergence. We first assume that the potential  $u$  is stable.

**Assumption 1:** *There exists a non-negative function  $b$  on  $\mathbb{X}$  such that, for all  $n$  and almost all  $x_1, \dots, x_n \in \mathbb{X}$ ,*

$$\prod_{1 \leq i < j \leq n} |1 + \zeta(x_i, x_j)| \leq \prod_{i=1}^n e^{b(x_i)}.$$

In other words, we assume the lower bound

$$\sum_{1 \leq i < j \leq n} \operatorname{Re} u(x_i, x_j) \geq - \sum_{i=1}^n b(x_i). \quad (2.4)$$

When the function  $b$  is constant, this is the usual definition of stability. “Almost all” means that, for given  $n$ , the set of points where the condition fails has measure zero with respect to the product measure  $\otimes^n \mu$ . If  $\mathbb{X}$  is countable, the condition must be satisfied for all  $x_1, \dots, x_n$  such that  $\mu(x_i) \neq 0$ . The second condition deals with the strength of interactions.

**Assumption 2:** *There exists a non-negative function  $a$  on  $\mathbb{X}$  such that for almost all  $x \in \mathbb{X}$ ,*

$$\int d|\mu|(y) |\zeta(x, y)| e^{a(y)+2b(y)} \leq a(x).$$

In order to guess the correct form of  $a$ , one should consider the left side of the equation above with  $a(y) \equiv 0$ . The integral may depend on  $x$ ; a typical situation is that  $x$  is characterized by a length  $\ell(x)$ , which is a positive number, so that the left side is roughly proportional to  $\ell(x)$ . This suggests to try  $a(x) = c\ell(x)$ , and one can then optimize on the value of  $c$ .

We also consider an alternate criterion that involves  $u$  rather than  $\zeta$ . It is inspired by the recent work of Procacci.<sup>22</sup> Let

$$\bar{u}(x, y) = \begin{cases} u(x, y) & \text{if } \operatorname{Re} u(x, y) \neq \infty \\ 1 & \text{if } \operatorname{Re} u(x, y) = \infty. \end{cases} \quad (2.5)$$

**Assumption 2'**: There exists a non-negative function  $a$  on  $\mathbb{X}$  such that for almost all  $x \in \mathbb{X}$ ,

$$\int d|\mu|(y) |\bar{u}(x, y)| e^{a(y)+b(y)} \leq a(x).$$

For positive  $u$  we can take  $b(x) \equiv 0$  and since  $1 - e^{-u} \leq u$ , Assumption 2 is always better than Assumption 2'. We actually conjecture that, together with Assumption 1, a sufficient condition is

$$\int d|\mu|(y) \min(|\zeta(x, y)|, |u(x, y)|) e^{a(y)+b(y)} \leq a(x). \quad (2.6)$$

That is, it should be possible to combine the best of both assumptions. In this respect Assumption 2 is optimal in the case of positive potentials, and Assumption 2' is optimal in the case of hard core plus negative potentials.

We denote by  $\mathcal{G}_n$  the set of all graphs with  $n$  vertices (unoriented, no loops) and  $\mathcal{C}_n \subset \mathcal{G}_n$  the set of connected graphs with  $n$  vertices. We introduce the following combinatorial function on finite sequences  $(x_1, \dots, x_n)$  of elements of  $\mathbb{X}$ :

$$\varphi(x_1, \dots, x_n) = \begin{cases} 1 & \text{if } n = 1 \\ \frac{1}{n!} \sum_{G \in \mathcal{C}_n} \prod_{\{i, j\} \in G} \zeta(x_i, x_j) & \text{if } n \geq 2. \end{cases} \quad (2.7)$$

The product is over edges of  $G$ .

**Theorem 2.1 (Cluster expansions)**: Suppose that Assumptions 1 and 2, or 1 and 2', hold true. We also suppose that  $\int d|\mu|(y) e^{a(y)+2b(y)} < \infty$ . Then we have

$$Z = \exp \left\{ \sum_{n \geq 1} \int d\mu(x_1) \cdots d\mu(x_n) \varphi(x_1, \dots, x_n) \right\}. \quad (2.8)$$

The term in the exponential converges absolutely. Furthermore, for almost all  $x_1 \in \mathbb{X}$ , we have the following estimate:

$$\sum_{n \geq 2} n \int d|\mu|(x_2) \cdots \int d|\mu|(x_n) |\varphi(x_1, \dots, x_n)| \leq (e^{a(x_1)} - 1) e^{2b(x_1)}. \quad (2.9)$$

[Under Assumption 2', Eq. (2.9) holds with  $e^{b(x_1)}$  instead of  $e^{2b(x_1)}$ .] Let us turn to correlation functions. We only consider one-point and two-point correlation functions since these are the most useful and expressions become more transparent. We refer to Ref. 26 for more general functions. First, we define the *un-normalized one-point correlation function* by

$$Z(x_1) = \sum_{n \geq 1} \frac{1}{(n-1)!} \int d\mu(x_2) \cdots \int d\mu(x_n) \prod_{1 \leq i < j \leq n} (1 + \zeta(x_i, x_j)) \quad (2.10)$$

(the term  $n=1$  is 1 by definition). And we define the *un-normalized two-point correlation function* by

$$Z(x_1, x_2) = \sum_{n \geq 2} \frac{1}{(n-2)!} \int d\mu(x_3) \cdots \int d\mu(x_n) \prod_{1 \leq i < j \leq n} (1 + \zeta(x_i, x_j)) \quad (2.11)$$

[the term  $n=2$  is equal to  $1 + \zeta(x_1, x_2)$ ]. Notice that  $Z(x_1)$  can be viewed as a regular partition function, given by Eq. (2.2), but with the modified measure  $(1 + \zeta(x_1, x))\mu(x)$  instead of  $\mu(x)$ . The

normalized correlation functions are  $Z(x_1)/Z$  and  $Z(x_1, x_2)/Z$ . As is shown in Theorem 2.2, they can be expressed using the “cluster functions”

$$\hat{Z}(x_1) = \sum_{n \geq 1} n \int d\mu(x_2) \cdots \int d\mu(x_n) \varphi(x_1, \dots, x_n) \quad (2.12)$$

and

$$\hat{Z}(x_1, x_2) = \sum_{n \geq 2} n(n-1) \int d\mu(x_3) \cdots \int d\mu(x_n) \varphi(x_1, \dots, x_n). \quad (2.13)$$

Notice that  $|\hat{Z}(x_1)| \leq e^{a(x_1)+2b(x_1)}$  by (2.9).

**Theorem 2.2 (Correlation functions):** *Under the same assumptions as in Theorem 2.1, we have*

$$\frac{Z(x_1)}{Z} = \hat{Z}(x_1),$$

$$\frac{Z(x_1, x_2)}{Z} = \hat{Z}(x_1)\hat{Z}(x_2) + \hat{Z}(x_1, x_2).$$

In statistical mechanics, the relevant expression is the *truncated two-point correlation function*

$$\frac{Z(x_1, x_2)}{Z} - \frac{Z(x_1)Z(x_2)}{Z^2}.$$

When the cluster expansion converges, it is equal to  $\hat{Z}(x_1, x_2)$  by the theorem above. This function usually provides an order parameter for phase transitions and it is useful to estimate its decay properties.

**Theorem 2.3 (Decay of correlations):** *If Assumptions 1 and 2 hold true, we have for almost all  $x, y \in \mathbb{X}$ ,*

$$|\hat{Z}(x, y)| \leq e^{a(y)+2b(y)} \left[ |\zeta(x, y)| e^{a(x)+2b(x)} + \sum_{m \geq 1} \int d|\mu|(x_1) \cdots \int d|\mu|(x_m) \prod_{i=0}^m |\zeta(x_i, x_{i+1})| e^{a(x_i)+2b(x_i)} \right]$$

(with  $x_0 \equiv x$  and  $x_{m+1} \equiv y$ ). If Assumptions 1 and 2' hold true, we have the same bound but with  $|\bar{u}(\cdot, \cdot)|$  instead of  $|\zeta(\cdot, \cdot)|$  and  $e^{b(\cdot)}$  instead of  $e^{2b(\cdot)}$ .

In many applications the functions  $\zeta(x, y)$  and  $u(x, y)$  depend on the difference  $x-y$  (this assumes that  $\mathbb{X}$  has additional structure, namely, that of a group). The estimates for  $|\hat{Z}(x, y)|$  are given by convolutions. The theorems of this section are proved in Sec. VII.

### III. CLASSICAL GAS

We consider a gas of point particles that interact with a pair potential. We work in the grand-canonical ensemble where the parameters are the fugacity  $z$  and the inverse temperature  $\beta$  (both are real and positive numbers). The set  $\mathbb{X}$  is an open bounded subset of  $\mathbb{R}^d$  and  $\mu(x) = z dx$  with  $dx$  the Lebesgue measure. We actually write  $\Lambda = \mathbb{X}$  so as to have more traditional notation. The interaction is given by a function  $U: \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\infty\}$  which we take to be piecewise continuous;  $u(x, y) = \beta U(x-y)$ . We suppose that  $U$  is stable, i.e., that there exists a constant  $B \geq 0$  such that for any  $n$  and any  $x_1, \dots, x_n \in \mathbb{R}^d$ :

$$\sum_{1 \leq i < j \leq n} U(x_i - x_j) \geq -Bn. \tag{3.1}$$

Our Assumption 1 holds with  $b(x) \equiv \beta B$ . The system is translation invariant so all  $x \in \mathbb{R}^d$  are equivalent. The function of Assumptions 2 and 2' can then be taken to be a constant,  $a(x) \equiv a$ . We seek a condition that does not depend on the size of the system. Then integrals over  $y$  are on  $\mathbb{R}^d$  instead of  $\Lambda$ . By translation invariance we can take  $x=0$ .

Assumption 2 gives the condition

$$ze^{2\beta B} \int_{\mathbb{R}^d} |e^{-\beta U(y)} - 1| dy \leq ae^{-a}. \tag{3.2}$$

We obviously choose the constant  $a$  that maximizes the right side, which is  $a=1$ . This condition is the one in Ref. 23. Let us now assume that  $U$  consists of a hard core of radius  $r$  and that it is otherwise integrable. Again with  $a=1$ , Assumption 2' gives the condition

$$ze^{\beta B} \left[ |\mathbb{B}|r^d + \beta \int_{|y|>r} |U(y)| dy \right] \leq e^{-1}. \tag{3.3}$$

Here,  $|\mathbb{B}| = \pi^{d/2} / \Gamma(d/2 + 1)$  is the volume of the ball in  $d$  dimensions. This condition is often better than (3.2). Without hard core it is the one in Ref. 5. The domains of parameters where these conditions hold correspond to low fugacities and high temperatures.

The thermodynamic pressure is defined as the infinite volume limit of

$$p_\Lambda(\beta, z) = \frac{1}{|\Lambda|} \log Z. \tag{3.4}$$

Using Theorem 2.1, we have

$$p_\Lambda(\beta, z) = \frac{1}{|\Lambda|} \int_\Lambda dx_1 \left[ \sum_{n \geq 1} z^n \int_\Lambda dx_2 \cdots \int_\Lambda dx_n \varphi(x_1, \dots, x_n) \right]. \tag{3.5}$$

Consider now any sequence of increasing domains  $\Lambda_1 \subset \Lambda_2 \subset \dots$  such that  $\Lambda_n \rightarrow \mathbb{R}^d$ . Thanks to the estimate (2.9) and using translation invariance, we get

$$p(\beta, z) \equiv \lim_{n \rightarrow \infty} p_{\Lambda_n}(\beta, z) = \sum_{n \geq 1} z^n \int_{\mathbb{R}^d} dx_2 \cdots \int_{\mathbb{R}^d} dx_n \varphi(0, x_2, \dots, x_n). \tag{3.6}$$

(The term with  $n=1$  is equal to  $z$ .) This expression for the infinite volume pressure  $p(\beta, z)$  should be viewed as a convergent series of analytic functions of  $\beta, z$ . Then  $p(\beta, z)$  is analytic in  $\beta$  and  $z$  by Vitali's theorem and no phase transition takes place in the domain of parameters where the cluster expansion is convergent.

The truncated two-point correlation function  $\sigma(x)$  is given by  $\hat{Z}(0, x)$ . We consider only the case of Assumption 2 but a similar claim can be obtained with Assumption 2'. Let  $c(x)$  be a function that satisfies the triangle inequality. The estimate of Theorem 2.3 yields

$$e^{c(x)} \sigma(x) \leq e^{2+4\beta B} \sum_{m \geq 0} z^m e^{m+2\beta Bm} (e^{c(\cdot)} |e^{-\beta U(\cdot)} - 1|)^{*m}(x) \tag{3.7}$$

(with  $f^{*0} \equiv f$ ). Recall that  $\|f^{*n}\|_\infty \leq \|f\|_\infty \|f\|_1^{n-1}$ , and let

$$C_p = \|e^{c(\cdot)} |e^{-\beta U(\cdot)} - 1|\|_p. \tag{3.8}$$

Then we get

$$\sigma(x) \leq e^{-c(x)} e^{2+4\beta B} \frac{C_\infty}{C_1} (1 - ze^{1+2\beta B} C_1)^{-1}. \quad (3.9)$$

If the inequality (3.2) is strict, one can usually find a function  $c(x)$  that satisfies the triangle inequality and such that  $C_1 \leq (ze^{1+2\beta B})^{-1}$ ; the truncated two-point correlation function then decays faster than  $e^{-c(x)}$ .

#### IV. POLYMER SYSTEMS

Polymer systems are discrete, which is technically simpler, but they also have internal structure. The first application of cluster expansions to polymer systems is due to Gruber and Kunz.<sup>13</sup> Among the many articles devoted to this subject, let us mention Refs. 15, 7, and 9. The main goal of this section is to illustrate our setting; we therefore restrict ourselves to a specific model of polymers with both repulsive and attractive interactions.

Our space  $\mathbb{X}$  is the set of all finite connected subsets of  $\mathbb{Z}^d$ . The measure  $\mu$  is the counting measure multiplied by the *activity*  $z(x)$  (a function  $\mathbb{X} \rightarrow \mathbb{R}_+$ ). We choose  $z(x) = e^{-\gamma|x|}$  with  $\gamma > 0$ . The interaction is hard core when polymers overlap and it is attractive when they touch:

$$u(x, y) = \begin{cases} \infty & \text{if } x \cap y \neq \emptyset \\ -\eta c(x, y) & \text{if } x \cap y = \emptyset. \end{cases} \quad (4.1)$$

Here,  $c(x, y)$  is the number of “contacts” between  $x$  and  $y$ , i.e., the number of bonds between sites of  $x$  and  $y$ ;  $\eta > 0$  is a parameter. The interaction is zero when the distance between polymers is greater than 1.

The stability condition can be written as

$$\frac{1}{2} \sum_{i=1}^n \sum_{j \neq i} u(x_i, x_j) \geq - \sum_{i=1}^n b(x_i). \quad (4.2)$$

Only disjoint polymers need to be considered; the left side is infinite otherwise. The sum over  $j$  is always larger than  $-\eta$  times the number of bonds connecting  $x_j$  with its exterior. Thus we can take  $b(x) = \eta d|x|$ .

The function  $a$  in Assumption 2' grows like  $|x|$ , so it is natural to choose  $a(x) = a|x|$  for some constant  $a$ . A sufficient condition is that

$$\sum_{y, y \cap x \neq \emptyset} z(y) e^{a|y| + \eta d|y|} + \sum_{y, \text{dist}(x, y) = 1} \eta z(y) c(x, y) e^{a|y| + \eta d|y|} \leq a|x|. \quad (4.3)$$

We can bound  $\eta c(x, y)$  by  $2\eta d|y|$ . Summing over the sites of  $x$  and requiring that  $y$  contains the given site or comes at distance 1, we get

$$\sum_{y \geq 0} (1 + 2d\eta|y|) z(y) e^{a|y| + \eta d|y|} \leq a. \quad (4.4)$$

We used the fact that the activity is translation invariant. If  $x$  is a connected set, there exists a closed walk with nearest neighbor jumps whose support is  $x$ , and whose length is at most  $2(|x| - 1)$ . This can be seen by induction: knowing the walk for  $x$ , it is easy to construct one for  $x \cup \{i\}$ . The number of connected sets of cardinality  $n$  that contain the origin is therefore smaller than the number of walks of length  $2n - 3$  starting at the origin, which is equal to  $(2d)^{2n-3}$ . Then it suffices that

$$\sum_{n \geq 1} e^{-n(\gamma - a - 3d\eta - 2 \log 2d)} \leq (2d)^3 a. \quad (4.5)$$

This is equivalent to

$$\gamma \geq a + \log\left(1 + \frac{1}{(2d)^3 a}\right) + 3d\eta + 2 \log 2d. \quad (4.6)$$

Assumption 2' holds for any  $a$ . Using  $\log(1+t) \leq t$  and optimizing on  $a$ , we find the sufficient condition

$$\gamma \geq 2(2d)^{-3/2} + 3d\eta + 2 \log 2d \quad (4.7)$$

with  $a = (2d)^{-3/2}$ .

We have just established the existence of a low density phase provided the activity is small enough. The condition depends on the contact parameter  $\eta$ . For large  $\eta$  one should expect interesting phases with many contacts between the polymers.

## V. QUANTUM GAS

We follow a course that is similar to Ginibre<sup>11</sup> using the Feynman–Kac formula so as to get a gas of winding Brownian loops. We get comparable results, with a larger domain of convergence in the case of integrable potentials. Winding Brownian loops are kind of continuous polymers; they combine the difficulties of both cases above—the continuous nature and the internal structure.

### A. Feynman–Kac representation

The state space for  $N$  fermions (bosons) in a domain  $\Lambda \subset \mathbb{R}^d$  is the Hilbert space  $L^2_{\text{anti}}(\Lambda^N)$  [ $L^2_{\text{sym}}(\Lambda^N)$ ] of square-integrable complex functions that are antisymmetric (symmetric) with respect to their arguments. The Hamiltonian is

$$H = - \sum_{i=1}^N \Delta_i + \sum_{1 \leq i < j \leq N} U(q_i - q_j), \quad (5.1)$$

with  $\Delta_i$  the Laplacian for the  $i$ th variable and  $U(q)$  a multiplication operator. As in the classical case, we consider the grand-canonical ensemble whose parameters are the fugacity  $z$  and the inverse temperature  $\beta$ . The partition function is

$$Z = \sum_{N \geq 0} z^N \text{Tr} e^{-\beta H}. \quad (5.2)$$

We need to cast the partition function in the form (2.2), which can be done using the Feynman–Kac representation. Namely, we have

$$Z = \sum_{N \geq 0} \frac{z^N}{N!} \sum_{\pi \in \mathcal{S}_N} \varepsilon(\pi) \int_{\Lambda^N} dq_1 \cdots dq_N \int dW_{q_1, q_{\pi(1)}}^{2\beta}(\omega_1) \cdots \int dW_{q_N, q_{\pi(N)}}^{2\beta}(\omega_N) \left( \prod_{i=1}^N \chi_{\Lambda}(\omega_i) \right) \exp \left\{ - \frac{1}{2} \sum_{1 \leq i < j \leq N} \int_0^{2\beta} U(\omega_i(s) - \omega_j(s)) ds \right\}. \quad (5.3)$$

$\mathcal{S}_N$  is the symmetric group of  $N$  elements;  $\varepsilon(\pi)$  is equal to the signature of the permutation  $\pi$  for fermions and  $\varepsilon(\pi) \equiv 1$  for bosons;  $W_{q, q'}^t$  is the Wiener measure for the Brownian bridge from  $q$  to  $q'$  in time  $t$ —the normalization is chosen so that

$$\int dW_{q, q'}^t(\omega) = (2\pi t)^{-d/2} e^{-|q - q'|^2/2t}; \quad (5.4)$$

$\chi_{\Lambda}(\omega)$  is 1 if  $\omega(s) \in \Lambda$  for all  $0 \leq s \leq 2\beta$ ; it is zero otherwise. An introduction to the Feynman–Kac formula in this context can be found in the survey of Ginibre.<sup>11</sup>



The right side of Eq. (5.3) is well defined for a large class of functions  $U: \mathbb{R}^d \rightarrow \mathbb{R} \cup \{\infty\}$ , that includes all piecewise continuous functions. Thus we take (5.3) as the definition for  $Z$ . Under additional assumptions on  $U$ , (5.3) is equal to (5.2) with Hamiltonian (5.1) and with Dirichlet boundary conditions.

We now rewrite the grand-canonical partition function in terms of winding loops. Let  $\mathbb{X}_k$  be the set of continuous paths  $[0, 2\beta k] \rightarrow \mathbb{R}^d$  that are closed. Its elements are denoted  $x = (q, k, \omega)$ , with  $q \in \mathbb{R}^d$  the starting point,  $k$  the winding number, and  $\omega$  the path; we have  $\omega(0) = \omega(2\beta k) = q$ . We consider the measure  $\mu$  given by

$$\mu(dx) = \frac{\varepsilon^{k+1} z^k}{k} dq e^{-v(x)} \chi_\Lambda(\omega) W_{q,q}^{2\beta k}(d\omega). \tag{5.5}$$

Here,  $v(x)$  is a self-interaction term that is defined below in (5.7);  $\varepsilon = -1$  for fermions and  $+1$  for bosons. Let  $\mathbb{X} = \cup_{k \geq 1} \mathbb{X}_k$ ; the measure  $\mu$  naturally extends to a measure on  $\mathbb{X}$ . The grand-canonical partition function can then be written as

$$Z = \sum_{n \geq 0} \frac{1}{n!} \int_{\mathbb{X}^n} d\mu(x_1) \cdots d\mu(x_n) \exp \left\{ - \sum_{1 \leq i < j \leq n} u(x_i, x_j) \right\}. \tag{5.6}$$

Let  $x = (q, k, \omega)$  and  $x' = (q', k', \omega')$ . The self-interaction  $v(x)$  and the two-loop interaction  $u(x, x')$  are given by

$$v(x) = \sum_{0 \leq \ell < m \leq k-1} \frac{1}{2} \int_0^{2\beta} U(\omega(s + 2\beta\ell) - \omega(s + 2\beta m)) ds,$$

$$u(x, x') = \sum_{\ell=0}^{k-1} \sum_{\ell'=0}^{k'-1} \frac{1}{2} \int_0^{2\beta} U(\omega(s + 2\beta\ell) - \omega'(s + 2\beta\ell')) ds. \tag{5.7}$$

We now treat separately the case of integrable potentials and the case of more general potentials.

**B. Stable integrable potentials**

We suppose that  $U$  is stable with constant  $B$ , i.e., it satisfies Eq. (3.1). For given loops  $x_1, \dots, x_n$ , stability implies that

$$\sum_{i=1}^n v(x_i) + \sum_{1 \leq i < j \leq n} u(x_i, x_j) \geq -\beta B \sum_{i=1}^n k_i. \tag{5.8}$$

Then Assumption 1 holds with

$$b(x) = \beta B k + v(x). \tag{5.9}$$

Notice that  $b(x) \geq 0$ , again by the stability of  $U$ .

We use Assumption 2' and we choose a function  $a(x) = ak$  with a constant  $a$  to be determined later. Explicitly, the assumption is that for any  $x = (q, k, \omega)$

$$\sum_{k' \geq 1} \frac{z^{k'} e^{ak'}}{k'} \int_{\mathbb{R}^d} dq' \int dW_{q',q'}^{2k'\beta}(\omega') e^{-v(x') + b(x')} |u(x, x')| \leq ak. \tag{5.10}$$

We have lifted the restriction that  $\omega'(s) \in \Lambda$  because we want a condition that does not depend on  $\Lambda$ . Equation (5.10) is easier to handle than its appearance suggests. Notice that the term in the second exponential is just  $\beta B k'$ . Using  $\int dW_{qq}^t(\omega) f(\omega) = \int dW_{00}^t(\omega) f(\omega + q)$  and the definition (5.7) of  $u(x, x')$ , it is enough that

$$\sum_{\ell=0}^{k-1} \frac{1}{2} \int_0^{2\beta} ds \sum_{k' \geq 1} \frac{z^{k'} e^{(a+\beta B)k'} k'^{-1}}{k'} \sum_{\ell'=0} \int_{\mathbb{R}^d} dq' \int dW_{0,0}^{2k'\beta}(\omega') |U(\omega(s+2\beta\ell) - \omega'(s+2\beta\ell') - q')| \leq ak. \tag{5.11}$$

We can immediately integrate over  $q'$ , which yields  $\|U\|_1$ . The Wiener integral then gives  $(4\pi k'\beta)^{-d/2}$  and we get the equivalent condition

$$\frac{\beta \|U\|_1}{(4\pi\beta)^{d/2}} \sum_{k' \geq 1} \frac{z^{k'} e^{(a+\beta B)k'}}{(k')^{d/2}} \leq a. \tag{5.12}$$

For any  $d, \beta, \|U\|_1$ , and  $a$ , the inequality holds for  $z$  small enough. Notice that  $z < 1$  in any case. One can get a more explicit condition for  $d \geq 3$  by choosing  $a$  such that  $ze^{a+\beta B} = 1$ . This yields

$$z \leq \exp \left\{ -\beta \left[ \frac{\|U\|_1 \zeta(d/2)}{(4\pi\beta)^{d/2}} + B \right] \right\}. \tag{5.13}$$

Here,  $\zeta(d/2) = \sum_{n \geq 1} n^{-d/2}$  is the Riemann zeta function.

When  $d=3$  and when the potential is repulsive, one can rewrite (5.13) in a more transparent way. Let  $a_0 = (1/8\pi)\|U\|_1$  denote the Born approximation to the scattering length. The condition is then

$$z \leq \exp \left\{ -\frac{\zeta(3/2) a_0}{\sqrt{\pi} \sqrt{\beta}} \right\}. \tag{5.14}$$

The critical fugacity is expected to be greater than 1. The present result helps nonetheless to obtain a range of densities where the pressure is analytic. In the bosonic case it compares well with physicists' expectations.<sup>24</sup>

### C. Stable potentials with hard core

The presence of a hard core makes the situation more complicated; we only sketch the argument in this section without trying to get explicit bounds. Our aim is to show that, using Theorem 2.1, the problem of convergence of the cluster expansion reduces to estimates of Wiener sausages.

We consider an interaction  $U = U' + U''$ . We assume that  $U' \geq 0$  is a repulsive potential of radius  $r$ , with a hard core of radius  $0 < r' \leq r$ , that  $U''(q) = 0$  for  $|q| < r$ , and that  $U''$  is integrable otherwise. We suppose that the stability condition takes the slightly stronger form

$$\sum_{i=1}^n U(q_0 - q_i) \geq -B \tag{5.15}$$

for any  $q_0, \dots, q_n$  such that  $|q_i - q_j| > r'$ . For potentials with a hard core this is equivalent to the property (3.1), possibly with a different constant  $B$ . Then one has<sup>10</sup>

$$\sum_{i=1}^n v(x_i) + 2 \sum_{1 \leq i < j \leq n} u(x_i, x_j) \geq -2\beta B \sum_{i=1}^n k_i. \tag{5.16}$$

Then Assumption 1 holds with

$$b(x) = \beta B k + \frac{1}{2} v(x). \tag{5.17}$$

Notice that the stability condition also holds with  $b$  given in (5.9) (and with a better constant  $B$ ). The advantage of (5.17) is the factor  $\frac{1}{2}$  in front of  $v(x)$ . Then  $e^{2b(x)}$  involves a term that cancels the self-interactions of  $x$ .

Given a loop  $x=(q, k, \omega)$ , let  $S(x)$  be the Wiener sausage generated by a ball of radius  $r$  when its center moves along the trajectory  $\omega$ :

$$S(x) = \{q \in \mathbb{R}^d : |\omega(s) - q| \leq r \text{ for some } s \in [0, 2k\beta]\}. \quad (5.18)$$

We also define the Wiener sausage that corresponds to the difference of two winding loops  $x=(q, k, \omega)$  and  $x'=(q', k', \omega')$ :

$$S(x, x') = \{q'' : |\omega(s + 2\beta\ell) - \omega'(s + 2\beta\ell') - q''| \leq r \text{ for some } \ell = 0, \dots, k-1; \ell' = 0, \dots, k'-1; s \in [0, 2\beta]\}. \quad (5.19)$$

We denote the volume of a Wiener sausage  $S(\cdot)$  by  $|S(\cdot)|$ . One can check that

$$|S(x, y)| \leq \frac{|S(x)||S(y)|}{r^d |\mathbb{B}|}, \quad (5.20)$$

with  $|\mathbb{B}|$  the volume of the unit ball (see Appendix 2 in Ref. 10).

We choose  $a(x) = |S(x)| + k$  in Assumption 2. Then a sufficient condition is that for any  $x \in \mathbb{X}$ ,

$$\sum_{k' \geq 1} \frac{z^{k'} e^{(2\beta B+1)k'}}{k'} \int_{\mathbb{R}^d} dq' \int dW_{0,0}^{2k'\beta}(\omega') |\zeta(x, x' + q')| e^{|S(x')|} \leq |S(x)| + k. \quad (5.21)$$

We consider separately the cases where  $q'$  belongs or not to  $S(x, x')$ . First,

$$\int_{S(x, x')} |\zeta(x, x' + q')| dq' \leq |S(x, x')|, \quad (5.22)$$

which we bound using (5.20). Second, using  $|e^{-u(x, x')} - 1| \leq |u(x, x')| e^{\beta B k'}$ ,

$$\int_{\mathbb{R}^d \setminus S(x, x')} |\zeta(x, x' + q')| dq' \leq \beta \|U''\|_1 k k' e^{\beta B k'}. \quad (5.23)$$

We certainly get (5.21) if we have the two inequalities

$$\begin{aligned} \sum_{k' \geq 1} \frac{z^{k'} e^{(2\beta B+1)k'}}{k'} \int dW_{0,0}^{2k'\beta}(\omega') e^{|S(x')|} |S(x')| &\leq r^d |\mathbb{B}|, \\ \sum_{k' \geq 1} z^{k'} e^{(3\beta B+1)k'} \int dW_{0,0}^{2k'\beta}(\omega') e^{|S(x')|} &\leq \frac{1}{\beta \|U''\|_1}. \end{aligned} \quad (5.24)$$

One can estimate the integrals of Wiener sausages (see Ref. 10), so that both conditions hold if  $z$  is small enough.

Now that the cluster expansion is known to converge, it is possible to write the pressure as an absolutely convergent series of analytic functions in  $\beta$  and  $z$ . It is also possible to study the decay of correlation functions. In the case of potentials that consist of hard core plus integrable part, one can apply Assumption 2' instead. This may give better results, especially if the integrable part is mostly attractive.

## VI. TREE ESTIMATES

In this section we obtain estimates of sums of connected graphs in terms of sums of trees. Our main result is Proposition 6.1 below. Such estimates seem to have been introduced by Penrose<sup>19</sup> and they have often been considered in the past.<sup>5,2,4,6,20,1,8</sup> We introduce a minimal setting that

clarifies its role in the cluster expansion. Namely, we fix the polymers so we only deal with the numbers that represent their interactions,  $\zeta$  or  $u$ , and the stability function  $b$ . Assumption 1 is vital here, but Assumptions 2 and 2' are not used in this section.

Let  $\mathcal{T}_n \subset \mathcal{C}_n$  denote the set of trees with  $n$  vertices. Let  $n$  be an integer,  $b_1, \dots, b_n$  be real non-negative numbers, and  $\zeta_{ij} = \zeta_{ji}$ ,  $1 \leq i, j \leq n$ , be complex numbers. We assume that the following bound holds for any subset  $I \subset \{1, \dots, n\}$ :

$$\prod_{i,j \in I, i < j} |1 + \zeta_{ij}| \leq \prod_{i \in I} e^{b_i}, \tag{6.1}$$

Let  $u_{ij}$  be such that  $\zeta_{ij} = e^{-u_{ij}} - 1$ ; let  $\bar{u}_{ij} = 1$  if  $\zeta_{ij} = -1$ , and  $\bar{u}_{ij} = u_{ij}$  otherwise. We state two distinct tree estimates, the first one involving  $|\zeta_{ij}|$  and the second one involving  $|\bar{u}_{ij}|$ . These bounds will allow to prove the convergence under either Assumption 2 or Assumption 2'.

**Proposition 6.1:** *If (6.1) holds true, we have the two bounds*

(a)

$$\left| \sum_{G \in \mathcal{C}_n} \prod_{\{i,j\} \in G} \zeta_{ij} \right| \leq \left( \prod_{i=1}^n e^{2b_i} \right) \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} |\zeta_{ij}|;$$

(b)

$$\left| \sum_{G \in \mathcal{C}_n} \prod_{\{i,j\} \in G} \zeta_{ij} \right| \leq \left( \prod_{i=1}^n e^{b_i} \right) \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} |\bar{u}_{ij}|.$$

We actually conjecture that the following estimate holds under the same hypotheses:

$$\left| \sum_{G \in \mathcal{C}_n} \prod_{\{i,j\} \in G} \zeta_{ij} \right| \leq \left( \prod_{i=1}^n e^{b_i} \right) \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} \min(|u_{ij}|, |\zeta_{ij}|). \tag{6.2}$$

We prove Proposition 6.1 (a) below using Ruelle's algebraic approach.<sup>23</sup> This method is usually combined with a Banach fixed point argument for correlation functions. However, we use it differently so as to get a tree estimate. Proposition 6.1 (b) follows from a tree identity due to Brydges, Battle, and Federbush,<sup>5,2,4</sup> combined with an argument due to Procacci;<sup>22</sup> its proof can be found at the end of this section.

Let  $\mathcal{A}$  be the set of complex functions on the power set  $\mathcal{P}(\{1, \dots, n\})$ . We introduce the following multiplication operation for  $f, g \in \mathcal{A}$ :

$$f * g(I) = \sum_{J \subset I} f(J)g(I \setminus J). \tag{6.3}$$

We use the standard conventions for sums and products, namely, that the empty sum is zero and the empty product is 1. Together with the addition,  $\mathcal{A}$  is a commutative algebra with unit  $1_{\mathcal{A}}(I) = \delta_{I,\emptyset}$ . It is possible to check that each  $f \in \mathcal{A} \setminus \{0\}$  has a unique inverse, which we denote  $f^{*(-1)}$ . We have

$$f^{*k}(I) = \sum_{\substack{J_1, \dots, J_k \subset I \\ J_i \cap J_j = \emptyset, \cup J_i = I}} f(J_1) \cdots f(J_k). \tag{6.4}$$

Let  $\mathcal{A}_0$  be the subset of functions  $f$  such that  $f(\emptyset) = 0$  ( $\mathcal{A}_0$  is an ideal of  $\mathcal{A}$ ). Notice that  $f^{*k} = 0$  for any  $k > n$  when  $f \in \mathcal{A}_0$ . We define the exponential mapping  $\exp_{\mathcal{A}}: \mathcal{A}_0 \rightarrow \mathcal{A}_0 + 1_{\mathcal{A}}$  by

$$\exp_{\mathcal{A}} f = 1_{\mathcal{A}} + f + \frac{1}{2}f^{*2} + \cdots + \frac{1}{n!}f^{*n}. \quad (6.5)$$

Let  $\Phi$  and  $\Psi$  be the functions defined by

$$\Phi(I) = \sum_{G \in \mathcal{C}(I)} \prod_{\{i,j\} \in G} \zeta_{ij},$$

$$\Psi(I) = \prod_{i,j \in I, i < j} (1 + \zeta_{ij}) = \sum_{G \in \mathcal{G}(I)} \prod_{\{i,j\} \in G} \zeta_{ij}. \quad (6.6)$$

Here,  $\mathcal{G}(I)$  [ $\mathcal{C}(I)$ ] is the set of graphs (connected graphs) on  $I$ . We have the relation

$$\Psi = \exp_{\mathcal{A}} \Phi. \quad (6.7)$$

We also introduce an operation that is reminiscent of differentiation:

$$D_I f(I) = \begin{cases} f(I \cup J) & \text{if } I \cap J = \emptyset \\ 0 & \text{otherwise.} \end{cases} \quad (6.8)$$

One can check that  $D_{\{i\}} \exp_{\mathcal{A}} f = (\exp_{\mathcal{A}} f) * D_{\{i\}} f$ .

For disjoint  $I, J \subset \{1, \dots, n\}$ , we define

$$g(I, J) = (\Psi^{*(-1)} * D_I \Psi)(J). \quad (6.9)$$

Let  $I \subset \{1, \dots, n\}$ . The assumption of Proposition 6.1 implies that

$$\prod_{i \in I} \prod_{j \in I \setminus \{i\}} |1 + \zeta_{ij}| \leq \prod_{i \in I} e^{2b_i}. \quad (6.10)$$

Then there exists  $i \in I$  such that

$$\prod_{j \in I \setminus \{i\}} |1 + \zeta_{ij}| \leq e^{2b_i}. \quad (6.11)$$

Such  $i$  is not unique in general but it does not matter. We consider a function  $\iota$  that assigns one of the indices  $i = \iota(I)$  above to each nonempty subset  $I \subset \{1, \dots, n\}$ . Notice that  $\iota(I) \in I$  for any subset  $I$ . It is also useful to introduce the notation  $I' = I \setminus \{\iota(I)\}$ .

**Lemma 6.2:** *The function  $g(I, J)$  of Eq. (6.9) is solution of the following equation:*

$$\begin{cases} g(\emptyset, J) = \delta_{\emptyset, J}, \\ g(I, J) = \left( \prod_{i \in I'} (1 + \zeta_{i, \iota(I)}) \right) \sum_{K \subset J} \left( \prod_{i \in K} \zeta_{i, \iota(I)} \right) g(I' \cup K, J \setminus K) \quad \text{if } I \neq \emptyset. \end{cases}$$

Since the equation gives  $g(I, J)$  in terms of  $g(K, L)$  with  $|K| + |L| = |I| + |J| - 1$ , it is well defined inductively and it has a unique solution. Notice that  $g(\emptyset, \emptyset) = 1$ , and that  $g(\{i\}, \emptyset) = 1$  for any index  $i$ .

*Proof:* Recall the definition (6.6) of  $\Psi$ . For disjoint  $I, K$  we have

$$\Psi(I \cup K) = \left( \prod_{j \in I' \cup K} (1 + \zeta_{j, \iota(I)}) \right) \Psi(I' \cup K) = \left( \prod_{j \in I'} (1 + \zeta_{j, \iota(I)}) \right) \left( \sum_{L \subset K} \prod_{k \in L} \zeta_{k, \iota(I)} \right) \Psi(I' \cup K). \quad (6.12)$$

Then

$$\begin{aligned}
 g(I, J) &= \sum_{K \subset J} \Psi^{*(-1)}(J \setminus K) \Psi(I \cup K) = \left( \prod_{j \in I'} (1 + \zeta_{j, u(I)}) \right) \sum_{L \subset K \subset J} \left( \prod_{k \in L} \zeta_{k, u(L)} \right) \Psi^{*(-1)}(J \setminus K) \Psi(I' \cup K) \\
 &= \left( \prod_{j \in I'} (1 + \zeta_{j, u(I)}) \right) \sum_{L \subset J} \left( \prod_{k \in L} \zeta_{k, u(L)} \right) \sum_{K' \subset J \setminus L} \Psi^{*(-1)}(J \setminus L \setminus K') \Psi(I' \cup L \cup K'). \tag{6.13}
 \end{aligned}$$

The last sum is equal to  $g(I' \cup L, J \setminus L)$ . One recognizes the equation of Lemma 6.2.  $\square$

We now estimate the function  $g$  using another function  $h$  that satisfies an equation that is similar to that of Lemma 6.2.

$$\begin{cases} h(\emptyset, J) = \delta_{\emptyset, J}, \\ h(I, J) = e^{2b, u(I)} \sum_{K \subset J} \left( \prod_{i \in K} |\zeta_{i, u(I)}| \right) h(I' \cup K, J \setminus K) \quad \text{if } I \neq \emptyset. \end{cases} \tag{6.14}$$

It also has a unique solution. Since  $\prod_{i \in I'} |1 + \zeta_{i, u(I)}| \leq e^{2b, u(I)}$ , we can check inductively that

$$|g(I, J)| \leq h(I, J) \tag{6.15}$$

for any sets  $I, J$  (with  $I \cap J = \emptyset$ ). Now the function  $h$  can be written explicitly.<sup>17,21</sup> Let  $\mathcal{F}_I(J)$  be the set of forests on  $I \cup J$  rooted in  $I$ . That is, a graph  $G \in \mathcal{F}_I(J)$  is a forest such that each tree contains exactly one element of  $I$ .

**Lemma 6.3:** *The solution of Eq. (6.14) is*

$$h(I, J) = \left( \prod_{i \in I \cup J} e^{2b_i} \right) \sum_{G \in \mathcal{F}_I(J)} \prod_{\{i, j\} \in G} |\zeta_{ij}|.$$

*Proof:* Since the solution of Eq. (6.14) is unique, it is enough to check that the ansatz of the lemma satisfies the equation. First, let us observe that both sides are multiplied by  $\prod_{i \in I \cup J} e^{2b_i}$ . Thus it is enough to consider the case  $b_i \equiv 0$ .

The sum over graphs in  $\mathcal{F}_I(J)$  can be realized by first summing over the set  $K$  of indices (necessarily in  $J$ ) that are connected to  $u(I)$  and then over sets of trees in  $J \setminus K$  and over connections to  $I' \cup K$ . Explicitly,

$$\sum_{G \in \mathcal{F}_I(J)} \prod_{\{i, j\}} |\zeta_{ij}| = \sum_{K \subset J} \left( \prod_{i \in K} |\zeta_{i, u(I)}| \right) \sum_{G \in \mathcal{F}_{I' \cup K}(J \setminus K)} \prod_{\{i, j\} \in G} |\zeta_{ij}|. \tag{6.16}$$

This equation is precisely (6.14).  $\square$

*Proof of Proposition 6.1, bound (a):* When  $I$  has a single element, the function  $g$  is equal to

$$g(\{1\}, \{2, \dots, n\}) = \sum_{G \in \mathcal{C}_n} \prod_{\{i, j\} \in G} \zeta_{ij}. \tag{6.17}$$

This is the left side of Proposition 6.1, bound (a). We have  $\mathcal{F}_{\{1\}}(\{2, \dots, n\}) = \mathcal{T}_n$ , the set of trees with  $n$  vertices. Thus  $h(\{1\}, \{2, \dots, n\})$  is equal to the right side of Proposition 6.1, bound (a), and the proof follows from Eq. (6.15).  $\square$

We now turn to the proof of Proposition 6.1, bound (b). Notice that in the absence of ‘‘hard cores,’’ i.e., when  $\zeta_{ij} \neq -1$ , our claim is just a reformulation of Corollary 3.2 (a) of Ref. 4. The present proof follows Ref. 22.

*Proof of Proposition 6.1, bound (b):* Let  $P$  be the set of  $\{i, j\}$  such that  $\zeta_{ij} = -1$ , i.e., such that  $\text{Re } u_{ij} = \infty$ . We regularize those numbers by setting

$$v_{ij}^{(m)} = \begin{cases} m & \text{if } \{i, j\} \in P \\ u_{ij} & \text{if } \{i, j\} \notin P. \end{cases} \tag{6.18}$$

This allows to use the tree identity of Refs. 5, 2, and 4; we will eventually take  $m$  to infinity. The tree identity can be written as

$$\sum_{G \in \mathcal{C}_n} \prod_{\{i,j\} \in G} (e^{-v_{ij}^{(m)}} - 1) = \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} (-v_{ij}^{(m)}) \int d\lambda_G(\{s_{ij}\}) e^{-\sum_{i < j} s_{ij} v_{ij}^{(m)}}. \tag{6.19}$$

The full definition of the measure  $\lambda_G$  can be found in Ref. 4; here we only mention its relevant properties.  $\lambda_G$  depends on the tree  $G$  and it is a probability measure supported on a subset  $L \subset \{s_{ij} \in [0, 1]: 1 \leq i < j \leq n\}$ . If  $v_{ij}^{(m)}$  satisfies the stability condition, then

$$\sum_{i,j \in I, i < j} s_{ij} \operatorname{Re} u_{ij} \geq - \sum_{i \in I} b_i \tag{6.20}$$

for all  $\{s_{ij}\} \in L$  and all  $I \subset \{1, \dots, n\}$ .

If  $\operatorname{Re} u_{ij} \neq \infty$  for any  $i, j$ , we immediately get the claim by using the stability condition above, since  $\int d\lambda_G(\{s_{ij}\}) = 1$ . The extension to possibly infinite numbers can be obtained using a trick due to Procacci.<sup>22</sup> Let  $G$  be a fixed tree and  $\varepsilon > 0$ . We introduce

$$w_{ij}^{(m,\varepsilon)} = \begin{cases} (1 - \varepsilon)m & \text{if } \{i,j\} \in G \cap P \\ \varepsilon & \text{if } \{i,j\} \in G \setminus P \\ 0 & \text{otherwise.} \end{cases} \tag{6.21}$$

Then

$$\sum_{1 \leq i < j \leq n} s_{ij} \operatorname{Re} v_{ij}^{(m)} \geq \sum_{1 \leq i < j \leq n} s_{ij} \operatorname{Re} v_{ij}^{(\varepsilon m)} + \sum_{1 \leq i < j \leq n} s_{ij} w_{ij}^{(m,\varepsilon)} - \varepsilon |G \setminus P|. \tag{6.22}$$

If  $m$  is large enough (depending on  $\varepsilon$ ),  $\{v_{ij}^{(\varepsilon m)}\}$  is stable and the first term of the right side is bounded below by  $-\sum b_i$ . Then

$$\begin{aligned} \left| \prod_{\{i,j\} \in G} (-v_{ij}^{(m)}) \int d\lambda_G(\{s_{ij}\}) e^{-\sum_{i < j} s_{ij} v_{ij}^{(m)}} \right| &\leq \left( \prod_{i=1}^n e^{b_i} \right) \left( \prod_{\{i,j\} \in G \cap P} \frac{w_{ij}^{(m,\varepsilon)}}{1 - \varepsilon} \right) \\ &\times \left( \prod_{\{i,j\} \in G \setminus P} \frac{w_{ij}^{(m,\varepsilon)} e^\varepsilon}{\varepsilon} |u_{ij}| \right) \int d\lambda_G(\{s_{ij}\}) e^{-\sum_{i < j} s_{ij} w_{ij}^{(m,\varepsilon)}}. \end{aligned} \tag{6.23}$$

A special case of the tree identity (6.19) is

$$\prod_{\{i,j\} \in G} w_{ij}^{(m,\varepsilon)} \int d\lambda_G(\{s_{ij}\}) e^{-\sum_{i < j} s_{ij} w_{ij}^{(m,\varepsilon)}} = \prod_{\{i,j\} \in G} (1 - e^{-w_{ij}^{(m,\varepsilon)}}) = (1 - e^{-(1-\varepsilon)m})^{|G \cap P|} (1 - e^{-\varepsilon})^{|G \setminus P|}. \tag{6.24}$$

We get

$$\left| \prod_{\{i,j\} \in G} (-v_{ij}^{(m)}) \int d\lambda_G(\{s_{ij}\}) e^{-\sum_{i < j} s_{ij} v_{ij}^{(m)}} \right| \leq \left( \prod_{i=1}^n e^{b_i} \right) \left( \frac{1 - e^{-(1-\varepsilon)m}}{1 - \varepsilon} \right)^{|G \cap P|} \left( \prod_{\{i,j\} \in G \setminus P} |u_{ij}| \frac{e^\varepsilon - 1}{\varepsilon} \right). \tag{6.25}$$

We can insert this estimate into Eq. (6.19). Letting  $m \rightarrow \infty$  and then  $\varepsilon \rightarrow 0$ , we get Proposition 6.1, bound (b). □

**VII. PROOFS OF THE THEOREMS**

In this section we prove the theorems of Sec. II. We consider only the case where Assumption 2 holds true—the case with Assumption 2' is entirely the same; one only needs to replace all  $|\zeta(x, y)|$  with  $|u(x, y)|$  and all  $e^{2b(\cdot)}$  with  $e^{b(\cdot)}$ . The proofs are based on the following tree estimate, which is a direct consequence of Proposition 6.1: for almost all  $x_1, \dots, x_n \in \mathbb{X}$ ,

$$|\varphi(x_1, \dots, x_n)| \leq \frac{1}{n!} \prod_{i=1}^n e^{2b(x_i)} \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} |\zeta(x_i, x_j)|. \tag{7.1}$$

*Proof of Theorem 2.1:* We start by proving the bound (2.9). Let us introduce

$$K_N(x_1) = \sum_{n=1}^N \frac{1}{(n-1)!} \int d|\mu|(x_2) \cdots \int d|\mu|(x_n) \prod_{i=1}^n e^{2b(x_i)} \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} |\zeta(x_i, x_j)|,$$

$$K(x) = \lim_{N \rightarrow \infty} K_N(x). \tag{7.2}$$

(The term  $n=1$  is equal to  $e^{2b(x_1)}$  by definition.) We show by induction that

$$K_N(x) \leq e^{a(x)+2b(x)} \tag{7.3}$$

for any  $N$ . Then  $K(x) \leq e^{a(x)+2b(x)}$  for almost all  $x$ , and using (7.1) we get (2.9).

The case  $N=1$  reduces to  $1 \leq e^{a(x)}$  and it is clear. The sum over trees with  $n$  vertices can be written as a sum over forests on  $\{2, \dots, n\}$  and a sum over edges between 1 and each tree of the forest. Explicitly,

$$K_N(x_1) = \sum_{n=1}^N \frac{1}{(n-1)!} \sum_{m \geq 1} \sum_{\substack{\{V_1, \dots, V_m\} \\ \text{partition of } \{2, \dots, n\}}} \int d|\mu|(x_2) \cdots \int d|\mu|(x_n) \\ \times \prod_{i=1}^n e^{2b(x_i)} \prod_{k=1}^m \left( \sum_{\ell \in V_k} |\zeta(x_1, x_\ell)| \sum_{G \in \mathcal{T}(V_k)} \prod_{\{i,j\} \in G} |\zeta(x_i, x_j)| \right). \tag{7.4}$$

Here,  $\mathcal{T}(V)$  denotes the set of trees with  $V$  as the set of vertices. If  $|V_k|=1$  the sum over  $G \in \mathcal{T}(V_k)$  is 1 by definition. The term after the sum over partitions depends on the cardinalities of the  $V_k$  but not on the actual labeling. Also, each  $\ell \in V_k$  gives the same contribution. We get

$$K_N(x_1) = e^{2b(x_1)} \sum_{n=1}^N \sum_{m \geq 1} \frac{1}{m!} \sum_{\substack{n_1, \dots, n_m \geq 1 \\ n_1 + \dots + n_m = n-1}} \prod_{k=1}^m \left( \frac{1}{(n_k-1)!} \int d|\mu|(y_1) \cdots \int d|\mu|(y_{n_k}) \right. \\ \left. \times |\zeta(x_1, y_1)| \prod_{i=1}^{n_k} e^{2b(y_i)} \sum_{G \in \mathcal{T}_{n_k}} \prod_{\{i,j\} \in G} |\zeta(y_i, y_j)| \right). \tag{7.5}$$

We obtain an upper bound by releasing the constraint  $n_1 + \dots + n_m \leq N-1$  to  $n_k \leq N-1, 1 \leq k \leq m$ . We then get

$$K_N(x_1) \leq e^{2b(x_1)} \exp \left\{ \sum_{n=1}^{N-1} \frac{1}{(n-1)!} \int d|\mu|(y_1) \cdots \int d|\mu|(y_n) |\zeta(x_1, y_1)| \prod_{i=1}^n e^{2b(y_i)} \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} |\zeta(y_i, y_j)| \right\} \\ = e^{2b(x_1)} \exp \left\{ \int d|\mu|(y_1) |\zeta(x_1, y_1)| K_{N-1}(y_1) \right\}. \tag{7.6}$$

We have  $K_{N-1}(y_1) \leq e^{a(y_1)+2b(y_1)}$  by the induction hypothesis. Equation (7.3) follows from Assumption 2.

The rest of the proof is standard combinatorics. The partition function can be expanded so as to recognize the exponential of connected graphs. Namely, we start with



$$Z = 1 + \sum_{n \geq 1} \frac{1}{n!} \int d\mu(x_1) \cdots \int d\mu(x_n) \sum_{G \in \mathcal{G}_n} \prod_{\{i,j\} \in G} \zeta(x_i, x_j). \quad (7.7)$$

The graph  $G \in \mathcal{G}_n$  can be decomposed into  $k$  connected graphs whose sets of vertices form a partition of  $\{1, \dots, n\}$ . Summing first over the number  $m_i$  of vertices for each set of the partition, we get

$$\begin{aligned} Z &= 1 + \sum_{n \geq 1} \sum_{k \geq 1} \frac{1}{k!} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} \frac{1}{m_1! \cdots m_k!} \prod_{\ell=1}^k \left\{ \int d\mu(x_1) \cdots \int d\mu(x_{m_\ell}) \sum_{G \in \mathcal{C}_{m_\ell}} \prod_{\{i,j\} \in G} \zeta(x_i, x_j) \right\} \\ &= 1 + \sum_{n \geq 1} \sum_{k \geq 1} \frac{1}{k!} \sum_{\substack{m_1, \dots, m_k \geq 1 \\ m_1 + \dots + m_k = n}} \prod_{\ell=1}^k \left\{ \int d\mu(x_1) \cdots \int d\mu(x_{m_\ell}) \varphi(x_1, \dots, x_{m_\ell}) \right\}. \end{aligned} \quad (7.8)$$

The triple sum is absolutely convergent, thanks to the estimate (2.9) that we have just established. One can then interchange the sums by the dominated convergence theorem. This removes the sum over  $n$ , and this completes the proof of Theorem 2.1.  $\square$

Next we prove Theorems 2.2 and 2.3 in reverse order, since we will use the convergence properties in the latter theorem to get the former.

*Proof of Theorem 2.3:* From the definition (2.13) and the tree estimate (7.1), we have

$$|\hat{Z}(x_1, x_2)| \leq \sum_{n \geq 2} \frac{1}{(n-2)!} \int d|\mu|(x_3) \cdots \int d|\mu|(x_n) \prod_{i=1}^n e^{2b(x_i)} \sum_{G \in \mathcal{T}_n} \prod_{\{i,j\} \in G} |\zeta(x_i, x_j)|. \quad (7.9)$$

The expression above involves a sum over trees of arbitrary size that connect 1 and 2. Any such tree decomposes into a line of  $m+1$  edges that connect 1 and 2 ( $m \geq 0$ ) and  $m+2$  trees rooted in the vertices of the connecting line. Taking into account the combinatorial factors, we obtain

$$|\hat{Z}(x, y)| \leq |\zeta(x, y)| K(x) K(y) + \sum_{m \geq 1} \int d|\mu|(x_1) \cdots \int d|\mu|(x_m) \left( \prod_{i=0}^m |\zeta(x_i, x_{i+1})| K(x_i) \right) K(y),$$

with  $x_0 \equiv x$  and  $x_{m+1} \equiv y$ . The result follows from the bound (7.3) for  $K$ .  $\square$

*Proof of Theorem 2.2:* It is actually similar to the end of the proof of Theorem 2.1.  $Z(x)$  can be expanded as a sum over graphs, that can be decomposed into a connected graph that contains 1, and other connected graphs. Taking into account the combinatorial factors, the contribution of connected graphs containing 1 yields  $\hat{Z}(x)$ , and the contribution of the others yields the expression (2.8) for  $Z$ . Thus  $Z(x) = \hat{Z}(x)Z$ . One step involved interchanging unbounded sums, which is justified because everything is absolutely convergent, thanks to (2.9) and Theorem 2.3.

In the graph expansion for  $Z(x, y)$ , the terms where 1 and 2 belong to the same connected graph yield  $\hat{Z}(x, y)Z$ , and the terms where 1 and 2 belong to different connected graphs yield  $\hat{Z}(x)\hat{Z}(y)Z$ . The detailed argument is the same as above. We then obtained the desired expression.  $\square$

## ACKNOWLEDGMENTS

We thank the referee for helpful comments. D.U. is grateful to Bill Faris, Roberto Fernández, Roman Kotecký, Charles-Édouard Pfister, Alan Sokal, and Miloš Zahradník for many useful discussions. He also acknowledges the hospitality of the Armenian Academy of Science, CNRS Marseille, the University of Geneva, ETH Zürich, the Center of Theoretical Studies of Prague, and the University of Arizona, where parts of this project were carried forward. D.U. is supported in part by Grant No. DMS-0601075 of the U.S. National Science Foundation.

- <sup>1</sup> Abdesselam, A. and Rivasseau, V., “Trees, forests, and jungles: a botanical garden for cluster expansions,” *Lect. Notes Phys.* **446**, 7 (1994); e-print arXiv:hep-th/9409094.
- <sup>2</sup> Battle, G. and Federbush, P., “A phase cell cluster expansion for Euclidean field theories,” *Ann. Phys.* **142**, 95 (1982).
- <sup>3</sup> Bovier, A. and Zahradník, M., “A simple approach to the problem of convergence of cluster expansions of polymer models,” *J. Stat. Phys.* **100**, 765 (2000).
- <sup>4</sup> Brydges, D. C., “A short course on cluster expansions,” in *Phénomènes critiques, systèmes aléatoires, théories de jauge, Les Houches*, 1984 (Elsevier/North Holland, Amsterdam, 1986), pp. 129–183.
- <sup>5</sup> Brydges, D. C. and Federbush, P., “A new form of the Mayer expansion in classical statistical mechanics,” *J. Math. Phys.* **19**, 2064 (1978).
- <sup>6</sup> Brydges, D. C. and Kennedy, T., “Mayer expansions and the Hamilton-Jacobi equation,” *J. Stat. Phys.* **48**, 19 (1987).
- <sup>7</sup> Dobrushin, R. L., “Estimates of semi-invariants for the Ising model at low temperatures,” in *Topics in Statistical and Theoretical Physics*, AMS Translational Series 2 (American Mathematical Society, Providence, RI, 1996), Vol. 177, pp. 59–81.
- <sup>8</sup> Faris, W. G., “A connected graph identity and convergence of cluster expansions,” *J. Math. Phys.* **49**, 113302 (2008).
- <sup>9</sup> Fernández, R. and Procacci, A., “Cluster expansion for abstract polymer models. New bounds from an old approach,” *Commun. Math. Phys.* **274**, 123 (2007).
- <sup>10</sup> Ginibre, J., “Reduced density matrices of quantum gases. III. Hard-core potentials,” *J. Math. Phys.* **6**, 1432 (1965).
- <sup>11</sup> Ginibre, J., “Some applications of functional integration in statistical mechanics,” in *Mécanique statistique et théorie quantique des champs, Les Houches*, 1970 (Gordon and Breach, New York, 1971), pp. 327–427.
- <sup>12</sup> Glimm, J., Jaffe, A., and Spencer, T., “The Wightman axioms and particle structure in the  $\mathcal{P}(\phi)_2$  quantum field model,” *Ann. Math.* **100**, 585 (1974).
- <sup>13</sup> Gruber, C. and Kunz, H., “General properties of polymer systems,” *Commun. Math. Phys.* **22**, 133 (1971).
- <sup>14</sup> Jackson, B., Procacci, A., and Sokal, A. D., “Complex zero-free regions at large  $|q|$  for multivariate Tutte polynomials (alias Potts-model partition functions) with general complex edge weights,” e-print arXiv:0810.4703.
- <sup>15</sup> Kotecký, R. and Preiss, D., “Cluster expansion for abstract polymer models,” *Commun. Math. Phys.* **103**, 491 (1986).
- <sup>16</sup> Malyshev, V. A., “Cluster expansions in lattice models of statistical physics and quantum field theory,” *Russ. Math. Surveys* **35**, 1 (1980).
- <sup>17</sup> Minlos, R. A. and Poghosyan, S., “Estimates of Ursell functions, group functions, and their derivatives,” *Theor. Math. Phys.* **31**, 408 (1977).
- <sup>18</sup> Miracle-Solé, S., “On the convergence of cluster expansions,” *Physica A* **279**, 244 (2000).
- <sup>19</sup> Penrose, O., *Statistical Mechanics, Foundations and Applications* (Benjamin, New York, 1967).
- <sup>20</sup> Pfister, Ch.-E., “Large deviations and phase separation in the two-dimensional Ising model,” *Helv. Phys. Acta* **64**, 953 (1991).
- <sup>21</sup> Poghosyan, S. and Zessin, H., “Decay of correlations of the Ginibre gas obeying Maxwell-Boltzmann statistics,” *Markov Processes Relat. Fields* **7**, 581 (2001).
- <sup>22</sup> Procacci, A., “Abstract polymer models with general pair interactions,” e-print arXiv:0707.0016 (version 2 of 26 November 2008).
- <sup>23</sup> Ruelle, D., *Statistical Mechanics: Rigorous Results* (World Scientific, Singapore, 1999).
- <sup>24</sup> Seiringer, R. and Ueltschi, D., “Rigorous upper bounds on the critical temperature of dilute Bose gases,” e-print arXiv:0904.0050.
- <sup>25</sup> Sokal, A. D., “Bounds on the complex zeros of (di)chromatic polynomials and Potts-model partition functions,” *Combinatorics, Probab. Comput.* **10**, 41 (2001).
- <sup>26</sup> Ueltschi, D., “Cluster expansions and correlation functions,” *Mosc. Math. J.* **4**, 511 (2004); e-print arXiv:math-ph/0304003.