

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Nazanin Haseli-Mashhadi, Tony Dadd, An Pan, Zhijie Yu, Xu Lin and Oscar H Franco

Article Title: Sleep quality in middle-aged and elderly Chinese: distribution, associated factors and associations with cardio-metabolic risk factors

Year of publication: 2009

Link to published version: <http://dx.doi.org/10.1186/1471-2458-9-130>

Publisher statement: None

Research article

Open Access

Sleep quality in middle-aged and elderly Chinese: distribution, associated factors and associations with cardio-metabolic risk factors

Nazanin Haseli-Mashhadi¹, Tony Dadd¹, An Pan², Zhijie Yu², Xu Lin² and Oscar H Franco*^{1,3}

Address: ¹Unilever R&D, Colworth Science Park, Sharnbrook, Bedfordshire, MK441LQ, UK, ²Key Laboratory of Nutrition and Metabolism, Institute for Nutritional Sciences, Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences and Graduate School of the Chinese Academy of Sciences, Shanghai, PR China and ³Health Sciences Research Institute, Warwick Medical School, The University of Warwick, Coventry, CV47AL, UK

Email: Nazanin Haseli-Mashhadi - hasseli57@yahoo.com; Tony Dadd - tony.dadd@unilever.com; An Pan - apan@sibs.ac.cn; Zhijie Yu - zjyu@sibs.ac.cn; Xu Lin - xlin@sibs.ac.cn; Oscar H Franco* - o.H.Franco@warwick.ac.uk

* Corresponding author

Published: 9 May 2009

Received: 22 December 2008

BMC Public Health 2009, 9:130 doi:10.1186/1471-2458-9-130

Accepted: 9 May 2009

This article is available from: <http://www.biomedcentral.com/1471-2458/9/130>

© 2009 Haseli-Mashhadi et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Poor sleep quality has been associated with increased risk of heart disease, diabetes and mortality. However, limited information exists on the distribution and determinants of sleep quality and its associations with cardio-metabolic risk factors in Chinese populations. We aimed to evaluate this in the current study.

Methods: A cross-sectional survey conducted in 2005 of 1,458 men and 1,831 women aged 50–70 years from urban and rural areas of Beijing and Shanghai. Using a questionnaire, sleep quality was measured in levels of well, common and poor. Comprehensive measures of socio-demographical and health factors and biomarkers of cardio-metabolic disease were recorded. These were evaluated in association with sleep quality using logistic regression models.

Results: Half of the population reported good sleep quality. After adjusting for potential confounders, women and Beijing residents had almost half the probability to report good sleep quality. Good physical and mental health (good levels of self-rated health (OR 2.48; 95%CI 2.08 to 2.96) and no depression (OR 4.05; 95%CI 3.12 to 5.26)) related to an increased chance of reporting good sleep quality, whereas short sleep duration (<7 hrs OR 0.10; 95%CI 0.07 to 0.14) decreased it substantially. There were significant associations between levels of sleep quality and concentrations of plasma insulin, total and LDL cholesterol, and index of insulin resistance.

Conclusion: Levels of good sleep quality in middle-age and elderly Chinese were low. Gender, geographical location, self-rated health, depression and sleep quantity were major factors associated with sleep quality. Prospective studies are required to distil the factors that determine sleep quality and the effects that sleep patterns exert on cardio-metabolic health.

Background

Sleep disturbances, including chronic insomnia (difficulty in maintaining sleep, having intermittent awakening and early awakening) and sleep apnea are increasingly common in modern societies with reported prevalences varying between 4 and 22 percent in different populations.[1] Sleep disturbances are common in later life, with loss of nocturnal sleep continuity and depth. [2-4] This could be due to the high prevalence of diseases accompanied by pain and stress, the side effects of taking medications and the inherent changes in sleep architecture that tend to occur with age. Aside from age and gender, physical and mental health problems are reported to be significantly associated with sleep disturbances. [5-8] Across different populations, several studies have found a significant deleterious effect of sleep disturbances on self-rated health, incidence of cardio-metabolic diseases (coronary heart disease, metabolic syndrome and type 2 diabetes), quality of life and mortality. [9-25] Various mechanisms have been explored and suggested as the potential link between poor sleep patterns and negative health outcomes (e.g. impaired glucose regulation and subsequent increased risk of type 2 diabetes[24]). Nevertheless the existing information is limited and the pathophysiological pathways between poor sleep patterns and cardio-metabolic disease remain unclear. Furthermore, although ample data exist for different populations, there is limited information on the levels of sleep disturbances among Chinese populations, the possible factors associated and the potential association of sleep quantity and quality with biomarkers of early cardio-metabolic disorders.

Therefore, in order to evaluate the levels of sleep patterns in Chinese and contribute to complement its related gap of epidemiological evidence and knowledge, in the current study we aimed to evaluate: (i) the distribution of sleep quality; (ii) factors that determine sleep quality; and (iii) whether self-reported sleep quality is associated with biomarkers of cardio-metabolic disease in middle-aged and elderly Chinese.

Methods

Study population

We used data from "Nutrition and Health of Aging Population in China" study. [26-28] This is a cross-sectional survey conducted between March and June 2005, among 3289 subjects (1,458 men and 1,831 women) aged 50 to 70 years from urban and rural areas of Beijing and Shanghai (metropolitan cities of north and south of China). The sampling was conducted with an emphasis on recruiting at least 40% of men, a similar number of rural/urban residents of Beijing and Shanghai and a representation of people from all levels of education and income. Trained physicians or public health workers completed the sampling through a face-to-face interview using a standard-

ized questionnaire to collect socio-demographical, lifestyle and health related information. Further details about the questionnaire can be found elsewhere.[29]

The study protocol was approved by the Institutional Review Board of the Institute for Nutritional Sciences. All participants provided written informed consents.

Assessment of sleep and related factors

Subjective sleep quality during the last month was recorded in 3 levels (well, common or poor). Sleep quantity was measured as self-reported average total hours of daily sleep during the last month, including both night and nap hours sleep and was categorized as under, normal and over (<7, 7 to 9 and >9 hours per day respectively).

Socio-demographic variables included age (50–59 and 60–70 yrs), gender, geographic region (Beijing or Shanghai), residence of living (urban or rural), educational attainment (low, moderate and high, based on the number of years of education as 0 to 6, 7 to 9, and ≥10 years respectively), total household income (low, moderate and high on the basis of less than 10,000, 10,000 to 29,999 and more than 30,000 Yuan annually), marital status (married or without spouse), employment status (grouped as currently employed, unemployed/on welfare system and retired) and social activity ('yes' or 'no' on the basis of participating in at least one of a list of social activities such as painting, playing chess, singing and dancing).

Lifestyle variables included smoking habit, defined as never, current and former (cessation of smoking >6 months); alcohol drinking (grouped into "yes" or "no") and physical activity level (low, moderate, or high) according to the IPAQ (International Physical Activity Questionnaire).

Self rated health (SRH) was initially recorded in 5 levels (very good, good, fair, poor and very poor) and then combined into two categories: good (very good and good) and poor (fair, poor and very poor).[30,31] Pulmonary disease was defined as existence of at least one of the following: asthma, chronic bronchitis, emphysema or chronic cor-pulmonale diseases. Presence of diabetes was defined as either measured fasting glucose greater than 7.0 mmol/L or being diagnosed with type 2 diabetes or taking anti-diabetes drugs or insulin. Presence of cardiovascular disease (CVD) was defined as one or more definite manifestations of coronary heart disease or stroke.

The Center for Epidemiologic Studies Depression Scale (CES-D)[32], which has been previously validated in Chinese populations[33,34], was used to measure the presence of clinically relevant depressive symptoms. Using the accepted cut-off point of 16 for the sum of scores, a CES-

D score of 0–15 was defined as no or minimal depression and a CES-D score ≥ 16 was defined as minor depression.[32] Following a physical examination, BMI levels were measured and categorized as under weight (<18.5 kg/m²), normal weight (18.5 to <24.0 kg/m²) and overweight/obese (≥ 24.0 kg/m²).[35] Three measurements of blood pressure were taken using an electronic blood pressure monitor and the mean of the last two measurements was used for analysis.

Laboratory methods

Overnight fasting blood samples were collected in tubes containing liquid ethylenediaminetetraacetic acid (EDTA), centrifuged at 4°C, and stored at -80°C until analysis. Total cholesterol, high-density lipoprotein (HDL) cholesterol, low-density lipoprotein (LDL) cholesterol, triglycerides (TG), and glucose were measured enzymatically (Hitachi 7080, Japan). Plasma C-reactive protein (CRP) was measured by a particle-enhanced immunoturbidimetric assay (Ultrasensitive CRP kit, Orion Diagnostica, Espoo, Finland). Plasma adiponectin concentrations were measured by Luminex xMAP™ Technology (Linco Research Inc, St Charles, Mo). All procedures followed the manufacture's instruction with quality controls in the expected ranges for each assay. Human Retinol Binding Protein 4 (RBP4) was measured by a sandwich ELISA developed in-house.[36] Interleukin 6 (IL-6) was measured with high-sensitivity enzyme-linked immuno-sorbent assay (ELISA) (Quantikine HS IL-6 Immunoassay, R&D Systems, Inc., Minneapolis, Minnesota). Plasma adiponectin, resistin and active plasminogen activator inhibitor-1 (PAI-1) concentrations were measured by Luminex xMAP™ Technology (Linco Research Inc, St Charles, Mo).[27] Fasting glucose was measured enzymatically on an automatic analyzer (Hitachi 7080, Japan). Fasting insulin was determined by radioimmunoassay (Linco Research, MO). Index of insulin resistance was calculated using updated homeostasis model assessment methods (HOMA2-IR, using the HOMA2 calculator, <http://www.dtu.ox.ac.uk>).[37]

Statistical analyses

We evaluated the distribution of sleep quality in the total population and how it varied by gender, geographic region and area of residence. Using a chi-square test, analysis of variance or Wilcoxon rank-sum test and univariate analysis, we evaluated the associations between sleep quality and potential associated factors (selected based on previous literature or scientific knowledge). Factors that significantly modified sleep quality or those that were consistently reported as significant determinants in previous publications were selected to be included in multivariate analysis applying an ordinal regression model. Multivariate analysis was repeated stratifying by gender,

region and residence one at a time and by stratifying for two of these factors simultaneously.

The associations between sleep quality and biomarkers of cardio-metabolic disease were tested on the total population and on a sub-sample of participants whom were free from CVD and diabetes with parametric t-tests (after logarithmic transformation of the data to satisfy normality assumptions) or non-parametric Wilcoxon rank-sum tests.

Data management and statistical analyses were performed with the SAS statistical package version 9.1 (SAS Institute, Cary, NC). Statistical tests were two-sided and at a 5% level of significance.

Results

Characteristics of the participants

The mean age of the study sample was 58.6 years (Table 1). More than half of the participants were women (55.7%), 72% were non-smokers (94% women and 44% men), 72% were non alcohol drinkers (90% women and 48% men), 92% were involved in moderate or high levels of physical activity and 53% were overweight or obese. Three percent of the participants reported taking sleeping pills (4% women and 1% men), 42% reported taking a nap (56% in Beijing, 29% in Shanghai), 68% reported having 7–9 hours of daily sleep and 53% reported good sleep quality (Table 1). Thirty two percent of participants reported good/excellent self-rated health, 9% of participants suffered from depressive symptoms (15% in Beijing and 4% in Shanghai), 14% had diabetes (19% in Beijing and 9% in Shanghai), 10% had CVD (15% in Beijing and 6% in Shanghai), 42% had metabolic syndrome (50% in Beijing and 35% in Shanghai), 14% had hyperlipidemia, 7% had pulmonary disease, 13% arthritis and 11% prostatism (Table 1).

Distribution of sleep quality

More than half of the study population (53%) reported good sleep quality (Beijing 46% vs. Shanghai 60%, women 47% vs. 61% men). Almost a third of the population (31%) reported a 'common' level of sleep quality (Beijing 34% vs. 27% Shanghai, women 33% vs. 28% men) and the rest (16%) reported poor sleep quality (19% Beijing vs. 13% Shanghai, women 20% vs. men 11%) (Table 1, data for gender difference not shown).

Eighty-eight percent of the participants who reported good sleep quality reported seven or more hours of daily sleep. In participants that reported poor sleep quality, 84% had fair, poor or very poor SRH. Three per cent of those with good sleep quality reported depressive symptoms, compared to 27% of those who reported poor sleep quality. Additionally, higher prevalence of CVD (20% vs.

Table 1: Characteristics of the study participants by region

Variable, N (%)	Total	Beijing	Shanghai	P-value
Female gender	1819 (55.71)	894 (54.91)	925 (56.51)	0.3599
Rural residence	1638 (50.17)	808 (49.63)	830 (50.70)	0.5405
Age, mean(std)	58.59 (6.01)	58.30 (5.95)	58.87 (6.06)	0.0072
Age category, years				0.0134
50 ≤ age < 60	1845 (56.51)	955 (58.66)	890 (54.37)	
60 ≤ age ≤ 70	1420 (43.49)	673 (41.34)	747 (45.63)	
BMI, mean(std)	24.46 (3.59)	25.26 (3.65)	23.67 (3.35)	<.0001
BMI category				<.0001
Under weight	111 (3.40)	37 (2.27)	74 (4.52)	
Normal	1411 (43.22)	570 (35.01)	841 (51.37)	
Over weight/Obese	1743 (53.38)	1021 (62.71)	722 (44.11)	
Currently married (YES, %)	2862 (87.66)	1407 (86.43)	1455 (88.88)	0.0328
Living alone (YES, %)	185 (5.67)	93 (5.72)	92 (5.62)	0.9056
Smoking				<.0001
Current	910 (27.87)	508 (31.20)	402 (24.56)	
Former	326 (9.98)	206 (12.65)	120 (7.33)	
Never	2029 (62.14)	914 (56.14)	1115 (68.11)	
Current alcohol drinker	930 (28.48)	602 (36.98)	328 (20.04)	<.0001
Physical activity, level				<.0001
Low	242 (7.41)	128 (7.86)	114 (6.96)	
Moderate	1368 (41.90)	581 (35.69)	787 (48.08)	
High	1655 (50.69)	919 (56.45)	736 (44.96)	
Educational level, years in school				<.0001
0–6	1348 (41.29)	562 (34.52)	786 (48.01)	
7–9	1163 (35.62)	698 (42.87)	465 (28.41)	
≥10	754 (23.09)	368 (22.60)	386 (23.58)	
Medical insurance (YES, %)	2280 (70.05)	872 (53.83)	1408 (86.12)	<.0001
Annual income, Yuan				<.0001
<10000	903 (29.07)	396 (25.13)	507 (33.14)	
10000–29999	1409 (45.36)	797 (50.57)	612 (40.00)	
≥30000	794 (25.56)	383 (24.30)	411 (26.86)	
Employment				<.0001
Employed	778 (23.83)	386 (23.71)	392 (23.95)	
Retired	1803 (55.22)	842 (51.72)	961 (58.70)	
Unemployed/on welfare	684 (20.95)	400 (24.57)	284 (17.35)	
Social activities (YES, %)	1661 (50.87)	848 (52.09)	813 (49.66)	0.1659
Self-rated health status (good, %)	1041 (31.90)	483 (29.69)	558 (34.11)	0.0067
Diabetes (YES, %)	446 (14.04)	292 (18.83)	154 (9.48)	<.0001
Metabolic syndrome (YES, %)	1392 (42.63)	815 (50.06)	577 (35.25)	<.0001
Hyperlipidemia (YES, %)	430 (13.78)	257 (17.11)	173 (10.69)	<.0001
Cardiovascular disease (YES, %)	327 (10.24)	228 (14.53)	99 (6.10)	<.0001
Pulmonary disease (YES, %)	238 (7.33)	118 (7.30)	120 (7.37)	0.9399
Arthritis (YES, %)	431 (13.41)	227 (14.29)	204 (12.55)	0.1458
Prostatism (YES, %)	150 (10.81)	77 (11.21)	73 (10.43)	0.6402
Depressive symptoms (YES, %)	307 (9.40)	241 (14.80)	66 (4.03)	<.0001
Sleeping pills intake (YES, %)	97 (2.97)	59 (3.63)	38 (2.32)	0.0281
Having nap (YES, %)	1382 (42.33)	911 (55.96)	471 (28.77)	<.0001
Sleep quality				<.0001
Well	1736 (53.17)	757 (46.50)	979 (59.80)	
Common	1001 (30.66)	558 (34.28)	443 (27.06)	
Poor	528 (16.17)	313 (19.23)	215 (13.13)	
Sleep quantity, total hours of sleep/day				<.0001
<7	799 (24.47)	346 (21.25)	453 (27.67)	
7–9	2208 (67.63)	1119 (68.73)	1089 (66.52)	
>9	258 (7.90)	163 (10.01)	95 (5.80)	

Abbreviations: SD, standard deviation; BMI, Body Mass Index; CI, Confidence interval.

*Data are given as frequency (percentage) unless otherwise indicated.

Table 2: Characteristics of the study participants by quality of sleep

Variable, N (%)	Total	Well	Common	Poor	P-value
Female gender	1819(55.71)	849(48.91)	603(60.24)	367(69.51)	<.0001
Beijing region	1628(49.86)	757(43.61)	558(55.74)	313(59.28)	<.0001
Rural residence	1638(50.17)	956(55.07)	420(41.96)	262(49.62)	<.0001
Age, mean(std)	58.59 (6.01)	58.61 (6.05)	58.69(6.00)	58.34(5.90)	0.5617
Age category, years					0.2195
50≤ age <60	1845(56.51)	975(56.16)	554(55.34)	316(59.85)	
60≤ age ≤70	1420(43.49)	761(43.84)	447(44.66)	212(40.15)	
BMI, mean(std)	24.46 (3.59)	24.33 (3.50)	24.66(3.68)	24.53(3.66)	0.056
BMI category					0.1373
Under weight	111 (3.40)	61 (3.51)	27 (2.70)	23 (4.36)	
Normal	1411(43.22)	776 (44.70)	423 (42.26)	212 (40.15)	
Over weight/Obese	1743(53.38)	889 (51.79)	551 (55.04)	293 (55.49)	
Currently married (YES, %)	2862(87.66)	1547(89.11)	858(85.71)	457(86.55)	0.0237
Living alone (YES, %)	185(5.67)	89(5.13)	63(6.29)	33(6.25)	0.3664
Smoking					<.0001
Current	910(27.87)	550(31.68)	241(24.08)	119(22.54)	
Former	326(9.98)	184(10.60)	98(9.79)	44(8.33)	
Never	2029(62.14)	1002(57.72)	662(66.13)	365(69.13)	
Current alcohol drinker	930(28.48)	524(30.18)	293(29.27)	113(21.40)	0.0004
Physical activity, level					0.0025
Low	242(7.41)	120(6.91)	74(7.39)	48(9.09)	
Moderate	1368(41.90)	681(39.23)	455(45.45)	232(43.94)	
High	1655(50.69)	935(53.86)	472(47.15)	248(46.97)	
Educational level, years in school					0.0026
0–6	1348(41.29)	762(43.89)	366(36.56)	220(41.67)	
7–9	1163(35.62)	599(34.50)	371(37.06)	193(36.55)	
≥10	754(23.09)	375(21.60)	264(26.37)	115(21.78)	
Medical insurance (YES, %)	2280(70.05)	1236(71.40)	711(71.24)	333(63.31)	0.0011
Annual income, Yuan					0.0116
<10000	903(29.07)	512(31.05)	239(24.82)	152(30.77)	
10000–29999	1409(45.36)	718(43.54)	469(48.70)	222(44.94)	
≥30000	794(25.56)	419(25.41)	255(26.48)	120(24.29)	
Employment					<.0001
Employed	778(23.83)	473(27.25)	207(20.68)	98(18.56)	
Retired	1803(55.22)	905(52.13)	598(59.74)	300(56.82)	
Unemployed/on welfare	684(20.95)	358(20.62)	196(19.58)	130(24.62)	
Social activities (YES, %)	1661(50.87)	910(52.42)	508(50.75)	243(46.02)	0.0362
Self-rated health status (good, %)	1041(31.90)	754(43.46)	203(20.30)	84(15.91)	<.0001
Diabetes (YES, %)	446(14.04)	211(12.48)	154(15.81)	81(15.85)	0.0255
Metabolic syndrome (YES, %)	1392(42.63)	694(39.98)	451(45.05)	247(46.78)	0.0038
Hyperlipidemia (YES, %)	430(13.78)	175(10.45)	169(17.81)	86(17.30)	<.0001
Cardiovascular disease (YES, %)	327(10.24)	122(7.17)	103(10.59)	102(19.69)	<.0001
Pulmonary disease (YES, %)	238(7.33)	111(6.43)	77(7.73)	50(9.52)	0.0501
Arthritis (YES, %)	431(13.41)	175(10.21)	144(14.69)	112(21.54)	<.0001
Prostatism (YES, %)	150(10.81)	66(7.78)	55(14.36)	29(18.59)	<.0001
Depressive symptoms (YES, %)	307(9.40)	49(2.82)	116(11.59)	142(26.89)	<.0001
Sleeping pills intake (YES, %)	97(2.97)	12(0.69)	21(2.10)	64(12.12)	<.0001
Having nap (YES, %)	1382(42.33)	739(42.57)	428(42.76)	215(40.72)	0.7131
Sleep quantity, total hours of sleep/day					<.0001
<7	799(24.47)	209(12.04)	273(27.27)	317(60.04)	
7–9	2208(67.63)	1348(77.65)	668(66.73)	192(36.36)	
>9	258(7.90)	179(10.31)	60(5.99)	19(3.60)	

Abbreviations: SD, standard deviation; BMI, Body Mass Index; CI, Confidence interval.

*Data are given as frequency (percentage) unless otherwise indicated.

7%), arthritis (21% vs. 10%) and prostatism (19% vs. 8%) was observed among those that reported poor sleep quality compared to those with good sleep quality (Table 2).

Sleep quality: associated factors

Based on the results of the univariate analyses -and previous publications-, gender, region, residential status, BMI, alcohol consumption and smoking, physical activity, educational level, marital status, medical insurance coverage, social activity, SRH, diabetes, CVD, arthritis, prostatism, depressive symptoms and sleep quantity were chosen to be analysed in the multivariate analysis (Table 3).

The results of the multivariate analysis indicated that gender (Odds Ratio (OR) 0.51; 95% Confidence Interval (CI) 0.41 to 0.64), region (OR 0.61; 95% CI 0.51 to 0.73), medical insurance coverage (OR 0.73; 95% CI 0.59 to

0.90), social activity (OR 0.80; 95% CI 0.69 to 0.94), residential status (OR 1.27; 95% CI 1.01 to 1.60), SRH (OR 2.48; 95% CI 2.08 to 2.96), depressive symptoms (OR 4.05; 95% CI 3.12 to 5.26), CVD (OR 4.05; 95% CI 3.12 to 5.26) and sleep duration (<7 hours daily sleep (OR 0.10; 95% CI 0.07 to 0.14), 7–9 hours daily sleep (OR 0.58; 95% CI 0.42 to 0.78)) affected the quality of sleep independently from the effect of other variables in the model (Table 3).

When we stratified the analyses by region (Beijing/Shanghai) we found that gender, depressive symptoms, self-rated health and sleep duration still remained significantly associated with sleep quality in both north and south China. Residential status, marital status, social activity and present CVD only affected the sleep quality of participants living in Shanghai, while arthritis only affected those living in Beijing (Table 3).

Table 3: Adjusted associated factors of sleep quality for the total population and stratified by regional area of living

Variable	Association with sleep quality (OR with 95% CI)		
	Total	Beijing	Shanghai
Gender (female vs. male)	0.51 (0.41–0.64)*	0.55 (0.40–0.75)*	0.50 (0.36–0.71)*
Residential location (rural vs. urban)	1.27 (1.01–1.60)*	1.08 (0.77–1.52)	1.66 (1.13–2.43)*
Geographic location (Beijing vs. Shanghai)	0.61 (0.51–0.73)*		
Body Mass Index (BMI)			
normal vs. over weight	0.92 (0.79–1.07)	1.05 (0.84–1.30)	0.81 (0.65–1.01)
under weight vs. over weight	0.74 (0.49–1.12)	0.35 (0.17–0.72)	1.03 (0.60–1.76)
Smoking			
current vs. never smoker	0.90 (0.71–1.14)	0.79 (0.58–1.08)	1.05 (0.72–1.55)
former vs. never smoker	0.98 (0.72–1.32)	1.15 (0.78–1.70)	0.69 (0.42–1.13)
Alcohol drinking (No vs. yes)	0.97 (0.80–1.18)	0.95 (0.74–1.23)	0.89 (0.65–1.22)
Physical activity			
high vs. low	1.09 (0.81–1.46)	0.93 (0.62–1.39)	1.26 (0.81–1.95)
moderate vs. low	0.94 (0.70–1.28)	0.81 (0.53–1.23)	1.07 (0.68–1.68)
Educational level, years in school			
group 1 vs. 3	1.24 (0.97–1.59)	1.38 (0.98–1.94)	0.91 (0.61–1.35)
group 2 vs. 3	1.06 (0.86–1.30)	1.13 (0.84–1.53)	0.93 (0.69–1.26)
Marital status (no spouse vs. spouse)	1.25 (1.00–1.56)*	1.14 (0.84–1.55)	1.52 (1.07–2.15)*
Medical insurance (no vs. yes)	0.73 (0.59–0.90)*	0.84 (0.60–1.16)	0.77 (0.55–1.07)
Employment			
employed vs. retired	1.05 (0.86–1.28)	1.07 (0.80–1.43)	0.96 (0.72–1.28)
unemployed vs. retired	1.00 (0.81–1.25)	0.80 (0.59–1.08)	1.30 (0.93–1.81)
Social activity (no vs. yes)	0.80 (0.69–0.94)*	0.83 (0.66–1.03)	0.80 (0.64–1.00)*
Self rated health status (good vs. not good)	2.48 (2.08–2.96)*	2.25 (1.76–2.88)*	2.63 (2.03–3.41)*
Diabetes (no vs. yes)	1.01 (0.82–1.26)	1.10 (0.84–1.44)	0.98 (0.68–1.41)
Cardiovascular disease (no vs. yes)	1.47 (1.15–1.88)*	1.25 (0.92–1.71)	2.14 (1.41–3.26)*
Arthritis (no vs. yes)	1.23 (0.99–1.53)	1.40 (1.04–1.89)*	1.11 (0.81–1.53)
Depressive Symptoms (no vs. yes)	4.05 (3.12–5.26)*	3.91 (2.88–5.29)*	5.00 (2.94–8.52)*
Total hours of sleep/day	*	*	*
<7 vs. 9< hrs	0.10 (0.07–0.14)*	0.10 (0.07–0.15)*	0.10 (0.05–0.17)*
7–9 vs. 9< hrs	0.58 (0.42–0.78)*	0.64 (0.44–0.92)*	0.53 (0.31–0.92)*

Abbreviations: CES-D, Center for Epidemiologic Studies Depression Scale; OR, Odds ratio; CI, Confidence interval. *P < 0.05. Group 1, 2 and 3 for educational level are 0–6, 7–9 and ≥10 years in school respectively.

After stratifying the analyses by gender and region simultaneously (Table 4), the effect of SRH, presence of depressive symptoms and sleep duration remained significant among all subgroups. Marital status, social activity, prostatism and residential status significantly affected Shanghai men and CVD significantly affected Beijing men and Shanghai women.

Sleep quality and cardio-metabolic biomarkers

Significant differences in the level of resistin, PAI-1, insulin, HOMA-IR, HOMA2-IR, LDL, TCH and TG were observed between different levels of sleep quality (Table 5). When we analyzed these associations among a subsample of the population who were free from CVD and diabetes, we found that significant associations remained for insulin, HOMA-IR, HOMA2-IR, LDL, TCH and TG (data not shown).

Discussion

Only half of middle aged and elderly Chinese in our study reported good sleep quality. Quality of sleep varied substantially by gender and geographical location and was affected by level of self-rated health, presence of depressive symptoms, sleep quantity, chronic diseases and social activity. Significant associations between levels of sleep quality and resistin, PAI-1, insulin, HOMA-IR, HOMA2-IR, LDL, TCH and TG were observed. These associations remained significant for insulin, HOMA-IR, HOMA2-IR, LDL and TCH among a subgroup of participants who were free of CVD and diabetes.

Sixteen percent of the elderly Chinese assessed their sleep quality as poor in this study. They showed higher prevalence of mental and physical diseases (depression, CVD, arthritis, prostatism (men only)) and shorter daily sleep time compared to participants that reported a good level

Table 4: Adjusted associated factors of sleep quality stratified by region and gender

Variable	Association with sleep quality (OR with 95% CI)			
	Beijing-Male	Shanghai-Male	Beijing-Female	Shanghai-Female
Residential location (rural vs. urban)	1.50 (0.92–2.45)	3.90 (2.06–7.39)*	0.68 (0.41–1.14)	0.86 (0.51–1.45)
Body Mass Index (BMI)				
normal vs. over weight	1.03 (0.73–1.47)	0.77 (0.53–1.11)	0.98 (0.73–1.32)	0.79 (0.60–1.05)
under weight vs. over weight	0.21 (0.08–0.57)*	1.27 (0.47–3.48)	0.42 (0.14–1.28)	0.91 (0.46–1.79)
Smoking				
current vs. never smoker	0.81 (0.52–1.27)	0.96 (0.62–1.48)	0.75 (0.47–1.22)	0.80 (0.24–2.66)
former vs. never smoker	1.16 (0.69–1.95)	0.63 (0.36–1.08)	1.06 (0.52–2.18)	0.17 (0.01–2.75)
Alcohol drinking (no vs. yes)	0.84 (0.59–1.19)	0.86 (0.58–1.26)	1.20 (0.81–1.77)	0.88 (0.44–1.75)
Physical activity				
high vs. low	0.99 (0.50–1.98)	2.09 (1.02–4.30)*	0.77 (0.46–1.30)	0.98 (0.55–1.75)
moderate vs. low	0.70 (0.34–1.42)	1.57 (0.75–3.31)	0.77 (0.45–1.32)	0.83 (0.45–1.50)
Educational level, years in school ‡				
group 1 vs. 3	1.70 (0.98–2.96)	0.74 (0.37–1.50)	1.26 (0.79–1.99)	0.91 (0.54–1.52)
group 2 vs. 3	1.05 (0.68–1.62)	0.82 (0.50–1.35)	1.18 (0.78–1.81)	0.82 (0.55–1.22)
Marital status (no spouse vs. spouse)	1.12 (0.64–1.96)	4.87 (2.12–11.19)*	1.30 (0.89–1.89)	1.08 (0.72–1.62)
Medical insurance (no vs. yes)	0.67 (0.42–1.09)	0.75 (0.44–1.28)	1.06 (0.66–1.71)	0.75 (0.48–1.16)
Employment				
employed vs. retired	0.82 (0.55–1.23)	1.10 (0.71–1.70)	1.34 (0.86–2.11)	0.93 (0.62–1.41)
unemployed vs. retired	0.77 (0.46–1.27)	1.29 (0.75–2.19)	0.96 (0.64–1.44)	1.68 (1.05–2.68)*
Social activity (no vs. yes)	0.86 (0.60–1.22)	0.64 (0.44–0.93)*	0.82 (0.61–1.10)	0.99 (0.74–1.33)
Self rated health status (good vs. not good)	2.18 (1.51–3.16)*	3.11 (2.00–4.83)*	2.36 (1.68–3.31)*	2.43 (1.75–3.39)*
Diabetes (no vs. yes)	1.05 (0.68–1.61)	0.78 (0.45–1.34)	1.15 (0.80–1.65)	1.11 (0.66–1.86)
CVD (no vs. yes)	1.78 (1.12–2.83)*	1.58 (0.85–2.96)	0.82 (0.53–1.25)	2.72 (1.47–5.04)*
Arthritis (no vs. yes)	1.74 (0.96–3.15)	0.56 (0.29–1.09)	1.35 (0.94–1.94)	1.31 (0.89–1.93)
Prostatism (no vs. yes)	1.17 (0.69–1.99)	1.72 (1.00–2.97)*		
Depressive Symptoms (no vs. yes)	3.13 (1.82–5.38)*	9.06 (3.50–23.49)*	4.34 (2.95–6.36)*	3.44 (1.77–6.66)*
Total hours of sleep/day	*	*	*	*
<7 vs. 9< hrs	0.12 (0.06–0.23)*	0.11 (0.04–0.29)*	0.08 (0.04–0.14)*	0.08 (0.04–0.16)*
7–9 vs. 9< hrs	0.66 (0.37–1.16)	0.61 (0.23–1.62)	0.64 (0.39–1.06)	0.45 (0.22–0.90)*

Abbreviations: CES-D, Center for Epidemiologic Studies Depression Scale; CVD, Cardio-Vascular Disease; OR, Odds ratio; CI, Confidence interval. * $p < 0.05$.

‡ Group 1, 2 and 3 for educational level are 0–6, 7–9 and ≥ 10 years in school respectively.

Table 5: Cardio-metabolic factors and biomarkers by levels of sleep quality

Variable, mean (SD)	Total	well	common	poor	p-value
Resistin	11.47(9.37)	11.26 (8.92)	11.24 (8.92)	12.57 (11.34)	0.0428*
PAI-I	14.74(18.17)	14.35 (18.71)	15.44 (17.56)	14.71 (17.50)	0.0011*
RBP4	40.11(11.74)	39.78 (11.59)	40.48 (11.58)	40.47 (12.48)	0.2924
Adiponectin	16.48(11.62)	16.47 (11.58)	16.06 (11.41)	17.29 (12.11)	0.1914
IL-6	1.55(2.78)	1.46 (2.13)	1.68 (3.63)	1.61 (2.84)	0.4680
CRP	1.58(4.30)	1.42 (2.29)	1.72 (5.54)	1.82 (6.24)	0.9505
Insulin	15.46(9.50)	15.00 (9.37)	16.07 (8.86)	15.81 (10.93)	<0.0001*
HOMA-IR	4.10(3.36)	3.95 (3.36)	4.32 (3.25)	4.14 (3.55)	<0.0001*
HOMA2-IR	0.31(0.22)	0.29 (0.19)	0.32 (0.27)	0.31 (0.22)	<0.0001*
SBP	140.11(22.49)	139.5 (22.06)	140.7 (22.94)	141.1 (22.99)	0.2338
DBP	80.16(10.79)	80.18 (10.74)	80.15 (10.82)	80.11 (10.92)	0.9892
HDL	1.28(0.33)	1.27 (0.33)	1.27 (0.33)	1.30 (0.36)	0.5415
LDL	3.26(0.97)	3.19 (0.94)	3.36 (0.99)	3.30 (1.01)	<0.0001*
TCH	4.70(0.98)	4.63 (0.97)	4.79 (0.97)	4.75 (1.02)	0.0001*
TG	1.39(1.07)	1.36 (1.07)	1.44 (1.12)	1.39 (0.96)	0.0490*

Abbreviations: PAI-I, plasminogen activator inhibitor-I; RBP4, retinol binding protein 4; IL-6, interleukin-6; IL6, interleukin-6; CRP, C-reactive protein; HOMA-IR, homeostasis model assessment of insulin resistance (fasting glucose (mmol/L) × fasting insulin (μU/ml)/22.5); SBP, systolic blood pressure; DBP, diastolic blood pressure; HDL, high-density lipoprotein; LDL, low-density lipoprotein; TCH, total cholesterol; TG, triglycerides; SD, standard deviation;

of sleep quality. We also observed significant gender differences in those who reported poor sleep quality wherein Chinese women were almost 50% less likely to report good sleep quality than men. Similar to our findings, Li et al.[6] found a 12% prevalence of insomnia among Hong Kong Chinese adults. Li et al.[6] studied 9851 Hong Kong Chinese aged 18–65 years and found a 60% greater chance of insomnia being reported among women than men. Klink et al.[5] also found there was a 50% higher chance of women reporting insomnia. Furthermore, in a study of 400 adults aged 20–70 years, Reyner et al.[7] found women, reported more awakenings, more total time spent awake during the night and poorer sleep quality. This difference in sleep quality could be due to gender differences in the prevalence of psychiatric morbidities, socio-cultural factors and coping strategies. [5-7]

We also found that sleep quality related to geographical location. Those living in the north were 40% less likely to report good sleep quality compared to those living in the south. This could be due to differences between the two cities wherein compared to Shanghai residents, Beijing residents show almost twice the prevalence of chronic diseases (depression, CVD, diabetes and metabolic syndrome), higher levels of alcohol consumption (37% Beijing vs. 20% Shanghai) and smoking (31% Beijing vs. 25% Shanghai), higher prevalence of nap taking (55% Beijing vs. 29% Shanghai) and less medical insurance coverage (53% Beijing vs. 86% Shanghai). To our knowledge this is the first study investigating differences in sleep quality as a function of geographical and residential status in China.

The effect of SRH (which is considered to be a valid representative of physical and mental health[38-40]) was substantial on sleep quality for the total population and for gender and geographical subgroups. Those who reported good/excellent SRH had twice the chance of reporting good sleep quality compared to those that had fair, poor or very poor SRH.

In our study, we also found a strong association between sleep quality and presence of depressive symptoms, with the strongest effect among Shanghai men and Beijing women. The likelihood of reporting poor sleep quality was three times higher in the presence of depressive symptoms. Our findings are consistent with other studies that have shown significant associations between depression and insomnia[41] and depression and sleep disturbances.[42] The association between poor sleep and depression has also been observed in an elderly population.[2]

Currently, the number of hours required for optimal functioning in an elderly population is not known. Although it is commonly believed that 8 hours of sleep per night is optimal for good health, recent studies suggest that mortality risk is lower among those sleeping 7 hours.[21] The total hours of daily sleep includes night time sleep and nap time (nap taking is common mainly in north of China) and for that reason, 7–9 hours was categorized as normal daily sleep hours in our study. We found a substantial effect of sleep quantity on quality of sleep. Having a short sleep time significantly decreased the chance of reporting good quality of sleep. Even those reporting 7–9

hours of daily sleep were less likely to report good sleep quality than those reporting more than 9 hours of daily sleep. Presence of CVD also affected the levels of sleep quality among the total population but when we stratified by gender and residential location this effect remained significant only among Beijing men and Shanghai women.

We did not find any significant association between sleep quality and different lifestyle characteristics (smoking and alcohol drinking habits), age and BMI. This is consistent with Elwood et al. [38] who studied 1986 Welsh men aged 55–69 y and found no significant association between prevalence of insomnia and age, BMI, smoking and alcohol drinking habits. However, it is possible that the effect of smoking and alcohol drinking in our analysis might be confounded with gender as the majority of Chinese smokers and drinkers were men. Furthermore, this is affected by the limited number of participants in the extreme categories of alcohol consumption and BMI, which limits the statistical power to identify a potential association between these two variables and sleep patterns in our population.

Despite lower socio-economic status (lower average income, fewer years in full-time education and less medical insurance coverage) rural residents were more likely to report good levels of sleep quality compare to urban residents. This could be due to differences in lifestyle wherein those who live in rural areas tend to have higher levels of physical activity, higher age of retirement, less stressful lifestyles and a lower prevalence of both CVD (7.7% rural vs. 13% urban) and diabetes (11.4% rural vs. 17% urban).

In our analyses of the total sample and subgroups of the sample, we found no significant effects of income and education. This could be due to the fact that residence is a more comprehensive factor that might partially confound the effect of socio-economic factors (i.e. our sample of the Chinese population is skewed towards lower income and education and late retirement in rural areas).

Concentrations of inflammatory biomarkers, plasma insulin and index of insulin resistance were associated with sleep quality for the total population and among a subgroup of participants that were free from CVD and diabetes. However, no clear trends in the levels of biomarkers as a function of the different levels of sleep quality (good/common/poor) were observed. This is the first study to evaluate the associations between sleep quality and inflammatory markers and measures of insulin resistance in China and to evaluate the distribution and factors associated with sleep quality in such a vast country which has many geographical and residential differences

Although we considered the effect of comprehensive measurements of lifestyle, socio-demographic factors and presence of disease, our findings were limited. This could be due to the cross-sectional nature of our data and further longitudinal studies should be carried out to confirm our findings.

Furthermore, sleep patterns were measured using a questionnaire; a criticism of this type of measurement is that answers are provided by the participants and can be affected by recall bias. Furthermore, due to the size of the population and the extent of the general questionnaire, for logistic reasons the questions that we used are not as thorough as alternative validated questionnaires designed to evaluate sleep disturbances (e.g The Pittsburg Sleep Quality Inventory). Nevertheless, these factors could only generate a non-differential misclassification of the exposure (sleep patterns) that would attenuate the size of the effects hereby reported and the level of sleep disturbances in the population studied, but not necessarily our results and conclusions. Prospective evaluations and studies that provide more sensitive measures of sleep patterns in Chinese populations are required to clarify the determinants that affect sleep quality. This will also allow us to assess whether sleep disturbances are associated with early deviations from adequate cardio-metabolic health.

Conclusion

In general, only half of the population reported good sleep quality. Compared to their counterparts, men and people living in south China were twice as likely to report good sleep quality. Both physical and mental health significantly affected the quality of sleep in elderly Chinese and having enough daily sleep hours substantially increased the likelihood of reporting better sleep quality. Although we did not find any clear trends between levels of sleep quality and levels of inflammatory markers, we did find a significant association between levels of sleep quality and resistin, insulin, HOMA-IR, HOMA2-IR, LDL, and TCH. These associations were found among the total population and a subgroup of participants free from CVD and diabetes.

Considering the alarming increase of sleep disturbances that modern societies are experiencing and the deleterious consequences that a poor night sleep can exert on our metabolism and health, it is fundamental to clarify the factors that might generate sleep disturbances, how these disturbances can affect the biological homeostasis and identify potential interventions to prevent sleep disturbances at an early stage.

Abbreviations

ADIPO: adiponectin; ANOVA: analysis of variance; BMI: body mass index; CI: confidence interval; CRP: C-reactive

protein; CVD: cardiovascular disease; DBP: diastolic blood pressure; HDL: high-density lipoprotein; IL-6: interleukin-6; LDL: low-density lipoprotein; MetS: metabolic syndrome; OR: odds ratio; PAI-1: plasminogen activator inhibitor-1; RBP4: retinol-binding protein 4; SBP: systolic blood pressure; SD: standard deviation; SRH: self rated health; TG: triglycerides; TCH: total cholesterol.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors mentioned participated actively in all and each of the following aspects for this article: Conception and design, or analysis and interpretation of data, drafting the article or revising it critically for important intellectual content and final approval of the version to be published.

Acknowledgements

The authors want to express their sincere appreciation to the study participants and to the researchers and the healthcare professionals from the Centers for Disease Control and Prevention in Beijing and in Shanghai. The authors also want to thank Dr. Yong Liu, Dr. Huaixing Li, Wang Jing, Xing-wang Ye and Qibin Qi for their help in conducting laboratory assays for biomarkers and Dr Hilde Hendrickxs and Dr. Gabrielle Osborne for their valuable comments.

This study was funded by grants SIBS2008006 from the Chief Scientist Program of Shanghai Institutes for Biological Sciences, Chinese Academy of Sciences to Dr. Xu Lin, KSCXI-YW-02 and KSCXI-YW-R-116 from the Knowledge Innovation Program of the Chinese Academy of Sciences, CH-2006-0941 from the Shanghai-Unilever Research Development Fund. These funding organizations did not participate in the design and conduct of the study, collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

References

- Chevalier H, Los F, Boichut D: **Evaluation of severe insomnia in the general population: results of a European multinational survey.** *J Psychopharmacol* 1999, **13**:S21-S24.
- Dew MA, Reynolds CF III, Monk TH: **Psychosocial correlates and sequelae of electroencephalographic sleep in healthy elders.** *J Gerontol* 1994, **49**:P8-P18.
- Dew MA, Hoch CC, Buysse DJ: **Healthy older adults' sleep predicts all-cause mortality at 4 to 19 years of follow-up.** *Psychosom Med* 2003, **65**:63-73.
- Ohayon MM, Carskadon MA, Guilleminault C, Vitiello MV: **Meta-analysis of quantitative sleep parameters from childhood to old age in healthy individuals: developing normative sleep values across the human lifespan.** *Sleep* 2004, **27**:1255-1273.
- Klink ME, Quan SF, Kaltenborn WT, Lebowitz MD: **Risk factors associated with complaints of insomnia in a general adult population. Influence of previous complaints of insomnia.** *Arch Intern Med* 1992, **152**:1634-1637.
- Li RH, Wing YK, Ho SC, Fong SY: **Gender differences in insomnia – a study in the Hong Kong Chinese population.** *J Psychosom Res* 2002, **53**:601-609.
- Reyner LA, Horne JA, Reyner A: **Gender- and age-related differences in sleep determined by home-recorded sleep logs and actimetry from 400 adults.** *Sleep* 1995, **18**:127-134.
- Urpunen H, Vuori I, Hasan J, Partinen M: **Self-evaluations of factors promoting and disturbing sleep: an epidemiological survey in Finland.** *Soc Sci Med* 1988, **26**:443-450.
- Driscoll HC, Serody L, Patrick S: **Sleeping well, aging well: a descriptive and cross-sectional study of sleep in "successful agers" 75 and older.** *Am J Geriatr Psychiatry* 2008, **16**:74-82.
- Baldwin CM, Griffith KA, Nieto FJ, O'Connor GT, Walsleben JA, Redline S: **The association of sleep-disordered breathing and sleep symptoms with quality of life in the Sleep Heart Health Study.** *Sleep* 2001, **24**:96-105.
- Engleman HM, Douglas NJ: **Sleepiness, cognitive function, and quality of life in obstructive sleep apnoea/hypopnoea syndrome.** *Thorax* 2004, **59**:618-622.
- Grunstein RR, Stenlof K, Hedner J, Sjoström L: **Impact of obstructive sleep apnea and sleepiness on metabolic and cardiovascular risk factors in the Swedish Obese Subjects (SOS) Study.** *Int J Obes Relat Metab Disord* 1995, **19**:410-418.
- Koskenvuo M, Kaprio J, Telakivi T, Partinen M, Heikkilä K, Sarna S: **Snoring as a risk factor for ischaemic heart disease and stroke in men.** *Br Med J (Clin Res Ed)* 1987, **294**:16-19.
- Liu Y, Tanaka H: **Overtime work, insufficient sleep, and risk of non-fatal acute myocardial infarction in Japanese men.** *Occup Environ Med* 2002, **59**:447-451.
- Martin SE, Engleman HM, Deary IJ, Douglas NJ: **The effect of sleep fragmentation on daytime function.** *Am J Respir Crit Care Med* 1996, **153**:1328-1332.
- Olson LG, King MT, Hensley MJ, Saunders NA: **A community study of snoring and sleep-disordered breathing health outcomes.** *Am J Respir Crit Care Med* 1995, **152**:717-720.
- Shahar E, Whitney CW, Redline S: **Sleep-disordered breathing and cardiovascular disease: cross-sectional results of the Sleep Heart Health Study.** *Am J Respir Crit Care Med* 2001, **163**:19-25.
- Spriggs DA, French JM, Murdy JM, Curless RH, Bates D, James OF: **Snoring increases the risk of stroke and adversely affects prognosis.** *Q J Med* 1992, **83**:555-562.
- Kripke DF, Garfinkel L, Wingard DL, Klauber MR, Marler MR: **Mortality associated with sleep duration and insomnia.** *Arch Gen Psychiatry* 2002, **59**:131-136.
- Patel SR, Ayas NT, Malhotra MR: **A prospective study of sleep duration and mortality risk in women.** *Sleep* 2004, **27**:440-444.
- Tamakoshi A, Ohno Y: **Self-reported sleep duration as a predictor of all-cause mortality: results from the JACC study, Japan.** *Sleep* 2004, **27**:51-54.
- Newman AB, Spiekerman CF, Enright P: **Daytime sleepiness predicts mortality and cardiovascular disease in older adults. The Cardiovascular Health Study Research Group.** *J Am Geriatr Soc* 2000, **48**:115-123.
- Ayas NT, White DP, Al-Delaimy WK: **A prospective study of self-reported sleep duration and incident diabetes in women.** *Diabetes Care* 2003, **26**:380-384.
- Knutson KL, Ryden AM, Mander BA, Van CE: **Role of sleep duration and quality in the risk and severity of type 2 diabetes mellitus.** *Arch Intern Med* 2006, **166**:1768-1774.
- Yaggi HK, Araujo AB, McKinlay JB: **Sleep duration as a risk factor for the development of type 2 diabetes.** *Diabetes Care* 2006, **29**:657-661.
- Pan A, Ye X, Franco OH, Li H, Yu Z, Zou S, Zhang Z, Jiao S, Lin X: **Insulin resistance and depressive symptoms in middle-aged and elderly Chinese: Findings from the Nutrition and Health of Aging Population in China Study.** *J Affect Disord* 2008, **109**:75-82.
- Pan A, Ye X, Franco OH, Li H, Yu Z, Wang J, Qi Q, Gu W, Pang X, Liu H, Lin X: **The association of depressive symptoms with inflammatory factors and adipokines in middle-aged and older Chinese.** *PLoS ONE* 2008:e1392.
- Pan A, Franco OH, Wang YF, Yu ZJ, Ye XW, Lin X: **Prevalence and geographic disparity of depressive symptoms among middle-aged and elderly in China.** *J Affect Disord* 2008, **105**:167-175.
- Ye X, Yu Z, Li H, Franco OH, Liu Y, Lin X: **Distributions of C-reactive protein and its association with metabolic syndrome in middle-aged and older Chinese people.** *J Am Coll Cardiol* 2007, **49**:1798-1805.
- Lim WY, Ma S, Heng D, Bhalla V, Chew SK: **Gender, ethnicity, health behaviour & self-rated health in Singapore.** *BMC Public Health* 2007, **7**:184.
- Pei X, Rodriguez E: **Provincial income inequality and self-reported health status in China during 1991-7.** *J Epidemiol Community Health* 2006, **60**:1065-1069.

32. Radloff LS: **The CES-D scale: A self-report depression scale for research in the general population.** *Applied Psychological Measurement* 1977, **1**:385-401.
33. Lin N: **Measuring depressive symptomatology in China.** *J Nerv Ment Dis* 1989, **177**:121-131.
34. Zhang J, Norvilitis JM: **Measuring Chinese psychological well-being with Western developed instruments.** *J Pers Assess* 2002, **79**:492-511.
35. Wang J, Li H, Franco OH, Yu Z, Liu Y, Lin X: **Adiponectin and metabolic syndrome in middle-aged and elderly Chinese.** *Obesity* 2008, **16**:172-178.
36. Qi Q, Yu Z, Ye X, Zhao F, Huang P, Hu FB, Franco OH, Wang J, Li H, Liu Y, Lin X: **Elevated retinol-binding protein 4 levels are associated with metabolic syndrome in Chinese people.** *J Clin Endocrinol Metab* 2007, **92**:4827-4834.
37. Levy JC, Matthews DR, Hermans MP: **Correct homeostasis model assessment (HOMA) evaluation uses the computer program.** *Diabetes Care* 1998, **21**:2191-2192.
38. Elwood P, Hack M, Pickering J, Hughes J, Gallacher J: **Sleep disturbance, stroke, and heart disease events: evidence from the Caerphilly cohort.** *J Epidemiol Community Health* 2006, **60**:69-73.
39. Quesnel-Vallee A: **Self-rated health: caught in the crossfire of the quest for 'true' health?** *Int J Epidemiol* 2007, **36**:1161-1164.
40. Singh-Manoux A, Gueguen A, Martikainen P, Ferrie J, Marmot M, Shipley M: **Self-rated health and mortality: short- and long-term associations in the Whitehall II study.** *Psychosom Med* 2007, **69**:138-143.
41. Ford DE, Kamerow DB: **Epidemiologic study of sleep disturbances and psychiatric disorders. An opportunity for prevention?** *JAMA* 1989, **262**:1479-1484.
42. Rodin J, McAvay G, Timko C: **A longitudinal study of depressed mood and sleep disturbances in elderly adults.** *J Gerontol* 1988, **43**:P45-P53.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2458/9/130/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

