

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): G. ROWLANDS and M. A. ALLEN

Article Title: Nonlinear magnetoacoustic waves in a cold plasma

Year of publication: 2007

Link to published version: <http://dx.doi.org/10.1017/S0022377806004818>

Publisher statement: None

# Nonlinear magnetoacoustic waves in a cold plasma

G. ROWLANDS<sup>1</sup> and M. A. ALLEN<sup>2</sup>

<sup>1</sup>Department of Physics, University of Warwick, Coventry, CV4 7AL, UK

<sup>2</sup>Physics Department, Mahidol University, Rama 6 Road, Bangkok 10400 Thailand  
(frmaa@mahidol.ac.th)

(Received 6 June 2006 and accepted 16 June 2006)

**Abstract.** The equations describing planar magnetoacoustic waves of permanent form in a cold plasma are rewritten so as to highlight the presence of a naturally small parameter equal to the ratio of the electron and ion masses. If the magnetic field is not nearly perpendicular to the direction of wave propagation, this allows us to use a multiple-scale expansion to demonstrate the existence and nature of nonlinear wave solutions. Such solutions are found to have a rapid oscillation of constant amplitude superimposed on the underlying large-scale variation. The approximate equations for the large-scale variation are obtained by making an adiabatic approximation and in one limit, new explicit solitary pulse solutions are found. In the case of a perpendicular magnetic field, conditions for the existence of solitary pulses are derived. Our results are consistent with earlier studies which were restricted to waves having a velocity close to that of long-wavelength linear magnetoacoustic waves.

---

## 1. Introduction

For a plasma composed of cold electrons and a single species of cold ions, both collisions and Landau damping can be neglected with the result that a two-fluid model provides an accurate description (Kakutani et al. 1968). Such a model is governed by the continuity and momentum equations for electrons and ions, and Maxwell's equations. In the study of non-relativistic hydromagnetic waves with a frequency much less than the plasma frequency, these equations may be simplified somewhat by neglecting the displacement current and taking the number densities of electrons and ions to be equal, except in Poisson's equation (Kakutani et al. 1967). Then, taking all quantities to be independent of  $y$  and  $z$ , one arrives at a set of equations governing planar hydromagnetic waves. Il'ichev (1996) integrates these to obtain the following equations for a magnetoacoustic wave of permanent form propagating in the  $x$ -direction at a constant speed  $V$ :

$$\frac{dv}{d\xi} = -\frac{R_i \cos \theta}{V}nw - R_i B_z, \quad (1.1a)$$

$$\frac{dw}{d\xi} = \frac{R_i \cos \theta}{V}nv + R_i \hat{B}_y + R_i \sin \theta(1 - n), \quad (1.1b)$$

$$\frac{d\hat{B}_y}{d\xi} = R_e nw + \frac{R_e \cos \theta}{V}nB_z, \quad (1.1c)$$

$$\frac{dB_z}{d\xi} = -R_e n v - \frac{R_e \cos \theta}{V} n \hat{B}_y, \quad (1.1d)$$

where  $\xi = x - Vt$ ,

$$\frac{1}{n} = 1 - \frac{1}{2V^2} (\hat{B}_y^2 + 2\hat{B}_y \sin \theta + B_z^2), \quad (1.2)$$

$$\hat{B}_y = B_y - \sin \theta, \quad (1.3)$$

and  $\theta$  is the angle between the equilibrium magnetic field and the  $x$ -axis. In the above equations,  $\xi$ , the ion density  $n$ , the  $y$  and  $z$  components of the ion drift velocity  $v$  and  $w$ , and the magnetic field ( $B_x, B_y, B_z$ ) are normalized, respectively, by the characteristic length  $l$ , the equilibrium ion density, the Alfvén velocity  $V_A$ , and the equilibrium magnetic field strength. In such units the speed of a linear long-wavelength magnetoacoustic wave is unity. The remaining parameters are defined by  $R_i = \omega_{ci} l / V_A$  and  $R_e = \omega_{ce} l / V_A$  where  $\omega_{ci}$  and  $\omega_{ce}$  are the ion and electron cyclotron frequencies, respectively. The values of the dependent variables in the absence of a wave are zero for  $v, w, \hat{B}_y$ , and  $B_z$ , and unity in the case of  $n$ .

The results we present here originate from our observation that (1.1) may be rewritten so as to include a small parameter,  $\varepsilon$ . After introducing the variable  $s \equiv R_e \xi / \bar{V}$ , where  $\bar{V} = V \sec \theta$ , the equations take the form

$$\frac{dv}{ds} = -\varepsilon \{nw + \bar{V} B_z\}, \quad (1.4a)$$

$$\frac{dw}{ds} = \varepsilon \{nv + \bar{V} \hat{B}_y + V \tan \theta (1 - n)\}, \quad (1.4b)$$

$$\frac{d\hat{B}_y}{ds} = n \{\bar{V} w + B_z\}, \quad (1.4c)$$

$$\frac{dB_z}{ds} = -n \{\bar{V} v + \hat{B}_y\}, \quad (1.4d)$$

where  $\varepsilon \equiv R_i / R_e$  is simply the ratio of the electron and ion masses. With  $\varepsilon \sim 10^{-3}$  or smaller, any analytical treatment should make use of the smallness of this parameter and some form of perturbation theory is obviously called for. Furthermore, even a purely numerical method of solution should attempt to take advantage of the smallness of  $\varepsilon$  as otherwise it means integrating over many small-scale variations before the underlying variation on the large scale takes place.

From (1.4) it can be seen that generally  $v$  and  $w$  change on a much slower scale than  $\hat{B}_y$  and  $B_z$ . It is therefore natural to make an adiabatic approximation (see Haken (1983)) which is equivalent to equating the right-hand sides of (1.4c) and (1.4d) to zero. This gives

$$B_z = -\bar{V} w, \quad \hat{B}_y = -\bar{V} v. \quad (1.5)$$

These relationships can be used to eliminate  $\hat{B}_y$  and  $B_z$  and it is shown in Appendix A that all the dependent variables can be expressed in terms of a single variable which itself satisfies a Newtonian-type energy equation with a polynomial Sagdeev potential. This equation is then used to show the existence of solitary pulses and nonlinear waves on the larger scale. However, it should be stressed that the very nature of the adiabatic approximation as used here is to eliminate any variation

on the smaller scale. Furthermore, the class of solutions to (1.4) is restricted, in view of relationships (1.5), which implies that at some particular value of  $s$  the corresponding values of  $(B_z, w)$  and  $(\hat{B}_y, v)$  are not independent.

It is the purpose of this paper to construct a perturbation expansion based on the smallness of  $\varepsilon$  which allows one to put the adiabatic approximation in context and to allow for rapid oscillations on the small scale. These effects have been studied analytically in Il'ichev (1996) and later numerically by Bakholdin and Il'ichev (1998) and Bakholdin et al. (2002). However, those studies were only carried out for the case  $V = 1 + \mu/2$  where  $\mu$  is small. In the present paper, the analytic theory based on the smallness of  $\varepsilon$ , a naturally small parameter, clearly shows the origin of all the basic features of the solutions they obtain and is not restricted to small  $\mu$ .

The underlying form of the governing equations suggests that a multiple-scale perturbation expansion is appropriate and this is carried out in Sec. 2 where explicit equations are obtained that describe the evolution of  $v, w, \hat{B}_y$ , and  $B_z$  on both the small and large scale to lowest significant order in  $\varepsilon$ . This analysis is only valid when  $\cos \theta$  is not small. For  $\theta$  close to  $\pm\pi/2$  singularities develop in (1.4) and so an alternative set of variables and equations derived from (1.1) must be employed, as is shown in Sec. 3. The final section summarizes our results and the further applicability of our approach is discussed.

## 2. Multiple-scale perturbation expansion

The perturbation expansion is based on the implication from equations (1.4) that two distinct spatial scales exist—a large and a small scale. We formally proceed by introducing multiple scales,  $s, s_1 = \varepsilon s, s_2 = \varepsilon^2 s, \dots$  and treating them as independent variables so that

$$\frac{d}{ds} = \frac{\partial}{\partial s} + \varepsilon \frac{\partial}{\partial s_1} + \varepsilon^2 \frac{\partial^2}{\partial s_2^2} + \dots,$$

and allowing all dependent variables  $f$  to be functions of  $s, s_1, \dots$  and expressible in the form

$$f(s) = f_0(s, s_1, \dots) + \varepsilon f_1(s, s_1, \dots) + \dots,$$

although in the following it is not necessary to consider the scaled variables beyond  $s$  (the small scale) and  $s_1$  (the large scale). For a more general introduction to this form of perturbation theory see, for example, Nayfeh and Mook (1979) and Rowlands (1990).

Substituting the above form for the dependent variables into (1.4a) and (1.4b) gives, to lowest order,

$$\frac{\partial v_0}{\partial s} = 0, \quad \frac{\partial w_0}{\partial s} = 0,$$

and so  $v_0$  and  $w_0$  can only be functions of  $s_1$ , and not  $s$ . At lowest order, (1.4c) and (1.4d) reduce to

$$\frac{\partial \hat{B}_{y0}}{\partial s} = n_0 \{ \bar{V} w_0 + B_{z0} \}, \quad \frac{\partial B_{z0}}{\partial s} = n_0 \{ \bar{V} v_0 + \hat{B}_{y0} \}. \quad (2.1)$$

Since  $v_0$  and  $w_0$  are independent of  $s$ , the solution of these equations is

$$\hat{B}_{y0} = h_y - \bar{V} v_0, \quad B_{z0} = h_z - \bar{V} w_0, \quad (2.2)$$

where

$$\frac{\partial h_y}{\partial s} = n_0 h_z, \quad \frac{\partial h_z}{\partial s} = -n_0 h_y.$$

These equations have the solution

$$h_y = h(s_1) \sin \phi, \quad h_z = h(s_1) \cos \phi, \quad (2.3)$$

where

$$\frac{\partial \phi}{\partial s} = n_0 \quad (2.4)$$

and  $h(s_1)$  is an as yet to be determined function of  $s_1$ . Substituting the above results into (1.2) gives, to this approximation,

$$\frac{1}{n_0} = A + D(s_1) \cos \phi + E(s_1) \sin \phi \quad (2.5)$$

where

$$A = 1 + \frac{2V \tan \theta v_0 - \bar{V}^2(w_0^2 + v_0^2) - h^2}{2V^2} \quad (2.6)$$

and

$$D = \frac{\bar{V} h w_0}{V^2}, \quad E = \frac{(\bar{V} v_0 - \sin \theta) h}{V^2}.$$

Thus  $A$ ,  $D$ , and  $E$  are functions of  $s_1$  only and the  $s$  variation of  $n_0$  is through  $\phi$  only. Integrating the reciprocal of (2.4) after using (2.5) to express  $n_0$  in terms of  $\phi$  yields

$$A\phi + D \sin \phi - E \cos \phi = s - \tilde{s}(s_1) \quad (2.7)$$

where  $\tilde{s}$  is a function of  $s_1$ . A solution of (2.7) in the form  $\phi = \phi(s, s_1)$  is obtained in Appendix B.

We now proceed to the next order of approximation and find using (1.4a) that

$$\frac{\partial v_1}{\partial s} = -\frac{dv_0}{ds_1} - n_0 w_0 + \bar{V}^2 w_0 - \bar{V} h \cos \phi. \quad (2.8)$$

Since the right-hand side depends on  $s$  via  $\phi$  only, we replace  $\partial v_1 / \partial s$  by  $n_0 \partial v_1 / \partial \phi$ . Remembering that  $v_0$ ,  $w_0$  and  $h$  are independent of  $\phi$ , and using (2.5), (2.8) is readily integrated to give

$$\begin{aligned} v_1 = & \left( \bar{V}^2 w_0 - \frac{dv_0}{ds_1} \right) (A\phi + D \sin \phi - E \cos \phi) - w_0 \phi \\ & + \bar{V} h \left( \frac{E}{4} \cos 2\phi - \frac{D}{4} (2\phi + \sin 2\phi) - A \sin \phi \right) + \tilde{v}(s_1) \end{aligned} \quad (2.9)$$

where  $\tilde{v}(s_1)$  is an undetermined function. To ensure that  $v_1$  is a bounded function of  $s$ , we must remove terms proportional to  $\phi$ . After replacing  $D$  by its definition this results in the consistency condition

$$A \frac{dv_0}{ds_1} = \left( A \bar{V}^2 - 1 - \frac{\bar{V}^2 h^2}{2V^2} \right) w_0. \quad (2.10)$$

This is the equation for the variation of  $v_0$  on the slowly varying scale,  $s_1$ . Using (2.10) to simplify (2.9) leaves us with

$$v_1 = \frac{1}{A} \left( w_0 + \frac{\bar{V}hD}{2} \right) (D \sin \phi - E \cos \phi) + \bar{V}h \left( \frac{E}{4} \cos 2\phi - \frac{D}{4} \sin 2\phi - A \sin \phi \right), \quad (2.11)$$

in which  $\tilde{v}$  has been absorbed into  $v_0$ .

Similarly, to first order, (1.4b) yields

$$\frac{\partial w_1}{\partial s} = -\frac{dw_0}{ds_1} + n_0 v_0 - \bar{V}^2 v_0 + \bar{V}h \sin \phi + V \tan \theta (1 - n_0). \quad (2.12)$$

After again replacing  $\partial/\partial s$  by  $n_0 \partial/\partial \phi$  and integrating, to obtain a bounded  $w_1$  we require that

$$A \frac{dw_0}{ds_1} = \left( 1 - A\bar{V}^2 + \frac{\bar{V}^2 h^2}{2V^2} \right) v_0 + V \tan \theta \left( A - 1 - \frac{h^2}{2V^2} \right), \quad (2.13)$$

with the result that

$$w_1 = \frac{1}{A} \left( V \tan \theta - v_0 - \frac{\bar{V}hE}{2} \right) (D \sin \phi - E \cos \phi) - \bar{V}h \left( A \cos \phi + \frac{D}{4} \cos 2\phi + \frac{E}{4} \sin 2\phi \right). \quad (2.14)$$

To lowest order, the adiabatic approximation (1.5) is

$$B_{z0} = -\bar{V}w_0, \quad \hat{B}_{y0} = -\bar{V}v_0. \quad (2.15)$$

Comparison with (2.2) shows that this approximation is equivalent to setting  $h = 0$ . Equations (2.10) and (2.13) then form a complete set that can be integrated. The details are given in Appendix A. In particular, the existence of solitary pulses is proven.

If the adiabatic approximation is not made, it is necessary to obtain an equation for the variation of  $h$  on the  $s_1$  scale. This is achieved by considering the equations for  $\hat{B}_y$  and  $B_z$  to next order in  $\varepsilon$ . From (1.4c) and (1.4d) we may write, respectively,

$$\frac{\partial \hat{B}_{y1}}{\partial s} + \frac{\partial \hat{B}_{y0}}{\partial s_1} = n_0 \{ \bar{V}w_1 + B_{z1} \} + n_1 h \cos \phi,$$

$$\frac{\partial B_{z1}}{\partial s} + \frac{\partial B_{z0}}{\partial s_1} = -n_0 \{ \bar{V}v_1 + \hat{B}_{y1} \} - n_1 h \sin \phi.$$

Since the variation of all coefficients with  $s$  is through  $\phi$ , we replace  $\partial/\partial s$  by  $n_0 \partial/\partial \phi$  and rewrite the above equations as

$$\frac{\partial \hat{B}_{y1}}{\partial \phi} + \frac{1}{n_0} \frac{\partial \hat{B}_{y0}}{\partial s_1} = \bar{V}w_1 + B_{z1} + \frac{n_1 h}{n_0} \cos \phi, \quad (2.16a)$$

$$\frac{\partial B_{z1}}{\partial \phi} + \frac{1}{n_0} \frac{\partial B_{z0}}{\partial s_1} = -\bar{V}v_1 - \hat{B}_{y1} - \frac{n_1 h}{n_0} \sin \phi. \quad (2.16b)$$

We proceed by adding (2.16a) multiplied by  $\sin \phi$  to (2.16b) multiplied by  $\cos \phi$  and then integrating from 0 to  $2\pi$ . Insisting that  $\hat{B}_{y1}$  and  $B_{z1}$  are periodic functions of

$\phi$  means that

$$\left\langle \frac{\partial \hat{B}_{y1}}{\partial \phi} \sin \phi \right\rangle = -\langle \hat{B}_{y1} \cos \phi \rangle, \quad \left\langle \frac{\partial B_{z1}}{\partial \phi} \cos \phi \right\rangle = \langle B_{z1} \sin \phi \rangle,$$

where  $\langle \cdot \rangle$  denotes the average as  $\phi$  varies from 0 to  $2\pi$ . The combined equations then reduce to the following equation for the variation of  $h$ :

$$\left\langle \frac{1}{n_0} \right\rangle \frac{dh}{ds_1} = \bar{V} \left\{ \frac{dv_0}{ds_1} \left\langle \frac{\sin \phi}{n_0} \right\rangle + \frac{dw_0}{ds_1} \left\langle \frac{\cos \phi}{n_0} \right\rangle + \langle w_1 \sin \phi \rangle - \langle v_1 \cos \phi \rangle \right\}. \quad (2.17)$$

Using (2.5), (2.11), and (2.14) we have

$$\left\langle \frac{1}{n_0} \right\rangle = A, \quad \left\langle \frac{\sin \phi}{n_0} \right\rangle = \frac{E}{2}, \quad \left\langle \frac{\cos \phi}{n_0} \right\rangle = \frac{D}{2},$$

$$\langle w_1 \sin \phi \rangle = \left( V \tan \theta - v_0 - \frac{\bar{V} h E}{2} \right) \frac{D}{2A}, \quad \langle v_1 \cos \phi \rangle = -\left( 1 + \frac{\bar{V}^2 h^2}{2V^2} \right) \frac{E w_0}{2A}.$$

Then inserting the above expressions and results (2.10) and (2.13) into (2.17) and simplifying we find that the right-hand side of (2.17) is zero and hence that  $h$  is a constant. This is in agreement with the result obtained for  $V$  close to 1 in Il'ichev (1996).

In summary, we have seen that to lowest order, the ion drift velocity components  $v$  and  $w$  only show large-scale variation. Fast periodic variation occurs at the next order of approximation, as given by (2.11) and (2.14), but given the smallness of  $\varepsilon$ , these oscillations would be barely discernible. On the other hand, even to lowest order, the magnetic field components show rapid oscillations on top of the large-scale variation:

$$\hat{B}_{y0} = -\bar{V} v_0(s_1) + h \sin \phi(s, s_1), \quad B_{z0} = -\bar{V} w_0(s_1) + h \cos \phi(s, s_1),$$

where, as is shown in Appendix B,  $\sin \phi$  and  $\cos \phi$  are periodic functions of the variable  $S$  given by (B 3).

### 3. The small $\cos \theta$ and $\theta = \pm\pi/2$ limits

So far we have treated  $\cos \theta$  as finite but our treatment does not allow one to pass to the case  $\cos \theta = 0$  since in this limit  $\bar{V}$  becomes infinite. To consider this limit we write  $\theta = \pm\pi/2 \mp \sqrt{\varepsilon}\psi$  so that  $\cos \theta = \sin(\sqrt{\varepsilon}\psi) \simeq \sqrt{\varepsilon}\psi$ . (In the remainder of this section the upper and lower signs refer to the cases where  $\theta$  is in the neighbourhood of  $\pi/2$  and  $-\pi/2$ , respectively.) It is now necessary to go back to the original equations expressed in terms of  $\xi$  rather than  $s$  and define  $X \equiv \sqrt{R_i R_e} \xi$ . Also, to avoid singular solutions we need to use scaled versions of the ion drift velocities, defined by  $\bar{v} = v/\sqrt{\varepsilon}$  and  $\bar{w} = w/\sqrt{\varepsilon}$ . Then dividing (1.1) by  $\sqrt{R_i R_e}$  we obtain

$$\frac{d\bar{v}}{dX} = -B_z + O(\varepsilon\psi), \quad (3.1a)$$

$$\frac{d\bar{w}}{dX} = \hat{B}_y \pm 1 - n + O(\varepsilon\psi^2), \quad (3.1b)$$

$$\frac{d\hat{B}_y}{dX} = n\bar{w} + \frac{nB_z}{V}\psi + O(\varepsilon\psi^3), \quad (3.1c)$$

$$\frac{dB_z}{dX} = -n\bar{v} - \frac{n\hat{B}_y}{V}\psi + O(\varepsilon\psi^3) \tag{3.1d}$$

with

$$\frac{1}{n} = 1 - \frac{1}{2V^2}(\hat{B}_y^2 \pm 2\hat{B}_y + B_z^2) + O(\varepsilon\psi^2). \tag{3.2}$$

Any solution to the above equations will be such that  $\bar{v} \sim B$  and hence the real velocity  $v = \sqrt{\varepsilon}\bar{v}$  will be small compared to  $B$ .

For the  $\theta = \pm\pi/2$  limits, (3.1) reduces to

$$\frac{d\bar{v}}{dX} = -B_z, \quad \frac{dB_z}{dX} = -n\bar{v}, \tag{3.3a}$$

$$\frac{d\bar{w}}{dX} = \hat{B}_y \pm 1 - n, \quad \frac{d\hat{B}_y}{dX} = n\bar{w}. \tag{3.3b}$$

We now define an operator  $L$  by

$$Lf \equiv \frac{d}{dX} \left( \frac{1}{n} \frac{df}{dX} \right).$$

Then from (3.3a) and (3.3b) it can be seen that, respectively,

$$LB_z = B_z, \tag{3.4a}$$

$$LB_y = B_y - n, \tag{3.4b}$$

in which we have re-instated  $B_y$ , as given in (1.3). Combining (3.4a) and (3.4b) gives

$$B_yLB_z - B_zLB_y = B_zn.$$

Integrating this over one period (or all  $X$  if boundary conditions permit) yields

$$\langle B_zn \rangle = 0, \tag{3.5}$$

where  $\langle \cdot \rangle$  denotes the integral over  $X$ . Similarly, (3.4a) and (3.4b) imply that

$$\langle B_z \rangle = 0, \quad \langle B_y \rangle = \langle n \rangle. \tag{3.6}$$

Relations (3.5) and (3.6) are satisfied if  $n$  and  $B_y$  are even functions and  $B_z$  is an odd function of  $X$ .

We can demonstrate the existence of a non-trivial solution of (3.3) by taking  $B_z = 0$ . In this case (3.2) becomes

$$\frac{1}{n} = \alpha(1 - \beta B_y^2)$$

where

$$\alpha = 1 + \frac{1}{2V^2}, \quad \beta = \frac{1}{1 + 2V^2}, \tag{3.7}$$

with the result that (3.4b) can be re-expressed as

$$\alpha \frac{d}{dX} \left( (1 - \beta B_y^2) \frac{dB_y}{dX} \right) = B_y - \frac{1}{\alpha(1 - \beta B_y^2)}.$$

Letting  $p = dB_y/dX$ , this can be written as the following first-order differential equation for  $p^2$ :

$$\frac{dp^2}{dB_y} - \left( \frac{4\beta B_y}{1 - \beta B_y^2} \right) p^2 = \frac{2}{\alpha} \left( \frac{B_y}{1 - \beta B_y^2} - \frac{1}{\alpha(1 - \beta B_y^2)^2} \right),$$



which has an integrating factor of  $(1 - \beta B_y^2)^2$ . Hence the solution is given by

$$p^2 = \frac{2Q}{\alpha(1 - \beta B_y^2)^2} \quad (3.8)$$

where

$$Q = \frac{B_y^2}{2} - \frac{\beta B_y^4}{4} - \frac{B_y}{\alpha} + Q_0 \quad (3.9)$$

and  $Q_0$  is an integration constant. For a solitary pulse solution, the appropriate boundary conditions are  $B_y \rightarrow \pm 1$  and  $Q \rightarrow 0$  as  $|X| \rightarrow \infty$ . Using these allows us to determine  $Q_0$ . We then rewrite (3.9) as the following expansion in  $\hat{B}_y$ :

$$Q = \hat{B}_y^2 \left\{ \frac{1 - 3\beta}{2} \mp \beta \hat{B}_y - \frac{\beta}{4} \hat{B}_y^2 \right\}. \quad (3.10)$$

A necessary condition for the existence of solitary pulses is therefore that  $\beta < 1/3$ . In addition, from (3.8) it can be seen that  $p$  is singular at  $B_y^2 = 1/\beta$ . Hence, for a solitary pulse to exist, at least one of the two non-trivial zeros of the expression for  $Q$  given in (3.10) must lie within the range  $-1/\sqrt{\beta} \mp 1 < \hat{B}_y < 1/\sqrt{\beta} \mp 1$ . The zero that is larger in magnitude never satisfies this. The remaining zero at  $\pm\sqrt{2/\beta - 2} \mp 2$  satisfies the condition if  $\beta > 1/9$ . Using (3.7) we can now write the sufficient condition for the existence of a solitary pulse solution as  $1 < V^2 < 4$ .

As a check on our calculation we look at the case when  $V = 1 + \mu/2$  for small positive  $\mu$ . Then (3.8) reduces to

$$\left( \frac{d\hat{B}_y}{dX} \right)^2 = \frac{\hat{B}_y^2}{4} (\hat{B}_y \pm 4) (\pm\mu - \hat{B}_y)$$

to lowest order in  $\mu$ . The above equation has the solution

$$\hat{B}_y = \pm\mu \operatorname{sech}^2 \frac{1}{2} \sqrt{\mu} (X - X_0) + O(\mu^2)$$

where  $X_0$  is an arbitrary constant. This is in agreement with the result given in Bakholdin and Il'ichev (1998).

#### 4. Conclusions

We have studied a set of magnetohydrodynamic equations for planar magnetoacoustic waves of permanent form propagating in a two-component cold plasma and, by taking advantage of the smallness of the ratio of the electron to ion masses, have obtained a reduced set of equations that describe the large-scale variation of the magnetoacoustic wave solution of the full equations. Superimposed on the large-scale variation, multiple-scale perturbation analysis indicates that there is a rapid oscillation which is of constant amplitude in the case of the lowest-order magnetic field components. These results are consistent with the study of Il'ichev (1996), which was restricted to a narrow range of velocities. In addition, the approach expounded in this paper puts the adiabatic approximation into its true context.

In this work we have obtained various conditions for the existence of solitary pulses. Whether these solutions correspond to phenomena that could occur in nature depends on whether they are stable. The linear stability analysis of the solutions shown to exist in this paper is a challenging problem. However, the numerical solution of the full (time-dependent) system of equations obtained by

Bakholdin and Il'ichev (1998), show that for a range of initial conditions the solution relaxes to the type of solution shown to exist here. This suggests that our solutions are stable, at least to perturbations applied in the direction of propagation.

Although the equations studied here arise from a magnetohydrodynamics problem, the method is applicable to a more general set of nonlinear equations where two distinct scales are a basic feature. An advantage of the present study is that the equations obtained on the large scale can be investigated analytically and describe real physical processes. Although we have only looked at cold plasmas, an exactly analogous procedure can be applied to the case of warm plasmas, at the expense of some additional algebraic complexity. The relevant governing equations are given in Bakholdin et al. (2002).

*Acknowledgements*

G.R. thanks Mahidol University for their hospitality during his visit.

**Appendix A. Adiabatic approximation**

When  $h = 0$ , the coupled equations for  $v_0$  and  $w_0$ , namely (2.10) and (2.13), reduce to

$$\frac{dv_0}{ds_1} = \left( \bar{V}^2 - \frac{1}{A} \right) w_0 \tag{A 1}$$

and

$$\frac{dw_0}{ds_1} = \left( \frac{1}{A} - \bar{V}^2 \right) v_0 + V \tan \theta \left( 1 - \frac{1}{A} \right). \tag{A 2}$$

Multiplying (A 1) and (A 2) by  $2v_0$  and  $2w_0$ , respectively, and adding gives

$$\frac{d}{ds_1} (v_0^2 + w_0^2) = 2V \tan \theta \left( 1 - \frac{1}{A} \right) w_0. \tag{A 3}$$

Differentiating (2.6) with respect to  $s_1$  and using the above, one obtains

$$\frac{dA}{ds_1} = \frac{(\bar{V}^2 - 1) \tan \theta}{V} \frac{w_0}{A},$$

or, provided that  $\bar{V}^2 \neq 1$ ,

$$w_0 = \frac{A}{\kappa} \frac{dA}{ds_1}, \quad \kappa = \frac{(\bar{V}^2 - 1) \tan \theta}{V}. \tag{A 4}$$

Substituting (A 4) into (A 3) and (A 1) and integrating yields, respectively,


$$v_0^2 + w_0^2 = C_1 + V \kappa^{-1} \tan \theta (A^2 - 2A) \tag{A 5}$$

and

$$v_0 = C_2 + \kappa^{-1} \left( \frac{1}{2} \bar{V}^2 A^2 - A \right) \tag{A 6}$$

where  $C_1$  and  $C_2$  are integration constants. Finally, after combining (A 4), (A 5) and (A 6) one obtains

$$\left( A \frac{dA}{ds_1} \right)^2 = \sum_{m=0}^4 \gamma_m A^m \tag{A 7}$$



**Figure A.1.** Phase plane for (A 8) when conditions are such that both compressive and rarefactive solitary pulses occur.

in which

$$\begin{aligned} \gamma_0 &= (C_1 - C_2^2)\kappa^2, & \gamma_1 &= 2(C_2 - V \tan \theta)\kappa, & \gamma_2 &= (V \tan \theta - C_2\bar{V}^2)\kappa - 1, \\ \gamma_3 &= \bar{V}^2, & \gamma_4 &= -\frac{1}{4}\bar{V}^4. \end{aligned}$$

This is of the form of the energy equation of a particle with position  $A$  in a Sagdeev potential which is minus the right-hand side of (A 7). In general, nonlinear waves exist and, in particular, solitary pulses exist. The latter can occur when the boundary conditions are such that  $v \rightarrow 0$ ,  $w \rightarrow 0$ , and  $n \rightarrow 1$  as  $s_1 \rightarrow \pm\infty$ . Using the result that in this limit  $A \rightarrow 1$ , the boundary conditions allow us to use (A 5) and (A 6) to determine the integration constants which in this case are

$$C_1 = V\kappa^{-1} \tan \theta, \quad C_2 = (1 - \frac{1}{2}\bar{V}^2)\kappa^{-1}.$$

Then (A 7) reduces to

$$\left(A \frac{dA}{ds_1}\right)^2 = \frac{(A-1)^2}{4}(a + bA + cA^2) \tag{A 8}$$

where

$$a = 4(\bar{V}^2 - 1) \tan^2 \theta - (\bar{V}^2 - 2)^2, \quad b = 2\bar{V}^2(2 - \bar{V}^2), \quad c = -\bar{V}^4.$$

The requirement that (A 8) gives rise to a solitary pulse is that  $a + b + c > 0$ . Using the above renders this condition as  $(\bar{V}^2 - 1)(1 - V^2) > 0$  which on rearranging yields

$$\cos^2 \theta < V^2 < 1.$$

As illustrated in Fig. A.1, since  $c < 0$ , if the above condition is satisfied, compressive solitary pulses will always occur. However, rarefactive pulses are only possible if the smaller root of  $a + bA + cA^2 = 0$  is above zero. This will occur if  $a < 0$  and  $\bar{V}^2 < 2$ . These requirements are equivalent to the condition

$$V^2 < 2(1 - |\sin \theta|).$$

This implies that if the values of  $V$  and  $\theta$  are such that compressive pulses exist, then rarefactive pulses will also occur if  $|\theta| \leq \pi/6$ .

It is possible to integrate (A 8) to obtain the spatial variation of the solitary pulses implicitly. An approximate explicit solution can be obtained when  $\bar{V}^2$  is just above 1. Introducing  $U \equiv A - 1$ , (A 8) becomes

$$\left(\frac{dU}{ds_1}\right)^2 = \frac{\bar{V}^4 U^2 (U_+ - U)(U - U_-)}{(1 + U)^2} \tag{A 9}$$

where

$$U_{\pm} = \frac{2\sqrt{\bar{V}^2 - 1}}{\bar{V}^2} \left( \pm \tan \theta - \sqrt{\bar{V}^2 - 1} \right).$$

If  $\sqrt{\bar{V}^2 - 1} \ll |\tan \theta|$ , then  $U_{\pm} \simeq \pm \nu$  where

$$\nu = \frac{2\sqrt{\bar{V}^2 - 1} \tan \theta}{\bar{V}^2}.$$

For solitary pulse solutions,  $|U| < |U_{\pm}|$ , and so if  $\tan \theta$  is of order unity,  $U \ll 1$ . Hence (A 9) reduces to

$$\frac{dU}{ds_1} = U\sqrt{\nu^2 - U^2}$$

at lowest order and one obtains

$$A \simeq 1 \pm \nu \operatorname{sech} \nu s_1. \tag{A 10}$$

Using (A 4) and (A 6) we can then obtain the corresponding expressions for  $w_0$  and  $v_0$ :

$$w_0 \simeq \mp 4V \tan \theta \operatorname{sech} \nu s_1 \tanh \nu s_1, \quad v_0 \simeq 2V \tan \theta \operatorname{sech}^2 \nu s_1. \tag{A 11}$$

In this adiabatic approximation, the lowest-order components of the magnetic field are just multiples of these quantities, as given by (2.15).

We can now also use (A 10) and (A 11) to obtain the solution when  $\bar{V}^2 = 1$ . This corresponds to the limit  $\nu \rightarrow 0$  in which case  $A \rightarrow 1$ ,  $w_0 \rightarrow 0$ , and  $v_0 \rightarrow 2V \tan \theta$ . As a check on our calculation, we note that these results are consistent with the definition of  $A$  as given by (2.6) when  $h = 0$ .

### Appendix B. Explicit expression for $\phi(s, s_1)$

The variation of  $\phi$  with  $s$  is given implicitly by (2.7). An explicit expression can be obtained by writing the equation in the form

$$S + \psi = \phi + \psi - \sigma \sin(\phi + \psi) \tag{B 1}$$

where  $S = (s - \bar{s})/A$ ,  $\psi = \arg(-D + iE)$  and  $\sigma = \sqrt{D^2 + E^2}/A$ . The following explicit solution to (B 1) was first obtained by Jackson (1960) (although for a more transparent exposition see p.154 of Infeld and Rowlands (2000)):

$$\phi = S + 2 \sum_{m=1}^{\infty} \frac{J_m(m\sigma)}{m} \sin m(S + \psi). \tag{B 2}$$

It can be seen that  $\phi$  has a directed component,  $S$ , on which a periodic variation is superimposed. Since  $A$  and hence the period vary on the  $s_1$  scale, to the order in  $\epsilon$  to which (B 2) applies, it is more appropriate to redefine  $S$  by

$$S = \int_{\bar{s}}^s \frac{ds'}{A(\epsilon s')}. \tag{B 3}$$

Such a definition avoids secular terms at higher orders in the  $\epsilon$  expansion.

### References

Bakholdin, I. and Il'ichev, A. 1998 *J. Plasma Phys.* **60**, 569.

- Bakholdin, I., Il'ichev, A. and Zharkov, A. 2002 *J. Plasma Phys.* **67**, 1.
- Haken, H. 1983 *Synergetics*, 3rd edn. Berlin: Springer.
- Il'ichev, A. 1996 *J. Plasma Phys.* **55**, 181.
- Infeld, E. and Rowlands, G. 2000 *Nonlinear Waves, Solitons and Chaos*, 2nd edn. Cambridge: Cambridge University Press.
- Jackson, E. A. 1960 *Phys. Fluids* **3**, 831.
- Kakutani, T., Kawahara, T. and Taniuti, T. 1967 *J. Phys. Soc. Japan* **23**, 1138.
- Kakutani, T., Ono, H., Taniuti, T. and Wei, C.-C. 1968 *J. Phys. Soc. Japan* **24**, 1159.
- Nayfeh, A. H. and Mook, D. T. 1979 *Nonlinear Oscillations*. New York: Wiley.
- Rowlands, G. 1990 *Non-Linear Phenomena in Science and Engineering*. London: Ellis Horwood.