

University of Warwick institutional repository: <http://go.warwick.ac.uk/wrap>

This paper is made available online in accordance with publisher policies. Please scroll down to view the document itself. Please refer to the repository record for this item and our policy information available from the repository home page for further information.

To see the final version of this paper please visit the publisher's website. Access to the published version may require a subscription.

Author(s): Mary N. Heneghan, Claudine Porta, Cunjin Zhang, Kerry S. Burton, Michael P. Challen, Andy M. Bailey, and Gary D. Foster
Article Title: Characterization of Serine Proteinase Expression in *Agaricus bisporus* and *Coprinopsis cinerea* by Using Green Fluorescent Protein and the *A. bisporus* SPR1 Promoter
Year of publication: 2009
Link to published version: <http://dx.doi.org/10.1128/AEM.01897-08>
Publisher statement: None

1 Characterisation of serine proteinase expression in *Agaricus bisporus* and *Coprinopsis*
2 *cinerea* using GFP and the *A. bisporus SPRI* promoter.

3 Mary N. Heneghan¹, Claudine Porta², Cunjin Zhang², Kerry S. Burton², Michael P.
4 Challen², Andy M. Bailey¹ & Gary D. Foster.^{1*}

5

6 ¹ School of Biological Sciences, University of Bristol, Bristol BS8 1UG, UK

7 ² Warwick HRI, University of Warwick, Wellesbourne, Warwick CV35 9EF, UK

8 Author for correspondence Prof. G.D. Foster, email Gary.Foster@bristol.ac.uk

9 Phone: +44-117-928-7474

10 Fax : +44-117-331-7985

11

12 Running title: Serine proteinase expression in basidiomycetes

1 **Abstract**

2 The *Agaricus bisporus* serine proteinase 1 (SPR1) appears to be significant in both
3 mycelial nutrition and senescence of the fruiting body. We report on the construction
4 of an SPR promoter::GFP fusion cassette (pGreen_hph1_SPR_GFP) for the
5 investigation of temporal and developmental expression of SPR1 in
6 homobasidiomycetes, and to determine how expression is linked to physiological and
7 environmental stimuli. Monitoring of *A. bisporus* pGreen_hph1_SPR_GFP
8 transformants on media rich in ammonia or containing different nitrogen sources,
9 demonstrated that SPR1 is produced in response to available nitrogen. In *A. bisporus*
10 fruiting bodies, GFP activity was localised to the stipe of postharvest senescing
11 sporophores. pGreen_hph1_SPR_GFP was also transformed into the model
12 basidiomycete *Coprinopsis cinerea*. Endogenous *C. cinerea* proteinase activity was
13 profiled during liquid culture and fruiting body development. Maximum activity was
14 observed in the mature cap, while activity dropped during autolysis. Analysis of the
15 *C. cinerea* genome revealed 7 genes showing significant homology to the *A. bisporus*
16 *SPR1* and *SPR2* genes. These genes contain the aspartic acid, histidine and serine
17 residues common to serine proteinases. Analysis of the promoter regions revealed at
18 least one CreA and several AreA regulatory motifs in all sequences. Fruiting was
19 induced in *C. cinerea* dikaryons and fluorescence determined in different
20 developmental stages. GFP expression was observed throughout the life cycle,
21 demonstrating that serine proteinase can be active in all stages of *C. cinerea* fruiting
22 body development. Serine proteinase expression (GFP fluorescence) was most
23 concentrated during development of young tissue, which may be indicative of high
24 protein turnover during cell differentiation.

25

- 1 **Index descriptors:** Basidiomycetes, *Agaricus bisporus*, *Coprinopsis cinerea*, serine
- 2 proteinase, green fluorescent protein (GFP), humic protein.

1 **1. Introduction**

2 *Agaricus bisporus*, the cultivated mushroom has economic and biotechnological
3 significance. It is the most extensively cultivated mushroom worldwide, with an
4 annual production in the region of 5 million tonnes (33) and is a major protected crop
5 in the UK accounting for 10% of such horticultural production (20). In addition to its
6 value as a food crop there is considerable interest in *A. bisporus* as a host for
7 molecular pharming of heterologous proteins (21, 51, 58, 62), and it also appears to
8 produce a number of compounds of potential biomedical/nutraceutical importance
9 (13). Application of biotechnology to *A. bisporus* has greatly increased due to the
10 development of a transformation system (14, 26) and recently Burns *et al.*, (2006)
11 developed an *A. bisporus* 'molecular toolkit' which tested different promoters for
12 efficient gene expression. Despite these recent advances, developmental studies in
13 *Agaricus* have been hampered due to the time and containment issues that exist when
14 studying a genetically modified strain.

15 The ink-cap mushroom, *Coprinopsis cinerea* (formally *Coprinus cinereus*), is a well-
16 studied homobasidiomycete (12, 43, 47) that forms an excellent model system for
17 studies of gene expression at several levels of differentiation, particularly mushroom
18 development and meiotic processes (46, 59). It has been used as an object for studies
19 of development (32) mainly because of its relatively short life cycle, which can be
20 completed in the laboratory within 2 weeks (44). In addition, genetic studies and
21 experimental manipulation of all phases of its life cycle are simple and relatively
22 straightforward (63). The *C. cinerea* genome sequence was released in 2003
23 (<http://www.broad.mit.edu>) and recently gene silencing has been demonstrated in the
24 basidiomycete (24, 47). We have exploited these characteristics of *C. cinerea* for the
25 investigation of a serine proteinase from *Agaricus bisporus*.

1 A serine proteinase (SPR1) has been purified from senescent sporophore tissue of *A.*
2 *bisporus*, which has a molecular mass of 27 kDa and an isoelectric point of 9.0 (11).
3 The protease has a broad pH optimum, 6.5-11.5, and a narrow substrate specificity,
4 requiring both a hydrophobic amino acid in the P1 position and a minimum peptide
5 chain length (11). The most active proteolysis of *A. bisporus* culture filtrate was
6 observed with Suc-Ala-Ala-Pro-Phe-pNA at neutral pH (10). Serine proteinase was
7 found to be the major proteinase produced by *A. bisporus* in sporophores during
8 senescence (9), and extracellular to mycelium in colonized compost where nitrogen is
9 largely in the form of protein suggesting a nutritional role for this enzyme (10). The
10 serine proteinase extracellular to mycelium was produced to a greater degree in
11 response to protein associated with humic substances than other pure proteins,
12 suggesting factors additional to the protein are involved in its induction. The cDNA
13 for this proteinase has been cloned and sequenced (accession no Y13805), which
14 revealed that this serine proteinase (SPR) belongs to the 'proteinase K family' (31).
15 The *SPR1* gene expression was not detected in freshly harvested mushrooms, while
16 increased transcript levels were observed 1-3 days post harvest. Expression of *SPR1*
17 was strongest in post harvest stipe tissue (31) which correlated well with the increase
18 in enzyme activity and protein level detected in senescent stipe (9). The relatively
19 high transcriptional and translational levels of SPR in the stipe demonstrate that the
20 enzyme is important during the metabolism of senescing mushrooms.

21 This paper reports on the construction of a promoter::GFP fusion cassette for the
22 investigation of the temporal and developmental expression of *SPR1* in *A. bisporus*
23 and *C. cinerea* and to characterise expression in response to physiological and
24 environmental stimuli. This paper further investigates the utility of *C. cinereus* as a
25 model system for basidiomycete gene expression and fruiting body production, as

1 development of a model species for basidiomycetes research is vital for future
2 progress.

3 **2. Materials and methods**

4 **2.1 Strains and culture maintenance**

5 *Escherichia coli* strain DH5 α was the host strain for recombinant plasmids.
6 *Agrobacterium tumefaciens* AGL1 (34) was used for *A. bisporus* transformations and
7 cultured as previously described (17, 26). The *A. bisporus* commercial strain A15 (18)
8 was used for transformations. Mycelia were routinely maintained at 25°C on MPA
9 (35) agar plates and supplemented with 25 μgml^{-1} hygromycinB to select for
10 transformants. A tryptophan auxotroph, LT2 (*A6B6*, *trp1.1;1.6*) (4) was used for *C.*
11 *cinerea* transformations. *C. cinerea* strains AT8 (*A43B43*, *trp-3*, *ade-8*) and
12 AmutBmut (*A43mutB43mut*, *pab1*) (41, 56) were used for fruiting studies. *C. cinerea*
13 mycelia were routinely maintained at 37°C on YMG agar (4) supplemented when
14 appropriate with 100 μgml^{-1} L-tryptophan.

15

16 **2.2 Construct design**

17 An 877bp *A. bisporus* *SPR* putative promoter region (5'UTR) was amplified from a
18 cosmid clone template using primers *spr1-fwd*
19 (TCCCCGCGGCGGGCTCAGAAGGTTTCTAT) and *spr1(rev)m*
20 (AAATCCATGGTCGGTGAAGAGATC) that respectively introduced 5' *SacII* and 3'
21 *NcoI* restriction sites. The resulting amplicon was cloned using pGEM-Teasy
22 (Promega Corp.) and *SPR1* promoter integrity confirmed by dsDNA sequencing of
23 recombinants. The *SPR1* promoter was cloned into a pBluescriptII based GFP
24 expression construct (pBlue-SPR-GFP) following removal of the *A. bisporus* *GPDII*
25 promoter (*SacII-NcoI* restriction) from an intron-GFP expression vector p004iGM (6).

1 The 1884bp SPR::GFP expression unit was excised by *SacI-KpnI* restriction and
2 ligated into the *ClaI-KpnI* restricted binary pGreen_hph1 (18) by addition of a *ClaI-*
3 *SacI* oligolinker (CGAGCT) to yield pGreen_hph1_SPR_GFP.

4

5 **2.3 Fungal transformations**

6 Plasmid DNA for fungal transformation was prepared using QIAGEN Midi Prep Kits.
7 *C. cinerea* protoplast co-transformations were performed as previously described (4,
8 6, 22, 24) using *ca.* 1µg of pCc1001 (*trp1*) (54) with 5µg of plasmid
9 pGreen_hph1_SPR_GFP. Trp⁺ transformants were maintained on *Coprinus*
10 regeneration agar (6, 16, 24). Putative transformants of *C. cinerea* were cultured as
11 described above and genomic DNA extracted as previously described (36). PCR
12 screening of *C. cinerea* transformants was performed using Reddymix components
13 (Abgene) with a general thermal cycling program of 95°C for 3 min, (95°C for 30 sec,
14 50°C for 1min, 72°C for 30 sec,) 30 cycles, 72°C for 10 min.

15 *A. bisporus* was transformed using *Agrobacterium tumefaciens* mediated transfection
16 of gill tissue as previously described (6, 7, 14, 35, 42). Transformants of *A. bisporus*
17 were identified using previously published methods (18, 35) and transcription of both
18 *hph* and GFP transgenes confirmed using rtPCR and/or quantitative rtPCR (24). *A.*
19 *bisporus* transformants for fruiting were selected from a large sample set by
20 fluorometric quantification of GFP activity in mycelia (24) following induction with
21 humic fraction (10).

22

23 **2.4 Fruiting studies**

24 Dikaryons of *C. cinerea* were produced on YMG^T plates by placing mycelial blocks
25 of AT8 and LT2 trp⁺ transformants 5mm apart at 37°C; dikaryotization was

1 confirmed by the presence of clamp cells. For growth and induction of fruiting bodies,
2 dikaryons on YMGT plates were incubated at 12 hr light/12 hr dark, 25°C, 90%
3 humidity under standard fruiting conditions (22, 37). *C. cinerea* strain AmutBmut was
4 selected as a control strain for fruiting studies as it exhibits clamp formation and fruit
5 body development like a dikaryon and produces uninucleate oidia like a monokaryon
6 (56). GFP expression in fruiting bodies was examined using a Leica MZFL111
7 microscope with SPOT 2.2.1 (Diagnostic Instruments inc.) imaging software.

8 *A. bisporus* sporophores were produced in small-scale compost cultures at the
9 University of Warwick's transgenic mushroom containment facility, harvested and
10 stored as previously described (18). GFP activity was measured in detached
11 mushrooms, 3-days post-harvest, using a portable GFP meter (ADC BioScientific
12 Ltd., UK; excitation 450nm/ emission 530nm/ gain setting 55). Metered readings were
13 recorded for both cap and stipe tissues of whole mushrooms and the freshly cut face
14 of longitudinally bisected sporophores. A minimum of three replicate readings was
15 taken for each sample tissue from two replicate sporophores. Sectioned mushrooms
16 were also viewed using a blue LED floodlight (Inova X5™, Emissive Energy, RI)
17 with appropriate blue/yellow filter sets (57) and photographed using a Nikon Coolpix
18 990.

19

20 **2.5 Proteinase assays**

21 A proteinase plate assay was carried out by inoculating *C. cinerea* LT2 onto
22 ammonium free regeneration agar (RA) plates containing 0.5% (w/v) skimmed milk
23 powder. To assess proteinase activity, colony size was measured, as well as the
24 clearing zone around each colony, produced by degradation of the milk layer by

1 extracellular proteinase activity. LT2 was inoculated onto standard RA media as a
2 control. Five replicate plates were measured per assay.

3 Expression of serine proteinase activity in liquid culture was determined by
4 inoculating LT2 into ammonium free RA containing 0.5% (w/v) milk solution and
5 into standard RA. Cultures were grown for 264 h and samples (8 ml) were aseptically
6 removed every 24 h and assayed. Serine proteinase activity was measured in fruiting
7 body developmental stages by homogenising fungal tissue in 50mM Tris buffer, pH
8 8.0 and centrifuging at 10,000g to remove particulate material.

9 Serine proteinase activity was assayed spectrophotometrically by absorbance at 405nm
10 following the release of p-nitroaniline from the synthetic peptide Suc-Ala-Ala-Pro-
11 Phe-pNA (0.15mM) in 50mM Tris buffer, pH 8.0. Hydrolysis was performed for 30
12 min at 37°C. Inhibition of serine proteinase was performed by pre-incubation of 0.1M
13 PMSF (Fluka) inhibitor with the enzyme at 37°C for 30 min. Soluble protein
14 concentrations were measured by the dye-binding method of Bradford (5). Bovine
15 serum albumin was used as a standard. Biochemical assays were performed in
16 triplicate.

17

18 **2.6 Sequence analysis**

19 Sequences were analysed by BLAST (NCBI) (1) and aligned using ClustalW (25).
20 The sequence manipulation suite (55) performed molecular weight and isoelectric
21 point prediction. Prosite was used to identify motifs and signature sequences in the
22 deduced protein sequences (3) and signal sequences were identified using SignalP
23 (48). Structural classification of sequences was based on SCOP (45). Transcription
24 factor binding sites were predicted using MOTIF search on Genome Net
25 (<http://motif.genome.jp/>).

1

2 **3. Results**

3 **3.1 Analysis of *A. bisporus* pGreen_hph1_SPR_GFP transformants**

4 To investigate the temporal and spatial expression of the *A. bisporus* *SPR1* gene, a
5 promoter::GFP fusion cassette was constructed. This expression vector was
6 engineered to contain a 5' intron, which has previously been shown to be necessary
7 for GFP expression in *A. bisporus* and *C. cinerea* (6). Plasmid
8 pGreen_hph1_SPR_GFP was transformed into *A. bisporus* via *A. tumefaciens* and
9 transformants were recovered on hygromycin selection. Nine transformants were
10 selected for further analysis. The presence of the intact expression cassette,
11 pGreen_hph1_SPR_GFP, was confirmed via PCR. Primers SPR1Fwd (5'-
12 CCGCGCAACATATGTATGTGAGAG-3') and GFPprev (5'-
13 GTGGCGGATCTTGAAGTTCACCTTG-3'), which bind 256bp downstream from
14 the 5' end of the *SPR1* promoter and 234bp upstream from the 3' end of the GFP gene
15 respectively, resulted in a 1226bp PCR product. Primers GFPFwd (5'-
16 GGCGTGCAGTGCTTCAGCCGC-3') and TrpCRev (5'-
17 GCACTCTTTGCTGCTTGGAC-3') which bind 222bp downstream from the 5' end
18 of the GFP gene and 146bp upstream from the 3' end of the TrpC terminator resulted
19 in a 665bp PCR product. Positive amplification of both fragments confirmed the
20 presence of the intact expression cassette. *A. bisporus* pGreen_hph1_SPR_GFP
21 transformants, wild type *A. bisporus* A15, and an *A. bisporus* strain expressing the
22 plasmid pGR4-4GiGM3' (G26) which contains GFP under the *A. bisporus* *GPD II*
23 promoter (6), were inoculated onto a range of media to investigate if changes in
24 nutrient availability would alter the expression of the proteinase which is know to be
25 involved in nutrient acquisition. GFP expression was monitored on media rich in

1 ammonia (YMG, MMP, and regeneration agar (RA)), potato dextrose agar (PDA),
2 and ammonia free regeneration agar containing one of the following sole nitrogen
3 sources; 0.094% (w/v) humic fraction, 0.084% (w/v) glutamic acid (GA), or 0.5%
4 (w/v) skimmed milk power. GFP expression was observed in the
5 pGreen_hph1_SPR_GFP transformants grown on humic fraction, milk, GA and PDA
6 media, while no GFP expression was observed on YMG, MMP and RA (Table 1).
7 Fig. 1A shows expression of GFP in an *A. bisporus* SPR::GFP transformant (TP17)
8 on ammonia free regeneration agar containing 0.094% humic fraction, and its
9 repression on standard regeneration media. As expected the GPD::GFP control
10 transformant (G26) exhibited strong GFP expression on all media, whilst GFP
11 fluorescence was not observed on any media with the wild type strain (Table 1).

12

13 **3.2 Monitoring of *SPR1* expression in *A. bisporus* sporophore development**

14 Fruiting was induced in *A. bisporus* transformants and GFP expression was detected
15 using blue LED illumination of bisected mushrooms (Fig 1B). Transformants for
16 fruiting were selected from a large sample set by fluorometric quantification of GFP
17 activity in mycelia (23) following induction with humic fraction (10). TP196 was
18 selected as a typical phenotypic representative of transformants, which also exhibited
19 excellent culture and fruiting capabilities. GFP expression was clearly observed in
20 both the cap and stipe tissues of freshly harvested (day 0) *A. bisporus* G26 fruiting
21 bodies, expressing GFP under the control of the GPD promoter (Fig. 1B: Panel A). In
22 senescing mushrooms (3d post harvest) no GFP expression was observed in a
23 hygromycin resistant (control) transformant of A15hph (no GFP cassette, Fig. 1B:
24 Panel B: leftmost mushroom), while GFP expression was clearly detected in the stipe
25 tissue of SPR::GFP transformant TP196 (Fig. 1B: Panel B: rightmost mushroom).

1 Metered readings (relative fluorescence units (RFUs)) for cap and stipe tissues of
2 whole and longitudinally bisected mushrooms of TP196 (*SPR::GFP*), G26
3 (*GPD::GFP*) and A15hph (no GFP cassette) were recorded 3 days post harvest (Fig.
4 2). GFP activity was substantially elevated in the stipes of senescing mushrooms for
5 the *SPR::GFP* transformant TP196. The tissue (stipe) specific expression of GFP in
6 TP196 is consistent with earlier histochemical observations of SPR activity in
7 senescing mushrooms (9). RFUs recorded for G26 represent background fluorescence
8 of the fruiting body, while A15hph exhibits a slight increase in RFUs compared to
9 G26 due to autofluorescence.

10

11 **3.3 Expression profiles of serine proteinases in *C. cinerea* LT2**

12 Endogenous proteinase activity was assessed by inoculating LT2 onto ammonium free
13 regeneration agar (RA) with and without a 0.5% (w/v) milk solution. Clearing zones,
14 indicative of proteinase activity, were only produced on media containing the milk
15 overlay (Fig. 3A). A proteinase expression profile was developed for LT2 grown in
16 broth by measuring the hydrolysis of the synthetic peptide Suc-Ala-Ala-Pro-Phe-
17 pNA. Proteinase activity was observed in LT2 cultures grown in ammonium free RA
18 containing 0.5% (w/v) milk solution after 120 h (0.0259 AU/mL) and continued to
19 increase until 240 h (1.283 AU/mL) (Fig. 3B). A small decrease in activity was
20 observed at 264 h but increased again at 288 h. Pre-incubation of the crude enzyme
21 extracts with the serine proteinase inhibitor PMSF resulted in a large decrease in
22 activity (from 1.283 AU/mL to 0.17 AU/mL at 240 h), thus confirming that the
23 majority of proteinase activity detected was the serine mechanistic class. Little or no
24 proteinase activity was observed in LT2 cultures grown in standard RA media, which
25 is rich in ammonia (Fig. 3B).

1 Serine proteinase activity was measured during the primordium, karyogamy, meiosis,
2 immature, mature and autolysis stages of fruiting body development (Fig. 3C).
3 Activity increased slowly from the primordium (1.29 units/g) to the meiosis stage
4 (1.69 units/g) with a slight dip at immature (1.52 units/g) followed by a large increase
5 in activity during the mature development stage (6.32 units/g). Maximum activity was
6 detected in the mature cap (6.32 units/g) followed by a decrease in activity during
7 autolysis (3.45 units/g). Similarly, pre-incubation of the crude enzyme extracts with
8 the inhibitor PMSF resulted in a large decrease in activity (from 6.32 units/g to 0.34
9 units/g in the mature cap), demonstrating that the class of proteinase activity detected
10 was serine proteinase (Fig. 3C).

11

12 **3.4 Identification and sequence analysis of homobasidiomycete serine proteinases**

13 Following confirmation of endogenous serine proteinase activity in *C. cinerea*,
14 identification of the encoding genes was undertaken using the published *C. cinerea*
15 genome sequence. Two *A. bisporus* serine proteinases have been previously identified
16 (*SPR1*, *SPR2*) and their sequence deposited in public databases under accession
17 numbers Y13805 and AJ344211 respectively (30, 31). Predicted molecular weights
18 (Table 2), for full-length *SPR1* and *SPR2* are considerably larger than the ca. 27kDa
19 experimental estimates from SDS-PAGE, cDNA and N-terminal amino acid
20 sequencing. Mature proteins for *SPR1* (286 aa, 28.29kDa) and *SPR2* (275 aa,
21 27.70kDa) are much closer to the 27kDa estimate previously observed (11, 31). Blast
22 analysis (1) of the *A. bisporus SPR1* and *SPR2* genes against the *C. cinerea* database
23 revealed 7 genes (04562.1, 10592.1, 10615.1, 07792.1, 10606.1, 0.3122.1 & 04470.1)
24 showing significant homology to the serine proteinases. ClustalW alignments of these
25 *C. cinerea* genes with the *A. bisporus SPR1* revealed amino acid sequence identity

1 values ranging between 44% and 61% while homology of the *SPR2* with the *C.*
2 *cinerea* genes ranged between 42% and 55% (Table 3). *SPR1* and *SPR2* have an
3 amino acid identity value of 75% while the *C. cinerea* genes have homology ranging
4 between 31% and 77% (Table 3).

5 Three motifs were identified within the *C. cinerea* genes that are common to other
6 serine proteinases; the aspartic acid residue (consensus: [STAIV]-X-[LIVMF]-
7 [LIVM]-D-[DSTA]-G-[LIVMFC]-X(2,3)-[DNH], the histidine residue (consensus: H-
8 G-[STM]-X-[VIC]-[STAGC]-[GS]-X-[LIVMA]-[STAGCLV]-[SAGM]) and the
9 serine residue (consensus: G-T-S-X-[SA]-X-P-X(2)-[STAVC]-[AG]) (31). These
10 residues were conserved between the *C. cinerea* genes and the *A. bisporus SPR1* and
11 *SPR2*, with the exception of the *C. cinerea* gene 10606.1 that lacked the serine
12 residue. This suggests that the *C. cinerea* genes are serine proteinases and they appear
13 to belong to the subtilisin family.

14 The probable *C. cinerea* serine proteinase genes ranged between 346 and 500 amino
15 acids in length (Table 2), and all contained introns. Each intron began with GT and
16 ended with AG, which is a common feature of fungal introns and has been observed
17 in the serine proteinase genes from *Acremonium chrysogenum* (28), *Lecanicillium*
18 *psalliotae* (60) and *Arthrobotrys conoides* (61). The number of introns varied between
19 2 and 14 depending on the gene (Table 2), and some conservation of intron position
20 was observed between the *C. cinerea* genes and *SPR2*.

21 The theoretical molecular weight and isoelectric points for the *C. cinerea SPR* genes
22 range between 35kDa and 53kDa and 5.83 and 9.97 respectively (Table 2), while the
23 theoretical molecular mass and isoelectric points for *SPR1* and *SPR2* are 39.39 kDa
24 and 5.93 and 38.85 kDa and 5.53 respectively (Table 2). A predicted signal peptide
25 was observed in the *C. cinerea* and *A. bisporus* serine proteinases, with cleavage

1 occurring either between amino acids 19 and 20, 20 and 21 or 21 and 22, suggesting
2 that these enzymes are secreted. Using the highest homology sequences, the predicted
3 secondary structure of these genes is composed of between 20-30% helices, 16-35%
4 strands and 42-61% loops (Table 2) and analysis of the degree of protein globularity
5 suggests that these enzymes exist as compact (globular) domains.

6 One kb of sequence upstream from the ATG start codon of each gene was analysed
7 for the presence of regulatory motifs. At least one CreA and several Nit2/AreA
8 regulatory elements were identified in the promoter regions of the *C. cinerea* and *A.*
9 *bisporus* genes (Table 2). No other regions of homology were detected between the
10 promoters.

11

12 **3.5 Analysis of *C. cinerea* pGreen_hph1_SPR_GFP transformants**

13 From a preliminary screen of one hundred Trp+ co-transformants on RA media (rich
14 in ammonia), and on ammonia free regeneration agar containing 0.094% (w/v) humic
15 fraction as the sole nitrogen source, 32% of transformants were found to express GFP
16 on humic fraction, which correlates well with the reported rate of co-transformation
17 (30-49% (6)). However, GFP expression was not observed on RA media. Four GFP+
18 transformants, non-transformed LT2, and a *C. cinerea* strain (PG78Gr) expressing
19 GFP under the regulation of the *A. bisporus GPDII* promoter (24), were selected for
20 further studies. GFP expression was monitored on media rich in ammonia (YMG, and
21 regeneration agar (RA)), potato dextrose agar (PDA), and ammonia free regeneration
22 agar containing either 0.094% (w/v) humic fraction, 0.084% (w/v) glutamic acid
23 (GA), or 0.5% (w/v) milk as the sole nitrogen source. LT2 exhibited no fluorescence
24 on any media while PG78Gr expressed GFP on all the media. GFP fluorescence was
25 observed in transformants grown on humic fraction, milk, GA and potato dextrose

1 media while no GFP fluorescence was observed on RA media (Table 1). The only
2 transformant to exhibit fluorescence on YMG media was T47.

3 *C. cinerea* transformants were mated with AT8 and the dikaryons inoculated onto a
4 range of media and screened for GFP expression. Similar expression profiles were
5 observed for both the monokaryons and dikaryons (Table 1). Fig. 4A shows *C.*
6 *cinerea* transformant T47 monokaryon and dikaryon expression of GFP on ammonia
7 free regeneration agar containing 0.094% (w/v) humic fraction, and repression of GFP
8 on standard regeneration media.

9

10 **3.6 Monitoring of *SPRI* expression in *C. cinerea* fruiting body development**

11 Fruiting was induced in the *C. cinerea* dikaryon strains and the control strain
12 AmutBmut. Different developmental stages of the fruiting body were examined
13 microscopically for fluorescence. Low levels of fluorescence were observed in the
14 hyphal knot (Fig 4B: Panel A). Fluorescence was also observed in the primordium
15 stage but was not localised (Fig 4B: Panel B). A similar observation was made for
16 karogamy stage but GFP localisation began to occur at the edge of the forming gill
17 tissue (Fig 4B: Panel C). GFP appeared more localised at the forming gill tissue
18 during meiosis (Fig 4B: Panel D), while at the immature stage GFP was observed
19 high up in the stipe close to the cap (Fig 4B: Panel E). In mature sporophores,
20 fluorescence was observed in the cap but was most concentrated at the junction of the
21 stipe and cap (Fig 4B: Panel F), while fluorescence was reduced in the stipe (Fig 4B:
22 Panel G). During autolysis fluorescence was greatly reduced in the cap (Fig 4B: Panel
23 H) but was concentrated in the stipe tissue (Fig 4B: Panel I). In the control strain
24 AmutBmut some autofluorescence was observed throughout the different

1 developmental stages. Fig 4C depicts a schematic of GFP fluorescence under the
2 control of the *A. bisporus SPR1* promoter through out the *C. cinerea* life cycle.

3

4 **4. Discussion**

5 *A. bisporus* SPR1 has previously been shown to be significant in both mycelial
6 nutrition and senescence of the mushroom fruit body (8-10). This study used an
7 SPR::GFP fusion construct to investigate temporal and developmental expression of
8 *SPR1* in *A. bisporus* and a heterologous host *C. cinerea* in response to physiological
9 and environmental stimuli. Developmental studies in *A. bisporus* are still hindered due
10 to the time and containment issues that exist when studying a genetically modified
11 strain. *C. cinerea* provides a model system for the studies of gene expression
12 throughout mushroom development (47, 59) and heterologous expression of the *A.*
13 *bisporus* *SPR1* promoter fusion is a further demonstration of the inkcap host utility as
14 a model species.

15 GFP has been widely used as a reporter molecule or as a fluorescent tag for fusion
16 proteins (53) and is now a valuable tool in the molecular analysis of filamentous fungi
17 (38). The use of GFP in ascomycete fungi has been widely reported (2, 27, 49), and
18 recently expression in homobasidiomycetes has also been achieved (6, 39, 40).

19 The aim of this study was to carry out a comparative molecular analysis of serine
20 proteases in both *C. cinereus* and *A. bisporus*. To this end, identification of *C.*
21 *cinereus* SPR genomic sequences was performed to establish the homology between
22 Agaricus and Coprinus SPR genes. Bioinformatics was employed to help predict if the
23 genes would be regulated in a similar fashion, thus providing evidence for the
24 suitability of *C. cinereus* as a heterologous host for *A. bisporus* SPR1. Blast analysis
25 of the *A. bisporus* *SPR1* cDNA and *SPR2* genomic sequence against the *C. cinerea*

1 database revealed 7 genes showing significant homology. Conservation of the aspartic
2 acid, histidine and serine residues within the genes suggested that they are serine
3 proteinases belonging to the subtilisin family. However, lack of a serine residue at the
4 active site in 10606.1 suggests that some of these are 'pseudogenes' that would be
5 unable to code for active enzymes. Signal peptide analyses were indicative of
6 extracellular activity and protein globularity infers that the enzymes would exist as
7 compact globular domains. Sequencing of the *A. bisporus* genome is currently
8 underway (<http://www2.warwick.ac.uk/fac/sci/whri/research/agaricusgenome/>) which
9 may reveal further SPR homologues.

10 All the *C. cinerea* genes contained introns with numbers varying between 2 and 14;
11 only two genes had less than 10 introns, six of the seven analysed contained between
12 11-14 introns. Short exons and high intron density in basidiomycetes and the
13 comparatively poor conservation of intron splice sequences compared with other
14 fungi can result in some inaccuracies when using intron predictive software. This may
15 account for the low number of introns identified in CC1G_10615.1 and
16 CC1G_04470.1.

17 In *A. bisporus*, two serine proteinases (*SPR1* and *SPR2*) were isolated from the same
18 cosmid clone, within 30kb of each other (30). Similarly, three *C. cinerea* serine
19 proteinases (CC1G_10592.1, CC1G_10606.1 and CC1G_10615.1) lay within 50kb of
20 each other on the genome suggesting either local duplication or a common ancestor.
21 Conservation of intron positions observed in these genes is indicative of local
22 duplications.

23 Endogenous protease activity was investigated in *C. cinereus*. A preliminarily plate
24 based assay resulted in a clearing zone around the fungal colonies thus confirming the
25 presence of proteases in the basidiomycete. As previously demonstrated in *A. bisporus*

1 (10), little or no serine proteinase activity was detected in *C. cinerea* cultures grown
2 in ammonia rich media. Activity was observed in cultures grown on ammonia free RA
3 containing milk as the sole nitrogen source after 120 h and continued to increase until
4 240 h with a slight decrease at 264 h before increasing again at 288 h which may be
5 indicative of the onset of autolysis.

6 Expression of SPR1 in response to physiological and environmental stimuli was
7 examined by inoculating the *A. bisporus* and *C. cinerea* pGreen_hph1_SPR_GFP
8 transformants onto a range of media to investigate if changes in nutrient availability
9 would alter the expression of the proteinase. At least one CreA and several Nit2/AreA
10 transcription factor-binding sites were identified in both the *A. bisporus* and *C.*
11 *cinereus* SPR promoter sequences, signifying regulation by factors such as carbon and
12 nitrogen sources. Conservation of these sites was not observed across the promoters.

13 Experimental evidence for the regulation of serine proteinases in response to nitrogen
14 sources is provided from *C. cinerea* biochemical profiles in broth culture; serine
15 proteinase was not detected on ammonia rich RA media but was observed on
16 ammonia free RA supplemented with milk. GFP expression was observed in *A.*
17 *bisporus* and *C. cinereus* transformants grown on PDB and on ammonia free RA
18 containing humic fraction, milk, or glutamate as the sole nitrogen source. GFP
19 expression was not observed on YMG, MMP or regeneration media (rich in
20 ammonium), with the exception of *C. cinereus* transformant TP47. GFP expression
21 was observed in TP47 grown on YMG media, which may result from multiple
22 insertion events, however this expression profile was atypical of the population of *C.*
23 *cinereus* transformants analysed. Expression profiles were similar for both
24 monokaryons and dikaryons. Collectively these results suggest that both *C. cinerea*
25 and *A. bisporus* produce serine proteinases in response to available nitrogen.

1
2 Developmental regulation of serine proteinase expression was investigated. Serine
3 proteinase activity has previously been reported during fruiting body development of
4 *A. bisporus* (9). From stages 2 to 6 of development (23), activity was relatively low
5 and cap and stipe activities were similar. *A. bisporus* developmental stages 2-6
6 roughly correspond to the primordium, karyogamy, meiosis, immature and mature
7 stages of *C. cinereus* development. In our SPR biochemical assays, activity was
8 relatively low in the first four stages of *C. cinerea* development but increased rapidly
9 at the mature stage. GFP expression was ubiquitous in the primordium stage, which
10 may be the result of a higher density of cytoplasm in the developing primordium. GFP
11 expression was observed throughout the karyogamy and meiosis stages though
12 localization of fluorescence began to occur at the edge of the forming gill tissue at the
13 karyogamy stage and became more pronounced at the meiosis stage. *C. cinerea* is
14 described as having a ruphymenial mode of hymenophore development, where the
15 gill is envisaged as widening towards the periphery of the cap as a differentiating
16 front moves into, and differentiates from, the basidiocarp (50). As the widest part of
17 the gills is those at the cap margin, the differentiating front is also moving upwards
18 towards the apex of the cap (52). GFP fluorescence was most concentrated at the base
19 of the gills in the karyogamy stage and moved upwards towards the apex of the cap in
20 meiosis suggesting that *SPRI* promoter activity was enhanced during the development
21 of young tissue, which may be indicative of high protein turnover during cell
22 differentiation. This could also result from autolysis of connective tissue as the gills
23 begin to separate from the stipe i.e. creating an abscission zone. At the immature stage
24 GFP was observed high up in the stipe close to the cap. Studies of *C. cinerea* stipe
25 elongation have revealed that it is variable along its length and that elongation is

1 greatest at the mid-upper portion (the stipe that is enclosed by the developing cap); the
2 apex and base of the stipe shows little elongation (15, 29). The rapid increase in
3 length is chiefly due to cellular elongation (29) but divisions also contribute, with
4 cells doubling in number and increasing six to eight fold in length (19). The
5 fluorescence observed in the mid-upper stipe demonstrates that the *SPRI* promoter is
6 activity upregulated during elongation, and is likely to support the elongating stipe by
7 providing free amino acids via protein degradation. Highest activity was recorded in
8 the mature cap with slightly less activity in the mature stipe, contrasting to the levels
9 recorded for *A. bisporus*. In the mature fruiting body GFP fluorescence was observed
10 in the cap but was most concentrated at junction of the stipe and cap. This may result
11 from a high density of cells where younger tissue is still developing resulting in
12 elevated protein turnover. With *A. bisporus* developmental stage 7, a large increase in
13 activity in the stipe and a small increase in the cap occurs (9), and further increases
14 are observed as stage 7 mushrooms progress to senescence. During *C. cinerea*
15 autolysis serine proteinase activity decreased and fluorescence was greatly reduced in
16 the cap but was highly concentrated in stipe tissue. Accumulation of serine proteinase
17 in the stipe during autolysis would suggest a role in the export of nutrients from the
18 stipe to the cap tissue during senescence. Similarly in *A. bisporus* sporophores,
19 highest *SPR::GFP* activity was observed in senescing stipe tissues, suggesting that the
20 stipe may act as an 'active source' during the export of nutrients to reproductive spore-
21 bearing tissues.

22

23 The results reported here confirm that the *A. bisporus* (*SPRI*) promoter is able to
24 regulate mycelial serine proteinase production in response to specific nitrogen sources
25 and have demonstrated tissue specific (stipe localised) expression in detached

1 sporophores. Use of the SPR::GFP fusion construct, coupled with genome data-
2 mining, suggests that serine proteinases also play an integral part in the development
3 of *C. cinerea* sporophores. The approaches developed in this study should underpin
4 further promoter analysis in these homobasidiomycete mushrooms and may permit
5 characterisation of promoter elements that regulate differential expression and
6 nutritional regulation of serine proteinases. Furthermore *C. cinereus* has been
7 validated as a potential model for expression and regulation studies of *A. bisporus*
8 genes.

9

10

11

12 **Acknowledgements**

13 The authors acknowledge Chris Thorogood for production of the *C. cinerea* lifecycle
14 illustration. Research at Universities of Bristol and Warwick was funded by grants
15 from BBSRC and DEFRA. The authors thank Nicholas Royat who helped screen
16 numerous *Agaricus bisporus* transformants.

1 **Table 1.** Evaluation of GFP fluorescence in *A. bisporus* and *C. cinerea* monokaryon
2 and dikaryon pGreen_hph1_SPR_GFP transformants on a range of different media.
3 Control strains included non transformed *A. bisporus* (A15) and *C. cinereus* (LT2)
4 and *A. bisporus* (G26) and *C. cinereus* (PG78Gr) strains expressing GFP under the *A.*
5 *bisporus* *GPD II* promoter. GFP expression was monitored on media rich in ammonia
6 (YMG, MMP, RA), potato dextrose agar (PDA), and ammonia free regeneration agar
7 containing 0.094% (w/v) humic fraction, 0.084% (w/v) glutamic acid (GA), or 0.5%
8 (w/v) milk as sole sources of nitrogen.

9

10 **Table 2.** Sequence analysis of the *A. bisporus* and the predicted *C. cinerea* serine
11 proteinases and promoter regions. *A. bisporus* genes: *SPR1* & *SPR2*. *C. cinerea* genes:
12 04562.1, 10592.1, 10615.1, 07792.1, 10606.1, 0.3122.1 & 04470.1

13

14 **Table 3.** Percentages of amino acid sequence identity between the *A. bisporus* and the
15 predicted *C. cinerea* serine proteinases. *A. bisporus* genes: *SPR1* & *SPR2*. *C. cinerea*
16 genes: 04562.1, 10592.1, 10615.1, 07792.1, 10606.1, 0.3122.1 & 04470.1

17

1 **Fig 1. A:** GFP expression in the *A. bisporus* SPR::*GFP* transformant TP17 when
2 grown on regeneration agar with or without 0.094% humic fraction under phase
3 contrast microscopy and UV light. Actively growing mycelia were examined using
4 40× objective on a Leitz Dialux 20 research microscope with excitation filters at 450–
5 490 nm, dichroic filter at 510 nm, and emission filter at 515 nm. Images clearly show
6 GFP fluorescence in TP17 grown on humic fraction while no fluorescence was
7 observed when grown on RA. **B:** Stipe localised GFP fluorescence in *A. bisporus*
8 transformant TP196. Fruiting was induced in *A. bisporus* transformants and bisected
9 mushrooms viewed under white light (WL) and blue LED illumination (BL). Panel A:
10 Images clearly show fluorescence in both the cap and stipe tissues of freshly harvested
11 (day 0) *A. bisporus* G26 fruiting bodies, expressing GFP under the control of the GPD
12 promoter. Panel B: In senescing mushrooms (3d post harvest) no GFP expression was
13 observed in a hygromycin resistant transformant of A15 (no GFP cassette, leftmost
14 mushroom), while GFP expression was clearly detected in the stipe tissue of
15 SPR::*GFP* transformant TP196 (rightmost mushroom).

16

17 **Fig 2.** GFP activity in senescing *A. bisporus* sporophores. Metered readings (relative
18 fluorescence units) are presented for cap and stipe tissues of whole and longitudinally
19 bisected mushrooms 3 days post harvest. GFP activity was substantially elevated in
20 the stipes of senescing mushrooms for the SPR::*GFP* transformant TP196 compared
21 with the control transformants, GPD::*GFP* (G26) and A15hph (no GFP cassette).

22

23 **Fig 3.** Proteinase profiles in *C. cinerea*. A: Proteinase plate assay of *C. cinerea*. 7mm
24 plugs of *C. cinerea* LT2 were inoculated onto RA and ammonium free RA plates
25 containing 0.5% (w/v) milk solution. Proteinase production was measured by the

1 clearing zone produced around the colony. B: Expression of serine proteinase activity
2 in culture filtrates during growth of *C. cinerea* LT2 in RA and ammonium free RA
3 containing 0.5% (w/v) milk solution. Cultures were grown for 264 h and samples (8
4 ml) were aseptically removed every 24 h and assayed using the synthetic peptide Suc-
5 Ala-Ala-Pro-Phe-pNA. C: Proteinase activity during *C. cinerea* AmutBmut
6 sporophore development as determined using the Suc-Ala-Ala-Pro-Phe-pNA substrate
7 in the presence or absence of inhibitor.

8

9 **Fig 4. A:** Expression of GFP in *C. cinerea* T47 monokaryon and dikaryon on
10 ammonia free regeneration agar containing 0.094% (w/v) humic fraction and on
11 standard regeneration media (RA) viewed under phase contrast (PC) microscopy and
12 UV light. Mycelia on actively growing plates were examined microscopically using
13 40× objective on a Leitz Dialux 20 research microscope with excitation filters at 450–
14 490 nm, dichroic filter at 510 nm, and emission filter at 515 nm. Images clearly show
15 GFP fluorescence in both TP47 monokaryon and dikaryon grown on humic fraction
16 while no fluorescence was observed in transformants grown on RA. **B:** Expression of
17 GFP in the *C. cinerea* developing fruiting body. Fruiting was induced in the dikaryon
18 *C. cinerea* TP24 mated with AT8 and GFP fluorescence was monitored in the hyphal
19 knot, primordium, karyogamy, meiosis, immature, mature and autolysis stages of
20 development. Fruiting was induced in *C. cinerea* AmutBmut and fruiting body stages
21 were also screened for GFP expression as a control. Samples were viewed under
22 phase contrast (PC) microscopy and UV light. **C:** Schematic illustration of GFP
23 fluorescence under the control of the *A. bisporus* *SPR1* promoter through out the *C.*
24 *cinerea* life cycle.

1 References

- 2
- 3
- 4 1. **Altschul, S. F., T. L. Madden, A. A. Schaffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman.** 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* **25**:3389-402.
- 5
- 6
- 7
- 8 2. **Amey, R. C., A. Athey-Pollard, C. Burns, P. R. Mills, A. Bailey, and G. D. Foster.** 2002. PEG-mediated and Agrobacterium-mediated transformation in the mycopathogen *Verticillium fungicola*. *Mycol Res* **106**:4-11.
- 9
- 10
- 11 3. **Bairoch, A., P. Bucher, and K. Hofmann.** 1997. The PROSITE database, its status in 1997. *Nucleic Acids Res* **25**:217-21.
- 12
- 13 4. **Binnering, D. M., C. Skrzynia, P. J. Pukkila, and L. A. Casselton.** 1987. DNA-Mediated Transformation of the Basidiomycete *Coprinus cinereus*. *Embo J* **6**:835-840.
- 14
- 15
- 16 5. **Bradford, M. M., and W. L. Williams.** 1976. New, Rapid, Sensitive Method for Protein Determination. *Federation Proceedings* **35**:274-274.
- 17
- 18 6. **Burns, C., K. E. Gregory, M. Kirby, M. K. Cheung, M. Riquelme, T. J. Elliott, M. P. Challen, A. Bailey, and G. D. Foster.** 2005. Efficient GFP expression in the mushrooms *Agaricus bisporus* and *Coprinus cinereus* requires introns. *Fungal Genet. Biol.* **42**:191-199.
- 19
- 20
- 21
- 22 7. **Burns, C., K. M. Leach, T. J. Elliott, M. P. Challen, G. D. Foster, and A. Bailey.** 2006. Evaluation of Agrobacterium-Mediated Transformation of *Agaricus bisporus* Using a Range of Promoters Linked to Hygromycin Resistance. *Mol Biotechnol* **32**:129-138.
- 23
- 24
- 25
- 26 8. **Burton, K. S., M. E. Love, and J. F. Smith.** 1993. Biochemical Changes Associated with Mushroom Quality in *Agaricus* Spp. *Enzyme Microb. Technol.* **15**:736-741.
- 27
- 28
- 29 9. **Burton, K. S., M. D. Partis, D. A. Wood, and C. F. Thurston.** 1997a. Accumulation of serine proteinase in senescent sporophores of the cultivated mushroom, *Agaricus bisporus*. *Mycological Research* **101**:146-152.
- 30
- 31
- 32 10. **Burton, K. S., J. F. Smith, D. A. Wood, and C. F. Thurston.** 1997b. Extracellular proteinases from the mycelium of the cultivated mushroom *Agaricus bisporus*. *Mycol Res* **101**:1341-1347.
- 33
- 34
- 35 11. **Burton, K. S., D. A. Wood, C. F. Thurston, and P. J. Barker.** 1993. Purification and Characterization of a Serine Proteinase from Senescent Sporophores of the Commercial Mushroom *Agaricus bisporus*. *J Gen Microbiol* **139**:1379-1386.
- 36
- 37
- 38
- 39 12. **Casselton, L. A., and N. S. Olesnicky.** 1998. Molecular genetics of mating recognition in basidiomycete fungi. *Microbiol. Mol. Biol. Rev.* **62**:55-70.
- 40
- 41 13. **Chen, S.** 2004. Anti-cancer properties of white button mushrooms. *Mushroom Science* **16**:25-29.
- 42
- 43 14. **Chen, X., M. Stone, C. Schlaghauser, and C. P. Romaine.** 2000. A fruiting body tissue method for efficient Agrobacterium-mediated transformation of *Agaricus bisporus*. *Appl Environ Microbiol* **66**:4510-3.
- 44
- 45
- 46 15. **Cox, R. J., and D. J. Niederpruem.** 1975. Differentiation in *Coprinus lagopus*. III. Expansion of excised fruit-bodies. *Arch Microbiol* **105**:257-60.
- 47
- 48 16. **Cummings, W. J., M. Celerin, J. Crodian, L. K. Brunick, and M. E. Zolan.** 1999. Insertional mutagenesis in *Coprinus cinereus*: use of a dominant
- 49

- 1 selectable marker to generate tagged, sporulation-defective mutants. *Curr*
2 *Genet* **36**:371-82.
- 3 17. **de Groot, M. J., P. Bundock, P. J. Hooykaas, and A. G. Beijersbergen.**
4 1998. *Agrobacterium tumefaciens*-mediated transformation of filamentous
5 fungi. *Nat Biotechnol* **16**:839-42.
- 6 18. **Eastwood, D. C., M. P. Challen, C. Zhang, H. Jenkins, J. Henderson, and**
7 **K. S. Burton.** 2008. Hairpin-mediated down-regulation of the urea cycle
8 enzyme argininosuccinate lyase in *Agaricus bisporus*. *Mycol Res* **112**:708-16.
- 9 19. **Eilers, F. I.** 1974. Growth-Regulation in *Coprinus radiatus*. *Arch. Microbiol.*
10 **96**:353-364.
- 11 20. **Elliott, T. J.** 1997. Mushrooms, p. 8-9, Society for General Microbiology
12 Quarterly.
- 13 21. **Elliott, T. J., K. S. Burton, and M. P. Challen.** 2001. Presented at the
14 Molecular Farming, OECD workshop, La Grande Motte (France), 2001.
- 15 22. **Granado, J. D., K. KerteszChaloupkova, M. Aebi, and U. Kües.** 1997.
16 Restriction enzyme-mediated DNA integration in *Coprinus cinereus*.
17 *Molecular & General Genetics* **256**:28-36.
- 18 23. **Hammond, J. B. W., and R. Nichols.** 1976. Carbohydrate-Metabolism in
19 *Agaricus bisporus* (Lange) Sing - Changes in Soluble Carbohydrates during
20 Growth of Mycelium and Sporophore. *J Gen Microbiol* **93**:309-320.
- 21 24. **Heneghan, M. N., A. Costa, M. P. Challen, P. R. Mills, A. Bailey, and G.**
22 **D. Foster.** 2007. A comparison of methods for successful triggering of gene
23 silencing in *Coprinus cinereus*. *Mol Biotechnol* **35**:283-296.
- 24 25. **Higgins, D., J. Thompson, and T. Gibson.** 1994. CLUSTAL W: improving
25 the sensitivity of progressive multiple sequence alignment through sequence
26 weighting, position-specific gap penalties and weight matrix choice. *Nucleic*
27 *Acids Res* **22**:4673-4680.
- 28 26. **Hooykaas, P. J. J., C. Roobol, and R. A. Schilperoort.** 1979. Regulation of
29 the Transfer of Ti Plasmids of *Agrobacterium tumefaciens*. *J Gen Microbiol*
30 **110**:99-109.
- 31 27. **Horowitz, S., S. Freeman, and A. Sharon.** 2002. Use of green fluorescent
32 protein-transgenic strains to study pathogenic and nonpathogenic lifestyles in
33 *Colletotrichum acutatum*. *Phytopathology* **92**:743-749.
- 34 28. **Isogai, T., M. Fukagawa, H. Kojo, M. Kohsaka, H. Aoki, and H. Imanaka.**
35 1991. Cloning and nucleotide sequences of the complementary and genomic
36 DNAs for the alkaline protease from *Acremonium chrysogenum*. *Agric Biol*
37 *Chem* **55**:471-7.
- 38 29. **Kamada, T., and T. Takemaru.** 1977. Stipe Elongation during Basidiocarp
39 Maturation in *Coprinus macrorhizus* - Mechanical-Properties of Stipe Cell-
40 Wall. *Plant Cell Physiol.* **18**:831-840.
- 41 30. **Kingsnorth, C. S., and K. S. Burton.** 1998. Genetic regulation of sporophore
42 senescence., p. 94-98. *In* L. J. L. D. Van Griensven and J. Visser (ed.),
43 *Proceedings of the Fourth Meeting on Genetics & Cellular Biology of*
44 *Basidiomycetes.* Mushroom Experimental Station, Horst, Netherlands,
45 Nijmegen, The Netherlands.
- 46 31. **Kingsnorth, C. S., D. C. Eastwood, and K. S. Burton.** 2001. Cloning and
47 postharvest expression of serine proteinase transcripts in the cultivated
48 mushroom *Agaricus bisporus*. *Fungal Genet Biol* **32**:135-44.
- 49 32. **Kües, U.** 2000. Life history and developmental processes in the basidiomycete
50 *Coprinus cinereus*. *Microbiol. Mol. Biol. Rev* **64**:316-53.

- 1 33. **Kües, U., and Y. Liu.** 2000. Fruiting body production in Basidiomycetes.
2 Appl Microbiol Biotechnol **54**:141-52.
- 3 34. **Lazo, G. R., P. A. Stein, and R. A. Ludwig.** 1991. A DNA transformation-
4 competent Arabidopsis genomic library in Agrobacterium. Biotechnology (N
5 Y) **9**:963-7.
- 6 35. **Leach, K., V. Odon, C. Zhang, H. K. Kim, J. Henderson, P. Warner, M.
7 Challen, and T. Elliott.** 2004. Progress in *Agaricus bisporus* transformation:
8 Agrobacterium methodologies and development of novel marker genes.
9 Mushroom Science **16**:93-102.
- 10 36. **Liu, D., S. Coloe, R. Baird, and J. Pederson.** 2000. Rapid mini-preparation
11 of fungal DNA for PCR. J Clin Microbiol **38**:471.
- 12 37. **Liu, Y., P. Srivilai, S. Loos, M. Aebi, and U. Kües.** 2006. An essential gene
13 for fruiting body initiation in the basidiomycete *Coprinopsis cinerea* is
14 homologous to bacterial cyclopropane fatty acid synthase genes. Genetics
15 **172**:873-84.
- 16 38. **Lorang, J. M., R. P. Tuori, J. P. Martinez, T. L. Sawyer, R. S. Redman, J.
17 A. Rollins, T. J. Wolpert, K. B. Johnson, R. J. Rodriguez, M. B. Dickman,
18 and L. M. Ciuffetti.** 2001. Green fluorescent protein is lighting up fungal
19 biology. Applied and Environmental Microbiology **67**:1987-1994.
- 20 39. **Lugones, L. G., K. Scholtmeijer, R. Klootwijk, and J. G. H. Wessels.** 1999.
21 Introns are necessary for mRNA accumulation in *Schizophyllum commune*.
22 Molecular Microbiology **32**:681-689.
- 23 40. **Ma, B., M. B. Mayfield, and M. H. Gold.** 2001. The green fluorescent
24 protein gene functions as a reporter of gene expression in *Phanerochaete*
25 *chrysosporium*. Appl. Environ. Microbiol. **67**:948-955.
- 26 41. **May, G., L. Le Chevanton, and P. J. Pukkila.** 1991. Molecular analysis of
27 the *Coprinus cinereus* mating type A factor demonstrates an unexpectedly
28 complex structure. Genetics **128**:529-38.
- 29 42. **Mikosch, T. S., B. Lavrijssen, A. S. Sonnenberg, and L. J. van Griensven.**
30 2001. Transformation of the cultivated mushroom *Agaricus bisporus* (Lange)
31 using T-DNA from *Agrobacterium tumefaciens*. Curr Genet **39**:35-9.
- 32 43. **Money, N. P., and J. P. Ravishankar.** 2005. Biomechanics of stipe
33 elongation in the basidiomycete *Coprinopsis cinerea*. Mycol Res **109**:627-34.
- 34 44. **Moore, D., and P. J. Pukkila.** 1985. *Coprinus cinereus* - an Ideal Organism
35 for Studies of Genetics and Developmental Biology. J. Biol. Educ. **19**:31-40.
- 36 45. **Murzin, A. G., S. E. Brenner, T. Hubbard, and C. Chothia.** 1995. SCOP: a
37 structural classification of proteins database for the investigation of sequences
38 and structures. J Mol Biol **247**:536-40.
- 39 46. **Namekawa, S., F. Hamada, S. Ishii, Y. Ichijima, T. Yamaguchi, T. Nara,
40 S. Kimura, T. Ishizaki, K. Iwabata, A. Koshiyama, H. Teraoka, and K.
41 Sakaguchi.** 2003. *Coprinus cinereus* DNA ligase I during meiotic
42 development. Biochimica Et Biophysica Acta-Genes Structure and Expression
43 **1627**:47-55.
- 44 47. **Namekawa, S. H., K. Iwabata, H. Sugawara, F. N. Hamada, A.
45 Koshiyama, H. Chiku, T. Kamada, and K. Sakaguchi.** 2005. Knockdown
46 of LIM15/DMC1 in the mushroom *Coprinus cinereus* by double-stranded
47 RNA-mediated gene silencing. Microbiology-Sgm **151**:3669-3678.
- 48 48. **Nielsen, H., J. Engelbrecht, S. Brunak, and G. von Heijne.** 1997. A neural
49 network method for identification of prokaryotic and eukaryotic signal
50 peptides and prediction of their cleavage sites. Int J Neural Syst **8**:581-99.

- 1 49. **Poggeler, S., S. Masloff, B. Hoff, S. Mayrhofer, and U. Kuck.** 2003.
2 Versatile EGFP reporter plasmids for cellular localization of recombinant gene
3 products in filamentous fungi. *Curr. Genet.* **43**:54-61.
- 4 50. **Reijnders, A. F. M.** 1979. Developmental anatomy of *Coprinus*. *Persoonia*
5 **10**:383-424.
- 6 51. **Romaine, C. P.** 2002. Gene transfer technology for mushrooms: the power
7 and potential for significant crop improvement. *Mushroom News* **50**:4-14.
- 8 52. **Rosin, I. V., j. Horner, and d. Moore.** 1985. Differentiation and pattern
9 formation in the fruit body cap of *Coprinus cinereus*. In D. Moore, L. A.
10 Casselton, D. A. Wood, and J. C. Frankland (ed.), *Developmental biology of*
11 *higher fungi*. Cambridge University Press, Cambridge.
- 12 53. **Schmid, J. A., and H. Neumeier.** 2005. Evolutions in science triggered by
13 green fluorescent protein (GFP). *Chembiochem* **6**:1149-+.
- 14 54. **Skrzynia, C., D. M. Binnering, J. A. Alspaugh, and P. J. Pukkila.** 1989.
15 Molecular Characterization of Trp1, a Gene Coding for Tryptophan-
16 Synthetase in the Basidiomycete *Coprinus cinereus*. *Gene* **81**:73-82.
- 17 55. **Stothard, P.** 2000. The sequence manipulation suite: JavaScript programs for
18 analyzing and formatting protein and DNA sequences. *Biotechniques* **28**:1102,
19 1104.
- 20 56. **Swamy, S., I. Uno, and T. Ishikawa.** 1984. Morphogenetic Effects of
21 Mutations at the A and B Incompatibility Factors in *Coprinus cinereus*. *J Gen*
22 *Microbiol* **130**:3219-3224.
- 23 57. **Tyas, D. A., T. Pratt, T. I. Simpson, J. O. Mason, and D. J. Price.** 2003.
24 Identifying GFP-transgenic animals by flashlight. *Biotechniques* **34**:474-+.
- 25 58. **Velco, A. J., R. W. Kerrigan, L. A. MacDonald, M. P. Wach, C.**
26 **Schlaghauser, and C. P. Romaine.** 2004. Expression of novel genes in
27 *Agaricus bisporus* using an Agrobacterium-mediated transformation
28 technique. *Mushroom Science* **16**:591-598.
- 29 59. **Yamaguchi, T., S. H. Namekawa, F. N. Hamada, N. Kasai, T. Nara, K.**
30 **Watanabe, K. Iwabata, T. Ishizaki, S. Ishii, A. Koshiyama, S. Inagaki, S.**
31 **Kimura, and K. Sakaguchi.** 2004. Expression of flap endonuclease-1 during
32 meiosis in a basidiomycete, *Coprinus cinereus*. *Fungal Genet. Biol.* **41**:493-
33 500.
- 34 60. **Yang, J., X. Huang, B. Tian, H. Sun, J. Duan, W. Wu, and K. Zhang.**
35 2005. Characterization of an extracellular serine protease gene from the
36 nematophagous fungus *Lecanicillium psalliotae*. *Biotechnol Lett* **27**:1329-34.
- 37 61. **Yang, J., J. Li, L. Liang, B. Tian, Y. Zhang, C. Cheng, and K. Q. Zhang.**
38 2007. Cloning and characterization of an extracellular serine protease from the
39 nematode-trapping fungus *Arthrobotrys conoides*. *Arch Microbiol* **188**:167-
40 174.
- 41 62. **Zhang, C., Odon, V., Kim, H.K., Challen, M.P., Burton, K.S., Hartley D.,**
42 **Elliott T.J.** 2004. Mushrooms for molecular pharming. *Mushroom Science*
43 **16**:611-617.
- 44 63. **Zolan, M. E., and P. J. Pukkila.** 1986. Inheritance of DNA Methylation in
45 *Coprinus cinereus*. *Mol. Cell. Biol.* **6**:195-200.
- 46
47

Table 1.

Strain		Media						
		YMG	RA	MMP	PDA	Humic fraction	GA	Milk
<i>A. bisporus</i>	17	-	-	-	+	+	-	-
	18	-	-	-	-	+	-	-
	19	-	-	-	+	+	+	-
	21	-	-	-	+	+	+	+
	22	-	-	-	+	+	-	+
	23	-	-	-	+	+	+	+
	119	-	-	-	+	+	-	+
	120	-	-	-	+	+	+	+
	121	-	-	-	+	+	+	+
	A15	-	-	-	-	-	-	-
	G26	+	+	+	+	+	+	+
<i>C. cinereus</i> monokaryon	1	-	-	ND	+	+	+	+
	24	-	-	ND	+	+	+	+
	37	-	-	ND	+	+	+	+
	47	+	-	ND	+	+	+	+
	LT2	-	-	ND	-	-	-	-
	PG78Gr	+	+	ND	+	+	+	+
<i>C. cinereus</i> dikaryon	1	-	-	ND	+	+	+	+
	24	-	-	ND	+	+	+	+
	37	-	-	ND	-	+	+	+
	47	+	-	ND	+	+	+	+

Table 2.

Gene	Length (aa)	Predicted Mol Wt kDa	Predicted PI	Signal peptide	Cleavage point	Introns	Secondary structure			Regulatory sequences in promoter	
							% helices	% strands	% Loops	Nit 2/AreA	CreA
<i>Spr1</i>	387	39.39	5.93	+	19/20	nd	20.41	24.81	54.78	9	1
<i>Spr2</i>	377	38.85	5.53	+	19/20	11	29.44	24.93	45.62	9	1
CC1G_04562.1	394	40.093	7.75	+	20/21	14	27.99	24.94	47.07	6	2
CC1G_10592.1	387	39.332	9.69	+	21/22	12	36.53	21.24	42.23	10	1
CC1G_10615.1	394	40.193	9.97	+	21/22	6	30.53	20.61	48.85	13	3
CC1G_07792.1	395	40.683	6.73	+	20/21	13	28.43	22.84	48.73	7	3
CC1G_10606.1	346	35.359	5.83	+	19/20	13	21.16	35.07	43.77	7	4
CC1G_0.3122.1	421	44.354	7.55	+	20/21	12	20.71	24.29	55.00	4	1
CC1G_04470.1	500	53.18	6.84	+	21/22	2	22.24	16.63	61.12	6	8

Table 3.

Gene	<i>Spr1</i>	<i>Spr2</i>	04562.1	10592.1	10615.1	07792.1	10606.1	0.3122.1	04470.1
<i>Spr1</i>	100								
<i>Spr2</i>	75	100							
04562.1	61	55	100						
10592.1	57	54	66	100					
10615.1	55	53	62	77	100				
07792.1	55	50	70	58	56	100			
10606.1	47	48	50	54	51	47	100		
0.3122.1	46	45	43	45	41	40	40	100	
04470.1	44	42	41	39	40	39	35	31	100

A:

UV

PC

A. bisporus
TP17 on humic
fraction

0.5mm

0.5mm

A. bisporus
TP17 on RA

0.5mm

0.5mm

B:


Panel

WL


BL




A: *A. bisporus*
G26

B: *A. bisporus* G26
(left) and TP196
(right)



GFP activity: *A. bisporus* 3 days Postharvest








UV

PC



C. cinereus
TP47
monokaryon on
humic fraction



C. cinereus
TP47
monokaryon on
RA



C. cinereus
TP47 dikaryon
on humic
fraction



C. cinereus
TP47 dikaryon
on RA

