

Research

## Antagonism of the prostaglandin D<sub>2</sub> receptor CRTH2 attenuates asthma pathology in mouse eosinophilic airway inflammation

Lena Uller\*<sup>1</sup>, Jesper Mosolff Mathiesen<sup>2</sup>, Lisa Alenmyr<sup>3</sup>, Magnus Korsgren<sup>3</sup>, Trond Ulven<sup>2</sup>, Thomas Högberg<sup>2</sup>, Gunnar Andersson<sup>2</sup>, Carl GA Persson<sup>3</sup> and Evi Kostenis<sup>2</sup>

Address: <sup>1</sup>Dept. Experimental Medical Science, Lund University, Sweden, <sup>2</sup>TM Pharma A/S, Fremtidsvej 3, 2970 Hørsholm, Denmark and <sup>3</sup>Dept. Clinical Pharmacology, Lund University Hospital, Lund, Sweden

Email: Lena Uller\* - [lenu.uller@med.lu.se](mailto:lenu.uller@med.lu.se); Jesper Mosolff Mathiesen - [JMM@zp.dk](mailto:JMM@zp.dk); Lisa Alenmyr - [lisa.alenmyr@med.lu.se](mailto:lisa.alenmyr@med.lu.se); Magnus Korsgren - [magnus.korsgren@med.lu.se](mailto:magnus.korsgren@med.lu.se); Trond Ulven - [ulven@chem.sdu.dk](mailto:ulven@chem.sdu.dk); Thomas Högberg - [TH@7tm.com](mailto:TH@7tm.com); Gunnar Andersson - [ga@7tm.com](mailto:ga@7tm.com); Carl GA Persson - [carl.persson@med.lu.se](mailto:carl.persson@med.lu.se); Evi Kostenis - [kostenis@uni-bonn.de](mailto:kostenis@uni-bonn.de)

\* Corresponding author

Published: 28 February 2007

Received: 10 October 2006

*Respiratory Research* 2007, **8**:16 doi:10.1186/1465-9921-8-16

Accepted: 28 February 2007

This article is available from: <http://respiratory-research.com/content/8/1/16>

© 2007 Uller et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Mast cell-derived prostaglandin D<sub>2</sub> (PGD<sub>2</sub>), may contribute to eosinophilic inflammation and mucus production in allergic asthma. Chemoattractant receptor homologous molecule expressed on TH<sub>2</sub> cells (CRTH2), a high affinity receptor for prostaglandin D<sub>2</sub>, mediates trafficking of TH<sub>2</sub>-cells, mast cells, and eosinophils to inflammatory sites, and has recently attracted interest as target for treatment of allergic airway diseases. The present study involving mice explores the specificity of CRTH2 antagonism of TM30089, which is structurally closely related to the dual TP/CRTH2 antagonist ramatroban, and compares the ability of ramatroban and TM30089 to inhibit asthma-like pathology.

**Methods:** Affinity for and antagonistic potency of TM30089 on many mouse receptors including thromboxane A<sub>2</sub> receptor mTP, CRTH2 receptor, and selected anaphylatoxin and chemokines receptors were determined in recombinant expression systems *in vitro*. *In vivo* effects of TM30089 and ramatroban on tissue eosinophilia and mucus cell histopathology were examined in a mouse asthma model.

**Results:** TM30089, displayed high selectivity for and antagonistic potency on mouse CRTH2 but lacked affinity to TP and many other receptors including the related anaphylatoxin C3a and C5a receptors, selected chemokine receptors and the cyclooxygenase isoforms 1 and 2 which are all recognized players in allergic diseases. Furthermore, TM30089 and ramatroban, the latter used as a reference herein, similarly inhibited asthma pathology *in vivo* by reducing peribronchial eosinophilia and mucus cell hyperplasia.

**Conclusion:** This is the first report to demonstrate anti-allergic efficacy *in vivo* of a highly selective small molecule CRTH2 antagonist. Our data suggest that CRTH2 antagonism alone is effective in mouse allergic airway inflammation even to the extent that this mechanism can explain the efficacy of ramatroban.

## Background

The small lipid mediator prostaglandin D<sub>2</sub> (PGD<sub>2</sub>) is the major cyclooxygenase metabolite of arachidonic acid and is released by activated mast cells in response to allergen exposure [1]. PGD<sub>2</sub> has long been considered a potentially important mediator in several diseases such as asthma, allergic rhinitis, atopic dermatitis, and allergic conjunctivitis [2-5]. PGD<sub>2</sub> elicits biological responses by interaction with three specific seven-transmembrane receptors, referred to as DP/DP1, DP2/CRTH2, and TP (DP, D prostanoid receptor; CRTH2, chemoattractant receptor homologous molecule expressed on T helper type 2 cells; TP, thromboxane A<sub>2</sub> receptor) [6-8]. Via interaction with one (or a combination) of its three specific receptors PGD<sub>2</sub> may contribute to bronchoconstriction, eosinophilia and mucus production in allergic asthma. However, assessment of actual roles of PGD<sub>2</sub> in allergic diseases has been hampered by its very short biological half-life and the lack of specific receptor antagonists suitable to uncover how signaling of individual PGD<sub>2</sub> receptors contribute to disease processes *in vivo*. It is also possible that the contribution of PGD<sub>2</sub> to allergic airway inflammation is easily missed if the load of allergen in challenge studies is too large [9].

CRTH2 is expressed on eosinophils, TH<sub>2</sub> cells and basophils, which are all considered to contribute to the pathogenesis of allergic diseases [3,10-15]. Several lines of evidence suggest that activation of CRTH2 in response to PGD<sub>2</sub> mediates recruitment of inflammatory cells *in vitro* and *in vivo*. *In vitro*, activation of CRTH2 induces chemotaxis of TH<sub>2</sub> cells, eosinophils, and basophils [7,16]. *In vivo*, CRTH2 mediates mobilization of eosinophils from guinea-pig bone marrow [17], promotes eosinophilia and exacerbates pathology in mouse models of allergic asthma and atopic dermatitis [18], and induces eosinophil infiltration into the airways upon intratracheal administration of PGD<sub>2</sub> or a selective CRTH2 agonist [19-21]. Based on evidence supporting a pro-inflammatory role of CRTH2, this receptor has attracted great interest as a drug target for therapeutic intervention in allergic diseases. Confusingly, however, allergic mice that lack a functional CRTH2 receptor and hence are incapable to signal through CRTH2 have been reported to exhibit both increased [22] and reduced [23,24] allergic inflammation in models of asthma [22-24] and atopic dermatitis [23]. These diverging reports involving gene-deficient animals further underscore the need to use specific CRTH2 antagonists to explore the *in vivo* function of CRTH2.

It was recently reported that ramatroban, which was initially developed as a TP antagonist and is now used for treatment of allergic rhinitis in Japan, also displays potent CRTH2 antagonistic activity [25]. Consistent with this finding, ramatroban has been shown to abrogate blood

eosinophilia induced by a CRTH2-specific agonist in rats [19] and to inhibit PGD<sub>2</sub>-stimulated human eosinophil migration *in vitro* [25]. While these data are congruent with the notion that ramatroban acts through inhibition of CRTH2 receptors, it is not clear whether its clinical efficacy in allergic rhinitis is due to inhibition of TP, CRTH2 or both receptors. We have recently reported a ramatroban analog (given internal code number TM30089) with high antagonistic potency on and selectivity for human CRTH2 and devoid of affinity to the TP receptor [26,27].

Here we report that TM30089 is a highly potent antagonist on mouse CRTH2 and lacks affinity to mouse TP in agreement with its pharmacological properties on the human receptor orthologs. In addition, this study demonstrates that TM30089 does not display appreciable affinity to a range of other receptors and enzymes that have potentially important roles in allergic asthma. Furthermore, using a mouse model of allergic asthma involving two allergen challenges, we have discovered that TM30089 similar to ramatroban inhibits allergen challenge-induced airway tissue eosinophilia and mucus cell hyperplasia. These *in vivo* data support the promise of CRTH2 as a target for treatment of allergic airways diseases.

## Methods

### In vitro analysis

*Cloning and expression of the mouse CRTH2 (mCRTH2) and mouse thromboxane A<sub>2</sub> (mTP) receptor in HEK293 cells*

The coding sequence of mCRTH2 (genbank accession no. [AF054507](#)) was amplified by PCR from mouse hippocampus cDNA and inserted into the pcDNA3.1(+) expression vector (Invitrogen). A HEK293 cell line, stably expressing mCRTH2 (hereafter referred to as mCRTH2-HEK cells) was generated under G418 (Gibco #11811) selection and used as described below in whole cell binding and functional inositol phosphate accumulation assays. The coding sequence of mTP (genbank accession no. [D10849](#)) was amplified by PCR from mouse spleen cDNA and inserted into pcDNA3.1(+). HEK293 cells were transiently transfected with mTP as described [27] and assayed after 48 hr as outlined below.

### mCRTH2 and mTP whole cell binding

Binding assays were performed essentially as described previously [27]. In brief, mCRTH2-HEK cells or HEK cells transiently transfected with mTP were seeded into white 96 well plates (Costar #3917) at a density of 30,000 cells/well. About 18–24 hr later, whole cell competition binding experiments were performed using 0.8 nM [<sup>3</sup>H]PGD<sub>2</sub> (Amersham #TRK734, 166 Ci/mmol) or 0.7 nM [<sup>3</sup>H]SQ29548 (PerkinElmer #NET-936, 48.2 Ci/mmol), for mCRTH2 and mTP respectively. Total and nonspecific binding were determined in the absence and presence of

10  $\mu\text{M}$   $\text{PGD}_2$  for mCRTH2 and 10  $\mu\text{M}$  U46619 for mTXA<sub>2</sub> receptors.

#### *Inositol phosphate accumulation assays*

mCRTH2-HEK or HEK cells transiently transfected with mTP were seeded at a density of 30,000 cells/well in poly-D-lysine coated 96-well tissue culture plates and labeled by overnight incubation in medium containing 5  $\mu\text{Ci}$  *myo*- [2-<sup>3</sup>H]-inositol (TRK911, Amersham Biosciences). After washing, cells were stimulated with the respective ligands in the presence of 5 mM LiCl (HBSS, GIBCO cat. 14025-050, 45 min, 37°C) and the inositol phosphate fraction was quantified using a scintillation proximity assay (SPA) as described previously [28]. To facilitate inositol phosphate generation by the Gi-selective CRTH2 receptor, mCRTH2-HEK cells were transiently transfected with a promiscuous G $\alpha$  protein [29].

#### *TM30089 profiling panel*

TM30089 was tested at a final concentration of 10  $\mu\text{M}$  in either binding or functional assays on several 7TM receptors, and in enzymatic assays on selected enzymes including those involved in arachidonic acid metabolism. All assays were performed using recombinant human receptors and enzymes, and included the appropriate controls for assay validation. Enzymatic and binding assays, except for muscarinic binding, were performed by CEREP in vitro pharmacology profiling services. Full description of methods and references is available on the CEREP Web site [30]. Muscarinic binding assays were performed on mouse brain homogenate (non-selective) or porcine heart membranes (M2-selective) with 0.2 nM 3H-NMS as a tracer and atropine as control. Functional assays were performed using recombinant human receptors and enzymes, and included the appropriate controls for C3a, C5a, ChemR23 receptors by determining the inhibitory potency of TM30089 in InsP assays essentially as described above. For stimulation of C3a, C5a, and ChemR23 receptors in InsP assays the following agonists were used: human recombinant C3a (C3a receptor), human recombinant C5a (C5a receptor), and the non-peptide Chemerin (ChemR23).

#### *Data analysis*

was performed using the Prism 3.0 software (GraphPad Software, San Diego) as described previously [27] with the following exceptions: Binding data from competition binding assays were normalized to percent of specific binding of [<sup>3</sup>H]PGD<sub>2</sub> or [<sup>3</sup>H]SQ29548 and the K<sub>i</sub> was then determined using the Cheng-Prusoff equation  $K_i = IC_{50} / (1 + L/K_d)$ , where L is the concentration of radioactive ligand and K<sub>d</sub> is its dissociation constant. For Schild analysis, global fitting using the Gaddum/Schild competitive interaction model was applied after the functional concentration-response curves for the agonists PGD<sub>2</sub> or U46619 in

the absence or presence of various antagonist concentrations were normalized to percent of the maximal response obtained by agonist alone.

#### **In vivo model of allergic asthma**

##### *Animals*

Female BALB/c mice about 6 weeks of age (MoB A/S; Ry, Denmark) were kept in well-controlled animal housing facilities and fed *ad libitum*. The study was approved by the Regional Ethics Committee in Malmö-Lund, Sweden.

##### *Allergen sensitization and challenge protocol*

All groups of mice (n = 10 in each group) were immunized by intraperitoneal (i.p) injections of ovalbumin (OVA, 10  $\mu\text{g}$  per injection; grade V, Sigma, St Louis, Mo) absorbed to alum adjuvant (Al(OH)<sub>3</sub>; Sigma) at day 0 as described previously [31,32]. Fourteen days after immunization mice were exposed to aerosolized OVA (1 % wt/vol) 30 minutes daily for 2 days. Control animals received saline challenge. Animals were sacrificed 24 hr after the last allergen challenge by pentobarbital i.p followed by bronchoalveolar lavage (BAL) and lung tissue sampling. In preliminary studies it was ascertained that daily allergen challenges for only a few days sufficed to produce a significant airway tissue eosinophilia as well as goblet cell hyperplasia.

##### *Treatment with CRTH2 antagonists in vivo*

On days 14 and 15 mice were treated twice daily with TM30089 and ramatroban (BAY3405), 5 mg/kg orally. Control animals received vehicle (PBS 10% Tween80). Drugs were administered 30 minutes before and 4 hr after each challenge. The dose of ramatroban was selected in accord with previous reports [20,23]. Furthermore, ramatroban and TM30089 differ chemically merely by the move of one carbon atom from the carboxyl side chain in ramatroban to the sulfonamide nitrogen in TM30089 – thus they have the same molecular weight and other important structural features in common – so it is highly reasonable to assume they will have very similar PK properties. Hence, based on the additional similarity regarding CRTH2 antagonism *in vitro* the same dose of the TM30089 was used in the in vivo part of this study to allow for a head-to-head comparison.


##### *Staining and quantification of lung tissue eosinophils and mucus-containing cells*

Eosinophils were detected by histochemical visualization of cyanide-resistant eosinophil peroxidase (EPO) activity [33]. In brief, 5  $\mu\text{m}$  cryo sections were incubated for 8 min at room temperature in PBS buffer (pH 7.4) supplemented with 3.3-diaminobenzidine tetrahydrochloride (60 mg/100 ml; Sigma), 30% H<sub>2</sub>O<sub>2</sub> (0.3 ml/100 ml), and NaCN (120 mg/100 ml). Slides were then rinsed in tap water and mounted in Kaisers medium (Merck, Darm-

stadt, Germany). Eosinophils were identified by their dark brown reaction product and eosinophils were counted around bronchi using a depth of 120  $\mu\text{m}$  from the epithelial basement membrane and expressed as cells/0.1  $\text{mm}^2$  tissue area as previously described [32]. 5  $\mu\text{m}$  cryo sections were stained with periodic acid-Schiff reagent (PAS) and the total number of PAS-positive cells counted and expressed as cells/mm basement membrane as previously described [32]. Although the focus in this study was on tissue pathology BAL was also carried out (see below).

**Bronchoalveolar lavage (BAL) and quantification of luminal cells**  
Although eosinophils, according to the preliminary studies, clearly had migrated to the airway tissue the brevity of present allergen challenge period until termination meant

that the tissue eosinophils had only started to be lost into the airway lumen. However, it was still of interest to examine BALF eosinophils to see if drug-induced effects on tissue eosinophilia in part could be due to increased loss of these cells through migration across the epithelial lining [34]. BAL was performed via a ligated tracheal cannula and 1 ml of PBS was allowed to passively enter the lungs at a pressure of 10  $\text{cm H}_2\text{O}$ , a procedure that is gentle to the lungs [32]. The obtained BAL-fluid (BALF) from each animal was immediately centrifuged and the supernatant frozen for ELISA analysis. The total number of cells was quantified using a cell sorter (NucleoCounter<sup>®</sup>, Chemometec A/S, Allerød, Denmark) and  $5 \times 10^5$  cells cytocentrifuged to microscope slides. Differential cell counts were performed on May-Grünwald Giemsa stained slides and



**Figure 1**

**In vitro characterization of compounds on mouse CRTH2 (mCRTH2) and mouse TP (mTP) receptors. A,** Competition binding analysis. **B,** inhibition of mCRTH2 and mTP receptor function. PGD<sub>2</sub>- or U-46619-concentration response curves in the absence and presence of the indicated compounds in mCRTH2- or mTP-expressing cells. Inserts: Schild plots. Schild analysis show potent antagonism of mCRTH2 by TM30089 ( $pA_2 = 9.15 \pm 0.11$ , Schild slope =  $1.45 \pm 0.08$ ) whereas it does not interfere with signaling of mTP. Ramatroban is a potent antagonist on mCRTH2 ( $pA_2 = 8.08 \pm 0.14$ , Schild slope =  $0.94 \pm 0.05$ ) and mTP ( $pA_2 = 9.36 \pm 0.10$ , Schild slope =  $1.35 \pm 0.06$ ). Experiments show mean  $\pm$  SE from 5–7 independent experiments.

percentage of eosinophils, lymphocytes, neutrophils, and macrophages determined by counting 200–300 cells in a blinded manner. To obtain the absolute number of each leukocyte subtype in each BALF, the percentage of cells was multiplied by the total number of cells recovered by the BAL.

#### Data analysis

Histology analyses were performed and quantified in a blinded manner. To compare histological data, the non-parametric Kruskal-Wallis test and followed by Conover's test was applied using Analyze It™ (Analyze-it software Ltd., Leeds, UK). Data are expressed as mean  $\pm$  SEM. A value of  $p < 0.05$  was considered statistically significant.

## Results

### *In vitro* pharmacological profile of TM30089 and ramatroban

We have recently reported that the ramatroban analog TM30089 displays high affinity to human CRTH2 (hCRTH2) but completely lacks affinity to the human thromboxane A<sub>2</sub> receptor hTP (TP, T prostanoid receptor), unlike ramatroban which antagonizes both receptors equally well [27]. To investigate whether TM30089 represents a suitable tool compound to explore the role of CRTH2 in a mouse model of allergic inflammation *in vivo*, we cloned the mouse orthologs of CRTH2 and TP and evaluated TM30089 and ramatroban (as a reference) in their ability to displace [<sup>3</sup>H]PGD<sub>2</sub> and [<sup>3</sup>H]SQ29548 from mCRTH2 and mTP, respectively, in competition binding

experiments using human embryonic kidney cells expressing the individual receptors. Both compounds displayed high affinity to mCRTH2 with  $\log pK_i = 8.96 \pm 0.05$  (1.1 nM), for TM30089 and  $pK_i = 8.38 \pm 0.05$  (4.2 nM) for ramatroban (FIG 1A). The affinity of ramatroban in our binding assay is in good agreement with the reported  $K_i$  value of 28 nM obtained by others in the same expression system [35]. In contrast to ramatroban which also displayed high affinity to mTP ( $pK_i = 8.92 \pm 0.05$  (1.2 nM)), TM30089 bound to mTP with only negligible affinity ( $pK_i = 5.30 \pm 0.03$  (50.000 nM)). Thus, both compounds retain their selectivity profile for CRTH2 versus TP on the mouse receptor orthologs: TM30089 is CRTH2-selective, while ramatroban is a dual CRTH2/TP ligand (FIG 1A).

To confirm antagonistic efficacy of the compounds their ability to inhibit cellular signaling of the receptors was evaluated in a functional second messenger assay (FIG 1B). In agreement with the binding data, TM30089 acts as a potent antagonist on mCRTH2 ( $pA_2 = 9.15 \pm 0.11$ , Schild slope =  $1.45 \pm 0.08$ ) but does not interfere with signaling of mTP, as compared with ramatroban which potently inhibits cellular signaling of both mCRTH2 ( $pA_2 = 8.08 \pm 0.14$ , Schild slope =  $0.94 \pm 0.05$ ) and mTP ( $pA_2 = 9.36 \pm 0.10$ , Schild slope =  $1.35 \pm 0.06$ ) receptors (FIG 1B). To ascertain that *in vivo* efficacy of TM30089 does not arise from inhibition of related 7TM receptors such as the anaphylatoxin receptors C3a and C5a, the chemokine receptors CCR3 and ChemR23, or the second high affinity PGD<sub>2</sub> receptor DP, it was tested for its affinity or antagonistic potency on these receptors (Table 1). Notably, TM30089 is very selective, exhibiting > 1000-fold preference for CRTH2 over DP [26], and lacks affinity to any of the other tested receptors and also to the two cyclooxygenase isoforms 1 and 2 which are recognized as important players in allergic airway inflammation [36,37]. Together, the affinity and selectivity profile of TM30089 suggests that it may be suited to uncover the *in vivo* contribution of CRTH2 in allergic airway inflammation.

**Table 1: Profiling of TM30089 at relevant 7TM receptors (binding or functional assays) and enzymes (enzymatic assays).**


Receptor	Assay type <sup>a</sup>	% of control <sup>b</sup>
AT <sub>1</sub>	B	90.1 $\pm$ 2.8
AT <sub>2</sub>	B	98.1 $\pm$ 1.4
BLT <sub>1</sub>	B	89.3 $\pm$ 3.5
C3a	F	88.7 $\pm$ 5.9
C5a	B	108.7 $\pm$ 6.3
ChemR23	F	96.6 $\pm$ 1.6
CXCR2	B	104.6 $\pm$ 6.1
CysLT <sub>1</sub>	B	116.9 $\pm$ 10.5
Glucocorticoid	B	98.9 $\pm$ 11.3
Muscarinic (non-selective)	B	106.7 $\pm$ 8.5
Muscarinic (M2-selective)	B	97.5 $\pm$ 1.9
Phospholipase A <sub>2</sub>	E	101.5 $\pm$ 1.0
COX <sub>1</sub>	E	109.6 $\pm$ 3.6
COX <sub>2</sub>	E	123.8 $\pm$ 13.4
iNOS	E	102.3 $\pm$ 3.6

<sup>a</sup> Assay type: B – radioligand binding, F – functional InsP, and E – enzymatic assays.

<sup>b</sup> Assays were performed in the absence (100% control) or presence of 10  $\mu$ M TM30089 (% of control). Results are given as % of control  $\pm$  SE (estimated) from 2–3 independent experiments.

### *In vivo* effects of CRTH2 antagonists on asthma-like airway histopathology

Mouse models of allergic asthma exhibit some of the cardinal histopathological signs of human asthmatic lungs, most notably eosinophilia and mucus cell hyperplasia [32,38]. These are also the key features investigated in this study. Mice were immunized with the allergen (OVA) and challenged two times by inhalation of aerosolized OVA. Lung histology was assessed 24 hr after the last challenge. OVA-challenged mice developed a marked peribronchial lung tissue eosinophilia (35.5 eosinophils  $\pm$  4.7/0.1 mm<sup>2</sup>)  $p < 0.0001$  compared to saline challenged animals (1.3  $\pm$  0.3 cells) (FIG 2a). Eosinophilia was accompanied by a significant increase of mucus cells in the airway mucosa (88.1  $\pm$  8.8; 3.2  $\pm$  1 cells/mm basement mem-

**Figure 2**

**CRTH2 antagonism in vivo attenuates airway tissue eosinophilia and mucus cell hyperplasia.** Above (A): OVA challenge produced marked eosinophilia and mucus cell hyperplasia (\*\*\*,  $p < 0.001$  compared to saline control). Treatment with ramatroban or the specific CRTH2 antagonist TM30089 significantly reduced airway tissue eosinophilia (black bars) and mucus production (white bars); #,  $p < 0.05$  compared to vehicle treatment. Below (B-G): Light micrographs showing effects in particularly well-responding animals. Lung tissue eosinophilia in normal saline treated lung (B), OVA/vehicle treated lung (C), and OVA/TM30089 treated lung (D). Airway mucus cells are shown in normal saline treated lung (E), OVA/vehicle treated lung (F), and OVA/TM30089 treated lung (G). Scale bar = 100  $\mu$ m.



brane), (FIG 2a). Treatment with TM30089 and ramatroban significantly diminished the allergen challenge-induced peribronchial lung tissue eosinophilia and mucus cell hyperplasia (FIG 2). Inhibition by both compounds of pathological changes was quantitative and did not alter the overall pattern of occurrence of eosinophils and mucus cells in the airway-lung tissues (FIG 2).

Elimination of the airway tissue eosinophils largely occurs through egression of these cells into the airway lumen [32,39,40] where they can be retrieved by BAL. As indicated by the BALF eosinophilia ( $1.0 \pm 0.2$ ;  $0.02 \pm 0.02$  % eosinophils in allergen and saline-challenged animals, respectively), eosinophils had already started to enter into the airway lumen about 24 hr after the second allergen challenge. Both drug treatments tended to reduce BALF eosinophilia (TM30089  $0.5 \pm 0.01$  %; ramatroban  $0.4 \pm 0.01$  % BALF eosinophils) indicating that the attenuation of tissue eosinophilia induced by these drugs did not reflect increased elimination of tissue eosinophils into the airway lumen.

## Discussion

Emerging evidence suggests that PGD<sub>2</sub>-activation of its high affinity receptor CRTH2 may be particularly critical in the pathogenesis of eosinophilic airway inflammation, since activation by PGD<sub>2</sub> of CRTH2 potently stimulates chemotaxis of eosinophils *in vitro* and *in vivo* [7,19-21,25,41-43]. Herein, we demonstrate that the small molecule TM30089 is a highly potent and selective inhibitor of mouse CRTH2 function *in vitro*, and by using this antagonist demonstrate for the first time that inhibition of CRTH2 signaling *in vivo* suppresses the development of certain key features characteristic for allergic asthma.

*In vitro*, TM30089, which is structurally closely related to ramatroban, was found to bind mouse CRTH2 with nanomolar affinity and potently inhibit its signaling in cells overexpressing the receptor. In addition, TM30089 completely lacks affinity to the mouse TP receptor, unlike ramatroban which represents a dual TP/CRTH2 antagonist on both human and mouse receptor orthologs [25,26] (this study). Furthermore TM30089 in concentrations up to 10  $\mu$ M did not show significant binding to or inhibition of selected chemokine receptors, anaphylatoxin receptors, the other high affinity PGD<sub>2</sub> receptor DP as well as the cyclooxygenases 1 and 2. Owing to its high selectivity over all other tested receptors and enzymes, and the fact that its chemical structure is closely related to ramatroban which has proven efficacious in various animal and human studies of allergic rhinitis and asthma [44-50], TM30089 emerges a suitable research tool to explore the contribution of CRTH2-signaling in allergic airway inflammation *in vivo*.

We found that administration of TM30089 displayed anti-inflammatory efficacy in a mouse model of allergic asthma that mimics some of the major histopathological features characteristic of human asthma such as eosinophilia and mucus cell hyperplasia [38]. In the present study we used allergen challenges during only two consecutive days which proved to be sufficient for causing goblet cell hyperplasia as well as a significant accumulation of eosinophils in airway-pulmonary tissues. More challenges for more days would have increased the eosinophilia further. However, the present model was preferred because it had been reported that efficacy of interference with PGD<sub>2</sub> is dependent on the allergen load [9]. Thus, the marked protection against allergen-induced asthma observed in DP<sup>-/-</sup> mice was completely lost when more than three allergen challenges were used [9]. It is of note that although the present number of provocations was low the load of allergen would still be far greater than the level of allergens human asthmatics are exposed to.

The present observation of increased BALF eosinophils agrees with previous data indicating that already during the build-up of tissue eosinophilia by allergen exposure these cells begin to be lost into the airway lumen [32]. Yet, studies resorting exclusively to the determination of BALF eosinophilia run the risk of drawing incorrect conclusions about changes in eosinophil numbers in the most important locale, the airway-pulmonary tissues. For example, we and others have previously observed in allergic mice that drug interventions may inhibit lumen eosinophilia whilst the tissue eosinophilia remains unchanged or is, indeed, increased as reviewed in [34]. It was, therefore, important to note in this study that both drug treatments tended to reduce BALF eosinophils demonstrating that increased elimination of these cells into the airway lumen could not explain the present inhibitory effects of these drugs on airway tissue eosinophilia. Using TM30089, we thus unravel that CRTH2-signaling appears integral to the recruitment of eosinophils to the airways *in vivo*. So far, it has only been demonstrated that externally administered PGD<sub>2</sub> is able to induce local eosinophilia in different models of inflammation [18-21,51], and that CRTH2 is the cellular mediator for this effect. However, whether CRTH2 signaling is relevant in a disease paradigm has not been established to date. The present finding that a small molecule inhibitor of CRTH2 is effectively attenuating eosinophil trafficking to the airway tissues hence suggests that CRTH2 is an important effector in this OVA-induced asthma model and regulates allergic inflammation *in vivo*. Since recruitment of eosinophils to inflammatory sites is considered a critical parameter in asthma and other allergic diseases [11-13], our findings also highlight the potential importance of CRTH2 as a novel therapeutic target.

Interestingly, the selective CRTH2 antagonist TM30089 equals the dual TP/CRTH2 antagonist ramatroban regarding inhibition of eosinophil recruitment *in vivo*. Ramatroban has previously not only been shown to attenuate airway inflammation in guinea pig and mouse asthma models [45], but also been effective in inhibiting eosinophil infiltration into the nasal mucosa in patients suffering from allergic rhinitis [48]. It remained elusive, however, whether inhibition of CRTH2 or TP or both receptors accounted for its anti-inflammatory efficacy in these studies. Potentially, inhibition by ramatroban of eosinophil trafficking may be explained by two different mechanisms: (i) blockade of TXA<sub>2</sub>-mediated expression of adhesion molecules on endothelial cells and/or (ii) direct inhibition of CRTH2-dependent eosinophil migration [7,52]. The present finding that both, a dual TP/CRTH2 antagonist as well as a selective CRTH2 antagonist are comparable in their ability to prevent peribronchial eosinophil infiltration in OVA-sensitized mice therefore is indicative of the notion that exclusive inhibition of CRTH2 may also be sufficient to prevent eosinophil infiltration into the airway tissues in allergic humans. Thus, it is tempting to speculate that efficacy of ramatroban is likely related to inhibition of CRTH2 rather than inhibition of TP both in rodent asthma models and in humans.

Other characteristic features of allergic asthma are mucus hypersecretion and airway remodeling [3,53]. Mice challenged by inhalation of aerosolized ovalbumin showed marked goblet cell hyperplasia, which was significantly attenuated in animals treated with the dual TP/CRTH2 antagonist ramatroban or the selective CRTH2 antagonist TM30089. Interestingly, exclusive inhibition of CRTH2 by TM30089 was equally effective as compared with dual TP/CRTH2 antagonism in ameliorating goblet cell hyperplasia. This finding is intriguing since corticosteroids, which now are mainstay asthma treatment, have variable effects on suppression of goblet cell hyperplasia [54], and offers the exciting perspective that selective inhibition of CRTH2 may be beneficial to achieve clinical improvement of allergic asthma. Further studies are warranted to investigate the molecular basis for the beneficial effects of CRTH2 antagonists in experimental allergic asthma in more detail.

Recent reports dealing with CRTH2-deficient mice have generated a highly inconsistent picture of receptor function. On one hand, OVA-sensitized mice lacking CRTH2 displayed enhanced occurrence of eosinophils in BALF [22]. Unfortunately, data on airway tissue eosinophils were not presented [22]. On the other hand and in apparent contrast, CRTH2-deficient mice from another laboratory were protected from bronchial hyperresponsiveness, and mucus production [24]. Furthermore, from a third laboratory IgE-mediated skin eosinophilic inflammation

was significantly reduced in CRTH2-deficient mice compared to wild type animals [23]. These latter authors [23] also reported that they could not confirm the previously reported increase in lung eosinophilia [22] and they stated that only further studies including CRTH2 antagonists will contribute to final conclusions in this field. Although the specific contribution of CRTH2 in allergic inflammation thus is inconsistent in studies using mice with targeted gene disruption, the findings of the present study are congruent with the notion that CRTH2 represents an eosinophilotactic receptor and provide strong support for the concept that CRTH2-signaling *in vivo* is an important molecular step in the pathogenesis of allergic asthma.

It is intriguing to note that CRTH2 is not only activated by PGD<sub>2</sub> and several of its metabolites the generation of which is dependent on the enzyme PGD synthase, but also by products of the arachidonic acid cascade that are generated independent of PGD<sub>2</sub> production. Among the latter are the TXA<sub>2</sub> metabolite 11-dehydro-TXB<sub>2</sub>, and PGF<sub>2α'</sub> that have recently been demonstrated to activate eosinophils and basophils [43, 55]. The possibility that CRTH2 can be activated in cellular contexts where PGD synthase is not present, i.e. in the absence of PGD<sub>2</sub> production, further corroborates its importance as a regulator of allergic inflammation and underscores the potential usefulness of CRTH2 antagonists as anti-asthmatic agents.

## Conclusion

Our study is the first to demonstrate efficacy of a small molecule selective CRTH2 antagonist in an experimental model of eosinophilic airway inflammation, and indicates that CRTH2-signaling is integral for orchestrating some of the pathological features characteristic for allergic asthma such as recruitment of eosinophils and mucus cell hyperplasia. Although we cannot rule out at present that inhibition of both TP and CRTH2 may be superior to obtain clinical efficacy in allergic diseases, our data certainly suggest that blockade of CRTH2 alone is sufficient to yield anti-inflammatory efficacy in experimental asthma, even to the extent that this mechanism may explain the efficacy of ramatroban. Although the OVA-induced asthma model does not reproduce all the features of the human disease, we propose that selective CRTH2 antagonists represent a novel and promising therapeutic approach to treat allergic asthma and related inflammatory diseases.

## Competing interests

The author(s) declare that they have no competing interests.



## Authors' contributions

LU, JMM, LA, MK, TU, TH, GA, CGAP, EK participated in the design and performance of the study and LU, JMM, TU, CGAP and EK drafted and wrote the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

Supported by grants from the Swedish Medical Research Council, The Medical Faculty Lund University, Asthma and Allergy Association, the European Community's Sixth Framework Programme and 7TM Pharma A/S.

## References

- Lewis RA, Soter NA, Diamond PT, Austen KF, Oates JA, Roberts LJ 2nd: **Prostaglandin D2 generation after activation of rat and human mast cells with anti-IgE.** *J Immunol* 1982, **129(4)**:1627-1631.
- Moore ML, Peebles RS Jr.: **Update on the role of prostaglandins in allergic lung inflammation: Separating friends from foes, harder than you might think.** *J Allergy Clin Immunol* 2006, **117(5)**:1036-1039.
- Luster AD, Tager AM: **T-cell trafficking in asthma: lipid mediators grease the way.** *Nat Rev Immunol* 2004, **4(9)**:711-724.
- Hata AN, Breyer RM: **Pharmacology and signaling of prostaglandin receptors: multiple roles in inflammation and immune modulation.** *Pharmacol Ther* 2004, **103(2)**:147-166.
- Fujishima H, Fukagawa K, Okada N, Takano Y, Tsubota K, Hirai H, Nagata K, Matsumoto K, Saito H: **Prostaglandin D2 induces chemotaxis in eosinophils via its receptor CRTH2 and eosinophils may cause severe ocular inflammation in patients with allergic conjunctivitis.** *Cornea* 2005, **24(8 Suppl)**:S66-S70.
- Boie Y, Sawyer N, Slipetz DM, Metters KM, Abramovitz M: **Molecular cloning and characterization of the human prostanoid DP receptor.** *J Biol Chem* 1995, **270(32)**:18910-18916.
- Hirai H, Tanaka K, Yoshie O, Ogawa K, Kenmotsu K, Takamori Y, Ichimasa M, Sugamura K, Nakamura M, Takano S, Nagata K: **Prostaglandin D2 selectively induces chemotaxis in T helper type 2 cells, eosinophils, and basophils via seven-transmembrane receptor CRTH2.** *J Exp Med* 2001, **193(2)**:255-261.
- Underwood DC, Muccitelli RM, Luttmann MA, Hay DW, Torphy TJ, Wasserman MA: **Differential antagonism of airway contractile responses to prostaglandin (PG)D2 and 9 alpha, 11 beta-PGF2 by atropine, SK&F 88046 and SQ 29,548 in the guinea pig.** *J Pharmacol Exp Ther* 1994, **268(1)**:304-310.
- Matsuoka T, Hirata M, Tanaka H, Takahashi Y, Murrata T, Kabashima K, Sugimoto Y, Kobayashi T, Ushikubi F, Aze Y, Eguchi N, Urade Y, Yoshida N, Kimura K, Mizoguchi A, Honda Y, Nagai H, Narumiya S: **Prostaglandin D2 as a mediator of allergic asthma.** *Science* 2000, **287(5460)**:2013-2017.
- Till S, Dickason R, Huston D, Humbert M, Robinson D, Larche M, Durham S, Kay AB, Corrigan C: **IL-5 secretion by allergen-stimulated CD4+ T cells in primary culture: relationship to expression of allergic disease.** *J Allergy Clin Immunol* 1997, **99(4)**:563-569.
- Humbles AA, Lloyd CM, McMillan SJ, Friend DS, Xanthou G, McKenna EE, Ghiran S, Gerard NP, Yu C, Orkin SH, Gerard C: **A critical role for eosinophils in allergic airways remodeling.** *Science* 2004, **305(5691)**:1776-1779.
- Lee JJ, Dimina D, Macias MP, Ochkur SI, McGarry MP, O'Neill KR, Protheroe C, Pero R, Nguyen T, Cormier SA, Lenkiewicz E, Colbert D, Rinaldi L, Ackerman SJ, Irvin CG, Lee NA: **Defining a link with asthma in mice congenitally deficient in eosinophils.** *Science* 2004, **305(5691)**:1773-1776.
- Rothenberg ME, Hogan SP: **The eosinophil.** *Annu Rev Immunol* 2006, **24**:147-174.
- Prussin C, Metcalfe DD: **5. IgE, mast cells, basophils, and eosinophils.** *J Allergy Clin Immunol* 2006, **117(2 Suppl Mini-Primer)**:S450-6.
- Marone G, Triggiani M, de Paulis A: **Mast cells and basophils: friends as well as foes in bronchial asthma?** *Trends Immunol* 2005, **26(1)**:25-31.
- Yoshimura-Uchiyama C, Iikura M, Yamaguchi M, Nagase H, Ishii A, Matsushima K, Yamamoto K, Shichijo M, Bacon KB, Hirai K: **Differential modulation of human basophil functions through prostaglandin D2 receptors DP and chemoattractant receptor-homologous molecule expressed on Th2 cells/DP2.** *Clin Exp Allergy* 2004, **34(8)**:1283-1290.
- Heinemann A, Schuligoi R, Sabroe I, Hartnell A, Peskar BA: **Delta 12-prostaglandin J2, a plasma metabolite of prostaglandin D2, causes eosinophil mobilization from the bone marrow and primes eosinophils for chemotaxis.** *J Immunol* 2003, **170(9)**:4752-4758.
- Spik I, Brenuchon C, Angeli V, Staumont D, Fleury S, Capron M, Trottein F, Dombrowicz D: **Activation of the prostaglandin D2 receptor DP2/CRTH2 increases allergic inflammation in mouse.** *J Immunol* 2005, **174(6)**:3703-3708.
- Shichijo M, Sugimoto H, Nagao K, Inbe H, Encinas JA, Takeshita K, Bacon KB, Gantner F: **Chemoattractant receptor-homologous molecule expressed on Th2 cells activation in vivo increases blood leukocyte counts and its blockade abrogates 13,14-dihydro-15-keto-prostaglandin D2-induced eosinophilia in rats.** *J Pharmacol Exp Ther* 2003, **307(2)**:518-525.
- Shiraishi Y, Asano K, Nakajima T, Oguma T, Suzuki Y, Shiomi T, Sayama K, Niimi K, Wakaki M, Kagyo J, Ikeda E, Hirai H, Yamaguchi K, Ishizaka A: **Prostaglandin D2-induced eosinophilic airway inflammation is mediated by CRTH2 receptor.** *J Pharmacol Exp Ther* 2005, **312(3)**:954-960.
- Almishri W, Cossette C, Rokach J, Martin JG, Hamid Q, Powell WS: **Effects of prostaglandin D2, 15-deoxy-Delta12,14-prostaglandin J2, and selective DPI and DP2 receptor agonists on pulmonary infiltration of eosinophils in Brown Norway rats.** *J Pharmacol Exp Ther* 2005, **313(1)**:64-69.
- Chevalier E, Stock J, Fisher T, Dupont M, Fric M, Fargeau H, Leport M, Soler S, Fabien S, Pruniaux MP, Fink M, Bertrand CP, McNeish J, Li B: **Cutting Edge: Chemoattractant Receptor-Homologous Molecule Expressed on TH2 Cells Plays a Restricting Role on IL-5 Production and Eosinophil Recruitment.** *J Immunol* 2005, **175(4)**:2056-2060.
- Satoh T, Moroi R, Aritake K, Urade Y, Kanai Y, Sumi K, Yokozeki H, Hirai H, Nagata K, Hara T, Utsuyama M, Hirokawa K, Sugamura K, Nishioka K, Nakamura M: **Prostaglandin D2 plays an essential role in chronic allergic inflammation of the skin via CRTH2 receptor.** *J Immunol* 2006, **177(4)**:2621-2629.
- Gonzalo J, Qiu Y, Coyle AJ, Hodge MR: **CRTH2(DP2) and not the DPI receptor mediate allergen induced mucus production and airway hyperresponsiveness.** *Am J Respir Crit Care Med* 2005, **163(5)**:A811.
- Sugimoto H, Shichijo M, Iino T, Manabe Y, Watanabe A, Shimazaki M, Gantner F, Bacon KB: **An orally bioavailable small molecule antagonist of CRTH2, ramatroban (BAY u3405), inhibits prostaglandin D2-induced eosinophil migration in vitro.** *J Pharmacol Exp Ther* 2003, **305(1)**:347-352.
- Ulven T, Kostenis E: **Minor structural modifications convert the dual TP/CRTH2 antagonist ramatroban into a highly selective and potent CRTH2 antagonist.** *J Med Chem* 2005, **48(4)**:897-900.
- Mathiesen JM, Christopoulos A, Ulven T, Royer JF, Campillo M, Heinemann A, Pardo L, Kostenis E: **On the mechanism of interaction of potent surmountable and insurmountable antagonists with the prostaglandin D2 receptor CRTH2.** *Mol Pharmacol* 2006, **69(4)**:1441-1453.
- Brandish PE, Hill LA, Zheng W, Scolnick EM: **Scintillation proximity assay of inositol phosphates in cell extracts: high-throughput measurement of G-protein-coupled receptor activation.** *Anal Biochem* 2003, **313(2)**:311-318.
- Kostenis E, Martini L, Ellis J, Waldhoer M, Heydorn A, Rosenkilde MM, Norregaard PK, Jorgensen R, Whistler JL, Milligan G: **A highly conserved glycine within linker I and the extreme C terminus of G protein alpha subunits interact cooperatively in switching G protein-coupled receptor-to-effector specificity.** *J Pharmacol Exp Ther* 2005, **313(1)**:78-87. (<http://www.cerep.fr/Cerep/Users/pages/ProductsServices/In-Vitro.asp>): . .
- Korsgren M, Erjefalt JS, Korsgren O, Sundler F, Persson CG: **Allergic eosinophil-rich inflammation develops in lungs and airways of B cell-deficient mice.** *J Exp Med* 1997, **185(5)**:885-892.
- Uller L, Lloyd CM, Rydell-Tormanen K, Persson CG, Erjefalt JS: **Effects of steroid treatment on lung CC chemokines, apoptosis and transepithelial cell clearance during development**

- and resolution of allergic airway inflammation. *Clin Exp Allergy* 2006, **36**(1):111-121.
33. Ten RM, Pease LR, McKean DJ, Bell MP, Gleich GJ: **Molecular cloning of the human eosinophil peroxidase. Evidence for the existence of a peroxidase multigene family.** *J Exp Med* 1989, **169**(5):1757-1769.
  34. Uller L, Lloyd CM, Rydell-Tormanen K, Persson CG, Erjefalt JS: **Resolution of airway disease: removal of inflammatory cells through apoptosis, egression or both?** *Trends Pharmacol Sci* 2006, **27**(9):461-466.
  35. Hata AN, Lybrand TP, Marnett LJ, Breyer RM: **Structural determinants of arylacetic acid nonsteroidal anti-inflammatory drugs necessary for binding and activation of the prostaglandin D2 receptor CRTH2.** *Mol Pharmacol* 2005, **67**(3):640-647.
  36. Peebles RS Jr., Hashimoto K, Morrow JD, Dworski R, Collins RD, Hashimoto Y, Christman JW, Kang KH, Jarzecka K, Furlong J, Mitchell DB, Talati M, Graham BS, Sheller JR: **Selective cyclooxygenase-1 and -2 inhibitors each increase allergic inflammation and airway hyperresponsiveness in mice.** *Am J Respir Crit Care Med* 2002, **165**(8):1154-1160.
  37. Gavett SH, Madison SL, Chulada PC, Scarborough PE, Qu W, Boyle JE, Tian HF, Lee CA, Langenbach R, Roggli VL, Zeldin DC: **Allergic lung responses are increased in prostaglandin H synthase-deficient mice.** *J Clin Invest* 1999, **104**(6):721-732.
  38. Persson CG, Erjefalt JS, Korsgren M, Sundler F: **The mouse trap.** *Trends Pharmacol Sci* 1997, **18**(12):465-467.
  39. Uller L, Persson CG, Kallstrom L, Erjefalt JS: **Lung tissue eosinophils may be cleared through luminal entry rather than apoptosis: effects of steroid treatment.** *Am J Respir Crit Care Med* 2001, **164**(10 Pt 1):1948-1956.
  40. Gervais FG, Cruz RP, Chateaufneuf A, Gale S, Sawyer N, Nantel F, Metters KM, O'Neill G P: **Selective modulation of chemokinesis, degranulation, and apoptosis in eosinophils through the PGD2 receptors CRTH2 and DP.** *J Allergy Clin Immunol* 2001, **108**(6):982-988.
  41. Monneret G, Gravel S, Diamond M, Rokach J, Powell WS: **Prostaglandin D2 is a potent chemoattractant for human eosinophils that acts via a novel DP receptor.** *Blood* 2001, **98**(6):1942-1948.
  42. Bohm E, Sturm GJ, Weighlofer I, Sandig H, Shichijo M, McNamee A, Pease JE, Kollrosier M, Peskar BA, Heinemann A: **11-Dehydrothromboxane B2, a stable thromboxane metabolite, is a full agonist of chemoattractant receptor-homologous molecule expressed on TH2 cells (CRTH2) in human eosinophils and basophils.** *J Biol Chem* 2004, **279**(9):7663-7670.
  43. Johnston SL, Bardin PG, Harrison J, Ritter W, Joubert JR, Holgate ST: **The effects of an oral thromboxane TP receptor antagonist BAY u 3405, on prostaglandin D2- and histamine-induced bronchoconstriction in asthma, and relationship to plasma drug concentrations.** *Br J Clin Pharmacol* 1992, **34**(5):402-408.
  44. Nagai H, Takeda H, Yamaguchi S, Tanaka H, Matsuo A, Inagaki N: **The effect of a thromboxane A2 receptor antagonist BAY-u-3405 on experimental allergic reactions.** *Prostaglandins* 1995, **50**(2):75-87.
  45. Aizawa H, Shigyo M, Nogami H, Hirose T, Hara N: **BAY u3405, a thromboxane A2 antagonist, reduces bronchial hyperresponsiveness in asthmatics.** *Chest* 1996, **109**(2):338-342.
  46. Narita S, Asakura K, Kataura A: **Effects of thromboxane A2 receptor antagonist (Bay u 3405) on nasal symptoms after antigen challenge in sensitized guinea pigs.** *Int Arch Allergy Immunol* 1996, **109**(2):161-166.
  47. Terada N, Yamakoshi T, Hasegawa M, Tanikawa H, Maesako K, Ishikawa K, Konno A: **The effect of ramatroban (BAY u 3405), a thromboxane A2 receptor antagonist, on nasal cavity volume and minimum cross-sectional area and nasal mucosal hemodynamics after nasal mucosal allergen challenge in patients with perennial allergic rhinitis.** *Acta Otolaryngol Suppl* 1998, **537**:32-37.
  48. Dogne JM, de Leval X, Benoit P, Delarge J, Masereel B: **Thromboxane A2 inhibition: therapeutic potential in bronchial asthma.** *Am J Respir Med* 2002, **1**(1):11-17.
  49. Kayasuga R, Iba Y, Hossen MA, Watanabe T, Kamei C: **The role of chemical mediators in eosinophil infiltration in allergic rhinitis in mice.** *Int Immunopharmacol* 2003, **3**(4):469-473.
  50. Honda K, Arima M, Cheng G, Taki S, Hirata H, Eda F, Fukushima F, Yamaguchi B, Hatano M, Tokuhisa T, Fukuda T: **Prostaglandin D2 reinforces Th2 type inflammatory responses of airways to low-dose antigen through bronchial expression of macrophage-derived chemokine.** *J Exp Med* 2003, **198**(4):533-543.
  51. Ishizuka T, Kawakami M, Hidaka T, Matsuki Y, Takamizawa M, Suzuki K, Kurita A, Nakamura H: **Stimulation with thromboxane A2 (TXA2) receptor agonist enhances ICAM-1, VCAM-1 or ELAM-1 expression by human vascular endothelial cells.** *Clin Exp Immunol* 1998, **112**(3):464-470.
  52. Spina D, Page CP: **Asthma -- a need for a rethink?** *Trends Pharmacol Sci* 2002, **23**(7):311-315.
  53. Ward C, Walters H: **Airway wall remodelling: the influence of corticosteroids.** *Curr Opin Allergy Clin Immunol* 2005, **5**(1):43-48.
  54. Sandig H, Andrew D, Barnes AA, Sabroe I, Pease J: **9alpha,11beta-PGF2 and its stereoisomer PGF2alpha are novel agonists of the chemoattractant receptor, CRTH2.** *FEBS Lett* 2006, **580**(2):373-379.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

