



**Syddansk Universitet**

## **Minimal Flavor Constraints for Technicolor**

Sakuma, Hidenori; Sannino, Francesco

*Published in:*

International Journal of Modern Physics A (IJMPA)

*DOI:*

[10.1142/S0217751X10050147](https://doi.org/10.1142/S0217751X10050147)

*Publication date:*

2010

*Document Version*

Submitted manuscript

[Link to publication](#)

*Citation for published version (APA):*

Sakuma, H., & Sannino, F. (2010). Minimal Flavor Constraints for Technicolor. *International Journal of Modern Physics A (IJMPA)*, 25(20), 3911-3932. DOI: 10.1142/S0217751X10050147

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

## **Minimal Flavor Constraints for Technicolor**

Hidenori S. Fukano <sup>♥\*</sup> and Francesco Sannino <sup>♥†</sup>

<sup>♥</sup> *CP<sup>3</sup>-Origins, Campusvej 55, DK-5230 Odense M, Denmark.*<sup>‡</sup>

(Dated: September 2, 2009)

### **Abstract**

We analyze the constraints on the the vacuum polarization of the standard model gauge bosons from a minimal set of flavor observables valid for a general class of models of dynamical electroweak symmetry breaking. We will show that the constraints have a strong impact on the self-coupling and masses of the lightest spin-one resonances. Our analysis is applicable to any four and higher dimensional extension of the standard model reducing to models of dynamical electroweak symmetry breaking.

arXiv:0908.2424v2 [hep-ph] 2 Sep 2009

---

<sup>‡</sup> Centre of Excellence for Particle Physics Phenomenology and the Origins of bright and dark mass.

\*E-mail: hidenori@cp3.sdu.dk

†E-mail: sannino@cp3.sdu.dk

## I. MINIMAL MODELS OF (EXTENDED) TECHNICOLOR

Dynamical electroweak symmetry breaking constitutes one of the best motivated extensions of the standard model (SM) of particle interactions.

Studies of the dynamics of gauge theories featuring fermions transforming according to higher dimensional representations of the new gauge group has led to several phenomenological possibilities [1, 2, 3, 4, 5] such as (Next) Minimal Walking Technicolor (MWT) [6] and Ultra Minimal Walking Technicolor (UMT) [7]. We will collectively refer to them as minimal models of technicolor. In [8] it was launched a coherent program to investigate different signals of minimal models of technicolor at the Large Hadron Collider experiment at CERN. Here, we also investigated in much detail, among other things also the production of the composite Higgs in association with a SM gauge boson suggested first in [9]. An interesting analysis relevant for the LHC phenomenology of low scale technicolor [10] has appeared [11].

Walking dynamics for breaking the electroweak symmetry was introduced in [12, 13, 14, 54]. It is worth noting that higher dimensional representations have been used earlier in particle physics phenomenology. Time honored examples are grand unified theories. The possibility of unifying the SM gauge interactions within a technicolor framework has been recently addressed within minimal technicolor models in [15]. The discovery [1] that theories with fermions transforming according to higher dimensional representations develop an infrared fixed point (IRFP) for an extremely small number of flavors and colors is intriguing. The dynamics of these theories is being investigated using several analytic methods not only for  $SU(N)$  gauge groups [1, 3, 16] but also for  $SO(N)$  and  $Sp(2N)$  gauge groups [17]. A better knowledge of the gauge dynamics of several nonsupersymmetric gauge theories has been useful to construct explicit UV-complete models able to break the electroweak symmetry dynamically while naturally featuring small contributions to the electroweak precision parameters [6, 18, 19, 20, 21]. These models are economical since they require the introduction of a very small number of underlying elementary fields and can feature a light composite Higgs [2, 3, 22]. Recent analyses lend further support to the latter observation [23, 24, 25]. The models feature also explicit dark matter candidates [7, 26, 27, 28] and associated interesting phenomenology [29, 30]. Moreover, extensions of the SM featuring a new underlying asymptotic free

gauge theory are naturally unitary at any arbitrary high energy scale. This strongly increases the theoretical appeal of these extensions. Another important aspect is that the underlying gauge theories can already be tested via first principle lattice computations [31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42]. Effective approaches, i.e. four and higher dimensional ones are to be considered as approximations of an underlying dynamics a la Technicolor or of an unspecified dynamics, see [43, 44, 45, 46] for recent efforts.

Whatever is the dynamical extension of the SM it will, in general, modify the vacuum polarizations of the SM gauge bosons. LEP I and II data provided direct constraints on these vacuum polarizations [47, 48, 49]. In this work we show that we can use flavor physics to provide stronger constraints than previously obtained for some of the precision observables. Our results are in agreement with the analysis made in [50, 51].

We are not attempting to provide a full theory of flavor but merely estimate the impact of a new dynamical sector, *per se*, on well known flavor observables. We will, however, assume that whatever is the correct mechanism behind the generation of the mass of the SM fermions it will lead to SM type Yukawa interactions [52]. This means that we will constrain models of technicolor with extended technicolor interactions [53, 54] entering in the general scheme of minimal flavor violation theories [55]. To be specific we will show that it is possible to provide strong constraints on the technirho and techniaxial self-couplings and masses for a general class of models of dynamical electroweak symmetry breaking. Our results can be readily applied to any extension of the SM featuring new heavy spin-one states. In particular it will severely limit the possibility to have very light spin-one resonances to occur at the LHC even if the underlying gauge theory has vanishing  $S$ -parameter.

## II. MINIMAL $\Delta F = 2$ FLAVOR CORRECTIONS FROM TECHNICOLOR

Our goal is to compute the minimal contributions, i.e. coming just from the technicolor sector, for processes in which the flavor number  $F$  changes by two units, i.e.  $\Delta F = 2$ . Here we consider  $F$  to be either the strange or the bottom number. Besides the intrinsic technicolor corrections to flavor processes one has also the corrections stemming out from extended technicolor models which are directly responsible for providing mass to the SM fermions. We will make the assumption, strongly supported by experiments, that if this

extended model exists it leads to a Yukawa sector similar to the SM one. It is, hence, by construction an extended technicolor model implementing the minimal flavor violation [55] idea. To be more specific we will determine the effects of heavy spin-one resonances mixing with the SM gauge bosons on flavor observables. We use the effective Lagrangian framework presented in [6] according to which the relevant interactions of the composite Higgs sector to the SM quarks up and down reads:

$$\mathcal{L}_{\text{yukawa}}^{\text{quark}} = \frac{\sqrt{2} m_{u_i}}{v} V_{ij} \cdot \bar{u}_{Ri} \pi^+ d_{Lj} - \frac{\sqrt{2} m_{d_i}}{v} V_{ji}^* \cdot \bar{d}_{Ri} \pi^- u_{Lj} + h.c., \quad (1)$$

where  $m_{u_i}$ , ( $u_i = u, c, t$ ) and  $m_{d_i}$ , ( $d_i = d, s, b$ ) are respectively the up and down quark masses of the  $i^{\text{th}}$  generation.  $V_{ij}$  is the  $i, j$  element of the Cabibbo-Kobayashi-Maskawa (CKM) matrix. This is our starting point which will allow us to compute the  $\Delta F = 2$  processes. We have also checked our results using the Hidden Local Gauge Symmetry [56] version of [6].




FIG. 1: Box diagrams for  $\Delta S = 2$  annihilation processes. To obtain the  $\Delta B = 2$  process, we should simply rename  $s$  with  $b$  and  $d$  with  $q$  ( $q = d, s$ ) in the various diagrams.

The diagrams contributing to the  $\Delta F = 2$  process are shown in Fig. 1. They amount to the *annihilation* process [80].

The final contribution to the  $\Delta F = 2$  amplitude is:

$$\begin{aligned} i\mathcal{M}(\Delta S = 2) &= 2 \times [i\mathcal{M}_{\square}^{(a)} + i\mathcal{M}_{\square}^{(b)} + i\mathcal{M}_{\square}^{(c)} + i\mathcal{M}_{\square}^{(d)}] \\ &= 2 \times \left( i \frac{g_{\text{EW}}}{\sqrt{2}} \right)^4 \times \frac{-i}{16\pi^2 M_W^2} \times \left[ \sum_{i,j=u,c,t} \lambda_i \lambda_j E(m_i, m_j, M_V, M_A) \right] \times Q_{\Delta F=2}, \end{aligned} \quad (2)$$

where  $m_i$ , ( $i = u, c, t$ ) indicates the  $u_i$  mass while  $M_V, M_A$  are respectively the mass of the lightest techni-vector meson and techni-axial vector one.  $Q_{\Delta F=2}$  is short for

$$Q_{\Delta F=2} = \begin{cases} (\bar{s}_L \gamma^\mu d_L)(\bar{s}_L \gamma_\mu d_L) & \text{for } F = S, \\ (\bar{b}_L \gamma^\mu q_L)(\bar{b}_L \gamma_\mu q_L) & \text{for } F = B. \end{cases} \quad (3)$$

We introduced the quantity  $\lambda_i$ :

$$\lambda_i = \begin{cases} V_{id} V_{is}^* & \text{for } K^0 - \bar{K}^0 \text{ system,} \\ V_{iq} V_{ib}^* & \text{for } B_q^0 - \bar{B}_q^0 \text{ system,} \end{cases} \quad (4)$$

encoding the information contained in the CKM matrix. Moreover,  $E(m_i, m_j, M_V, M_A)$  keeps track of the technicolor-modified gauge bosons propagators. Its cumbersome full expression is reported in the technical appendix.

It is convenient to rewrite the induced  $\Delta F = 2$  term of the Lagrangian as follows:

$$\mathcal{L}_{\text{eff}}^{\Delta F=2} = -\frac{G_F^2 M_W^2}{4\pi^2} \cdot A(a_V, a_A) \cdot Q_{\Delta F=2}, \quad (5)$$

with

$$A(a_V, a_A) \equiv \sum_{i,j=u,c,t} [\lambda_i \lambda_j \cdot E(a_i, a_j, a_V, a_A)]. \quad (6)$$

Here we have expressed all the quantities by means of the following ratios

$$a_\alpha \equiv m_\alpha^2 / M_W^2, (\alpha = i, j) \quad \text{and} \quad a_v \equiv M_v^2 / M_W^2, (v = V, A). \quad (7)$$

Indicating with  $g_{\text{EW}}$  the weak-coupling constant and  $\tilde{g}$  the coupling constant governing the massive spin-one self interactions and by expanding up to the order in  $\mathcal{O}(g_{\text{EW}}^4 / \tilde{g}^4)$  one can rewrite the previous expression as:

$$E(a_i, a_j, a_V, a_A) = E_0(a_i, a_j) + \frac{g_{\text{EW}}^2}{\tilde{g}^2} \Delta E(a_i, a_j, a_V, a_A). \quad (8)$$

The explicit expressions can be found in the appendix and are consistent with the results in [57]. The SM contribution is fully contained in  $E_0$  and the technicolor one appear first in  $\Delta E$ . The latter can be divided into a vector and an axial-vector contribution as follows:

$$\Delta E(a_i, a_j, a_V, a_A) = h(a_i, a_j, a_V) + (1 - \chi)^2 \cdot h(a_i, a_j, a_A), \quad (9)$$

where the expressions for  $h(a_i, a_j, a_v)$  are reported in the appendix. The quantity  $\chi$  was introduced first in [18, 58]. Subsequently the associated effective Lagrangian [18, 58] was extended to take into account terms involving the space-time  $\epsilon^{\mu\nu\rho\gamma}$  tensor, and topological terms, in [59] for any technicolor models for which the global symmetry group is either  $SU(N_f) \times SU(N_f)$  or  $SU(2N_f)$  breaking spontaneously respectively to  $SU(N_f)$  or  $Sp(2N_f)$ , and  $N_f$  is the number of techniflavors [81].

The axial-vector decay constant is directly proportional to the quantity  $(1 - \chi)^2$ . The vector and axial decay constant are:

$$f_V^2 = \frac{M_V^2}{\bar{g}^2} \quad f_A^2 = \frac{M_A^2}{\bar{g}^2} (1 - \chi)^2. \quad (10)$$

Note also that for  $\chi = 2$  and  $\chi = 0$  the vector and axial-vector meson contributions are identical while for  $\chi = 1$  only the direct technirho contribution survives. The limit  $\chi = 0$  and  $M_V = M_A = M$  corresponds to the *custodial technicolor* model introduced in [18, 58, 59]. In this limit the S-parameter vanishes identically because is protected by a new symmetry. We also write:

$$A(a_V, a_A) = A_0 + \frac{g_{EW}^2}{\bar{g}^2} \cdot \Delta A(a_V, a_A). \quad (11)$$

Upon taking into account the unitarity of the CKM matrix and setting  $a_u \rightarrow 0$  one has

$$A_0 = \eta_1 \cdot \lambda_c^2 \cdot \bar{E}_0(a_c) + \eta_2 \cdot \lambda_t^2 \cdot \bar{E}_0(a_t) + \eta_3 \cdot 2\lambda_c \lambda_t \cdot \bar{E}_0(a_c, a_t), \quad (12)$$

and

$$\Delta A(a_V, a_A) = \eta_1 \cdot \lambda_c^2 \cdot \Delta \bar{E}(a_c, a_V, a_A) + \eta_2 \cdot \lambda_t^2 \cdot \Delta \bar{E}(a_t, a_V, a_A) + \eta_3 \cdot 2\lambda_c \lambda_t \cdot \Delta \bar{E}(a_c, a_t, a_V, a_A), \quad (13)$$

where  $\eta_{1,2,3}$  are the QCD corrections to  $\bar{E}_0$  and  $\Delta \bar{E}$ . The explicit expressions for the functions  $\bar{E}$  and  $\Delta \bar{E}$  various expressions are provided in the appendix. The expressions simplify for

the  $\Delta\bar{E}$  in the relevant limit  $a_v \gg a_t, a_c$ :

$$\Delta\bar{E}(a_c, a_t, a_v, a_A) \simeq (7.28 \times 10^{-5}) \times \left[ \frac{1}{a_v} + \frac{(1-\chi)^2}{a_A} \right], \quad (14)$$

$$\Delta\bar{E}(a_c, a_v, a_A) \simeq (3.13 \times 10^{-8}) \times \left[ \frac{1}{a_v} + \frac{(1-\chi)^2}{a_A} \right] \quad (15)$$

$$\Delta\bar{E}(a_t, a_v, a_A) \simeq (-3.30) \times \left[ \frac{1}{a_v} + \frac{(1-\chi)^2}{a_A} \right]. \quad (16)$$

De facto, the formulae above are a reasonable approximation for  $M_{V,A} > 400$  GeV. The numerical prefactors are independent of the specific model of dynamical electroweak symmetry breaking but depend on the SM values for  $a_c$  and  $a_t$ .

### III. MINIMAL FLAVOR CONSTRAINTS

We can now compare the generic technicolor effects encoded in Eq.(5) and due primarily to the techni-vector and axial vector contributions with the CP-violation parameter  $\epsilon_K$  in the  $K^0$  meson system as well as the mass difference of the  $K^0 - \bar{K}^0$  and  $B_q^0 - \bar{B}_q^0$  mesons systems with  $q = d, s$ .

We recall that the absolute value of the CP-violation parameter in the  $K^0 - \bar{K}^0$  system is given by [61]:

$$(|\epsilon_K|)_{\text{full}} = \frac{G_F^2 M_W^2}{12 \sqrt{2} \pi^2} \times \left[ \frac{M_K}{\Delta M_K} \right]_{\text{exp.}} \times B_K f_K^2 \times [-\text{Im}A(a_v, a_A)]. \quad (17)$$

The meson mass difference in the  $Q^0 - \bar{Q}^0$ ,  $Q = (K, B_d, B_s)$  system is given by

$$\left( \Delta M_Q \right)_{\text{full}} \equiv 2 \cdot \left| \langle \bar{Q}^0 | - \mathcal{L}_{\text{eff}}^{\Delta F=2} | Q^0 \rangle \right| = \frac{G_F^2 M_W^2}{6 \pi^2} \cdot f_Q^2 \cdot M_Q \times B_Q \times |A(a_v, a_A)|, \quad (18)$$

where  $f_Q$  is the decay constant of the  $Q$ -meson and  $M_Q$  is its mass.  $B_Q$  is identified with the QCD bag parameter correcting for possible deviations of the true value of the matrix elements  $\langle \bar{Q}^0 | - \mathcal{L}_{\text{eff}}^{\Delta F=2} | Q^0 \rangle$  from its approximate value computed using the vacuum insertion approximation. This bag parameter is an intrinsic QCD contribution and we assume that the technicolor sector does not contribute to the bag parameter [82]. There are many estimates available for the bag parameters, such as the ones from the lattice [62, 63],  $1/N$ -approximation [64], etc. In this paper we use, for definitiveness, the values quoted



$G_F$	$1.1664 \times 10^{-5}$	$\text{GeV}^{-2}$	Ref. [66]
$M_W$	80.398	$\text{GeV}$	Ref. [66]
$m_t$	$161.3 \pm 1.8$	$\text{GeV}$	$\overline{\text{MS}}$ mass in Ref. [66]
$m_c$	$1.274^{+0.036}_{-0.045}$	$\text{GeV}$	$\overline{\text{MS}}$ mass in Ref. [66]
$M_K$	$497.61 \pm 0.02$	$\text{MeV}$	Ref. [66]
$\Delta M_K$	$5.292 \pm 0.0009$	$\text{ns}^{-1}$	Ref. [66]
$ \epsilon_K $	$(2.229 \pm 0.012) \times 10^{-3}$		Ref. [66]
$f_K$	155.5	$\text{MeV}$	Ref. [66]
$B_K$	$0.72 \pm 0.040$		Ref. [65]
$M_{B_d}$	$5279.5 \pm 0.3$	$\text{MeV}$	Ref. [66]
$\Delta M_{B_d}$	$0.507 \pm 0.005$	$\text{ps}^{-1}$	Ref. [66]
$f_{B_d} \sqrt{B_{B_d}}$	$225 \pm 35$	$\text{MeV}$	Ref. [65]
$M_{B_s}$	$5366.3 \pm 0.6$	$\text{MeV}$	Ref. [66]
$\Delta M_{B_s}$	$17.77 \pm 0.10$	$\text{ps}^{-1}$	Ref. [66]
$f_{B_s} \sqrt{B_{B_s}}$	$270 \pm 45$	$\text{MeV}$	Ref. [65]

TABLE I: Fermi constant ( $G_F$ ),  $W^\pm$  boson mass ( $M_W$ ), top quark and charm quark masses in the  $\overline{\text{MS}}$ -scheme ( $m_t, m_c$ ), meson masses ( $M_K, M_{B_q}$ ), indirect CP violation parameter ( $\epsilon_K$ ), meson mass difference ( $\Delta M_K, \Delta M_{B_q}$ ), decay constants ( $f_K, f_{B_q}$ ) and Bag parameters ( $B_K, B_{B_q}$ ). We show the central value for  $G_F, M_W, f_K$ .

in [65]. The experimental values of  $G_F, M_W, f_Q, M_Q, \Delta M_Q$  and the bag parameter  $B_Q$  are shown in Table I.

It is convenient to define the following quantities:

$$\delta_\epsilon \equiv \frac{g_{\text{EW}}^2}{\tilde{g}^2} \cdot \frac{\text{Im}\Delta A(a_V, a_A)}{\text{Im}A_0}, \quad (19)$$

$$\delta_{M_Q} \equiv \frac{g_{\text{EW}}^2}{\tilde{g}^2} \cdot \frac{\Delta A(a_V, a_A)}{A_0}. \quad (20)$$

Using these expressions we write  $(|\epsilon_K|)_{\text{full}}$  and  $(\Delta M_Q)_{\text{full}}$  as

$$(|\epsilon_K|)_{\text{full}} = (|\epsilon_K|)_{\text{SM}} \times (1 + \delta_\epsilon) \quad , \quad (\Delta M_Q)_{\text{full}} = (\Delta M_Q)_{\text{SM}} \times |1 + \delta_{M_Q}|. \quad (21)$$

Of course,  $(|\epsilon_K|)_{\text{SM}}$  and  $(\Delta M_Q)_{\text{SM}}$  are the SM expressions encoded in:

$$(|\epsilon_K|)_{\text{SM}} = \frac{G_F^2 M_W^2}{12 \sqrt{2} \pi^2} \times \left[ \frac{M_K}{\Delta M_K} \right]_{\text{exp}} \times B_K f_K^2 \times [-\text{Im}A_0], \quad (22)$$

$$(\Delta M_Q)_{\text{SM}} = \frac{G_F^2 M_W^2}{6 \pi^2} \cdot f_Q^2 \cdot M_Q \times B_Q \times |A_0|. \quad (23)$$

They assume the values:

$$(|\epsilon_K|)_{\text{SM}} = (2.08_{-0.13}^{+0.14}) \times 10^{-3}, \quad (24)$$

$$(\Delta M_K)_{\text{SM}} = (3.55_{-1.00}^{+1.09}) \text{ ns}^{-1}, \quad (25)$$

$$(\Delta M_{B_d})_{\text{SM}} = (0.56_{-0.16}^{+0.19}) \text{ ps}^{-1}, \quad (26)$$

$$(\Delta M_{B_s})_{\text{SM}} = (17.67_{-5.40}^{+6.38}) \text{ ps}^{-1}. \quad (27)$$

To evaluate the expressions above we used the values of  $G_F, M_W, M_Q, \Delta M_K, f_Q, B_Q$  in Table I. We also used the CKM matrix elements expressed in the Wolfenstein parameterization [67] and reported in Appendix B. We also need the QCD correcting factors  $\eta_{1,2,3}$  to evaluate  $A_0$ . Following [65] these are:

$$\eta_1 = (1.44 \pm 0.35) \cdot \left( \frac{1.3 \text{ GeV}}{m_c} \right)^{1.1}, \quad \eta_2 = 0.57, \quad \eta_3 = 0.47 \pm 0.05, \quad (28)$$

for the kaon system while we also need  $\eta_B = 0.55$  [65], corresponding to  $\eta_2$ , for the system containing a bottom quark.

Given that

$$\bar{E}_0(a_c) \simeq 2.51 \times 10^{-4}, \quad \bar{E}_0(a_t) \simeq 2.27, \quad \bar{E}_0(a_c, a_t) \simeq 2.22 \times 10^{-3}, \quad (29)$$

and that the CKM derived quantities  $\lambda_c^2, \lambda_t^2, \lambda_c \lambda_t$  are roughly of the same order for the  $B_q^0$  system we neglected the  $\eta_1 \bar{E}_0(a_c)$  and  $\eta_3 \bar{E}_0(a_c, a_t)$  terms when providing the estimates for this system. The uncertainty in Eqs (24) -(27) were deduced by propagating the theoretical ones plaguing  $\eta_{1,3}, B_K, f_{B_q}$  and  $\sqrt{B_{B_q}}$  [61, 65].

We are now ready to compare the SM value given in Eq.(24) with the experimental one in Table I and read off the constrain on  $\delta_\epsilon$  which is:

$$\delta_\epsilon = (7.05_{-7.07}^{+7.93}) \times 10^{-2} \quad (68\% \text{ C.L.}). \quad (30)$$

In order to compare the corrections associate to the kaon mass  $\Delta M_K$  we formally separate the short distance contribution from the long distance one and write

$$\Delta M_K = (\Delta M_K)_{\text{SD}} + (\Delta M_K)_{\text{LD}} . \quad (31)$$

Here  $(\Delta M_K)_{\text{SD}}$  encodes the short distance contribution which must be confronted with the technicolor one Eq.(20). The SM contribution to the short distance kaon mass difference evaluated in Eq.(25) is circa 70% of  $(\Delta M_K)_{\text{exp}}$ . The long distance contribution,  $(\Delta M_K)_{\text{LD}}$ , corresponds to the exchange of the light pseudoscalar mesons and its contribution may yield the remaining 30% of the experimental value  $(\Delta M_K)_{\text{exp}}$ . [61]. However, it is difficult to pin-point the  $(\Delta M_K)_{\text{LD}}$  contribution [61, 68] and hence we can only derive very weak constraints from  $\delta_{M_K}$ . In fact we simply require that

$$(\Delta M_K)_{\text{SD}} = (\Delta M_K)_{\text{SM}} |1 + \delta_{M_K}| \leq (\Delta M_K)_{\text{exp}} . \quad (32)$$

This means that:

$$|1 + \delta_{M_K}| \leq 2.08 \quad (68\% \text{ C.L.}) . \quad (33)$$

On the other hand the short distance contribution dominates the  $B_q^0 - \bar{B}_q^0$  mass difference [68] yielding the following constraints:

$$|1 + \delta_{M_{B_d}}| = 0.91_{-0.24}^{+0.38} \quad (68\% \text{ C.L.}) , \quad (34)$$

$$|1 + \delta_{M_{B_s}}| = 1.01_{-0.27}^{+0.44} \quad (68\% \text{ C.L.}) . \quad (35)$$

These constitute the *minimal* flavor constraints on any model of dynamical electroweak symmetry breaking. On the top of these corrections one has the ones coming from a given explicit extended technicolor model. Typically these models are hard to construct and, hence, to constrain. On the other hand assuming the existence of a successful extension, meaning that it provides the correct masses to the SM fermions and no direct flavor changing neutral currents effects, one has still to consider the experiment constraints above on the technicolor sector we have just computed.

Although the analytic formulae for  $\Delta A$  are valid for any value assumed by the vector meson masses they simplify considerably in the limit  $M_V^2, M_A^2 \gg m_t^2, m_c^2$ . We term it the

intermediate vector limit, iVL, and define the associated quantities with  $\delta_\epsilon^{(iVL)}, \delta_{M_Q}^{(iVL)}$ . They read:

$$\delta_\epsilon^{(iVL)} \simeq -2.14 \times W \quad , \quad \delta_{M_K}^{(iVL)} \simeq -0.026 \times W \quad , \quad \delta_{B_d}^{(iVL)} = \delta_{B_s}^{(iVL)} \simeq -2.90 \times W \quad , \quad (36)$$

where the numerical values depend on  $a_c, a_t, \eta_{1,2,3}, \eta_B, \lambda_i$  with

$$W = \frac{g_{EW}^2}{2\tilde{g}^2} \left[ \frac{1}{a_V} + \frac{(1-\chi)^2}{a_A} \right] \quad , \quad Y \equiv \frac{g_{EW}^2 M_W^2}{2} \left[ \Pi_{33}''(0) \right] \quad , \quad (37)$$

and  $\Pi_{33}$  the  $W_3 W_3$  corrections to the vacuum polarization due to the exchange of the new heavy vectors. The  $Y$  parameter is defined as [48]

$$Y \equiv \frac{g_{EW}^2 M_W^2}{2} \left[ \Pi_{BB}''(0) \right] \quad (38)$$

and for a generic minimal model of technicolor, i.e. in which the techniquarks are not charged under ordinary color interactions, we have:

$$Y = \frac{g_{EW}^2}{2\tilde{g}^2} \left[ \frac{1+4y^2}{a_V} + \frac{(1-\chi)^2}{a_A} \right] \quad . \quad (39)$$

The flavor constraints on the  $W$  parameter, as we shall see, are important and will provide tight constraints on the underlying technicolor dynamics, or alike models. We will then compare the limits with the ones deriving from LEP II data, i.e.  $W = (-0.2 \pm 0.8) \times 10^{-3}$  and  $Y = (0.0 \pm 1.2) \times 10^{-3}$  corresponding to the 68% C.L. constraints for a heavy Higgs in [48]. As a consistency check one can see that the expression for  $W$  and  $Y$  coincide with the ones derived in [69].

#### IV. CONSTRAINING MODELS OF DYNAMICAL ELECTROWEAK SYMMETRY BREAKING

We will now use the minimal flavor experimental information to reduce the parameter space of a general class of models of dynamical electroweak symmetry breaking.

### A. Scaled up version of QCD: The running case

If the underlying TC theory is QCD like we can impose the standard 1st and 2nd Weinberg's sum rules as shown in [6, 18, 58, 59]

$$\text{1st WSR : } f_V^2 - f_A^2 = f_\pi^2 = \left(\frac{v_{EW}}{\sqrt{2}}\right)^2, \quad (40)$$

$$\text{2nd WSR : } f_V^2 M_V^2 - f_A^2 M_A^2 = 0. \quad (41)$$

with  $f_V$  and  $f_A$  the vector and axial decay constants. One obtains exactly the expression in [6] via the re-definition  $F_i = \sqrt{2}f_i, (i = V, A, \pi)$ . Using the explicit expressions of the decay constants in terms of the coupling  $\tilde{g}$  and vector masses provided in [6, 18, 58, 59] and imposing the above sum rules we derive:

$$\frac{1}{a_V} = \frac{g_{EW}^2 S}{16\pi} - \frac{1}{a_A} \quad (\geq 0), \quad (42)$$

with the  $S$ -parameter [47] reading [6, 18, 58, 59]:

$$S \equiv 8\pi \left[ \frac{f_V^2}{M_V^2} - \frac{f_A^2}{M_A^2} \right] = \frac{8\pi}{\tilde{g}^2} [1 - (1 - \chi)^2]. \quad (43)$$

The condition above yields the following additional constraint for  $\tilde{g}$  by simply noting that the quantity  $(1 - \chi)^2$  is positive:

$$\tilde{g} < \sqrt{\frac{8\pi}{S}}. \quad (44)$$

The constraints on  $(M_A, \tilde{g})$  induced by (42) and (44) are stronger, for a given  $S$ , than the ones deriving from flavor experiments and expressed in (30)-(35). This is not surprising given that in an ordinary technicolor theory the spin-one states are very heavy. However the situation changes when allowing for a walking behavior.

### B. Walking Models

Besides the flavor constraints one has also the ones due to the electroweak precision measurements [69] as well as the unitarity constraint of  $W_L - W_L$  scattering [70]. We will consider all of them. As for the technicolor case we reduce the number of independent

parameters at the effective Lagrangian level via the 1st- and the second modified [18] Weinberg sum rules which now read [18]

$$\text{1st WSR : } f_V^2 - f_A^2 = f_\pi^2 = \left( \frac{v_{\text{EW}}}{\sqrt{2}} \right)^2, \quad (45)$$

$$\text{2nd MWSR : } f_V^2 M_V^2 - f_A^2 M_A^2 = a \cdot \frac{16\pi^2}{d(\mathbf{R})} f_\pi^4, \quad (46)$$

where  $a$  is a number expected to be positive and  $\mathcal{O}(1)$  [18].  $d(\mathbf{R})$  is the dimension of the representation of the underlying technifermions as shown in [6]. We have now:

$$M_A^2 < \frac{8\pi f_\pi^2}{S} \frac{2 - \chi}{1 - \chi} = \frac{8\pi f_\pi^2}{S} \left[ 1 + \frac{1}{\sqrt{1 - \frac{\tilde{g}^2 S}{8\pi}}} \right]. \quad (47)$$

In Figure 2, we show the allowed region in the  $(M_A, \tilde{g})$ -plane after having imposed the minimal flavor constraints due to the experimental values of  $|\epsilon_K|$  and  $\Delta M_Q$  obtained using Eqs. (30)-(35) together with the theoretical constraints for  $\tilde{g}, M_A^2$  obtained via Eqs. (44) and (47). To obtain Fig. 2, we used the expressions for  $A_0$  and  $\Delta A(a_V, a_A)$  shown respectively in (12) and (13) in which  $a_V = M_V^2/M_W^2$  reads:

$$a_V = \left[ 1 - \frac{\tilde{g}^2}{8\pi} S \right] \cdot a_A + \frac{2\tilde{g}^2}{g_{\text{EW}}^2}. \quad (48)$$

We obtained the last expression imposing the first Weinberg sum rule of Eq. (45). Given that the upper bound for  $\delta_{M_K}$  is always larger than the theoretical estimate in the region  $M_A > 200 \text{ GeV}$  we conclude that the  $\Delta M_K$  constraint is not yet very severe and hence it is not displayed in Fig. 2.

To make the plots we need also the value of the  $S$  parameters and hence we analyzed as explicit example minimal walking technicolor models.

### 1. Minimal Walking Technicolor

For definitiveness we use for  $S$  the naive MWT estimate, i.e.  $S = 1/(2\pi)$  [1, 3] while  $\tilde{g}$  is constrained via Eq. (44) to be  $\tilde{g} < 12.5$ .

We have plotted the various constraints on the  $(M_A, \tilde{g})$ -plane for MWT in the upper and lower left panel of Fig. 2. In the upper (lower) left figure we compare the 68% C.L.

(95% C.L.) allowed regions coming from the minimal flavor constraints (the darker region above the blue-dotted line) with the ones from LEP II data (region above the green-dashed line). It is clear that the flavor constraints are stronger for the 68% C.L. case but are weaker for the 95% C.L. one with respect to the constraints from LEP II data.

The region above the straight solid line is forbidden by the condition  $\tilde{g} < 12.5$  while the region below the black solid curve (on the right corner) by the condition (47) [83].

## 2. Next to Minimal Walking Technicolor (NMWT)

In this case the naive  $S$  is approximately  $1/\pi$  [1, 3] and the constraint on  $\tilde{g}$  from Eq. (44) yields  $\tilde{g} < 8.89$ . We have plotted the various constraints on the  $(M_A, \tilde{g})$ -plane for NMWT in the upper and lower right panel of Fig. 2 for the 68% C.L. and 95% C.L. constraints. We see again that the flavor constraints are stronger for the 68% C.L. case but are weaker for the 95% C.L. one.

## C. Custodial Technicolor

In the limit  $M_A = M_V = M$  and  $\chi = 0$  the effective theory acquires a new symmetry [58, 59]. This new symmetry relates a vector and an axial field and can be shown to work as a custodial symmetry for the  $S$  parameter [58, 59]. The only non-zero electroweak parameters are now:

$$W = \frac{g_{EW}^2}{\tilde{g}^2} \frac{M_W^2}{M^2}, \quad Y = \frac{g_{EW}'^2}{2\tilde{g}^2} \frac{M_W^2}{M^2} (2 + 4y^2). \quad (49)$$

It was already noted in [69] that a custodial technicolor model cannot be easily achieved via an underlying walking dynamics and should be interpreted as an independent framework. This is so since custodial technicolor models do not respect the Weinberg's sum rules [84]. This symmetry is also present in the BESS models [77, 78, 79] which will, therefore, be constrained as well. We directly compare in the Fig. 3 the constraints on the custodial technicolor parameter region  $(M, \tilde{g})$  coming from LEP II and flavor constraints and find a similar trend as for the other cases.




FIG. 2: The upper and lower left panels represent the allowed region in the  $(M_A, \tilde{g})$ -plane for MWT respectively for the 68% C.L. and 95% C.L.. A similar analysis is shown for NMWT in the right hand upper and lower panels. The region above the straight solid line is forbidden by the condition  $\tilde{g} < 12.5$  for MWT and  $\tilde{g} < 8.89$  for NMWT while the region below the black solid curve (on the right corner) by the condition (47). In the two upper (lower) plots the blue dotted lines correspond to the 68% C.L. (95% C.L.) flavor constraints while the green dashed lines are the 68% C.L. (95% C.L.) from LEP II data. The flavor constraints come only from  $\epsilon_K$  since the ones from  $\Delta M_{B_q}$  are not as strong.






FIG. 3: The left (right) panel represents the allowed region in the  $(M_A, \tilde{g})$ -plane for CT respectively for the 68% C.L. (95% C.L.). In the two upper (lower) plots the blue dotted lines correspond to the 68% C.L. (95% C.L.) flavor constraints while the green dashed lines are the 68% C.L. (95% C.L.) from LEP II data. The flavor constraints come only from  $\epsilon_K$  since the ones from  $\Delta M_{B_q}$  are not as strong.

## V. SUMMARY

Flavor constraints are relevant for models of dynamical electroweak symmetry breaking with light spin-one resonances, in fact, any model featuring spin-one resonances with the same quantum numbers of the SM gauge bosons will have to be confronted with these flavor constraints. Combining the flavor and LEP II data the new value for  $W$  at the one sigma level is  $W_{\text{avg}} \simeq (-1.6^{+3.7}_{-3.3}) \times 10^{-2}$ .

### Acknowledgments

We thank Mads T. Frandsen, Gino Isidori, Matti Järvinen, Maurizio Pierini, Luca Silvestrini, Koichi Yamawaki and Roman Zwicky for valuable comments and relevant discussions.

## APPENDIX A: RELEVANT EXPRESSIONS

We provided the explicit form of each quantity introduced in the main text starting with:

$$\begin{aligned}
E_0(a_i, a_j) = & -\frac{3}{4}h_0(a_i, a_j) - \frac{7}{4}\left(\frac{1}{a_i - 1} + \frac{1}{a_i - 1} + 1\right) \\
& + \left[\frac{a_i a_j}{a_i - a_j}\left(\frac{1}{4} - \frac{3}{2(a_i - 1)}\right) + \frac{7a_i}{4(a_i - 1)^2}\right] \ln a_i \\
& + \left[\frac{a_i a_j}{a_j - a_i}\left(\frac{1}{4} - \frac{3}{2(a_j - 1)}\right) + \frac{7a_j}{4(a_j - 1)^2}\right] \ln a_j, \tag{A1}
\end{aligned}$$

and

$$h_0(a_i, a_j) = \frac{1}{a_i - a_j} \left[ \left(\frac{a_i}{a_i - 1}\right)^2 \ln a_i - \left(\frac{a_j}{a_j - 1}\right)^2 \ln a_j - \frac{1}{a_i - 1} + \frac{1}{a_j - 1} \right]. \tag{A2}$$

We also have:

$$\Delta E(a_i, a_j, a_v, a_A) = h(a_i, a_j, a_v) + (1 - \chi)^2 \cdot h(a_i, a_j, a_A), \tag{A3}$$

where  $h(a_i, a_j, a_v)$  is given by

$$\begin{aligned}
h(a_i, a_j, a_v) = & \frac{a_i^2 \ln a_i}{(a_i - 1)^3(a_i - a_j)(a_i - a_v)} \cdot \left[ a_i^2 - \frac{3}{4}a_i^2 a_j - a_i a_j \right] \\
& + \frac{a_j^2 \ln a_j}{(a_j - 1)^3(a_j - a_i)(a_j - a_v)} \cdot \left[ a_j^2 - \frac{3}{4}a_i a_j^2 - a_i a_j \right] \\
& + \frac{a_v^2 \ln a_v}{(a_v - 1)^3(a_v - a_i)(a_v - a_j)} \cdot \left[ a_v^2 - \frac{3}{4}a_i a_j a_v - a_i a_j \right] \\
& - \frac{1}{(a_i - 1)(a_j - 1)(a_v - 1)} \cdot \left[ \frac{1}{a_i - 1} + \frac{1}{a_j - 1} + \frac{1}{a_v - 1} \right] \cdot \left[ 1 - \frac{7}{4}a_i a_j \right] \\
& - \frac{1}{(a_i - 1)(a_j - 1)(a_v - 1)} \cdot \left[ \frac{7}{2} - \frac{27}{8}a_i a_j \right]. \tag{A4}
\end{aligned}$$

We can now provide the full expression for  $A(a_v, a_A)$

$$A(a_v, a_A) = A_0 + \frac{g_{EW}^2}{\delta^2} \cdot \Delta A(a_v, a_A). \tag{A5}$$

Taking into account the unitarity constraint from the CKM matrix and setting  $a_u \rightarrow 0$  one finds:

$$A_0 = \eta_1 \cdot \lambda_c^2 \cdot \bar{E}_0(a_c) + \eta_2 \cdot \lambda_t^2 \cdot \bar{E}_0(a_t) + \eta_3 \cdot 2\lambda_c \lambda_t \cdot \bar{E}_0(a_c, a_t), \tag{A6}$$

and

$$\Delta A(a_V, a_A) = \eta_1 \cdot \lambda_c^2 \cdot \Delta \bar{E}(a_c, a_V, a_A) + \eta_2 \cdot \lambda_t^2 \cdot \Delta \bar{E}(a_t, a_V, a_A) + \eta_3 \cdot 2\lambda_c \lambda_t \cdot \Delta \bar{E}(a_c, a_t, a_V, a_A). \quad (\text{A7})$$

where  $\Delta \bar{E}(a_i, a_j, a_V, a_A) = \Delta \bar{E}(a_i, a_j, a_v) + (1 - \chi)^2 \Delta \bar{E}(a_i, a_j, a_A)$ , etc.  $\eta_{1,2,3}$  encodes the QCD corrections for  $\bar{E}_0$  and  $\Delta \bar{E}$ .

Here,  $\bar{E}_0(a_i, a_j)$  is given by

$$\begin{aligned} \bar{E}_0(a_i, a_j) &= \lim_{a_u \rightarrow 0} \left[ E_0(a_i, a_j) - E_0(a_u, a_j) - E_0(a_i, a_u) + E_0(a_u, a_u) \right] \\ &= \frac{a_i a_j}{a_i - a_j} \left[ K_0(a_i) - K_0(a_j) \right], \end{aligned} \quad (\text{A8})$$

with

$$K_0(x) = \left[ \frac{1}{4} - \frac{3}{2(x-1)} - \frac{3}{4(x-1)^2} \right] \ln x + \frac{3}{4(x-1)}. \quad (\text{A9})$$

We also have:

$$\begin{aligned} \Delta \bar{E}(a_i, a_j, a_v) &= \lim_{a_u \rightarrow 0} \left[ \Delta E(a_i, a_j, a_v) - \Delta E(a_u, a_j, a_v) - \Delta E(a_i, a_u, a_v) + \Delta E(a_u, a_u, a_v) \right] \\ &= \frac{a_i a_j \cdot \left[ (a_j - a_v)K(a_i) + (a_v - a_i)K(a_j) + (a_i - a_j)K(a_v) \right]}{(a_i - a_j)(a_j - a_v)(a_v - a_i)}, \end{aligned} \quad (\text{A10})$$

where

$$\begin{aligned} K(x) &= \frac{-\ln x}{(x-1)^3} \cdot \left[ x^2 - \frac{3}{4} a_i a_j \cdot x - a_i a_j \right] \\ &+ \frac{1}{x-1} \left[ 1 - \frac{7}{4} a_i a_j \right] \left[ \frac{1}{a_i - 1} + \frac{1}{a_j - 1} + \frac{1}{a_v - 1} \right] + \frac{1}{x-1} \left[ \frac{3}{2} + \frac{1}{8} a_i a_j \right]. \end{aligned} \quad (\text{A11})$$

Moreover

$$\begin{aligned} \bar{E}_0(a_i) &\equiv \lim_{a_j \rightarrow a_i} \bar{E}_0(a_i, a_j) \\ &= \frac{3}{2} \left( \frac{a_i}{a_i - 1} \right)^3 \ln a_i + \left[ \frac{1}{4} - \frac{9}{4(a_i - 1)} - \frac{3}{2(a_i - 1)^2} \right] a_i, \end{aligned} \quad (\text{A12})$$

$$\Delta \bar{E}(a_i, a_V, a_A) \equiv \lim_{a_j \rightarrow a_i} \Delta \bar{E}(a_i, a_j, a_V, a_A) = \Delta \bar{E}(a_i, a_v) + (1 - \chi)^2 \Delta \bar{E}(a_i, a_A), \quad (\text{A13})$$

and

$$\begin{aligned}
\Delta\bar{E}(a_i, a_v) &\equiv \lim_{a_j \rightarrow a_i} \Delta\bar{E}(a_i, a_j, a_v), \\
&= \frac{3}{4} \frac{a_i^2 a_v}{(a_i - a_v)^2} \left[ \frac{a_i^2 \ln a_i}{(a_i - 1)^3} - \frac{a_v^2 \ln a_v}{(a_v - 1)^3} \right] \\
&\quad + \frac{a_i^2}{a_i - a_v} \left[ \frac{a_i \ln a_i}{(a_i - 1)^3} - \frac{a_v \ln a_v}{(a_v - 1)^3} \right] + \frac{a_i^3}{a_i - a_v} \left[ \frac{\ln a_i}{(a_i - 1)^3} - \frac{\ln a_v}{(a_v - 1)^3} \right] \\
&\quad + \frac{9}{4} \frac{a_i^5 \ln a_i}{(a_i - a_v)(a_i - 1)^4} - \frac{3}{4} \frac{a_i^4}{(a_i - a_v)(a_i - 1)^3} \\
&\quad - \frac{a_i^2}{(a_i - 1)^2(a_v - 1)} \cdot \left[ 1 - \frac{7}{4} a_i^2 \right] \left[ \frac{2}{a_i - 1} + \frac{1}{a_v - 1} \right] \\
&\quad - \frac{a_i^2}{(a_i - 1)^2(a_v - 1)} \left[ \frac{3}{2} + \frac{1}{8} a_i^2 \right], \tag{A14}
\end{aligned}$$

Some of the formulae simplify considerably in the limit

$$a_v \gg a_c, a_t, \tag{A15}$$

yielding:

$$\Delta\bar{E}(a_c, a_t, a_v) \simeq -\frac{a_c}{a_v} \times \left\{ \frac{a_c^2 \ln a_c}{(1 - a_c)^3} - \frac{1}{1 - a_c} \left[ \frac{1}{a_t - 1} - \frac{1}{1 - a_c} + \frac{3}{2} \right] \right\}, \tag{A16}$$

$$\Delta\bar{E}(a_i, a_v) \simeq -\frac{a_i^2}{a_v} \times \left[ \frac{\frac{3}{2} a_i^3 + \frac{11}{4} a_i^2 - 2a_i}{(a_i - 1)^4} \ln a_i + \frac{\frac{1}{2} + \frac{3}{2} a_i - \frac{35}{8} a_i^2 + \frac{1}{8} a_i^3}{(a_i - 1)^2} \right]. \tag{A17}$$

## APPENDIX B: WOLFENSTEIN'S PARAMETRIZATION OF THE CKM MATRIX

The Wolfenstein parameterization [67] of the CKM matrix is:

$$V = \begin{pmatrix} 1 - \frac{1}{2} \lambda^2 - \frac{1}{8} \lambda^4 & \lambda & A \lambda^3 (\rho - i\eta) \\ -\lambda + \frac{1}{2} A^2 \lambda^5 [1 - 2(\rho + i\eta)] & 1 - \frac{1}{2} \lambda^2 - \frac{1}{8} \lambda^4 (1 + 4A^2) & A \lambda^2 \\ A \lambda^3 (1 - \bar{\rho} - i\bar{\eta}) & -A \lambda^2 + \frac{1}{2} A \lambda^4 [1 - 2(\rho + i\eta)] & 1 - \frac{1}{2} A^2 \lambda^2 \end{pmatrix} + \mathcal{O}(\lambda^6), \tag{B1}$$

where  $\lambda, A, \bar{\rho} = \rho(1 - \lambda^2/2), \bar{\eta} = \eta(1 - \lambda^2/2)$  are [66]

$$\lambda = 0.2257_{-0.001}^{+0.009}, \quad A = 0.814_{-0.022}^{+0.021}, \quad \bar{\rho} = 0.135_{-0.016}^{+0.031}, \quad \bar{\eta} = 0.349_{-0.017}^{+0.015}, \tag{B2}$$

### APPENDIX C: CORRELATION BETWEEN $|\epsilon_K|_{\text{SM}}$ AND $(\bar{\rho}, \bar{\eta})$

In our analysis, we have used the values of  $\bar{\rho}, \bar{\eta}$  in [66]. It is instructive, however, to show how  $|\epsilon_K|_{\text{SM}}$  modifies when the values  $(\bar{\rho}, \bar{\eta})$  change by a small amount. We indicate with  $[|\epsilon_K|_{\text{SM}}]_{\text{new}}$  the modified expression and with  $[|\epsilon_K|_{\text{SM}}]_{\text{old}}$  the initial value. We have then:

$$[|\epsilon_K|_{\text{SM}}]_{\text{new}} = [|\epsilon_K|_{\text{SM}}]_{\text{old}} \times \left(1 + \frac{\Delta\bar{\eta}}{\bar{\eta}_{\text{old}}}\right) - C_{\mathcal{E}} \times \left(A^4 \lambda^{10} \bar{\eta}\right)_{\text{old}} \times \Delta\bar{\rho}, \quad (\text{C1})$$

where  $\Delta_{\bar{\rho}(\bar{\eta})} \equiv \bar{\rho}(\bar{\eta})_{\text{new}} - \bar{\rho}(\bar{\eta})_{\text{old}}$  and

$$C_{\mathcal{E}} \equiv \frac{G_F^2 M_W^2}{12 \sqrt{2} \pi^2} \times \left[ \frac{M_K}{\Delta M_K} \right]_{\text{exp.}} \times B_K f_K^2 \times 2 \bar{E}_0(a_t) = 5.93 \times 10^4. \quad (\text{C2})$$

For example if we assume for  $(\bar{\rho}, \bar{\eta})_{\text{new}}$  the values  $(0.136 \pm 0.032, 0.340 \pm 0.016)$  [85] one has  $[|\epsilon_K|_{\text{SM}}]_{\text{new}} = (2.03^{+0.14}_{-0.12}) \times 10^{-3}$ . This leads to  $\delta_{\epsilon}$  is  $\delta_{\epsilon} = 0.100^{+0.076}_{-0.077}$  (68% C.L.). This shows that at the 68% C.L. we have always  $\delta_{\epsilon} > 0$ . However in models of dynamical electroweak symmetry breaking we analyzed here  $\delta_{\epsilon}^{(\text{iVL})}$  is always negative since  $W$  is always positive. Hence these value strongly reduce the allowed space of parameters  $(M_A, \tilde{g})$ . We need to go to two sigmas to allow for the introduction of vector states.

- 
- [1] F. Sannino and K. Tuominen, "Orientifold theory dynamics and symmetry breaking," Phys. Rev. D **71**, 051901 (2005) [arXiv:hep-ph/0405209].
  - [2] D. D. Dietrich, F. Sannino and K. Tuominen, "Light composite Higgs from higher representations versus electroweak precision measurements: Predictions for LHC," Phys. Rev. D **72**, 055001 (2005); *ibid* [arXiv:hep-ph/0505059]. "Light composite Higgs and precision electroweak measurements on the Z resonance: An update," Phys. Rev. D **73**, 037701 (2006) [arXiv:hep-ph/0510217].
  - [3] D. D. Dietrich and F. Sannino, "Conformal window of SU(N) gauge theories with fermions in higher dimensional representations." Phys. Rev. D **75**, 085018 (2007) [arXiv:hep-ph/0611341]. The section on the light composite Higgs is only on the archive version of the paper.
  - [4] T. A. Ryttov and F. Sannino, "Conformal Windows of SU(N) Gauge Theories, Higher Dimensional Representations and The Size of The Unparticle World," Phys. Rev. D **76**, 105004 (2007) [arXiv:0707.3166 [hep-th]].

- [5] N. D. Christensen and R. Shrock, "Technifermion representations and precision electroweak constraints," *Phys. Lett. B* **632**, 92 (2006) [arXiv:hep-ph/0509109].
- [6] R. Foadi, M. T. Frandsen, T. A. Ryttov and F. Sannino, "Minimal Walking Technicolor: Set Up for Collider Physics," *Phys. Rev. D* **76**, 055005 (2007) [arXiv:0706.1696 [hep-ph]].
- [7] T. A. Ryttov and F. Sannino, "Ultra Minimal Technicolor and its Dark Matter TIMP," *Phys. Rev. D* **78**, 115010 (2008) [arXiv:0809.0713 [hep-ph]].
- [8] A. Belyaev, R. Foadi, M. T. Frandsen, M. Jarvinen, F. Sannino and A. Pukhov, "Technicolor Walks at the LHC," *Phys. Rev. D* **79**, 035006 (2009) [arXiv:0809.0793 [hep-ph]].
- [9] A. R. Zerwekh, "Associate Higgs and gauge boson production at hadron colliders in a model with vector resonances," *Eur. Phys. J. C* **46**, 791 (2006) [arXiv:hep-ph/0512261].
- [10] E. Eichten and K. Lane, "Low-scale technicolor at the Tevatron and LHC," *Phys. Lett. B* **669**, 235 (2008) [arXiv:0706.2339 [hep-ph]].
- [11] K. Lane and A. Martin, "An Effective Lagrangian for Low-Scale Technicolor," arXiv:0907.3737 [hep-ph].
- [12] B. Holdom, "Raising The Sideways Scale," *Phys. Rev. D* **24**, 1441 (1981).
- [13] K. Yamawaki, M. Bando and K. i. Matumoto, "Scale Invariant Technicolor Model And A Technidilaton," *Phys. Rev. Lett.* **56**, 1335 (1986).
- [14] T. W. Appelquist, D. Karabali and L. C. R. Wijewardhana, "Chiral Hierarchies and the Flavor Changing Neutral Current Problem in Technicolor," *Phys. Rev. Lett.* **57**, 957 (1986).
- [15] S. B. Gudnason, T. A. Ryttov and F. Sannino, "Gauge coupling unification via a novel technicolor model," *Phys. Rev. D* **76**, 015005 (2007) [arXiv:hep-ph/0612230].
- [16] T. A. Ryttov and F. Sannino, "Supersymmetry Inspired QCD Beta Function," *Phys. Rev. D* **78**, 065001 (2008) [arXiv:0711.3745 [hep-th]].
- [17] F. Sannino, "Conformal Windows of  $SP(2N)$  and  $SO(N)$  Gauge Theories," arXiv:0902.3494 [hep-ph]. Published in *Physical Review D*.
- [18] T. Appelquist and F. Sannino, "The Physical Spectrum of Conformal  $SU(N)$  Gauge Theories," *Phys. Rev. D* **59**, 067702 (1999) [arXiv:hep-ph/9806409].
- [19] M. Harada, M. Kurachi and K. Yamawaki, " $\pi^+$  -  $\pi^0$  mass difference from the Bethe-Salpeter equation," *Phys. Rev. D* **70**, 033009 (2004) [arXiv:hep-ph/0403120].
- [20] M. Harada, M. Kurachi and K. Yamawaki, "Enhanced  $\pi^+$  -  $\pi^0$  mass difference and vanishing  $S$  in the large  $N(f)$  QCD," *Prog. Theor. Phys.* **115**, 765 (2006) [arXiv:hep-ph/0509193].

- [21] M. Kurachi and R. Shrock, “Behavior of the S parameter in the crossover region between walking and QCD-like regimes of an SU(N) gauge theory,” *Phys. Rev. D* **74**, 056003 (2006) [arXiv:hep-ph/0607231].
- [22] D. K. Hong, S. D. H. Hsu and F. Sannino, “Composite Higgs from higher representations,” *Phys. Lett. B* **597**, 89 (2004) [arXiv:hep-ph/0406200].
- [23] A. Doff and A. A. Natale, “Mass and width of a composite Higgs boson,” arXiv:0902.2379 [hep-ph].
- [24] A. Doff, A. A. Natale and P. S. Rodrigues da Silva, “Light composite Higgs from an effective action for technicolor,” *Phys. Rev. D* **77**, 075012 (2008) [arXiv:0802.1898 [hep-ph]].
- [25] A. Doff, A. A. Natale and P. S. R. da Silva, “Light composite Higgs boson from the normalized Bethe-Salpeter equation,” arXiv:0905.2981 [hep-ph].
- [26] S. B. Gudnason, C. Kouvaris and F. Sannino, “Dark Matter from new Technicolor Theories,” *Phys. Rev. D* **74**, 095008 (2006) [arXiv:hep-ph/0608055].
- [27] K. Kainulainen, K. Tuominen and J. Virkajarvi, “The WIMP of a minimal technicolor theory,” *Phys. Rev. D* **75**, 085003 (2007) [arXiv:hep-ph/0612247].
- [28] C. Kouvaris, “Dark Majorana Particles from the Minimal Walking Technicolor,” *Phys. Rev. D* **76**, 015011 (2007) [arXiv:hep-ph/0703266].
- [29] R. Foadi, M. T. Frandsen and F. Sannino, “Technicolor Dark Matter,” arXiv:0812.3406 [hep-ph]. Accepted for publication in *Brief Reports of Phys. Rev. D*.
- [30] E. Nardi, F. Sannino and A. Strumia, “Decaying Dark Matter can explain the electron/positron excesses,” *JCAP* **0901**, 043 (2009) [arXiv:0811.4153 [hep-ph]].
- [31] S. Catterall and F. Sannino, “Minimal walking on the lattice,” *Phys. Rev. D* **76**, 034504 (2007) [arXiv:0705.1664 [hep-lat]].
- [32] S. Catterall, J. Giedt, F. Sannino and J. Schneible, “Phase diagram of SU(2) with 2 flavors of dynamical adjoint quarks,” *JHEP* **0811**, 009 (2008) [arXiv:0807.0792 [hep-lat]].
- [33] Y. Shamir, B. Svetitsky and T. DeGrand, “Zero of the discrete beta function in SU(3) lattice gauge theory with color sextet fermions,” *Phys. Rev. D* **78**, 031502 (2008) [arXiv:0803.1707 [hep-lat]].
- [34] L. Del Debbio, M. T. Frandsen, H. Panagopoulos and F. Sannino, “Higher representations on the lattice: perturbative studies,” *JHEP* **0806**, 007 (2008) [arXiv:0802.0891 [hep-lat]].
- [35] L. Del Debbio, A. Patella and C. Pica, “Higher representations on the lattice: numerical

- simulations. SU(2) with adjoint fermions,” arXiv:0805.2058 [hep-lat].
- [36] T. Appelquist, G. T. Fleming and E. T. Neil, “Lattice Study of the Conformal Window in QCD-like Theories,” *Phys. Rev. Lett.* **100**, 171607 (2008) [arXiv:0712.0609 [hep-ph]].
- [37] A. Hietanen, J. Rantaharju, K. Rummukainen and K. Tuominen, “Spectrum of SU(2) gauge theory with two fermions in the adjoint representation,” *PoS LATTICE2008*, 065 (2008) [arXiv:0810.3722 [hep-lat]].
- [38] A. J. Hietanen, J. Rantaharju, K. Rummukainen and K. Tuominen, “Spectrum of SU(2) lattice gauge theory with two adjoint Dirac flavours,” *JHEP* **0905**, 025 (2009) [arXiv:0812.1467 [hep-lat]].
- [39] A. J. Hietanen, K. Rummukainen and K. Tuominen, “Evolution of the coupling constant in SU(2) lattice gauge theory with two adjoint fermions,” arXiv:0904.0864 [hep-lat].
- [40] A. Deuzeman, M. P. Lombardo and E. Pallante, “Evidence for a conformal phase in SU(N) gauge theories,” arXiv:0904.4662 [hep-ph].
- [41] L. Del Debbio, B. Lucini, A. Patella, C. Pica and A. Rago, “Conformal vs confining scenario in SU(2) with adjoint fermions,” arXiv:0907.3896 [hep-lat].
- [42] Z. Fodor, K. Holland, J. Kuti, D. Nogradi and C. Schroeder, “Nearly conformal gauge theories in finite volume,” arXiv:0907.4562 [hep-lat].
- [43] C. D. Carone, J. Erlich and M. Sher, “Holographic Electroweak Symmetry Breaking from D-branes,” *Phys. Rev. D* **76**, 015015 (2007) [arXiv:0704.3084 [hep-th]].
- [44] J. Hirn, A. Martin and V. Sanz, “Describing viable technicolor scenarios,” *Phys. Rev. D* **78**, 075026 (2008) [arXiv:0807.2465 [hep-ph]].
- [45] D. D. Dietrich and C. Kouvaris, “Constraining vectors and axial-vectors in walking technicolour by a holographic principle,” *Phys. Rev. D* **78**, 055005 (2008) [arXiv:0805.1503 [hep-ph]].
- [46] C. Nunez, I. Papadimitriou and M. Piai, “Walking Dynamics from String Duals,” arXiv:0812.3655 [hep-th].
- [47] M. E. Peskin and T. Takeuchi, “Estimation of oblique electroweak corrections,” *Phys. Rev. D* **46**, 381 (1992); M. E. Peskin and T. Takeuchi, *Phys. Rev. Lett.* **65**, 964 (1990).
- [48] R. Barbieri, A. Pomarol, R. Rattazzi and A. Strumia, “Electroweak symmetry breaking after LEP1 and LEP2,” *Nucl. Phys. B* **703**, 127 (2004) [arXiv:hep-ph/0405040].
- [49] A. J. Buras, M. E. Lautenbacher and G. Ostermaier, “Waiting For The Top Quark Mass,  $K^+ \rightarrow \pi^+ \text{ Neutrino Anti-Neutrino}$ ,  $B(S)0 - \text{Anti-}B(S)0$  Mixing And CP Asymmetries In B Decays,”



- Phys. Rev. D **50**, 3433 (1994) [arXiv:hep-ph/9403384].
- [50] M. Bona *et al.* [UTfit Collaboration], “Model-independent constraints on  $\Delta F=2$  operators and the scale of new physics,” JHEP **0803**, 049 (2008) [arXiv:0707.0636 [hep-ph]].
- [51] M. Antonelli *et al.*, “Flavor Physics in the Quark Sector,” arXiv:0907.5386 [hep-ph].
- [52] R. S. Chivukula and H. Georgi, “Composite Technicolor Standard Model,” Phys. Lett. B **188**, 99 (1987).
- [53] S. Dimopoulos and L. Susskind, “Mass Without Scalars,” Nucl. Phys. B **155**, 237 (1979).
- [54] E. Eichten and K. D. Lane, “Dynamical Breaking Of Weak Interaction Symmetries,” Phys. Lett. B **90**, 125 (1980).
- [55] G. D’Ambrosio, G. F. Giudice, G. Isidori and A. Strumia, “Minimal flavour violation: An effective field theory approach,” Nucl. Phys. B **645**, 155 (2002) [arXiv:hep-ph/0207036].
- [56] M. Bando, T. Kugo and K. Yamawaki, “Nonlinear Realization and Hidden Local Symmetries,” Phys. Rept. **164**, 217 (1988); M. Bando, T. Fujiwara and K. Yamawaki, “Generalized Hidden Local Symmetry and the A1 meson,” Prog. Theor. Phys. **79**, 1140 (1988).
- [57] T. Inami and C. S. Lim, “Effects Of Superheavy Quarks And Leptons In Low-Energy Weak Processes  $K(L) \rightarrow \text{Mu Anti-Mu}$ ,  $K^+ \rightarrow \text{Pi}^+ \text{ Neutrino Anti-Neutrino}$  And  $K^0 - \text{Anti-K}^0$ ,” Prog. Theor. Phys. **65**, 297 (1981) [Erratum-ibid. **65**, 1772 (1981)].
- [58] T. Appelquist, P. S. Rodrigues da Silva and F. Sannino, “Enhanced global symmetries and the chiral phase transition,” Phys. Rev. D **60**, 116007 (1999) [arXiv:hep-ph/9906555].
- [59] Z. y. Duan, P. S. Rodrigues da Silva and F. Sannino, “Enhanced global symmetry constraints on epsilon terms,” Nucl. Phys. B **592**, 371 (2001) [arXiv:hep-ph/0001303].
- [60] O. Kaymakcalan, S. Rajeev and J. Schechter, “Nonabelian Anomaly And Vector Meson Decays,” Phys. Rev. D **30**, 594 (1984).
- [61] G. Buchalla, A. J. Buras and M. E. Lautenbacher, “Weak decays beyond leading logarithms,” Rev. Mod. Phys. **68**, 1125 (1996) [arXiv:hep-ph/9512380].
- [62] Y. Aoki *et al.*, “The kaon B-parameter from quenched domain-wall QCD,” Phys. Rev. D **73**, 094507 (2006) [arXiv:hep-lat/0508011].
- [63] M. Okamoto, “Full determination of the CKM matrix using recent results from lattice PoS **LAT2005**, 013 (2006) [arXiv:hep-lat/0510113].
- [64] W. A. Bardeen, A. J. Buras and J. M. Gerard, “The B Parameter Beyond The Leading Order Of  $1/N$  Expansion,” Phys. Lett. B **211**, 343 (1988); J. M. Gerard, “Electroweak interactions of

- hadrons," *Acta Phys. Polon. B* **21**, 257 (1990).
- [65] U. Nierste, "Three Lectures on Meson Mixing and CKM phenomenology," arXiv:0904.1869 [hep-ph].
- [66] C. Amsler *et al.* [Particle Data Group], *Phys. Lett. B* **667**, 1 (2008).
- [67] L. Wolfenstein, "Parametrization Of The Kobayashi-Maskawa Matrix," *Phys. Rev. Lett.* **51**, 1945 (1983);
- [68] J. F. Donoghue, E. Golowich and B. R. Holstein, "Dynamics Of The Standard Model," *Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol.* **2**, 1 (1992).
- [69] R. Foadi, M. T. Frandsen and F. Sannino, "Constraining Walking and Custodial Technicolor," *Phys. Rev. D* **77**, 097702 (2008) [arXiv:0712.1948 [hep-ph]].
- [70] R. Foadi, M. Jarvinen and F. Sannino, "Unitarity in Technicolor," *Phys. Rev. D* **79**, 035010 (2009) [arXiv:0811.3719 [hep-ph]].
- [71] F. Sannino, "Dynamical Stabilization of the Fermi Scale: Phase Diagram of Strongly Coupled Theories for (Minimal) Walking Technicolor and Unparticles," arXiv:0804.0182 [hep-ph].
- [72] F. Sannino, "QCD Dual," arXiv:0907.1364 [hep-th].
- [73] E. Poppitz and M. Unsal, "Conformality or confinement: (IR)relevance of topological excitations," arXiv:0906.5156 [hep-th].
- [74] A. Armoni, "The Conformal Window from the Worldline Formalism," arXiv:0907.4091 [hep-ph].
- [75] O. Antipin, F. Sannino and K. Tuominen, work in progress.
- [76] F. Sannino and R. Zwicky, "Unparticle & Higgs as Composites," *Phys. Rev. D* **79**, 015016 (2009) [arXiv:0810.2686 [hep-ph]].
- [77] R. Casalbuoni, S. De Curtis, D. Dominici, F. Feruglio and R. Gatto, "Vector and axial vector bound states from a strongly interacting electroweak sector," *Int. J. Mod. Phys. A* **4**, 1065 (1989).
- [78] R. Casalbuoni, A. Deandrea, S. De Curtis, D. Dominici, F. Feruglio, R. Gatto and M. Grazzini, "Symmetries for vector and axial vector mesons," *Phys. Lett. B* **349**, 533 (1995) [arXiv:hep-ph/9502247].
- [79] R. Casalbuoni, F. Coradeschi, S. De Curtis and D. Dominici, "A linear moose model with pairs of degenerate gauge boson triplets," *Phys. Rev. D* **77**, 095005 (2008) [arXiv:0710.3057 [hep-ph]].

- [80] Note that the contribution of the *scattering* process to the invariant amplitude is equal to that of the *annihilation* one.
- [81] We also note that the approach used in [59] to add the  $\epsilon$  terms is the most general one given that preserves the *true* global symmetries of the underlying gauge theory. For example the interpretation of the new spin-one states as gauge bosons of a local version of the flavor symmetry allows to relate the coefficients of the  $\epsilon$  terms to the one of the gauged Wess-Zumin-Witten topological term [60]. This approach is, however, not justified on theoretical grounds.
- [82] This is a particularly good approximation when the technicolor sector does not have techni-quarks charged under ordinary color. The best examples are Minimal Walking Technicolor models.
- [83] It might be that the original theory behind Minimal Walking Technicolor, i.e. the  $SU(2)$  gauge theory with two Dirac flavors transforming according to the adjoint representation of the gauge group, develops an infrared fixed point. This case seems to be supported by recent lattice results [31, 32, 35, 38, 39]. Such a possibility was first predicted in [71] using the the beta function proposed in [16]. The Rytov-Sannino (RS) beta function results have, very recently, been shown to be consistent with new solutions of the 't Hooft anomaly matching conditions presented in [72]. Other analytic analysis further supporting the RS beta function have recently appeared [73, 74]. If the presence of such a nontrivial fixed point is confirmed this will not reduce the phenomenological interest for this theory but rather increase it both for constructing interesting new models of dynamical electroweak symmetry breaking [75] or to provide *minimal models of unparticle physics* similar to the ones suggested in [76].
- [84] One can, of course, imagine more complicated vector spectrum leading to such a symmetry.
- [85] We thank Maurizio Pierini for stressing that these are the best measured values for these quantities