



Syddansk Universitet

Fully Automated Radiosynthesis and Formulation of [11C]MeAIB Applied for In Vivo Imaging of Glioblastoma

Dam, Johan Hygum; Halle, Bo; Thisgaard, Helge; Hvidsten, Svend; Kristensen, Bjarne Winther; Någren, Kjell

Published in:
Journal of Labelled Compounds and Radiopharmaceuticals

DOI:
[10.1002/jlcr.3058](https://doi.org/10.1002/jlcr.3058)

Publication date:
2013

Document version
Publisher's PDF, also known as Version of record

Citation for published version (APA):

Dam, J. H., Halle, B., Thisgaard, H., Hvidsten, S., Kristensen, B. W., & Någren, K. (2013). Fully Automated Radiosynthesis and Formulation of [11C]MeAIB Applied for In Vivo Imaging of Glioblastoma. *Journal of Labelled Compounds and Radiopharmaceuticals*, 56(S1), S112. DOI: 10.1002/jlcr.3058

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

P-025 Fully automated radiosynthesis and formulation of [¹¹C]MeAIB applied for *in vivo* imaging of glioblastoma

Dam, Johan Hygum^{1*}; Halle, Bo²; Thisgaard, Helge¹; Hvidsten, Svend¹; Kristensen, Bjarne Winther²; Någren, Kjell¹

1 PET- & Cyclotron Unit, Department of Nuclear Medicine, Odense University Hospital, Odense, Denmark; 2 Department of Neurosurgery, Odense University Hospital, Odense, Denmark; 3 Department of Pathology, Odense University Hospital, Odense, Denmark

Objectives: Amino acids and analogues thereof play an important role in the growth of many tumor cells. The clinical value of radiolabelled amino acids such as methionine, fluoroethyltyrosine and fluorodopa has been confirmed in a plethora of studies [1]. We sought to prepare, formulate and apply a metabolically stable analogue of the amino acid alanine for system A transport, namely [*N*-methyl-¹¹C]α-methylamino-isobutyric acid, [¹¹C]MeAIB [2].

Methods: The labelling precursor, [¹¹C]MeOTf, was prepared on a Tracerlab FXc Pro by standard in-line gas-phase conversions from [¹¹C]CO₂. The [¹¹C]MeOTf was bubbled through a cooled solution of precursor in CH₃OH, CH₃CN and pentamethylpiperidine at 5 °C. The solution was warmed to 60 °C for 1 min, where 2 M NaOH was added. The heating was continued for 3 min. Shortly after, the mixture was automatically injected onto a semipreparative HPLC column via a 2 mL loop.




Figure 1: Synthesis of [¹¹C]MeAIB

The HPLC purification was performed in aqueous, buffered CH₃CN. Hence, the fraction containing the product was separated and diluted with 0.2 M H₃PO₄. This solution was loaded onto a Strata-X-C column, which was washed with sterile 0.1 M HCl and isotonic, sterile saline. The product was eluted with sterile 10 mL 2% Na₂HPO₄ through a sterile filter and into a sterile vial. Male athymic nude rats (HsdHanTM:RNU-*Foxn1^{tmu}*) were intracranially implanted in the right hemisphere with 300.000 low passage human glioblastoma single cells suspended in HBSS (Invitrogen) with 0.9% glucose (SAD 500 mg/mL). Four weeks post-implantation, rats with infiltrating glioblastomas were anesthetized and PET/CT scanned using a Siemens Inveon system after tail vein injection of [¹¹C]MeAIB, [¹¹C]methionine or [¹⁸F]FDG. Immediately after injection a dynamic acquisition was commenced.

Results: The full synthesis of [¹¹C]MeAIB including purification by prep. HPLC, solvent exchange by solid phase extraction and formulation takes about 35 min. delivering 2-5 GBq in 10 mL sterile phosphate-buffer. In all cases pH was maintained between 6.3-6.8 and the content of CH₃CN was below 350 ppm. We have produced and more than 40 productions in the described manner. After administration to orthotopically implanted nude rats the glioblastoma were visualized with a tumor to background (normal brain) ratio for [¹⁸F]FDG, [¹¹C]methionine and [¹¹C]MeAIB of 1.5, 2.3 or 8.2, respectively.

Conclusions: We have demonstrated the fast and reliable production of GMP-grade [¹¹C]MeAIB. Also, [¹¹C]MeAIB was successfully used in the imaging of human infiltrating glioblastoma in nude rats. [¹¹C]MeAIB was found to have a significantly higher tumor to background ratio than [¹¹C]methionine and [¹⁸F]FDG.

References: [1] McConathy J, et al. (2008) Cancer Metastasis Rev, 27, 555-73. [2] Någren K, et al. (2000) J Label Compd Radiopharm, 43, 1013-21.