



Syddansk Universitet

**Reduced CD300LG mRNA tissue expression, increased intramyocellular lipid content and impaired glucose metabolism in healthy male carriers of Arg82Cys in CD300LG a novel genometic cross-link between CD300LG and common metabolic phenotypes**

Støy, Julie; Kampmann, Ulla; Mengel, Annette; Magnusson, Nils E; Jessen, Niels; Grarup, Niels; Rungby, Jørgen; Stødkilde-Jørgensen, Hans; Brandslund, Ivan; Christensen, Cramer; Hansen, Torben; Pedersen, Oluf; Møller, Niels

*Published in:*

B M J Open Diabetes Research & Care

*DOI:*

[10.1136/bmjdr-2015-000095](https://doi.org/10.1136/bmjdr-2015-000095)

*Publication date:*

2015

*Document version*

Publisher's PDF, also known as Version of record

*Document license*

CC BY-NC

*Citation for published version (APA):*

Støy, J., Kampmann, U., Mengel, A., Magnusson, N. E., Jessen, N., Grarup, N., ... Møller, N. (2015). Reduced CD300LG mRNA tissue expression, increased intramyocellular lipid content and impaired glucose metabolism in healthy male carriers of Arg82Cys in CD300LG: a novel genometic cross-link between CD300LG and common metabolic phenotypes. *B M J Open Diabetes Research & Care*, 3(1), [e000095]. DOI: 10.1136/bmjdr-2015-000095

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

**Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 09. Sep. 2018

# Reduced *CD300LG* mRNA tissue expression, increased intramyocellular lipid content and impaired glucose metabolism in healthy male carriers of Arg82Cys in *CD300LG*: a novel genomemtabolic cross-link between *CD300LG* and common metabolic phenotypes

Julie Støy,<sup>1</sup> Ulla Kampmann,<sup>1</sup> Annette Mengel,<sup>2</sup> Nils E Magnusson,<sup>2</sup> Niels Jessen,<sup>3</sup> Niels Grarup,<sup>4</sup> Jørgen Rungby,<sup>5,6</sup> Hans Stødkilde-Jørgensen,<sup>7</sup> Ivan Brandslund,<sup>8</sup> Cramer Christensen,<sup>9</sup> Torben Hansen,<sup>4</sup> Oluf Pedersen,<sup>4</sup> Niels Møller<sup>1,2</sup>

**To cite:** Støy J, Kampmann U, Mengel A, et al. Reduced *CD300LG* mRNA tissue expression, increased intramyocellular lipid content and impaired glucose metabolism in healthy male carriers of Arg82Cys in *CD300LG*: a novel genomemtabolic cross-link between *CD300LG* and common metabolic phenotypes. *BMJ Open Diabetes Research and Care* 2015;**3**:e000095. doi:10.1136/bmjdr-2015-000095

Received 20 February 2015  
Revised 2 June 2015  
Accepted 26 June 2015



CrossMark

For numbered affiliations see end of article.

**Correspondence to**  
Dr Julie Støy; julistoe@rm.dk

## ABSTRACT

**Background:** *CD300LG* rs72836561 (c.313C>T, p. Arg82Cys) has in genetic-epidemiological studies been associated with the lipoprotein abnormalities of the metabolic syndrome. *CD300LG* belongs to the CD300-family of membrane-bound molecules which have the ability to recognize and interact with extracellular lipids. We tested whether this specific polymorphism results in abnormal lipid accumulation in skeletal muscle and liver and other indices of metabolic dysfunction.

**Methods:** 40 healthy men with a mean age of 55 years were characterized metabolically including assessment of insulin sensitivity by the hyperinsulinemic euglycemic clamp, intrahepatic lipid content (IHLC) and intramyocellular lipid content (IMCL) by MR spectroscopy, and  $\beta$ -cell function by an intravenous glucose tolerance test. Changes in insulin signaling and *CD300LG* mRNA expression were determined by western blotting and quantitative PCR in muscle and adipose tissue.

**Results:** Compared with the 20 controls (CC carriers), the 20 CT carriers (polymorphism carriers) had higher IMCL ( $p=0.045$ ), a reduced fasting forearm glucose uptake ( $p=0.011$ ), a trend toward lower M-values during the clamp; 6.0 mg/kg/min vs 7.1 ( $p=0.10$ ), and higher IHLC ( $p=0.10$ ). CT carriers had lower *CD300LG* mRNA expression and *CD300LG* expression in muscle correlated with IMCL ( $\beta=-0.35$ ,  $p=0.046$ ), forearm glucose uptake ( $\beta=0.37$ ,  $p=0.03$ ), and tended to correlate with the M-value ( $\beta=0.33$ ,  $p=0.06$ ), independently of *CD300LG* genotype.  $\beta$ -cell function was unaffected.

**Conclusions:** The *CD300LG* polymorphism was associated with decreased *CD300LG* mRNA expression in muscle and adipose tissue, increased IMCL, and

## Key messages

- An amino acid polymorphism in *CD300LG* (Arg82Cys), a protein which belongs to the CD300 family of lipid-binding surface glycoproteins in highly vascularized tissues, has in genetic-epidemiological studies been associated with the fasting lipoprotein abnormalities of the metabolic syndrome.
- We performed a comprehensive metabolic phenotype characterization of healthy male carriers of the identical polymorphism to test the hypothesis that the polymorphism may associate with additional metabolic phenotypes.
- The *CD300LG* polymorphism was associated with reduced expression of the gene in adipose tissue and skeletal muscle, accumulation of lipid in the skeletal muscle, and abnormalities in glucose metabolism. The findings of the study suggest a role for *CD300LG* in the regulation of common metabolic phenotypes.

abnormalities of glucose metabolism. *CD300LG* mRNA levels correlated with IMCL and forearm glucose uptake. These findings link a specific *CD300LG* polymorphism with features of the metabolic syndrome suggesting a role for *CD300LG* in the regulation of common metabolic traits.

**Trial registration number:** NCT01571609.

## BACKGROUND

The metabolic syndrome, comprising prediabetes, abdominal obesity, hypertension, and

dyslipidemia, is a risk factor for the future development of type 2 diabetes (T2DM) and cardiovascular diseases (CVD).<sup>1–3</sup> Accumulation of lipids in muscle and liver is associated with insulin resistance and is also a risk factor for T2DM and CVD.<sup>4–6</sup>

Epidemiological studies have shown that prenatal and postnatal environmental factors together with genetic factors exert major influences on the risk of metabolic disorders.<sup>7–15</sup> Each of the aspects of the metabolic syndrome has been studied in large-scale genetic association studies with a multitude of loci identified and each associated with a subtle effect on disease susceptibility.<sup>16–19</sup> The hypothesis that metabolic disorders on top of their phenotype-specific genes share overlapping causative genes was addressed in a three-stage genetic-epidemiological study combining whole exome sequencing of 2000 Danish individuals with genotyping and association studies in Danish and European individuals.<sup>20</sup> A low-frequency amino acid polymorphism in *CD300LG* (c.313 C>T, p.Arg82Cys) with a minor allele frequency of 3.5% was associated with a decreased fasting serum high-density lipoprotein (HDL) cholesterol level and an increased fasting serum triglyceride level, that is, components of the metabolic syndrome. The protein encoded by *CD300LG* belongs to the CD300 family of membrane-bound molecules, which have broad and diverse immunological actions, including the ability to recognize and interact with extracellular lipids.<sup>21–22</sup> The CD300LG protein is expressed in a broad range of tissues with highest expression in the placenta, adipose tissue, and skeletal muscle.<sup>20–23</sup> The biological functions of CD300LG are not well characterized, but the protein has been implicated in lymphocyte interaction with endothelial cells in lymph node capillaries,<sup>24</sup> endocytosis of immunoglobulins,<sup>23</sup> and binding of polar lipids.<sup>22</sup> The biological functions of CD300LG in adipose tissue and skeletal muscle are largely unknown, but with the present study we propose a role for CD300LG in lipid storage in skeletal muscle. It remains a distinct possibility that a genetically determined dysfunction of the CD300LG protein could induce abnormal lipid deposition in insulin-sensitive tissues, such as skeletal muscle contributing to insulin resistance. On the basis of these considerations, we performed a comprehensive metabolic phenotype characterization of male carriers of *CD300LG* rs72836561.

Our overall aim was to test if there are associations between the Arg82Cys polymorphism in *CD300LG* and metabolic phenotypes on whole body and tissue levels. More specifically, we aimed at testing whether *CD300LG* polymorphism carriers display reduced *CD300LG* mRNA expression in insulin-sensitive target tissues (muscle and fat) and whether any such genetic defects translate into excessive lipid accumulation in muscle and liver, features of insulin resistance, and metabolic dysfunction in a homogeneous group of male participants.

## METHODS

Study participants were recruited from a Danish biobank of nearly 10 000 participants.<sup>20</sup> Members of the biobank had been genotyped for the *CD300LG* polymorphism as part of the genetic-epidemiological study.<sup>20–25</sup> Potential participants were invited to participate in the study according to *CD300LG* genotype. Sixty male *CD300LG* CT carriers (polymorphism carriers) fulfilled inclusion criteria of age between 18 and 70 years, body mass index (BMI) <30 kg/m<sup>2</sup>, and normal glucose tolerance and were invited by letter to participate in the study; 20 CT carriers were included in the study, 31 did not respond to the invitation or did not wish to participate, and 9 had a medical condition that was incompatible with study participation. High-performance athletes were not included in the study (marathon runners, etc).


The 20 *CD300LG* CT carriers were age-matched and BMI-matched with 20 healthy male CC carriers (controls). The participants were examined over 2 days with the aim of assessing the effects of the *CD300LG* polymorphism on metabolic, cardiovascular, and bone health.<sup>25</sup> Only data on metabolic health are included in the present manuscript. Only men were included in the study to avoid the effect of the menstrual cycle in women on metabolic variables. The study participants were healthy as assessed by medical history, a physical examination, and a routine biochemical profile.

All participants provided written informed consent. The study protocol was approved by the Ethical Committee of Region Midt (protocol number 1-10-72-113-12) and the studies were conducted according to the principles of the Helsinki declaration. The study was registered at [clinicaltrials.gov](http://clinicaltrials.gov) (trial number NCT01571609).

## Design

After an overnight fast, the examination day started with a whole body DXA scan for evaluation of body composition (Hologic QDR 2000). This was followed by a 2 h basal period with infusion of a glucose and palmitate tracer to assess the rate of endogenous glucose production and free fatty acid (FFA) production. Ninety minutes into the basal period, blood was sampled for a biochemical profile. The basal period was followed by a Botnia clamp<sup>26</sup> which combines an intravenous glucose tolerance test (IVGTT; 60 min) with a hyperinsulinemic euglycemic clamp (HEC; for 120 min). Indirect calorimetry was performed during the last 30 min of the basal period and the clamp for assessment of resting energy expenditure (REE) and respiratory quotient (RQ) (Deltatrac monitor, Datex, Helsinki, Finland). Lipid and glucose oxidation was estimated after correction for protein oxidation, which was calculated from the urinary urea excretion during the study day. Muscle and subcutaneous adipose tissue (SAT) biopsies were collected 15 min into the basal period and repeated 50 min into the clamp (figure 1). MR spectroscopy of the liver for estimation of the intrahepatic lipid content (IHLIC) and

**Figure 1** Schematic overview of the examination day. The MR spectroscopy was performed on a separate day (IVGTT, intravenous glucose tolerance test).



of the left anterior tibial muscle for estimation of the intramyocellular lipid content (IMCL) was completed on the morning of a separate day. IHLC and IMCL are expressed as the arbitrary ratio between the lipid and water signal in the voxel. Study participants were non-fasting on the day of the MR spectroscopy. Study participants had refrained from major physical exercise for 48 h prior to both study days. The study participants arrived by transportation to the research facilities on the morning of the examination days.

### Intravenous catheters

A catheter was inserted retrogradely in a deep antecubital vein for sampling of venous blood from the forearm. Prior to sampling, a wrist cuff was inflated to suprasystolic blood pressure to ensure that only blood from the deep veins of the forearm was sampled. Regional blood flow was estimated by venous occlusion plethysmography. On the contralateral arm, an intravenous catheter was inserted in a heated superficial dorsal hand vein for sampling of arterialized venous blood and in an antecubital vein for infusions.<sup>27</sup>

### Muscle and adipose tissue biopsies

Muscle tissue was obtained from the lateral vastus muscle 15 cm proximal to the knee joint line by the use of a Bergstrom biopsy needle. SAT was obtained from the periumbilical region by liposuction with a 50 mL syringe. Tissue samples were cleaned from visible blood with sterile RNA-free saline, snap frozen in liquid nitrogen, and stored at  $-80^\circ\text{C}$ .

### Quantitative PCR

Real-time PCR assays were performed in duplicate for assessment of *CD300LG* mRNA tissue expression in skeletal muscle and SAT and included no-template controls, a calibrator sample, and a standard curve of five serial dilution points of the calibrator sample based on a pool

of the sampled cDNAs. Relative expressions were calculated from standard curves using software implemented in the light cycler. Target genes were normalized to endogenous controls, and expressions relative to the calibrator were calculated. For muscle tissue, cyclophilin A (CP) and  $\beta$ -actin were used as internal reference genes, whereas only CP was used in SAT. Further details on the RNA isolation and PCR are provided in the online supplementary material.

### Western blotting

Skeletal muscle biopsies obtained under basal conditions and during the clamp were used for western blotting for assessment of dynamic changes in insulin signaling downstream of the insulin receptor. Quantifications of protein phosphorylation are expressed as a ratio of phosphorylated protein to total protein measured on the same membrane for pAKT-threonine308/AKT2. For the ratios of pAS160/AS160, pAKT-serine473/AKT2, pmTOR/mTOR, and pGS/GS, the phosphorylated target and the total target were measured on separate membranes and the ratios were calculated after normalization for total protein of the membranes. Detailed descriptions of the applied methods are provided in the online supplementary material.

### Tracers

For the assessment of endogenous glucose production (EGP) during the basal period, a  $^3\text{-}^3\text{H}$ -glucose tracer (New England Nuclear Life Science Products, Boston, Massachusetts, USA) was infused for 120 min (0.15  $\mu\text{Ci}/\text{min}$ ). A priming dose  $^3\text{-}^3\text{H}$ -glucose (15  $\mu\text{Ci}$ ) was given as a bolus prior to the constant infusion. Glucose rate of appearance (Ra) was calculated from Steele's equation for non-steady state.<sup>28-30</sup> For evaluation of endogenous palmitate production during the basal period,  $9.10\text{-}^3\text{-H}$ -palmitate was infused for 60 min (0.3  $\mu\text{Ci}/\text{min}$ )



## Metabolism

from time 60 to 120. Blood was sampled for measurement of palmitate concentration and specific activity (SA) at time 0, and at time 100, 110, and 120 for steady-state analyses. Palmitate concentration and SA were analyzed by high performance liquid chromatography using  $^2\text{H}_{31}$ -palmitate as an internal standard.<sup>31</sup> Systemic palmitate flux ( $\mu\text{mol}/\text{min}$ ) was calculated using the  $9.10\text{-}^3\text{H}$ -palmitate infusion rate (dpm/min) divided by the steady-state palmitate SA (dpm/ $\mu\text{mol}$ ).<sup>32</sup> Regional palmitate balances were estimated using blood flow, SA, and palmitate concentrations from arterial and venous samples and calculated as previously described.<sup>33</sup>

### IVGTT and HEC

The Botnia clamp model was used to assess  $\beta$ -cell function and insulin sensitivity on the same study day.<sup>26</sup> A bolus of 25 g glucose from a 50% glucose solution was given at time 120 min followed by blood sampling at time 120, 122, 124, 126, 128, 130 for estimation of first-phase insulin release (FPIR) and C-peptide concentrations. Blood was sampled at time 140, 160, and 180 min for assessment of second-phase insulin release (SPIR) and C-peptide concentrations. FPIR was quantified as the time 120–130 and SPIR as the time 130–180 incremental area under the curve (iAUC) insulin concentration.<sup>26</sup> The clamp was started at time 180 with an insulin infusion rate of 1 mU/kg/min (Insulin Actrapid, Novo-Nordisk, Copenhagen, Denmark) until time 300. Plasma glucose was clamped at 5 mmol/L by adjusting an intravenous infusion of 20% glucose (200 g/L, SAD, Copenhagen, Denmark) according to plasma glucose measurements every 10 min (Beckmann Instruments, Palo Alto, California, USA). Steady-state serum insulin concentrations were measured at time 280, 290, and 300 min during the clamp. Insulin sensitivity in the two genotype groups was calculated as the mean glucose infusion rates (mg/kg/min) during the last 30 min of the clamp, the M-value.<sup>34</sup>

### MR spectroscopy

$^1\text{H}$ -MR spectroscopy was used to assess IHLC and IMCL using a Signa Excite 1.5 Tesla Twin Speed scanner (GE Medical systems, Milwaukee, Wisconsin, USA). The MR spectroscopy of the liver and muscle included point resolved spectroscopy sequences as described earlier.<sup>35</sup> Spectrum quality was checked by visual inspection during each examination and the mean full width at half maximum was 10.0 (95% CI 9.5 to 10.5) Hz for liver spectra and 10.8 (95% CI 10.0 to 11.6) Hz for muscle spectra. Data processing provided an estimate of the lipid-to-water ratio in the voxel (LC Model, Dr Steven Provencher, Ontario, Canada).<sup>36</sup> Three CT carriers were excluded from the analysis of IHLC due to technical difficulties with the liver spectroscopy.

### Assays

Plasma lipoproteins and procalcitonin were analyzed in the Department of Clinical Biochemistry, Aarhus

University Hospital. Proinsulin (intact proinsulin kit, Dakocytomation, Glostrup, Denmark), insulin (DAKO, Glostrup, Denmark), C-peptide (ALPCO, Salem, New Hampshire, USA), and cortisol (EIA-1887, AH-Diagnostic) were analyzed using ELISA-based kits. FFAs were analyzed with a commercial kit (Wako chemicals, Neuss, Germany). High-sensitive C reactive protein (BAM 17072 and MAB 17071, R&D Systems Europe Ltd, Abingdon, UK) and osteoprotegerin (OPG/TNFRSF11B, R&D Systems Europe Ltd, Abingdon, UK) were analyzed using an in-house ELISA assay. Adiponectin was analyzed using a commercial time-resolved immunofluorometric assay (TR-IFMAs; Autodelphia, PerkinElmer, Turku, Finland). Glucagon was analyzed by an in-house radioimmunoassay.<sup>37</sup> Plasma mannose-binding lectin was determined by a TR-IFMA as previously described.<sup>38</sup> Blood for analyses of proinsulin was sampled at time 90 min (basal period) and for glucagon at time 100, 110, 120, 180, 280, 290, and 300.

### Statistical analysis

Statistical analyses were performed using SPSS V.21.0. Results are presented as mean $\pm$ 95% CI or median $\pm$ IQR. A Student t test was applied to compare the two genotype groups. Non-parametric data were log-transformed to obtain normality. For data that did not achieve normal distribution on log-transformation, between-group comparisons were performed using a Wilcoxon-Mann-Whitney U test. p-values <0.05 were considered significant. A multiple regression model was applied to adjust the M-value for steady-state serum insulin concentration (clamp). The analyses of FPIR and SPIR included adjustment for the M-value. Linear regression with the log-transformed *CD300LG* mRNA level as the independent variable was used to test for correlation with relevant outcome measures. A Fisher's exact test was used to test for association between dichotomous variables. A mixed analysis of variance with repeated measurements was used for analysis of data available for both the basal period and the clamp. The between-participant factor was genotype and the within-participant factor was condition (fasting or insulin-stimulated). The within-participant effect of insulin stimulation was evaluated by a paired t test. Main outcomes were also adjusted for mean arterial pressure. Power calculations were based on the glucose infusion rates during the clamp, and with 20 study participants in each genotype group, the statistical power to detect a 20% difference in glucose infusion rates during the clamp was >80%. Box plots were generated using a web tool for generation of box plots (<http://boxplot.tyerslab.com/>).<sup>39</sup>

## RESULTS

### Anthropometrics

Clinical characteristics of the study participants are presented in [table 1](#).

**Table 1** Anthropometric data on the 40 male study participants according to *CD300LG* rs72836561 CC and CT genotype

Variable	CC (n=20)	CT (n=20)	p-Value
<b>Anthropometry</b>			
Age (years)	55.1 (50.8 to 59.2)	55.0 (50.8 to 59.3)	0.99
BMI (kg/m <sup>2</sup> )	24.6 (23.8 to 25.5)	24.5 (23.4 to 25.6)	0.79
Systolic blood pressure (mm Hg)	130 (126 to 133)	140 (132 to 147)	0.02
Diastolic blood pressure (mm Hg)	78 (75 to 82)	84 (81 to 87)	0.02
Waist circumference (cm)	91.9 (88.5 to 95.4)	92.5 (89.0 to 95.9)	0.82
Current smokers (yes/no)	1/19	5/15	0.18
<b>Body composition from DXA scan</b>			
Lean body mass (kg)	59.0 (56.7 to 61.3)	58.3 (55.3 to 61.4)	0.72
Fat mass (kg)	16.9 (15.1 to 18.6)	16.9 (15.0 to 18.8)	0.99
Fat per cent (%)	21.4 (19.6 to 23.2)	21.5 (19.9 to 23.1)	0.90
<b>Fasting blood samples</b>			
HbA1c (IFCC, mmol/mol)	35.2 (33.8 to 36.5)	35.7 (34.5 to 36.9)	0.56
HbA1c (%)	5.4 (5.2 to 5.5)	5.4 (5.3 to 5.5)	0.65
Glucose (mmol/L)	5.4 (5.3 to 5.6)	5.5 (5.2 to 5.7)	0.81
Total cholesterol (mmol/L)	5.3 (5.0 to 5.6)	5.1 (4.7 to 5.4)	0.18
LDL (mmol/L)	3.4 (3.1 to 3.7)	3.1 (2.8 to 3.4)	0.19
HDL (mmol/L)*	1.4 (1.1 to 1.7)	1.2 (1.0 to 1.5)	0.42
Triglyceride (mmol/L)*	1.1 (0.8 to 1.4)	1.3 (0.7 to 1.7)	0.49
Insulin (pmol/L)*	34.7 (24.5 to 50.0)	39.7 (26.8 to 54.0)	0.61
Proinsulin (pmol/L)*	3.41 (2.71 to 4.80)	4.00 (2.73 to 5.44)	0.26
Proinsulin/insulin ratio	0.12 (0.09 to 0.15)	0.13 (0.10 to 0.15)	0.67
C-peptide (pmol/L)*	362 (283 to 384)	375 (292 to 544)	0.22
Proinsulin/C-peptide	0.011 (0.009 to 0.013)	0.012 (0.009 to 0.014)	0.78
HOMA-B (%)	55.3 (44.7 to 65.9)	61.8 (47.4 to 76.1)	0.45
HOMA-IR	1.27 (1.03 to 1.52)	1.55 (1.07 to 1.99)	0.28

Mean and 95% CI.

\*Median and IQR.

BMI, body mass index; CC, control; CT, polymorphism carriers; HbA1c, glycated haemoglobin; HDL, high-density lipoprotein; IFCC, international federation of clinical chemistry; LDL, low-density lipoprotein.

### Substrate metabolism and insulin sensitivity

Forearm metabolism was evaluated in the basal period by assessing arteriovenous differences of glucose and FFA. Fasting EGP and palmitate kinetics were evaluated using a 3-<sup>3</sup>H-glucose tracer and a 9.10-<sup>3</sup>H-palmitate tracer, respectively (table 2). The glucose uptake in the forearm was reduced in CT carriers under basal conditions, but no differences were detected between groups in rates of EGP or in palmitate kinetics.

Insulin sensitivity was assessed by use of an HEC. Arteriovenous substance differences were also evaluated during insulin stimulation. A trend toward lower insulin sensitivity in CT carriers was detected; the M-value in the CT group was 5.96 mg/kg/min (95% CI 4.99 to 6.93) compared with 7.07 (6.08 to 8.05) in the CC group (p=0.10; figure 2). Steady-state insulin concentrations during the clamp was 404 pmol/L (95% CI 358 to 450) in CT carriers compared with 418 (391 to 444) in CC carriers (p=0.58). Adjusting the M-value for serum insulin concentration during the steady state of the clamp had some effect on the results (p=0.08). There was no effect of genotype on arteriovenous balances of glucose or FFA during the HEC. There were no between-group differences in forearm blood flow during the basal period or during the clamp (data not shown).

### β-cell function

β-cell function was assessed by FPIR and SPIR in response to intravenous administration of 25 g glucose (figure 3). There were no genotype-group differences in the iAUC of glucose, insulin, or C-peptide from time 120–130 (glucose, p=0.97; FPIR, p=0.37; C-peptide, p=0.18) or time 130–180 (glucose, p=0.93; SPIR, p=0.20; C-peptide, p=0.28). Adjusting the data for the M-value did not change the results.

### Energy expenditure

Fasting and insulin-stimulated REE and RQ were obtained by indirect calorimetry. The genotype of the study participants did not affect basal and insulin-stimulated energy expenditure and substrate oxidation (table 2).

### MR spectroscopy

IMCL was 0.005 au (95% CI 0.0038 to 0.0062) in the CT carriers compared with 0.004 in the controls (0.0029 to 0.0044), p=0.045. The median IHLC in CT-carriers was 0.07 (IQR 0.036–0.18) compared with 0.043 (0.024–0.068) in the controls (p=0.10; figure 4A, B).

**Table 2** Various measures of insulin sensitivity and energy expenditure in the 40 male study participants according to *CD300LG* rs72836561 CC and CT genotype

	CC (n=20)		CT (n=20)		p-value
	Basal	Clamp	Basal	Clamp	
Indirect calorimetry					
Respiratory quotient	0.81 (0.79 to 0.83)	0.92 (0.90 to 0.94)	0.82 (0.80 to 0.84)	0.92 (0.90 to 0.94)	<0.01*/0.24†/0.39‡
Energy expenditure (kcal/24 h)	1617 (1555 to 1679)	1713 (1650 to 1776)	1675 (1581 to 1769)	1721 (1637 to 1804)	<0.01*/0.29†/0.12‡
Glucose oxidation (mg/kg/min)	1.03 (0.75 to 1.32)	2.46 (2.15 to 2.79)	1.28 (0.98 to 1.59)	2.46 (2.16 to 2.76)	<0.01*/0.49†/0.32‡
Protein oxidation (mg/kg/min)	0.49 (0.39 to 0.59)	0.45 (0.35 to 0.56)	0.55 (0.40 to 0.70)	0.65 (0.52 to 0.79)	0.65*/0.56†/0.11‡
Lipid oxidation (mg/kg/min)	0.81 (0.69 to 0.93)	0.33 (0.22 to 0.44)	0.76 (0.61 to 0.91)	0.26 (0.16 to 0.37)	<0.01*/0.20†/0.77‡
Forearm glucose uptake (mmol/100 mL/min)/10 <sup>3</sup>	0.27 (0.18 to 0.36)	2.43 (1.75 to 3.12)	0.11 (0.02 to 0.20)	2.37 (1.38 to 3.35)	<0.01*/0.01†/0.86‡
FFA (mmol/L)					
FFA, arterial	1.07 (0.90 to 1.25)	0.06 (0.04 to 0.07)	0.96 (0.82 to 1.1)	0.07 (0.05 to 0.08)	<0.01*/0.32†/0.29‡
FFA, venous	1.07 (0.93 to 1.21)	0.06 (0.04 to 0.07)	0.95 (0.77 to 1.13)	0.07 (0.05 to 0.09)	<0.01*/0.39†/0.33‡
Endogenous glucose production (mg/kg/min)	1.72 (1.35 to 1.84)	–	1.62 (1.43 to 2.05)	–	0.94†
Palmitate flux (µmol/min)	173 (145 to 201)	–	182 (151 to 213)	–	0.80†
Forearm palmitate kinetics (µmol/100 mL/min)					
Palmitate uptake	0.080 (0.063 to 0.098)	–	0.073 (0.058 to 0.088)	–	0.53†
Palmitate net release	0.0006 (–0.024 to 0.015)	–	–0.011 (–0.028 to 0.005)	–	0.53†
Palmitate release	0.0734 (0.053 to 0.108)	–	0.057 (0.034 to 0.112)	–	0.20†
Miscellaneous					
Glucagon (pg/mL)	48.9 (40.7 to 57.1)	35.0 (27.0)	49.5 (41.9 to 57.0)	34.8 (29.2 to 40.4)	<0.01*/0.84†/0.83‡
Adiponectin (mg/L)	9.5 (7.7 to 11.3)		8.5 (6.9 to 10.0)		0.37†
Procalcitonin (µg/L)	0.04 (0.02 to 0.04)		0.04 (0.02 to 0.04)		0.95¶
Osteoprotegerin (pg/mL)	669 (550 to 948)		758 (566 to 878)		0.75§
Cortisol (ng/mL)	66 (52 to 91)		66 (52 to 85)		0.71¶
HsCRP (mg/L)	1.35 (0.39 to 3.63)		1.24 (0.66 to 2.96)		0.91§
Mannan-binding lectin (nm/mL)	1084 (355 to 1400)		404 (161 to 1874)		0.59¶

Mean and 95% CI unless otherwise indicated.

\*p-value from paired t test comparing basal to insulin-stimulated values.

†p-value from Student t test (comparison of mean basal values).

‡Mixed ANOVA where data are available from the basal period and from the clamp. The factors genotype and condition (basal/clamp) were included in the model. p-values represent interaction between genotype and condition.

§Median and IQR. p-value from Student t test of log-transformed data.

¶Median and IQR. p-value from Wilcoxon-Mann-Whitney U test.


ANOVA, analysis of variance; CC, control; CT, polymorphism carriers; FFA, free fatty acid; HsCRP, high-sensitive C reactive protein.





## Metabolism

**Figure 3** (A–B) Plasma glucose concentration (mmol/L) by *CD300LG* genotype during the IVGTT (time 120–180) and during the HEC (time 180–300) (A) and FPIR and SPIR (pmol/L) by *CD300LG* genotype during the IVGTT (time 120–180) and insulin concentrations during the HEC (time 180–300) (B). The CT group is represented by black triangles and the CC group is represented by black circles. The whiskers correspond to 95% CI. AUC, area under the curve; CC, control; CT, polymorphism carriers; FPIR, first-phase insulin release; HEC, hyperinsulinemic euglycemic clamp; SPIR, second-phase insulin release.




insights into dynamic measures of glucose and lipid turnover on whole body and tissue level and provided in-depth characterization of the metabolic phenotypes of the study participants. In the genetic-epidemiological gene discovery study, despite using a large study cohort, no signal was detected for additional metabolic traits or phenotypes for the *CD300LG* polymorphism (other than the observed changes in the lipoprotein profile).<sup>20</sup> Furthermore, the study participants in this study were heterogeneous with respect to gender and metabolic health, and associations between the *CD300LG* polymorphism and metabolic traits and phenotypes were tested on fasting levels of glucose and insulin, or using dichotomized data with study participants grouped into diabetic/non-diabetic. Potentially, the effects of the polymorphism on dynamic measures of glucose and lipid metabolism (or on unmeasured variables such as IMCL)

could have been missed in the large-scale study. In the present study we aimed at detecting novel associations between the *CD300LG* polymorphism and metabolic phenotypes through a detailed phenotype characterization of the study participants.

The CT polymorphism induced a significant reduction in *CD300LG* mRNA expression in adipose tissue and muscle. In SAT, large amounts of tissue were required to extract sufficient mRNA for cDNA synthesis reducing the number of participants available for the analyses. Owing to low yields of RNA from SAT, we were unable to inspect the integrity of the RNA on an agarose gel. On the other hand, yields of mRNA from muscle tissue were high and we were able to evaluate RNA integrity. Consequently, gene expression data from skeletal muscle seem solid, whereas the findings in AT should be interpreted more cautiously. Whether the reduced *CD300LG* mRNA

**Figure 4** (A–B) Box plot representing the median IMCL (A) and IHLC (B) as assessed by MR spectroscopy by *CD300LG* genotype. Other definitions of the plot are described in the legend to [figure 2](#). CC, control; CT, polymorphism carriers; IHCL, intrahepatic lipid content; IMCL, intramyocellular lipid content.




expression in skeletal muscle and adipose tissue in polymorphism carriers translates into quantitative and/or structural changes in the *CD300LG* protein is currently unknown. Future studies of the impact of the *CD300LG* polymorphism on human metabolism could include studies of the effect of the polymorphism on *CD300LG* protein expression and structure.

Our data show that CT carriers had a higher IMCL and tended to have higher IHLC than controls as evaluated by MR spectroscopy. A considerable body of evidence suggests that excessive accumulation of lipids in muscle and liver in sedentary persons is associated with insulin resistance,<sup>4–6</sup> possibly mediated by specific intracellular lipid fractions such as diacylglycerol (DAG) and ceramide.<sup>40</sup> It has been shown that in obesity, increased IMCL is associated with increased intracellular DAG and ceramide contents.<sup>41–42</sup> However, the cause and effect relationship of

these associations remains to be finally established. Our data support and expand the concept of IMCL-associated insulin resistance by providing evidence that in humans a specific polymorphism in *CD300LG* leads to the combination of high IMCL and indices of abnormal glucose turnover. The biological roles of the CD300 group of transmembrane molecules are not well characterized, but it appears that the CD300 receptors exist in activating and inhibitory forms, which engage in specific interactions with major polar lipids of cell membranes;<sup>21–22</sup> whether and, if so, how *CD300LG* is involved in lipid turnover in skeletal muscle cells remains elusive. Furthermore, the effect of dysfunction or reduced amounts of the *CD300LG* protein in polymorphism carriers on this proposed functional relationship is unknown, but it may include intracellular lipid accumulation and insulin resistance as suggested by results from this study.

**Figure 5** (A–B) Box plot representing the median relative expression of *CD300LG* mRNA in skeletal muscle (A) and in SAT (B) by *CD300LG* genotype. Other definitions of the plot are described in the legend to [figure 2](#). CC, control; CT, polymorphism carriers; SAT, subcutaneous adipose tissue.



Although we only saw a tendency towards lower M-values during the HEC in CT-carriers, we observed a lower fasting glucose uptake in the forearm when compared with the controls and skeletal muscle *CD300LG* mRNA expression correlated well with IMCL, M-value, and forearm glucose uptake when all participants were assessed as a whole. Taken together, these findings strongly suggest the existence of insulin resistance, although the fasting glucose uptake is also determined by insulin-independent mechanisms. We observed no effect of *CD300LG* genotype on forearm glucose uptake during the clamp, but a trend toward reduced insulin sensitivity in CT carriers. A decreased suppression of EGP during the clamp due to hepatic insulin resistance could have affected the results, but in healthy participants the relatively high insulin dose (1 mU/kg/min) has been shown to suppress the EGP by 98%.<sup>43</sup> Unfortunately, EGP was only determined under fasting conditions as the Botnia clamp model has not been

validated for simultaneous use of a glucose tracer. Consequently, with the present data, we are unable to assess the role of the liver in glucose turnover during the clamp. In this context, it should be noticed that the experimental procedures of the current investigation are relatively demanding and complex, therefore only allowing for a limited number of participants to be investigated. In addition, many genetic and lifestyle-related environmental factors may introduce biological noise and affect the outcome in an investigation such as ours.

The effect of the *CD300LG* polymorphism on clinical measures of insulin sensitivity did not translate into changes that were detectable using western blot analysis of signaling pathways downstream of the insulin receptor in skeletal muscle. We did detect a clear dynamic response to insulin when comparing basal conditions with insulin-stimulated conditions, but this response did not differ between genotype groups. We obtained a broad biochemical profile to explore the mechanisms

underlying the metabolic changes in *CD300LG* CT carriers. No differences between genotype groups were detected for the measured endocrine hormones and inflammatory markers.

For practical purposes, we used the Botnia clamp model which combines measures of  $\beta$ -cell function and insulin sensitivity on a single study day.<sup>26</sup> Whereas the glucose clamp offers a sensitive and reproducible way of evaluating insulin sensitivity, the IVGTT may be less accurate for determining  $\beta$ -cell function.<sup>44</sup> We are not aware of any data linking the *CD300LG* polymorphism to  $\beta$ -cell function, and we did not detect any effects of *CD300LG* on any measures of insulin secretion.

As mentioned, the relatively low number of study participants was a limitation to our study as it may reduce the statistical power to draw solid conclusions. The finding of a correlation between *CD300LG* expression and measures of insulin sensitivity despite the low number of participants therefore suggests the existence of a relatively robust signal and indicates that the findings are not merely reflecting a type I error. With the low frequency of the *CD300LG* polymorphism, it required a large biobank of 10 000 participants to include 20 suitable male CT carriers in the study. We did not include women in the study due to the use of radioactive tracers and the impact of menstrual cycle on metabolic measurements. It would, however, be of interest to include women in a future *CD300LG* phenotype study as the effects on lipid traits were more pronounced in women in the original genetic-epidemiological study.<sup>20</sup>

Although the effect of the *CD300LG* polymorphism on the fasting serum HDL cholesterol and triglyceride level was discrete in the genetic-epidemiological study,<sup>20</sup> the effect of the polymorphism on fasting serum HDL cholesterol was marked when compared with previously genome wide association study (GWAS)-identified HDL cholesterol-associated variants.<sup>16</sup> The association between the *CD300LG* polymorphism and lipid traits was detected using several thousand study participants, suggesting that we were underpowered to detect a significant effect on fasting serum HDL cholesterol and triglyceride levels in this study.

## CONCLUSIONS

Our data point to an effect of the *CD300LG* rs72836561 polymorphism on IMCL, fasting forearm glucose uptake, and on expression of *CD300LG* mRNA in SAT and muscle. Independently of the genotype, *CD300LG* mRNA expression was found to associate with IMCL and measures of glucose turnover. Conclusively, our data link a specific *CD300LG* polymorphism with features of the metabolic syndrome suggesting a role for *CD300LG* in the regulation of common metabolic traits.

### Author affiliations

<sup>1</sup>Department of Internal Medicine and Endocrinology, Aarhus University Hospital, Aarhus, Denmark

<sup>2</sup>Medical Research Laboratories, Institute for Clinical Medicine, Aarhus University, Aarhus, Denmark

<sup>3</sup>Research Laboratory for Biochemical Pathology, Institute for Clinical Medicine, Aarhus University, Aarhus, Denmark

<sup>4</sup>Faculty of Health and Medical Sciences, The Novo Nordisk Foundation Center for Basic Metabolic Research, University of Copenhagen, Copenhagen, Denmark

<sup>5</sup>Center for Diabetes Research and Department of Clinical Pharmacology, University Hospital of Copenhagen, Copenhagen, Denmark

<sup>6</sup>Department of Biomedicine, Aarhus University, Aarhus, Denmark

<sup>7</sup>The MR Research Center, Aarhus University Hospital, Aarhus, Denmark

<sup>8</sup>Department of Clinical Biochemistry, Vejle Hospital, Vejle, Denmark

<sup>9</sup>Department of Medicine, Vejle Hospital, Vejle, Denmark

**Acknowledgements** The authors wish to acknowledge laboratory technicians Lisa Buus, Merete Møller, and Lone Kvist from the Medical research laboratories, Institute for Clinical Medicine, Aarhus University, for invaluable assistance with biochemical analyses and assistance during the study days. They would like to acknowledge laboratory technician Helle Zibrandtsen from the Research laboratory for biochemical pathology, Institute for Clinical Medicine, Aarhus University, for introducing JS to western blotting. They would also like to acknowledge laboratory technician Dorthe Bødker Jensen from the Centre for carbohydrate recognition and signaling, Department of molecular biology and genetics, Aarhus University, for assisting with the gene expression analyses and for the use of laboratory equipment for gene expression analyses. They are grateful to the 40 study participants for their participation in the study.

**Contributors** All authors were responsible for the conception and design of the study described in the manuscript, drafting of the manuscript, and its critical revision for intellectual content. All authors approved the version to be published.

**Funding** The study was conducted by LuCamp investigators (<http://www.lucamp.org>) funded by the Lundbeck Foundation and a full scholarship for JS from Aarhus University. The Novo Nordisk Foundation Center for Basic Metabolic Research is an independent Research Center at the University of Copenhagen partially funded by an unrestricted donation from the Novo Nordisk Foundation.

**Competing interests** None declared.

**Ethics approval** The study protocol was approved by the Ethical Committee of Region Midt (protocol number 1-10-72-113-12), Denmark.

**Provenance and peer review** Not commissioned; externally peer reviewed.

► Additional material is available. To view please visit the journal online (<http://dx.doi.org/10.1136/bmjdr-2015-000095>).

**Open Access** This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## REFERENCES

- Mottillo S, Filion KB, Genest J, *et al*. The metabolic syndrome and cardiovascular risk a systematic review and meta-analysis. *J Am Coll Cardiol* 2010;56:1113–32.
- Isomaa B, Almgren P, Tuomi T, *et al*. Cardiovascular morbidity and mortality associated with the metabolic syndrome. *Diabetes Care* 2001;24:683–9.
- Kelly T, Yang W, Chen CS, *et al*. Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond)* 2008;32:1431–7.
- Pan DA, Lillioja S, Milner MR, *et al*. Skeletal muscle membrane lipid composition is related to adiposity and insulin action. *J Clin Invest* 1995;96:2802–8.
- Bachmann OP, Dahl DB, Brechtel K, *et al*. Effects of intravenous and dietary lipid challenge on intramyocellular lipid content and the relation with insulin sensitivity in humans. *Diabetes* 2001;50:2579–84.
- Korenblat KM, Fabbrini E, Mohammed BS, *et al*. Liver, muscle, and adipose tissue insulin action is directly related to intrahepatic



- triglyceride content in obese subjects. *Gastroenterology* 2008;134:1369–75.
7. Park YW, Zhu S, Palaniappan L, *et al*. The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988–1994. *Arch Intern Med* 2003;163:427–36.
  8. Boney CM, Verma A, Tucker R, *et al*. Metabolic syndrome in childhood: association with birth weight, maternal obesity, and gestational diabetes mellitus. *Pediatrics* 2005;115:e290–6.
  9. Hales CN, Barker DJ. Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Int J Epidemiol* 2013;42:1215–22.
  10. Almgren P, Lehtovirta M, Isomaa B, *et al*. Heritability and familiarity of type 2 diabetes and related quantitative traits in the Botnia Study. *Diabetologia* 2011;54:2811–19.
  11. Andersen G, Hansen T, Pedersen O. Genetics of common forms of glycaemia with pathological impact on vascular biology: are we on the right track? *Curr Mol Med* 2005;5:261–74.
  12. Köbberling J, Tillil H. *Empirical risk figures for first-degree relatives of non-insulin dependent diabetic patients. The genetics of diabetes mellitus*. London: Academic Press, 1982:201–9.
  13. Pedersen O, Echwald SM. Genetics of human obesity. In: LeRoith D, Olefsky JM, Taylor SI, eds. *Diabetes—a fundamental and clinical text*. 3rd edn. Lippincott, Williams & Williams, 2004:827–38.
  14. Kotchen TA, Kotchen JM, Grim CE, *et al*. Genetic determinants of hypertension: identification of candidate phenotypes. *Hypertension* 2000;36:7–13.
  15. Tuomi T, Santoro N, Caprio S, *et al*. The many faces of diabetes: a disease with increasing heterogeneity. *Lancet* 2014;383:1084–94.
  16. Teslovich TM, Musunuru K, Smith AV, *et al*. Biological, clinical and population relevance of 95 loci for blood lipids. *Nature* 2010;466:707–13.
  17. Scott RA, Lagou V, Welch RP, *et al*. Large-scale association analyses identify new loci influencing glycemic traits and provide insight into the underlying biological pathways. *Nat Genet* 2012;44:991–1005.
  18. Ehret GB, Munroe PB, Rice KM, *et al*. Genetic variants in novel pathways influence blood pressure and cardiovascular disease risk. *Nature* 2011;478:103–9.
  19. Randall JC, Winkler TW, Kutalik Z, *et al*. Sex-stratified genome-wide association studies including 270,000 individuals show sexual dimorphism in genetic loci for anthropometric traits. *PLoS Genet* 2013;9:e1003500.
  20. Albrechtsen A, Grarup N, Li Y, *et al*. Exome sequencing-driven discovery of coding polymorphisms associated with common metabolic phenotypes. *Diabetologia* 2013;56:298–310.
  21. Borrego F. The CD300 molecules: an emerging family of regulators of the immune system. *Blood* 2013;121:1951–60.
  22. Cannon JP, O'Driscoll M, Litman GW. Specific lipid recognition is a general feature of CD300 and TREM molecules. *Immunogenetics* 2012;64:39–47.
  23. Takatsu H, Hase K, Ohmae M, *et al*. CD300 antigen like family member G: a novel Ig receptor like protein exclusively expressed on capillary endothelium. *Biochem Biophys Res Commun* 2006;348:183–91.
  24. Umemoto E, Hayasaka H, Bai Z, *et al*. Novel regulators of lymphocyte trafficking across high endothelial venules. *Crit Rev Immunol* 2011;31:147–69.
  25. Stoy J, Grarup N, Horlyck A, *et al*. Blood pressure levels in male carriers of Arg82Cys in CD300LG. *PLoS ONE* 2014;9:e109646.
  26. Tripathy D, Wessman Y, Gullstrom M, *et al*. Importance of obtaining independent measures of insulin secretion and insulin sensitivity during the same test: results with the Botnia clamp. *Diabetes Care* 2003;26:1395–401.
  27. Moller N, Butler PC, Antsiferov MA, *et al*. Effects of growth hormone on insulin sensitivity and forearm metabolism in normal man. *Diabetologia* 1989;32:105–10.
  28. Steele R. Influences of glucose loading and of injected insulin on hepatic glucose output. *Ann N Y Acad Sci* 1959;82:420–30.
  29. Debedo RC, Steele R, Altszuler N, *et al*. On the hormonal regulation of carbohydrate metabolism; studies with C14 GLUCOSE. *Recent Prog Horm Res* 1963;19:445–88.
  30. Radziuk J, Norwich KH, Vranic M. Experimental validation of measurements of glucose turnover in nonsteady state. *Am J Physiol* 1978;234:E84–93.
  31. Nielsen S, Guo Z, Johnson CM, *et al*. Splanchnic lipolysis in human obesity. *J Clin Invest* 2004;113:1582–8.
  32. Jensen MD, Rogers PJ, Ellman MG, *et al*. Choice of infusion-sampling mode for tracer studies of free fatty acid metabolism. *Am J Physiol* 1988;254(5 Pt 1):E562–5.
  33. Jensen MD. Regulation of forearm lipolysis in different types of obesity. In vivo evidence for adipocyte heterogeneity. *J Clin Invest* 1991;87:187–93.
  34. DeFronzo RA, Tobin JD, Andres R. Glucose clamp technique: a method for quantifying insulin secretion and resistance. *Am J Physiol* 1979;237:E214–23.
  35. Maersk M, Belza A, Stodkilde-Jorgensen H, *et al*. Sucrose-sweetened beverages increase fat storage in the liver, muscle, and visceral fat depot: a 6-mo randomized intervention study. *Am J Clin Nutr* 2012;95:283–9.
  36. Provencher SW. Estimation of metabolite concentrations from localized in vivo proton NMR spectra. *Magn Reson Med* 1993;30:672–9.
  37. Orskov H, Thomsen HG, Yde H. Wick chromatography for rapid and reliable immunoassay of insulin, glucagon and growth hormone. *Nature* 1968;219:193–5.
  38. Thiel S, Moller-Kristensen M, Jensen L, *et al*. Assays for the functional activity of the mannan-binding lectin pathway of complement activation. *Immunobiology* 2002;205:446–54.
  39. Krzywinski M, Altman N. Visualizing samples with box plots. *Nat Methods* 2014;11:119–20.
  40. Shulman GI. Ectopic fat in insulin resistance, dyslipidemia, and cardiometabolic disease. *N Engl J Med* 2014;371:1131–41.
  41. Dube JJ, Amati F, Toledo FG, *et al*. Effects of weight loss and exercise on insulin resistance, and intramyocellular triacylglycerol, diacylglycerol and ceramide. *Diabetologia* 2011;54:1147–56.
  42. Moro C, Galgani JE, Luu L, *et al*. Influence of gender, obesity, and muscle lipase activity on intramyocellular lipids in sedentary individuals. *J Clin Endocrinol Metab* 2009;94:3440–7.
  43. Bergman RN, Finegood DT, Ader M. Assessment of insulin sensitivity in vivo. *Endocr Rev* 1985;6:45–86.
  44. Hermans MP, Levy JC, Morris RJ, *et al*. Comparison of tests of beta-cell function across a range of glucose tolerance from normal to diabetes. *Diabetes* 1999;48:1779–86.

# Reduced *CD300LG* mRNA tissue expression, increased intramyocellular lipid content and impaired glucose metabolism in healthy male carriers of Arg82Cys in *CD300LG*: a novel genometic cross-link between *CD300LG* and common metabolic phenotypes

Julie Støy, Ulla Kampmann, Annette Mengel, Nils E Magnusson, Niels Jessen, Niels Grarup, Jørgen Rungby, Hans Stødkilde-Jørgensen, Ivan Brandslund, Cramer Christensen, Torben Hansen, Oluf Pedersen and Niels Møller

*BMJ Open Diab Res Care* 2015 3:  
doi: 10.1136/bmjdr-2015-000095

---

Updated information and services can be found at:  
<http://drc.bmj.com/content/3/1/e000095>

---

*These include:*

## Supplementary Material

Supplementary material can be found at:  
<http://drc.bmj.com/content/suppl/2015/08/26/3.1.e000095.DC1>

## References

This article cites 42 articles, 7 of which you can access for free at:  
<http://drc.bmj.com/content/3/1/e000095#BIBL>

## Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

## Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

---

## Topic Collections

Articles on similar topics can be found in the following collections  
[Metabolism](#) (10)

---

## Notes

---

To request permissions go to:  
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:  
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:  
<http://group.bmj.com/subscribe/>