



# Syddansk Universitet

## Draft Genome Sequence of Parabacteroides goldsteinii with Putative Novel Metallo--Lactamases Isolated from a Blood Culture from a Human Patient

Krogh, Thøger Jensen; Agergaard, Charlotte Nielsen; Møller-Jensen, Jakob; Justesen, Ulrik Stenz

Published in: Genome Announcements

DOI:

10.1128/genomeA.00937-15

Publication date: 2015

Document version Publisher's PDF, also known as Version of record

Citation for pulished version (APA):

Krogh, T. J., Agergaard, C. N., Møller-Jensen, J., & Justesen, U. S. (2015). Draft Genome Sequence of Parabacteroides goldsteinii with Putative Novel Metallo--Lactamases Isolated from a Blood Culture from a Human Patient. Genome Announcements, 3(4), [e00937-15]. DOI: 10.1128/genomeA.00937-15

**General rights** 

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
  You may not further distribute the material or use it for any profit-making activity or commercial gain
  You may freely distribute the URL identifying the publication in the public portal?

Take down policy
If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.





# Draft Genome Sequence of *Parabacteroides goldsteinii* with Putative Novel Metallo- $\beta$ -Lactamases Isolated from a Blood Culture from a Human Patient

Thøger Jensen Krogh,<sup>a</sup> Charlotte Nielsen Agergaard,<sup>b</sup> Jakob Møller-Jensen,<sup>a</sup> Ulrik Stenz Justesen<sup>b</sup>

Department of Biochemistry and Molecular Biology, University of Southern Denmark, Odense, Denmark<sup>a</sup>; Department of Clinical Microbiology, Odense University Hospital, Odense, Denmark<sup>b</sup>

Parabacteroides goldsteinii was isolated from a blood culture. Genomic DNA was sequenced using a MiSeq sequencer and assembled using the SPAdes genome assembler. The draft genome sequence was 6,851,868 bp, spanning 282 contigs of 5,253 coding sequences, 66 tRNAs, and 5 rRNAs. Several putative novel metallo-β-lactamases were discovered.

Received 13 July 2015 Accepted 14 July 2015 Published 20 August 2015

**Citation** Krogh TJ, Agergaard CN, Møller-Jensen J, Justesen US. 2015. Draft genome sequence of *Parabacteroides goldsteinii* with putative novel metallo- $\beta$ -lactamases isolated from a blood culture from a human patient. Genome Announc 3(4):e00937-15. doi:10.1128/genomeA.00937-15.

Copyright © 2015 Krogh et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Address correspondence to Ulrik Stenz Justesen, ulrik.stenz.justesen@rsyd.dk.

*parabacteroides goldsteinii* is an anaerobic, Gram-negative rod belonging to the microbiota of the human gut (1, 2).

Here, we report the draft genome sequence of a *P. goldsteinii* strain isolated from a blood culture of a human patient. Antimicrobial susceptibility testing with gradient MIC strips demonstrated resistance toward clindamycin (>256 mg/liter), piperacillin-tazobactam (>256 mg/liter), and meropenem (>32 mg/liter) but susceptibility toward metronidazole (0.38 mg/liter). A meropenem-EDTA double-ended gradient strip was positive, indicating the presence of a metallo- $\beta$ -lactamase (3). In anaerobic bacteria, metallo- $\beta$ -lactamases are usually demonstrated only in *Bacteroides fragilis* and encoded by the *cfiA* gene.

The genomic DNA of the *P. goldsteinii* strain was purified using the DNeasy blood and tissue kit (Qiagen), according to protocol.

Paired-end libraries with an insert size averaging 350 bp were generated using the Illumina Nextera DNA sample preparation kit. The DNA was sequenced via an Illumina MiSeq benchtop sequencer with 150-bp reads at a 30× theoretical coverage. Reads were merged using PEAR version 0.9.5 (4), and *de novo* genome assembly was done via SPAdes version 3.0 (5) available through Illumina BaseSpace.

The final assembly consisted of 282 contigs with an  $N_{50}$  of 60,312, totaling 6,851,868 bp, and with a GC content of 43.46%. For comparison, the three available sequenced *P. goldsteinii* strains listed on the NCBI genome database (strains dnLKV18, CL02T12C30, and DSM 19448) are 6.49 Mb to 7.09 Mb with a GC content of 43.30% to 43.50%.

Annotation was carried out via the NCBI Prokaryotic Genome Annotation Pipeline, identifying 5 rRNAs, 66 tRNAs, and 5,253 coding sequences (CDS), of which 2,268 of the latter code for purely hypothetical proteins with no obvious homology-inferred function. A search for resistance-associated genes via the Res-Finder tool (version 2.1, accessed 2 June 2015) (settings, 30% ID, 20% length—lowest possible) (6) identified genes that accounted

for clindamycin (*ermF*) and tetracycline (*tetX* and *tetQ*) resistance, but no genes that account for resistance toward piperacillin-tazobactam or meropenem.

Genome analysis led to the discovery of nine genes coding for putative novel  $\beta$ -lactamases, of which six are putative novel metallo- $\beta$ -lactamases. All of the nine genes are located on different contigs with low to no internal primary sequence homology and low to no primary sequence homology for any of the  $\beta$ -lactamases in the ResFinder, Brenda, and NCBI databases (accessed 2 June 2015). Tertiary structure predictions via Phyre2 predict a general  $\beta$ -lactamase-like  $\alpha\beta\beta\alpha$  fold for all nine CDS (7, 8). These findings may represent several new subclasses of  $\beta$ -lactamases.

**Nucleotide sequence accession numbers.** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number LFJV000000000. The version described in this paper is version LFJV01000000.

## ACKNOWLEDGMENT

This research was supported by the involved departments.

### **REFERENCES**

- 1. Song Y, Liu C, Lee J, Bolanos M, Vaisanen ML, Finegold SM. 2005. "Bacteroides goldsteinii sp. nov." isolated from clinical specimens of human intestinal origin. J Clin Microbiol 43:4522–4527. http://dx.doi.org/10.1128/JCM.43.9.4522-4527.2005.
- Sakamoto M, Benno Y. 2006. Reclassification of Bacteroides distasonis, Bacteroides goldsteinii and Bacteroides merdae as Parabacteroides distasonis gen. nov., comb. nov., Parabacteroides goldsteinii comb. nov. and Parabacteroides merdae comb. nov. Int J Syst Evol Microbiol 56:1599–1605. http://dx.doi.org/10.1099/ijs.0.64192-0.
- 3. Bogaerts P, Engelhardt A, Berhin C, Bylund L, Ho P, Yusof A, Glupczynski Y. 2008. Evaluation of a new meropenem-EDTA double-ended Etest strip for the detection of the cfiA metallo-beta-lactamase gene in clinical isolates of *Bacteroides fragilis*. Clin Microbiol Infect 14:973–977. http://dx.doi.org/10.1111/j.1469-0691.2008.02065.x.

- 4. Zhang J, Kobert K, Flouri T, Stamatakis A. 2014. PEAR: a fast and accurate Illumina PairedEnd reAd mergeR. Bioinformatics 30:614–620. http://dx.doi.org/10.1093/bioinformatics/btt593.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol 19:455–477. http://dx.doi.org/10.1089/cmb.2012.0021.
- 6. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund
- O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. J Antimicrob Chemother 67:2640–2644. http://dx.doi.org/10.1093/jac/dks261.
- 7. Carfi A, Pares S, Duée E, Galleni M, Duez C, Frère JM, Dideberg O. 1995. The 3-D structure of a zinc metallo-beta-lactamase from *Bacillus cereus* reveals a new type of protein fold. EMBO J 14:4914–4921.
- Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. 2015. The Phyre2 Web portal for protein modeling, prediction and analysis. Nat Protoc 10:845–858. http://dx.doi.org/10.1038/nprot.2015.053.