



Syddansk Universitet

ImmunoChip analysis identifies association of the RAD50/IL13 region with human longevity

Flachsbart, Friederike; Ellinghaus, David; Gentschew, Liljana; Heinsen, Femke-Anouska; Caliebe, Amke; Christiansen, Lene; Nygaard, Marianne; Christensen, Kaare; Blanché, Hélène; Deleuze, Jean-François; Derbois, Céline; Galan, Pilar; Büning, Carsten; Brand, Stephan; Peters, Anette; Strauch, Konstantin; Müller-Nurasyid, Martina; Hoffmann, Per; Nöthen, Markus M; Lieb, Wolfgang; Franke, Andre; Schreiber, Stefan; Nebel, Almut

Published in:
Aging Cell

DOI:
[10.1111/accel.12471](https://doi.org/10.1111/accel.12471)

Publication date:
2016

Document version
Publisher's PDF, also known as Version of record

Document license
CC BY

Citation for published version (APA):

Flachsbart, F., Ellinghaus, D., Gentschew, L., Heinsen, F-A., Caliebe, A., Christiansen, L., ... Nebel, A. (2016). ImmunoChip analysis identifies association of the RAD50/IL13 region with human longevity. *Aging Cell*, 15(3), 585-588. DOI: 10.1111/accel.12471

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 09. Sep. 2018



SHORT TAKE

ImmunoChip analysis identifies association of the *RAD50/IL13* region with human longevity

Friederike Flachsbart,^{1*} David Ellinghaus,^{1*} Liljana Gentschew,¹ Femke-Anouska Heinsen,¹ Amke Caliebe,² Lene Christiansen,³ Marianne Nygaard,^{3,4} Kaare Christensen,^{3,4,5} Hélène Blanché,⁶ Jean-François Deleuze,^{6,7} Céline Derbois,⁷ Pilar Galan,⁸ Carsten Büning,⁹ Stephan Brand,¹⁰ Anette Peters,^{11,12,13} Konstantin Strauch,^{14,15} Martina Müller-Nurasyid,^{12,14,16} Per Hoffmann,^{17,18,19} Markus M. Nöthen,^{17,18} Wolfgang Lieb,²⁰ Andre Franke,¹ Stefan Schreiber^{1,21} and Almut Nebel¹

¹Institute of Clinical Molecular Biology, Kiel University, Kiel, Germany

²Institute of Medical Informatics and Statistics, Kiel University, Kiel, Germany

³Epidemiology, Biostatistics and Biodemography, Department of Public Health, University of Southern Denmark, Odense, Denmark

⁴Department of Clinical Genetics, Odense University Hospital, Odense, Denmark

⁵Department of Clinical Biochemistry and Pharmacology, Odense University Hospital, Odense, Denmark

⁶Fondation Jean Dausset-Centre du Polymorphisme Humain (CEPH), Paris, France

⁷Centre National de Génotypage CNG-IG-CEA, Evry, France

⁸Université Sorbonne Paris Cité-UREN, Unité de Recherche en Epidémiologie Nutritionnelle; U557 Inserm; U1125 Inra; Cnam; Université Paris 13, CRNH IdF, Bobigny, France

⁹Department of Gastroenterology, Hepatology and Endocrinology, Charité, Campus Mitte, Berlin, Germany

¹⁰Department of Medicine II – Grosshadern, Ludwig-Maximilians-University Munich, Munich, Germany

¹¹Institute of Epidemiology II, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany

¹²DZHK (German Centre for Cardiovascular Research), Partner Site Munich Heart Alliance, Munich, Germany

¹³German Center for Diabetes Research, Neuherberg, Germany

¹⁴Institute of Genetic Epidemiology, Helmholtz Zentrum München – German Research Center for Environmental Health, Neuherberg, Germany

¹⁵Institute of Medical Informatics, Biometry and Epidemiology, Chair of Genetic Epidemiology, Ludwig-Maximilians-University Munich, Munich, Germany

¹⁶Department of Medicine I, Ludwig-Maximilians-University Munich, Munich, Germany

¹⁷Institute of Human Genetics, University of Bonn, Bonn, Germany

¹⁸Department of Genomics, Life and Brain Center, University of Bonn, Bonn, Germany

¹⁹Division of Medical Genetics, University Hospital Basel and Department of Biomedicine, University of Basel, Basel, Switzerland

²⁰Institute of Epidemiology and Popgen Biobank, Kiel University, Kiel, Germany

²¹Clinic for Internal Medicine I, University Hospital of Schleswig-Holstein, Kiel, Germany

Correspondence

Almut Nebel, Institute of Clinical Molecular Biology, Kiel University, Schittenhelmstraße 12, 24105 Kiel, Germany. Tel.: +49-431-597-1373; fax: +49-431-597-2196; e-mail: a.nebel@mucosa.de

*Equal contribution of authors.

Accepted for publication 22 February 2016

Summary

Human longevity is characterized by a remarkable lack of confirmed genetic associations. Here, we report on the identification of a novel locus for longevity in the *RAD50/IL13* region on chromosome 5q31.1 using a combined European sample of 3208 long-lived individuals (LLI) and 8919 younger controls. First, we performed a large-scale association study on 1458 German LLI (mean age 99.0 years) and 6368 controls (mean age 57.2 years) by targeting known immune-associated loci covered by the ImmunoChip. The analysis of 142 136 autosomal single nucleotide polymorphisms (SNPs) revealed an ImmunoChip-wide significant signal ($P_{\text{ImmunoChip}} = 7.01 \times 10^{-9}$) for the SNP rs2075650 in the *TOMM40/APOE* region, which has been previously described in the context of human longevity. To identify novel susceptibility loci, we selected 15 markers with $P_{\text{ImmunoChip}} < 5 \times 10^{-4}$ for replication in two samples from France (1257 LLI, mean age 102.4 years; 1811 controls, mean age 49.1 years) and Denmark (493 LLI, mean age 96.2 years; 740 controls, mean age 63.1 years). The association at SNP rs2706372 replicated in the French study collection and showed a similar trend in the Danish participants and was also significant in a meta-analysis of the combined French and Danish data after adjusting for multiple testing. In a meta-analysis of all three samples, rs2706372 reached a *P*-value of $P_{\text{ImmunoChip+Repl}} = 5.42 \times 10^{-7}$ (OR = 1.20; 95% CI = 1.12–1.28). SNP rs2706372 is located in the extended *RAD50/IL13* region. *RAD50* seems a plausible longevity candidate due to its involvement in DNA repair and inflammation. Further studies are needed to identify the functional variant(s) that predispose(s) to a long and healthy life.

Key words: 5q31.1; genetic association; human longevity; *IL13*; ImmunoChip; *RAD50*.

Despite more than 20 years of research into the genetic basis of human longevity, only alleles in the *APOE* and *FOXO3* genes have repeatedly been shown to be associated with survival to very advanced ages (Schächter *et al.*, 1994; Willcox *et al.*, 2008; Flachsbart *et al.*, 2009; Soerensen *et al.*, 2010; Deelen *et al.*, 2013). *APOE* and *FOXO3* were initially detected in candidate-driven case-control investigations, but *APOE* has since then been confirmed in a number of genome-wide association studies (GWAS). In addition, a single nucleotide polymorphism (SNP) on chromosome 5q33.3 was recently identified in a GWAS meta-analysis on long-lived individuals (LLI) aged ≥ 90 years (Deelen *et al.*, 2014). Besides the detection of *APOE* and the 5q33.3 locus, longevity GWAS have been relatively unsuccessful and have failed to reveal novel associations with genome-wide significance or sufficient reproducibility (Deelen *et al.*, 2011, 2014; Nebel *et al.*, 2011). Here, we performed a large-scale candidate gene study by targeting established immune-associated loci present on the ImmunoChip (Trynka *et al.*, 2011). The ImmunoChip was designed to perform fine-mapping of GWAS loci of major immune-mediated diseases using data from the

1000 Genomes Project and other sequencing initiatives. The application of the array in this study is based on the hypothesis that a well-functioning immune system and efficient anti-inflammatory networks are potent longevity-assurance mechanisms (Franceschi *et al.*, 2007). We employed the ImmunoChip to screen 1458 German LLI and 6368 younger controls in a discovery phase (panel A in Table S1) for novel longevity loci followed by replication in 1750 LLI and 2551 younger controls from France and Denmark (panel B in Table S1).

After applying conservative and established quality filters to the German longevity sample, 142 136 autosomal SNPs were available for association analysis (see Appendix S1). We used a predefined threshold of $P = 6.15 \times 10^{-7}$ to define statistical significance for the ImmunoChip-wide analysis, based on the Bonferroni correction for the number of linkage disequilibrium (LD)-independent markers on the ImmunoChip (see Appendix S1). The comparison of the case-control frequencies yielded an ImmunoChip-wide significant association signal for the SNP rs2075650 in the *TOMM40/APOE* region ($P_{\text{ImmunoChip}} = 7.01 \times 10^{-9}$; OR = 0.69; 95% CI = 0.60–0.78; Table 1, Figs S1 and S2a). This SNP is in moderate LD ($r^2 = 0.52$) with SNP rs429358 that defines the well-established *APOE* $\epsilon 4$ allele (Deelen *et al.*, 2011), and hence, rs2075650 was not considered for replication. With regard to the *FOXO3* and the 5q33.3 loci, the ImmunoChip is uninformative due to poor coverage (Fig. S2b,c). Eighty-four other SNPs showed $P_{\text{ImmunoChip}} < 5 \times 10^{-4}$ in the discovery ImmunoChip analysis (Table S2). Of these, we selected 15 markers for replication, each one representing the best SNP for a specific associated region defined by the clumping procedure (see Appendix S1, Table S3). The replication analysis was performed in two longevity samples from France and Denmark (France: 1257 LLI and 1811 controls; Denmark: 493 LLI and 740 controls; panel B in Table S1). The signal at SNP rs2706372, located in a region encompassing *RAD50* (radiation sensitive, *Saccharomyces cerevisiae* homolog) and *IL13* (Interleukin 13), replicated in the French sample ($P_{\text{Repl-France}} = 2.69 \times 10^{-3}$; OR = 1.21; 95% CI = 1.07–1.38; $P_{\text{Repl-France; adj}} = 0.04$ (corrected for 15 tests)) (Table S4). In the smaller Danish sample, the allelic effect of rs2706372 showed a similar trend ($P_{\text{Repl-Denmark}} = 0.08$; OR = 1.19; 95% CI = 0.98–1.45; $P_{\text{Repl-Denmark; adj}} = 1$ (corrected for 15 tests)). In the combined French-Danish replication sample, meta-analysis association analysis yielded a P -value of 4.95×10^{-4} (OR = 1.21; 95% CI = 1.09–1.34; $P_{\text{Repl-France-Denmark; adj}} = 0.0074$ (corrected for 15 tests)). In a meta-analysis of the German discovery and French-Danish replication samples, rs2706372 reached a P -value of $P_{\text{ImmunoChip+Repl}} = 5.42 \times 10^{-7}$ (Table 1). Estimates of odds ratios for rs2706372 were consistent across all three studies (OR_{Germany} = 1.19; OR_{Denmark} = 1.19; OR_{France} = 1.21; statistical metric of heterogeneity $I^2 = 0.0$), supporting the validity of the association finding.

Our targeted immune gene approach on a combined European sample of 3208 LLI and 8919 controls resulted in the identification of a novel association for longevity in the *RAD50/IL13* region on chromosome 5q31.1. The lead SNP rs2706372 is located in the intronic region of the *RAD50* gene and is in strong LD with other associated SNPs close to *IL13* and *IL5*. The actual association signal extends even further to include additional genes (Fig. 1). At this point, this observation renders it difficult to assess which gene is actually affected by the association, although *RAD50* is a plausible candidate. The protein encoded by *RAD50* is highly similar to *Saccharomyces cerevisiae* Rad50 which is involved in repairing DNA double-strand breaks. Similarly, the human *RAD50* is integrated in a functional DNA-binding complex (Kinoshita *et al.*, 2015) that is important for recombination, repair, and genomic stability (Trujillo *et al.*, 1998). Hence, it is conceivable that variation in *RAD50* could positively influence longevity by increasing DNA stability.

Table 1 ImmunoChip loci associated with human longevity. 5q31.1 (*RAD50/IL13*) is a newly associated locus.

Chr.	Association boundaries (kb)	dbSNP ID	A1	A2	AF cases	AF controls	Functional annotation	Key genes (n additional genes within locus)	Discovery ImmunoChip (1458/6368)		Replication (1750/2551)		ImmunoChip + Replication (3208/8919)		I^2
									P	OR (95% CI)	P	OR (95% CI)	P _{combined}	OR (95% CI)	
19q13.32	45357–45444	rs2075650	G	A	0.1080	0.1488	<i>TOMM40</i> (intronic)	<i>TOMM40/APOE</i> (3)	7.01×10^{-9}	0.69 (0.60–0.78)	*	–	–	–	–
5q31.1	131784–132143	rs2706372	T	C	0.2565	0.2241	<i>RAD50</i> (intronic)	<i>RAD50/IL13</i> (7)	3.11×10^{-4}	1.19 (1.08–1.31)	4.95×10^{-4}	1.21 (1.09–1.34)	5.42×10^{-7}	1.20 (1.12–1.28)	0.0

*SNP rs2075650 was not considered for replication in this study, because the association at 19q31.32 was already established in the French sample (Schächter *et al.*, 1994) and parts of the Danish sample (Soerensen *et al.*, 2010). Chr: chromosome of marker; **association boundaries**: association boundaries for each index SNP (see Appendix S1). Genomic positions were retrieved from NCBI's dbSNP build v141 (genome build hg19); **dbSNP id**: rs ID of the index SNP; **A1**: minor allele; **A2**: major allele; **AF**: allele frequency of A1 estimated from ImmunoChip (German population); **key genes**: candidate gene(s) in the region; **P/OR**: P -value and corresponding allelic odds ratio and 95% confidence interval with respect to A1. We used a significance level of 6.15×10^{-7} for the statistical association analysis of the ImmunoChip in the discovery experiment and in the combined experiment of discovery and replication based on Bonferroni correction (see Appendix S1). For each panel, numbers of LLI/controls are displayed in parentheses; **I^2** : statistical metric of heterogeneity; I^2 ranges from 0 to 100% and is considered low for values 0–25%.

Associations with other traits: Overlaps with other disease phenotypes (listed if anywhere within association boundaries, see Appendix S1).

rs2075650: age-related macular degeneration; Alzheimer's disease; apolipoprotein levels; blood metabolite; brain imaging; C-reactive protein; cardiovascular disease; carotid intima media thickness; cerebrospinal AB1-42 levels; cholesterol total; cognitive decline; HDL cholesterol; LDL cholesterol; lipid traits; lipid metabolism; lipoprotein-associated phospholipase A2 activity and mass; metabolic syndrome; metabolic levels; quantitative traits; response to statin therapy (LDL-C); sphingolipid levels; triglycerides. rs2706372: asthma; asthma (childhood, severe); asthma (sex interaction); atopic dermatitis; Crohn's disease; eosinophil counts; fibrinogen; Hodgkin's lymphoma; IgE levels; platelet counts; psoriasis; self-reported allergy.




Fig. 1 Association plot for 5q31.1 (*RAD50/IL13*). Blue shaded region corresponds to locus association boundaries (Table 1 and Appendix S1). Shown are the $-\log_{10}$ P -values from the Immunochip analysis ($P_{\text{Immunochip}}$) of the German longevity discovery panel (panel A in Table S1) with regard to the physical location of marker. **Purple diamond:** lead SNP; **filled circles:** analyzed SNPs where the fill color corresponds to the strength of linkage disequilibrium (r^2) with the lead SNP (for color coding see legend in the upper right corner of the plot); **blue line:** recombination intensity (cM/Mb). Positions and gene annotations are according to NCBI's build 37 (hg19). Plot was generated using LocusZoom (Pruim *et al.*, 2010).

Alternatively, it could exert its effect via the direct modulation of cytokine expression; recent evidence suggests at least two possible avenues. First, in dendritic cells *RAD15* was found to activate—upon sensing viral DNA—the transcription factor NF- κ B, thus leading to the production of pro-inflammatory IL-1 β (Roth *et al.*, 2014). Second, the *RAD50* gene harbors at its 3' end an evolutionarily highly conserved locus control region (LCR; Lee *et al.*, 2003; Li *et al.*, 2010) that regulates the expression of the neighboring cytokine genes *IL-4*, *IL-13*, and *IL-5* (Fig. 1) in Th2 cells (Kelly & Locksley, 2000). Variants in the LCR were found to be associated with asthma (Li *et al.*, 2010). Taken together, these findings indicate that the *RAD50* locus may very well contribute to longevity via its role in inflammation and immunity. Nevertheless, it is still possible that the *RAD50* signal is a result of its LD with other markers within the observed association boundaries. Multiple SNPs in the extended *RAD50/IL13* region were previously identified as susceptibility factors for various chronic inflammatory diseases such as Crohn's disease, psoriasis, asthma, and atopic dermatitis (Rioux *et al.*, 2000; Li *et al.*, 2008, 2010; Paternoster *et al.*, 2011). Further studies are therefore needed to identify the functional variant(s) and the underlying molecular mechanisms that predispose(s) to a long and healthy life.

Acknowledgments

The authors wish to thank all individuals for their participation.

Funding

This study was supported by the RESOLVE project (grant FP7-HEALTH-F4-2008-202047 to S.S., A.N., and F.F.); the Deutsche Forschungsgemeinschaft (DFG) Cluster of Excellence 'Inflammation at Interfaces' (grant to A.N. with fellowship for L.G.); the German Federal Ministry of Education and Research (BMBF) within the framework of the e:Med research and funding concept (SysInflame grant O1ZX1306A); the biobank popgen (the popgen 2.0 network is financed by the German Ministry for Education and Research (grant O1EY1103 to W.L.)); the German Federal Ministry for

Education and Research within the context of the National Genome Research Network 2 (NGFN-2), the National Genome Research Network plus (NGFNplus) and the Integrated Genome Research Network (IG) MoodS (grant O1GS08144 to M.M.N., grant O1GS08147); the INTERREG4A program Syddanmark-Schleswig-K.E.R.N (with EU funds from the European Regional Development Fund, grant to K.C., A.N., L.C., and F.F.); the VELUX Foundation; the Danish National Research Foundation; the US National Institutes of Health—National Institute on Aging (grant number P01 AG08761 to K.C. and L.C.); the Danish Agency for Science, Technology and Innovation (grant number 09-070081 to K.C. and L.C.). The CEPH centenarian cohort was supported by the "Ministère de l'Enseignement supérieur et de la Recherche". The SU.VI.MAX cohort was funded by the French Institut National de la Santé et de la Recherche Médicale, the Institut National de la Recherche Agronomique, the Université Paris 13. The genotyping of French samples was supported by the Commissariat à l'Énergie Atomique—Centre National de Génotypage. The KORA research platform (KORA, Cooperative Research in the Region of Augsburg) was initiated and financed by the Helmholtz Zentrum München—German Research Center for Environmental Health, which is funded by the German Federal Ministry of Education and Research and by the State of Bavaria. Furthermore, KORA research was supported within the Munich Center of Health Sciences (MC Health), Ludwig-Maximilians-University Munich, as part of LMUinnovativ. S. B. is supported by DFG grant BR1912/6-1 and the Else-Kröner-Fresenius-Stiftung (stipend 2010_EKES.32). M. M. N. received support from the Alfried Krupp von Bohlen und Halbach-Stiftung. The HNR study is supported by the Heinz Nixdorf Foundation (Germany). Additionally, the study is funded by the German Ministry of Education and Science and the German Research Council (DFG; Project SI 236/8-1, SI236/9-1, ER 155/6-1). A. F. receives an endowment professorship by the Foundation for Experimental Medicine (Zurich, Switzerland).

Conflict of interest

The authors declare no competing financial interests.

Author contributions

F.F., A.N., A.F., D.E., and S.S. designed research; W.L., A.F., S.S., C.B., S.B., A.P., K.S., M.M.-N., P.H., and M.M.N. were involved in recruitment of German study subjects and assembling of phenotypic data; F.F. and A.N. organized chip genotyping of German long-lived individuals; H.B., J.D., C.D., P.G., L.C., M.N., and K.C. performed replication experiments; F.-A.H., M.N., and C.D. helped with the experimental work; D.E., L.G., A.C., and F.F. analyzed data; D.E., F.F., and A.N. interpreted the data and wrote the manuscript; all authors performed critical revision and approved the final version of the manuscript.

References

- Deelen J, Beekman M, Uh HW, Helmer Q, Kuningas M, Christiansen L, Kremer D, van der Breggen R, Suchiman HE, Lakenberg N, van den Akker EB, Passtoors WM, Tiemeier H, van Heemst D, de Craen AJ, Rivadeneira F, de Geus EJ, Perola M, van der Ouderaa FJ, Gunn DA, Boomsma DI, Uitterlinden AG, Christensen K, van Duijn CM, Heijmans BT, Houwing-Duistermaat JJ, Westendorp RG, Slagboom PE (2011) Genome-wide association study identifies a single major locus contributing to survival into old age; the APOE locus revisited. *Aging Cell* **10**, 686–698.
- Deelen J, Beekman M, Capri M, Franceschi C, Slagboom PE (2013) Identifying the genomic determinants of aging and longevity in human population studies: progress and challenges. *BioEssays* **35**, 386–396.
- Deelen J, Beekman M, Uh HW, Broer L, Ayers KL, Tan Q, Kamatani Y, Bennet AM, Tamm R, Trompet S, Guethbjartsson DF, Flachsbarth F, Rose G, Viktorin A, Fischer

- K, Nygaard M, Cordell HJ, Crocco P, van den Akker EB, Bohringer S, Helmer Q, Nelson CP, Saunders GI, Alver M, Andersen-Ranberg K, Breen ME, van der Breggen R, Caliebe A, Capri M, Cevenini E, Collerton JC, Dato S, Davies K, Ford I, Gampe J, Garagnani P, de Geus EJ, Harrow J, van Heemst D, Heijmans BT, Heinsen FA, Hottenga JJ, Hofman A, Jeune B, Jonsson PV, Lathrop M, Lechner D, Martin-Ruiz C, Mc Nerlan SE, Mihailov E, Montesanto A, Mooijaart SP, Murphy A, Nohr EA, Paternoster L, Postmus I, Rivadeneira F, Ross OA, Salvioli S, Sattar N, Schreiber S, Stefansson H, Stott DJ, Tiemeier H, Uitterlinden AG, Westendorp RG, Willemssen G, Samani NJ, Galan P, Sorensen TI, Boomsma DI, Jukema JW, Rea IM, Passarino G, de Craen AJ, Christensen K, Nebel A, Stefansson K, Metspalu A, Magnusson P, Blanche H, Christiansen L, Kirkwood TB, van Duijn CM, Franceschi C, Houwing-Duistermaat JJ, Slagboom PE (2014) Genome-wide association meta-analysis of human longevity identifies a novel locus conferring survival beyond 90 years of age. *Hum. Mol. Genet.* **23**, 4420–4432.
- Flachsbart F, Caliebe A, Kleindorp R, Blanche H, von Eller-Eberstein H, Nikolaus S, Schreiber S, Nebel A (2009) Association of FOXO3A variation with human longevity confirmed in German centenarians. *Proc. Natl Acad. Sci. USA* **106**, 2700–2705.
- Franceschi C, Capri M, Monti D, Giunta S, Olivieri F, Sevini F, Panourgia MP, Invidia L, Celani L, Scurti M, Cevenini E, Castellani GC, Salvioli S (2007) Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mech. Ageing Dev.* **128**, 92–105.
- Kelly BL, Locksley RM (2000) Coordinate regulation of the IL-4, IL-13, and IL-5 cytokine cluster in Th2 clones revealed by allelic expression patterns. *J. Immunol.* **165**, 2982–2986.
- Kinoshita E, van Rossum-Fikkert S, Sanchez H, Kertokallio A, Wyman C (2015) Human RAD50 makes a functional DNA-binding complex. *Biochimie* **113**, 47–53.
- Lee GR, Fields PE, Griffin TJ, Flavell RA (2003) Regulation of the Th2 cytokine locus by a locus control region. *Immunity* **19**, 145–153.
- Li Y, Chang M, Schrodli SJ, Callis-Duffin KP, Matsunami N, Civello D, Bui N, Catanese JJ, Leppert MF, Krueger GG, Begovich AB (2008) The 5q31 variants associated with psoriasis and Crohn's disease are distinct. *Hum. Mol. Genet.* **17**, 2978–2985.
- Li X, Howard TD, Zheng SL, Haselkorn T, Peters SP, Meyers DA, Bleecker ER (2010) Genome-wide association study of asthma identifies RAD50-IL13 and HLA-DR/DQ regions. *J. Allergy Clin. Immunol.* **125**, 328–335 e311.
- Nebel A, Kleindorp R, Caliebe A, Nothnagel M, Blanche H, Junge O, Wittig M, Ellinghaus D, Flachsbart F, Wichmann HE, Meitinger T, Nikolaus S, Franke A, Krawczak M, Lathrop M, Schreiber S (2011) A genome-wide association study confirms APOE as the major gene influencing survival in long-lived individuals. *Mech. Ageing Dev.* **132**, 324–330.
- Paternoster L, Standl M, Chen CM, Ramasamy A, Bonnelykke K, Duijts L, Ferreira MA, Alves AC, Thyssen JP, Albrecht E, Baurecht H, Feenstra B, Sleiman PM, Hysi P, Warrington NM, Curjuric I, Myhre R, Curtin JA, Groen-Blokhuis MM, Kerkhof M, Saaf A, Franke A, Ellinghaus D, Folster-Holst R, Dermitzakis E, Montgomery SB, Prokisch H, Heim K, Hartikainen AL, Pouta A, Pekkanen J, Blakemore AJ, Buxton JL, Kaakinen M, Duffy DL, Madden PA, Heath AC, Montgomery GW, Thompson PJ, Matheson MC, Le Souef P, Pourcain BS, Smith GD, Henderson J, Kemp JP, Timpson NJ, Deloukas P, Ring SM, Wichmann HE, Muller-Nurasyid M, Novak N, Klopp N, Rodriguez E, McArdle W, Linneberg A, Menne T, Nohr EA, Hofman A, Uitterlinden AG, van Duijn CM, Rivadeneira F, de Jongste JC, van der Valk RJ, Wjst M, Jogi R, Geller F, Boyd HA, Murray JC, Kim C, Mentch F, March M, Mangino M, Spector TD, Bataille V, Pennell CE, Holt PG, Sly P, Tiesler CM, Thiering E, Illig T, Imboden M, Nystad W, Simpson A, Hottenga JJ, Postma D, Koppelman GH, Smit HA, Soderhall C, Chawes B, Kreiner-Moller E, Bisgaard H, Melen E, Boomsma DI, Custovic A, Jacobsson B, Probst-Hensch NM, Palmer LJ, Glass D, Hakonarson H, Melbye M, Jarvis DL, Jaddoe VW, Gieger C, Strachan DP, Martin NG, Jarvelin MR, Heinrich J, Evans DM, Weidinger S (2011) Meta-analysis of genome-wide association studies identifies three new risk loci for atopic dermatitis. *Nat. Genet.* **44**, 187–192.
- Pruim RJ, Welch RP, Sanna S, Teslovich TM, Chines PS, Glied TP, Boehnke M, Abecasis GR, Willer CJ (2010) LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* **26**, 2336–2337.
- Rioux JD, Silverberg MS, Daly MJ, Steinert AH, McLeod RS, Griffiths AM, Green T, Brettin TS, Stone V, Bull SB, Bitton A, Williams CN, Greenberg GR, Cohen Z, Lander ES, Hudson TJ, Siminovitch KA (2000) Genomewide search in Canadian families with inflammatory bowel disease reveals two novel susceptibility loci. *Am. J. Hum. Genet.* **66**, 1863–1870.
- Roth S, Rottach A, Lotz-Havla AS, Laux V, Muschwack A, Gersting SW, Muntau AC, Hopfner KP, Jin L, Vanness K, Petrini JH, Drexler I, Leonhardt H, Ruland J (2014) Rad50-CARD9 interactions link cytosolic DNA sensing to IL-1 β production. *Nat. Immunol.* **15**, 538–545.
- Schächter F, Faure-Delanef L, Guenot F, Rouger H, Froguel P, Lesueur-Ginot L, Cohen D (1994) Genetic associations with human longevity at the APOE and ACE loci. *Nat. Genet.* **6**, 29–32.
- Soerensen M, Dato S, Christensen K, McGue M, Stevnsner T, Bohr VA, Christiansen L (2010) Replication of an association of variation in the FOXO3A gene with human longevity using both case-control and longitudinal data. *Ageing Cell* **9**, 1010–1017.
- Trujillo KM, Yuan SS, Lee EY, Sung P (1998) Nuclease activities in a complex of human recombination and DNA repair factors Rad50, Mre11, and p95. *J. Biol. Chem.* **273**, 21447–21450.
- Trynka G, Hunt KA, Bockett NA, Romanos J, Mistry V, Szperl A, Bakker SF, Bardella MT, Bhaw-Rosun L, Castillejo G, de la Concha EG, de Almeida RC, Dias KR, van Diemen CC, Dubois PC, Duerr RH, Edkins S, Franke L, Fransen K, Gutierrez J, Heap GA, Hrdlickova B, Hunt S, Plaza Izurieta L, Izzo V, Joosten LA, Langford C, Mazzilli MC, Mein CA, Midah V, Mitrovic M, Mora B, Morelli M, Nutland S, Nunez C, Onengut-Gumuscu S, Pearce K, Platteel M, Polanco I, Potter S, Ribes-Koninckx C, Ricano-Ponce I, Rich SS, Rybak A, Santiago JL, Senapati S, Sood A, Szajewska H, Troncone R, Varade J, Wallace C, Wolters VM, Zhernakova A, Thelma BK, Cukrowska B, Urcelay E, Bilbao JR, Mearin ML, Barisani D, Barrett JC, Plagnol V, Deloukas P, Wijmenga C, van Heel DA (2011) Dense genotyping identifies and localizes multiple common and rare variant association signals in celiac disease. *Nat. Genet.* **43**, 1193–1201.
- Willcox BJ, Donlon TA, He Q, Chen R, Grove JS, Yano K, Masaki KH, Willcox DC, Rodriguez B, Curb JD (2008) FOXO3A genotype is strongly associated with human longevity. *Proc. Natl Acad. Sci. USA* **105**, 13987–13992.

Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article.

Appendix S1 Experimental procedure.

Table S1 LLI/control panels used in the analysis.

Table S2 ImmunoChIP association statistics in German panel (panel A in Supplementary Table 1) for 84 SNPs with $P_{\text{ImmunoChIP}} < 5 \times 10^{-4}$.

Table S3 ImmunoChIP association statistics in German panel (panel A in Supplementary Table 1) for the 15 SNPs selected for replication.

Table S4 Association statistics in French and Danish samples (panel B in Supplementary Table 1) for the 15 SNPs selected for replication.

Fig. S1 Manhattan plot of ImmunoChIP association statistics of 142 136 SNPs.

Fig. S2 Regional association plots (from ImmunoChIP analysis; panel A in Supplementary Table 1) of established longevity susceptibility loci.

Fig. S3 Principal component analysis of QCed ImmunoChIP data.

Fig. S4 Quantile-quantile (Q-Q) plot for the discovery panel (panel A in Supplementary Table 1).