

This is a repository copy of *Pot, atom and step economic (PASE) synthesis of highly functionalized piperidines: a five-component condensation*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/3384/>

---

**Article:**

Clarke, Paul A. [orcid.org/0000-0003-3952-359X](http://orcid.org/0000-0003-3952-359X), Zaytzev, Andrey V. and Whitwood, Adrian C. (2007) Pot, atom and step economic (PASE) synthesis of highly functionalized piperidines: a five-component condensation. *Tetrahedron Letters*. pp. 5209-5212. ISSN 0040-4039

<https://doi.org/10.1016/j.tetlet.2007.05.141>

---

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

*promoting access to White Rose research papers*



**Universities of Leeds, Sheffield and York**  
**<http://eprints.whiterose.ac.uk/>**

---

This paper was subsequently published in **Tetrahedon Letters**.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/3384>

---

**Published paper**

Clarke, Paul A., Zaytzev, Andrey V. and Whitwood, Adrian C. (2007). Pot, atom and step economic (PASE) synthesis of highly functionalized piperidines: a five-component condensation. *Tetrahedron Letters*, 48 (30). 5209-5212.

---

## Accepted Manuscript

Pot, atom and step economic (PASE) synthesis of highly functionalized piperidines: a five-component condensation

Paul A. Clarke, Andrey V. Zaytzev, Adrian C. Whitwood

PII: S0040-4039(07)01037-4  
DOI: [10.1016/j.tetlet.2007.05.141](https://doi.org/10.1016/j.tetlet.2007.05.141)  
Reference: TETL 32247

To appear in: *Tetrahedron Letters*

Received Date: 19 April 2007  
Revised Date: 15 May 2007  
Accepted Date: 24 May 2007



Please cite this article as: Clarke, P.A., Zaytzev, A.V., Whitwood, A.C., Pot, atom and step economic (PASE) synthesis of highly functionalized piperidines: a five-component condensation, *Tetrahedron Letters* (2007), doi: [10.1016/j.tetlet.2007.05.141](https://doi.org/10.1016/j.tetlet.2007.05.141)


This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Graphical Abstract

To create your abstract, type over the instructions in the template box below.  
Fonts or abstract dimensions should not be changed or altered.

**Pot, atom and step economic (PASE)  
synthesis of highly functionalised  
piperidines: a five component condensation**

Paul A. Clarke,\* Andrey V. Zaytzev and Adrian C. Whitwood



Leave this area blank for abstract info.



Pergamon

---



---

TETRAHEDRON  
LETTERS

---



---

## Pot, atom and step economic (PASE) synthesis of highly functionalized piperidines: a five-component condensation

Paul A. Clarke,<sup>\*</sup> Andrey V. Zaytzev and Adrian C. Whitwood<sup>†</sup>


*Department of Chemistry, University of York, Heslington, York, YO10 5DD, UK.*

**Abstract**—: The diastereoselective pot, atom and step economic (PASE) synthesis of highly functionalized piperidines has been realized. The procedure simply involves mixing methyl acetoacetate, 2 eq. aldehyde and 2 eq. of aniline together in the presence of  $\text{InCl}_3$ . In most cases the piperidine precipitates out of solution.

Functionalized piperidines occur with great regularity in the natural product arena and as important units in pharmaceuticals. Consequently, a huge amount of effort has been directed towards developing ever more efficient methods to synthesize them.<sup>1</sup> For example, over the last ten years thousands of piperidine-containing compounds have been entered into preclinical and clinical trials.<sup>2</sup> We have recently become interested in the development of ‘Greener’ methods for the synthesis of organic molecules of medium complexity, and have reported the pot, atom and step economic (PASE) synthesis of highly substituted tetrahydropyran-4-ones.<sup>3</sup> Pot, atom<sup>4</sup> and step<sup>5</sup> economy aims to combine as many transformations as possible into a single reaction vessel, without the need for work-up and isolation of intermediate compounds. Ideally, all the reagents should also be incorporated into the final product. This should lead to a reduction in the amount of waste generated by a synthetic route due to the minimisation of by-products, reaction and extraction solvents, silica gel for chromatographic purification of intermediate compounds and a like. Process chemists in industry have long been applying these criteria to synthetic routes and during our discussions with them we became aware of a need to extend our PASE synthesis of THPs to the synthesis of functionalised piperidines. This Letter reports our success at developing a *five-component*, pot, atom and step

economic, diastereoselective synthesis of highly functionalised piperidines.

Our initial strategy was similar to the one we had developed for our synthesis of THPs,<sup>3, 6</sup> that is, disconnection of the target piperidine to a  $\beta$ -ketoester derivative and two aldehyde derived reaction partners (Scheme 1). In this instance we envisaged using an imine as one of the reaction partners and an aldehyde as the other, giving us the potential for diversification at the greatest number of positions around the piperidine ring.




**Scheme 1.**

We initially considered using diketene as the  $\beta$ -ketoester derivative, however, problems with supply and the hazards associated with its use prompted us to consider other alternatives. One interesting possibility was to use a  $\beta$ -

**Keywords:** piperidines, PASE, one-pot, multi-component.

<sup>\*</sup> Corresponding author. Phone: +44 01904 432614. E-mail: [pac507@york.ac.uk](mailto:pac507@york.ac.uk).

<sup>†</sup> Corresponding author for information concerning X-ray crystallography.



Scheme 2.

ketoester itself and to increase the nucleophilicity of its  $\alpha$ -carbon by formation of the enamine derivative. We rationalized that it may then be possible to develop a multi-component reaction which coupled the *in situ* formation of the enamine of the  $\beta$ -ketoester, with formation of an imine and the sequential condensation of the enamine to the aldehyde and then the imine, forming the desired piperidine in a pot, atom and step economic fashion. To assess the feasibility of this approach we conducted a few investigative studies on the reactivity of the potential reacting partners (Scheme 2).


We elected to use  $\text{InCl}_3$  as a Lewis acid promoter as it had been used previously to catalyze Knoevenagel reactions,<sup>7</sup> Mannich-like reactions and imine formation<sup>8</sup> and Michael reactions.<sup>9</sup> Initially we pre-formed Knoevenagel adduct **1** and imine **2** and mixed them with aniline in the presence of  $\text{InCl}_3$  and we were pleased that the desired piperidine was formed in a reasonable yield (**A**). However, when this reaction was repeated in the presence of anhydrous potassium carbonate no reaction took place (**B**). This led us to speculate that the reaction was actually promoted by trace amounts of HCl present in the  $\text{InCl}_3$ , although when we ran the reaction using a solution of HCl in 1,4-dioxane no product was formed (**C**). We next examined the reaction of pre-formed enamine **3** with benzaldehyde and imine **2** in the presence of  $\text{InCl}_3$ . In this case the desired piperidine **4** was again formed in a moderate yield (**D**). The Brønsted acid promoted reaction was also investigated, and in this instance a 19% yield of piperidine **4i** was recovered

from the reaction (**E**). Given the success of routes **A** and **D**, we investigated the *in-situ* formation of both the enamine and the imine and their condensation/cyclisation reactions, which is in effect a *five-component condensation reaction*. We were delighted that not only did this reaction yield piperidine **4i**, it did so in an increased yield of 60% (**F**). The relative stereochemistry of the 2,6-phenyl groups in **4i** was confirmed as *trans*- by single crystal X-ray analysis (Fig 1). These studies provided proof of concept that a *five-component condensation reaction* was indeed a viable approach to the PASE synthesis of highly functionalised piperidine rings.



Fig.1: ORTEP diagram of the X-ray Crystal Structure of **4i** (ADP shown at 50%)

We suggest that a plausible mechanism for the formation of piperidine **4i** involves the Lewis acid promoted formation of imine **2** and enamine **3**, enamine **3** can then undergo a 'Knoevenagel-like' condensation with benzaldehyde to form the iminium ion–Knoevenagel product **5**. Loss of a proton and tautomerisation of the imine to the enamine generates a diene **6** which can undergo either an aza-Diels-Alder cyclisation or a tandem Mannich-Michael reaction to furnish piperidine **4** (Scheme 3).



**Scheme 3.**


Given the success of these preliminary studies we decided to investigate the scope of the reaction and the results are displayed in Table 1.<sup>10</sup> Some interesting observations were made during these studies. *p*-Anisidine reacted faster than aniline, and in general reactions were complete within 24 h rather than 48 h. While 4-chloroaniline reacted very slowly and took 7 days to generate a 52% yield of the piperidine product, *p*-nitroaniline did not react at all. This is undoubtedly due to the reduced nucleophilicity of these anilines compared to *p*-anisidine and aniline. A wide range of aromatic aldehydes could be used in the reaction. Yields were found to be lower when 2-substituted aldehydes were employed and this is probably due to steric effects. In the cases of aniline/4-methylbenzaldehyde **4g** and aniline/benzaldehyde **4i**, hydrolysis of the final enamine-piperidine occurred to some extent yielding the enol-piperidines in 3% and 11%, respectively. It was also found that certain aldehyde–aniline combinations did not form any piperidine, as the reaction stopped due to the precipitation of an insoluble imine (Table 2).

We next investigated the use of the aliphatic aldehydes benzoylacetdehyde, butanal and *iso*-butyl aldehyde with both aniline and *p*-anisidine, but in all cases multiple products arose. This can be attributed to a number of reasons: (i) the propensity for aliphatic aldehydes to favour enamine formation rather than imine formation and, (ii) the enamines of the aliphatic aldehydes forming preferentially to the enamine of the  $\beta$ -ketoester and then condensing with any remaining aldehyde before the desired reaction occurs. Additionally, we investigated the use of benzylamine instead of aniline, however, only the HCl salt of the amine was isolated, presumably due to the more basic nature of aliphatic amines compared to anilines.


In summary, we have developed a five-component condensation reaction for the formation of highly substituted piperidines which is pot, atom and step economic. In general the yields are good and the piperidine product precipitates from the reaction allowing for easy isolation. Work is underway to extend the procedure to the synthesis of piperidines using different aldehyde and imine

components in order to form piperidines with different 2,6-substituents and also to the asymmetric synthesis of such compounds. These studies will be reported in due course.

**Table 1.**



	Ar <sup>1</sup> CHO	Ar <sup>2</sup> NH <sub>2</sub>	Yield (%)
<b>a</b>	4-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	45
<b>b</b>	3-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	48
<b>c</b>	Ph	4-MeOC <sub>6</sub> H <sub>4</sub>	74
<b>d</b>	2-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	27
<b>e</b>	3-CF <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	57
<b>f</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	Ph	52
<b>g</b>	4-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Ph	50
<b>h</b>	3-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Ph	64
<b>i</b>	Ph	Ph	60
<b>j</b>	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	Ph	52
<b>k</b>	2-CH <sub>3</sub> C <sub>6</sub> H <sub>4</sub>	Ph	16
<b>l</b>	4-MeO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub>	Ph	24
<b>m</b>	4-MeOC <sub>6</sub> H <sub>4</sub>	4-ClC <sub>6</sub> H <sub>4</sub>	52



	Ar <sup>1</sup> CHO	Ar <sup>2</sup> NH <sub>2</sub>	Yield (%)
	4-CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	59
	4-NO <sub>2</sub> C <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	42
	2-naphth	4-MeOC <sub>6</sub> H <sub>4</sub>	58
	4-MeO <sub>2</sub> CC <sub>6</sub> H <sub>4</sub>	4-MeOC <sub>6</sub> H <sub>4</sub>	83

**Table 2.**

### Acknowledgements

We thank the EPSRC for funding under the 'Greener' chemistry initiative (EP/C523970/1) and AstraZeneca for an unrestricted research award (PAC).

### References and Notes

1. For recent reviews on the synthesis of piperidines see: Laschat, S.; Dickner, T. *Synthesis* **2000**, 1781. Weintraub, P. M.; Sabol, J. S.; Kane, J. M.; Borcharding, D. R. *Tetrahedron* **2003**, *59*, 2953. Buffat, M. G. P. *Tetrahedron* **2004**, *60*, 1701.
2. Watson, P. S.; Jiang, B.; and Scott, B. *Org. Lett.* **2000**, *2*, 3679.
3. Clarke, P. A.; Santos, S.; Martin, W. H. C. *Green Chem.* **2007**, *9*, 438.
4. (a) Trost, B. M.; *Science* **1991**, *254*, 1471. (b) Trost, B. M. *Angew. Chem. Int. Ed.* **1995**, *34*, 259.
5. (a) Wender, P. A.; Bi, F. C.; Gamber, G. G.; Gosselein, F.; Hubbard, R. D.; Scanio, M. J. C.; Sun, R.; Williams, T. J.; Zhang, L. *Pure Appl. Chem.* **2002**, *74*, 25. (b) Wender, P. A.; Handy, S. T.; and Wright, D. L. *Chem. Ind. (London)* **1997**, 765. (c) Wender, P. A.; Miller, B. L. *Organic Synthesis: Theory and Applications*; Hudlicky, T. Ed.; JAI: Greenwich, 1993; Vol. 2, pp 27-66.
6. Clarke, P. A.; Martin, W. H. C.; Hargreaves, J. M.; Wilson, C.; Blake, A. J. *Chem. Commun.* **2005**, 1061. Clarke, P. A.; Martin, W. H. C.; Hargreaves, J. M.; Wilson, C.; Blake, A. J. *Org. Biomol. Chem.* **2005**, *3*, 3551.
7. Prajapati, D.; Gohain, M. *Beilstein J. Org. Chem.* **2006**, *2*:11.
8. Loh, T.-P.; Wei, L.-L. *Tetrahedron Lett.* **1998**, *39*, 323.
9. Loh, T.-P.; Wei, L.-L. *Synlett* **1998**, 975.
10. General procedure for the PASE synthesis of piperidines:  
Methyl acetoacetate (2 mmol), aldehyde (4 mmol) and aniline (4 mmol) were dissolved in acetonitrile (4 ml) and  $\text{InCl}_3$  (0.147 g, 0.67 mmol) was introduced. The reaction mixture was stirred at room temperature for 24 or 48 h. The product was isolated by filtration of the precipitate and washing with a small amount of acetonitrile or, if no precipitation occurred, the solvent was evaporated and the residue was purified by flash column chromatography on silica gel ( $\text{Et}_2\text{O}$  – Petrol).  
Compound **4i**  $\nu_{\text{max}}/\text{cm}^{-1}$  (KBr) 3446, 2918, 2850, 1662, 1588, 1503, 1254, 1078;  $\delta_{\text{H}}$  (500 MHz,  $\text{CDCl}_3$ ) 10.26 (1H, s, NH), 7.33-7.25 (7H, m, Ar), 7.21 (1H, m, Ar), 7.17 (2H, d,  $J$  7.5, Ar), 7.10-7.05 (5H, m, Ar), 6.61 (1H, t,  $J$  7.0, Ar), 6.53 (2H, d,  $J$  8.5, Ar), 6.46 (1H, s, 2-H), 6.30-6.28 (2H, m, Ar), 5.15 (1H, m, 6-H), 2.88 (1H, dd,  $J$  15.3 and 5.8, 5- $\text{H}_a$ ), 2.77 (1H, dd,  $J$  15.3 and 2.3, 5- $\text{H}_b$ );  $\delta_{\text{C}}$  (125 MHz,  $\text{CDCl}_3$ ) 168.6 (C,  $\text{COOCH}_3$ ), 156.3 (C), 146.9 (C), 143.9 (C), 142.7 (C), 137.8 (C), 128.9 (CH), 128.8 (CH), 128.6 (CH), 128.2 (CH), 127.1 (CH), 126.6 (CH), 126.4 (CH), 126.3 (CH), 125.9 (CH), 125.8 (CH), 116.1 (CH), 112.9 (CH), 97.9 (C, C-3), 58.2 (CH, C-6), 55.1 (CH, C-2), 51.0 ( $\text{CH}_3$ ,  $\text{COOCH}_3$ ), 33.6 ( $\text{CH}_2$ , C-5); MS (ES) 461 ( $\text{M}+\text{H}^+$ , 12), 386 (8), 280 (34), 182 (100), 150 (10), 121 (23), 106 (36), 94 (31); HRMS (ES) 461.2224 ( $\text{M} + \text{H}^+$ ,  $\text{C}_{31}\text{H}_{29}\text{N}_2\text{O}_2$  requires 461.2224).