



Modulating absorption and postprandial handling of dietary fatty acids by structuring fat in the meal: a randomized crossover clinical trial.

Cécile Vors, Gaëlle Pineau, Laure Gabert, Jocelyne Draï, Corinne Louche-Pélissier, Catherine Defoort, Denis Lairon, Michel Désage, Sabine Danthine, Stéphanie Lambert-Porcheron, et al.

► To cite this version:

Cécile Vors, Gaëlle Pineau, Laure Gabert, Jocelyne Draï, Corinne Louche-Pélissier, et al.. Modulating absorption and postprandial handling of dietary fatty acids by structuring fat in the meal: a randomized crossover clinical trial.: Fat structuring modifies postprandial metabolism. American Journal of Clinical Nutrition, American Society for Nutrition, 2013, 97 (1), pp.23-36. <10.3945/ajcn.112.043976>. <inserm-00766661>

HAL Id: inserm-00766661

<http://www.hal.inserm.fr/inserm-00766661>

Submitted on 18 Dec 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Modulating absorption and postprandial handling of dietary fatty acids
by structuring fat in the meal: a randomized cross-over clinical trial¹⁻³**

Cécile Vors, Gaëlle Pineau, Laure Gabert, Jocelyne Draï,
Corinne Louche-Pélissier, Catherine Defoort, Denis Lairon,
Michel Désage, Sabine Danthine, Stéphanie Lambert-Porcheron,
Hubert Vidal, Martine Laville, and Marie-Caroline Michalski

¹ From INRA USC1362, CarMeN, Cardiovascular Metabolism Diabetes and Nutrition laboratory, F-69621 Villeurbanne, France (CV, MCM); INSA-Lyon, IMBL, F-69621 Villeurbanne, France (CV, MCM); Université de Lyon, F-69100 Villeurbanne, France (LG, CLP, MD); INSERM U1060, CarMeN, F-69921 Oullins, France (GP, HV); CRNH Rhône-Alpes, CENS, F-69921 Oullins, France (LG, JD, CLP,MD, SLP, ML, MCM); INSERM UMR1062, Nutrition, Obesity and Risk of Thrombosis laboratory, F-13385, Marseille, France (CD, DL); INRA, UMR1260, F-13385 Marseille, France (CD, DL); Aix-Marseille Univ, Faculté de Médecine, F-13385 Marseille, France (CD, DL); Université de Liège, GxABT, Department of Food Science and Formulation, Gembloux, Belgique (SD); Hospices Civils de Lyon, F-69008 Lyon, France (JD, SLP); Laboratoire de Biochimie, Centre Hospitalier Lyon Sud (JD).

Authors' last names: Vors, Pineau, Gabert, Draï, Louche-Pélissier, Defoort, Lairon, Désage, Danthine, Lambert-Porcheron, Vidal, Laville, Michalski.

² This work was funded by the Centre National Interprofessionnel de l'Economie Laitière (CNIEL). CV received funding for PhD study by INRA and CNIEL. CV received a Research Prize from the Foundation Nestlé France. HV received financial support from Ezus-Lyon1. The funding agencies had no role in the data analysis.

³ Corresponding author, to whom reprints may be requested:

INRA U1235, Cardiovasculaire Métabolisme diabEtologie et Nutrition, CarMeN, Bâtiment IMBL, INSA-Lyon, 11 avenue Jean Capelle, 69621 VILLEURBANNE cedex, France

E-mail: marie-caroline.michalski@insa-lyon.fr

Phone: +33 4 72 43 81 12 - Fax: +33 4 72 43 85 24

Abbreviations: AMF: anhydrous milk fat; AP: atom percent; APE: atom percent excess; AUC: area under curve; iAUC: incremental area under curve; CMRF: chylomicron-rich fraction; CRNH-RA: Human Nutrition Research Center Rhône-Alpes; d_{32} : volume-surface mean diameter; dARF: acetate recovery factor; DSC: differential scanning calorimetry; FA: fatty acid; FAMES: fatty acid methyl esters; NEFA: non-esterified fatty acid; NW: normal-weight; TAG: triacylglycerol; XRD: X-ray diffraction.

Running title: Fat structuring modifies postprandial metabolism.

Clinical trial registry: Clinical Trials, #NCT01249378

1 ABSTRACT

2 **Background:** Prolonged postprandial hypertriglyceridemia is a potential risk factor for
3 cardiovascular diseases. In the context of obesity, this is associated with a chronic imbalance
4 of lipid partitioning oriented towards storage and not towards β -oxidation.

5 **Objectives:** We tested the hypothesis that the physical structure of fat in a meal can modify
6 absorption, chylomicron transport and further metabolic handling of dietary fatty acids.

7 **Design:** 9 normal-weight and 9 obese subjects were fed 40g of milkfat (+ ^{13}C -
8 triacylglycerols), either emulsified or not, in breakfasts of identical composition. We
9 measured the postprandial triglyceride content and size of the chylomicron-rich fraction,
10 plasma kinetics of ^{13}C -fatty acids, exogenous lipid oxidation using breath-test/indirect
11 calorimetry, and fecal excretion.

12 **Results:** The emulsified fat resulted in earlier (>1h) and sharper chylomicron and ^{13}C -fatty
13 acids peaks in plasma compared to spread fat in both groups ($P<0.0001$). After 2h, the
14 emulsified fat increased ApoB48 concentration (9.7 ± 0.7 vs 7.1 ± 0.9 mg/L; $P<0.05$) in the
15 normal-weight subjects compared to the spread fat. For the obese subjects, emulsified fat
16 resulted in 3-fold larger chylomicrons (218 ± 24 nm) compared to the spread fat ($P<0.05$). The
17 emulsified fat induced higher dietary fatty acid spillover in plasma and sharper $^{13}\text{CO}_2$
18 appearance, provoking increased exogenous lipid oxidation in each group: from 45% to 52%
19 in normal-weight subjects ($P<0.05$), 40% to 57% in obese ($P<0.01$).

20 **Conclusions:** This study supports a new concept of “slow vs fast fat” whereby intestinal
21 absorption can be modulated by structuring of dietary fat to modulate postprandial lipemia
22 and lipid β -oxidation in humans of different BMI.

23
24 **Key words:** intestinal absorption; chylomicron; emulsion; stable isotopes; obesity.

25

26 INTRODUCTION

27 The metabolic importance of intestinal absorption and transport of nutrients in the
28 postprandial period is recognized as important in the context of metabolic diseases such as
29 obesity and type 2 diabetes (1). Regarding lipid metabolism, plasma kinetics, timing of peak
30 of lipemia as well as chylomicron size are recognized as factors determining metabolic
31 complications that are still an open field of research (2-4). For this reason, control of intestinal
32 lipid absorption, the resulting chylomicron transport dynamics and ultimate dietary lipid fate
33 may be an effective tool in the management of metabolic diseases. Recent studies have shown
34 differential effects of oral sensory stimulation with high vs low amounts of dietary fat on
35 intestinal lipid absorption (5). The possible effects of fatty acid (FA) profile of an oral fat load
36 on chylomicron size have been suggested (6-10). While both fat load and composition can
37 affect postprandial lipid absorption, few studies have investigated the effects of fat structure
38 on the postprandial metabolism of an identical lipid load. In diabetes, the concept of
39 “slow/low glycemic index carbohydrates and fast/high glycemic index carbohydrates” is well
40 established and has facilitated the development of specific foods and/or cooking methods to
41 control postprandial glycemia (11). We thus raised the question of whether a similar concept
42 may be applicable to dietary fat according to the way it is structured in the meal.

43 Dietary lipids are incorporated in food products with different physicochemical structures,
44 e.g., in dispersed lipid droplets in oil-in-water emulsions like ice cream or in a continuous
45 lipid phase in butter and margarine. Emulsions are the most widespread fat structures, in
46 processed foods and enteral formulas, and are therefore of interest regarding their role in lipid
47 digestion and absorption (12). Indeed, we have previously demonstrated in rodents the
48 importance of lipid emulsified structure on FA absorption and β -oxidation (13-15). However,
49 the impact of fat structure on the kinetics of lipid absorption and dietary FA handling in
50 humans remains to be elucidated. Such effects also deserve to be elucidated in obese subjects
51 that present altered storage function of dietary FA in the postprandial state (16).

52 We therefore hypothesized that the physicochemical structure of the fat in a meal could
53 modulate postprandial lipemia and fat partitioning: storage *vs* oxidation, and that the effect
54 would be more pronounced in obese subjects. We investigated the metabolic response to fat-
55 containing meals (40 g) differing only in the structuring of fat, emulsified or not, in healthy
56 young normal weight and obese men. Measurements included chylomicron number and size,
57 FA β -oxidation and FA excretion in feces. The aim of this study was to define the
58 contribution of fat structure and subject BMI on the postprandial lipemia and metabolism of
59 dietary FA.

60

61 **SUBJECTS AND METHODS**

62 **Study design**

63 The study was an open label trial with a cross-over randomized controlled design
64 involving 2 days of metabolic testing separated by at least three weeks (Supplemental Figure
65 1). It was conducted at the Human Nutrition Research Center Rhône-Alpes (CRNH-RA;
66 Lyon, France) according to the Second Declaration of Helsinki and the French Huriet-
67 Serusclat law. The LIPINFLOX study was approved by the Scientific Ethics Committee of
68 Lyon Sud-Est-II and AFSSAPS and registered at Clinical Trials (#NCT01249378).
69 Volunteers received written and oral information and their medical history was reviewed. In
70 addition they underwent a physical examination and fasting clinical analysis were assessed
71 before enrolment. Informed written consent was obtained from all subjects. Volunteers
72 performed trial in the period from April 2010 to July 2011. During the protocol, all subjects
73 were asked to continue their regular diet and activity except for the week before and the 3-d
74 period following each test day. Subjects were told to avoid foods naturally rich in ^{13}C and
75 were given a list. For 48 h prior to testing subjects were asked to refrain from consuming
76 alcohol and to avoid exercise. In addition, subjects were provided with a standardized dinner

77 the evening prior to testing. Compliance was checked through diet records, 5 days before and
78 3 days after each test day.

79 After an overnight of fast, subjects ingested one of the two test breakfasts. The primary
80 outcome measured was the effect of fat structure on postprandial lipemia. Secondary
81 outcomes measured were the effect of fat structure and BMI on postprandial lipid metabolism.
82 Previous studies on lipemia (17) and lipid oxidation (18) were used for the power analysis: a
83 minimum sample size of 8 subjects per BMI group was calculated to be necessary to detect
84 significant changes in these parameters. The treatments were randomized according to a
85 random allocation sequence performed by a CRNH-RA biostatistician using Stat® v.11; two
86 randomization lists were generated and stratified over BMI. Subjects were anonymized using
87 a number corresponding to randomization sequence order.

88

89 **Subjects**

90 Twenty-two healthy men were recruited, 11 normal-weight (NW) and 11 obese, and 20
91 completed the study (see Flow Diagram online). One subject in each group was not included
92 in data analyses due to abnormal postprandial lipid metabolism; therefore 18 healthy subjects
93 divided in two groups, 9 NW and 9 obese with comparable mean age were finally tested for
94 the primary outcome, (**Table 1**). Volunteers were required to be non-smokers, sedentary or
95 having <4h per week of physical activity and non-claustrophobic. We excluded persons under
96 medication interfering with lipid metabolism, with psychological illness, or those with
97 eating/metabolic disorders. In addition, subjects were required to have had stable weight, to
98 be free of diabetes and to have not made blood donation for 3 months prior to the start of the
99 study. Data were collected at CRNH-RA.

100

101 **Test meals**

102 The test breakfasts were isoenergetic, equal in nutrient composition (**Table 2**) and both
103 consisted of bread (50 g), skimmed milk (160 mL) and anhydrous milk fat (AMF; 40 g),
104 containing 600 mg of tracers, either spread on bread or emulsified in skim milk. Both meals
105 had the same composition as no additional emulsifier was added because milk proteins are
106 sufficient to provide a submicronic milk fat emulsion. Prior to the test day, a mixture of
107 labeled triacylglycerols (TAG) proportionally representing each FA type present in test fat
108 was first incorporated into melted milk fat: 300 mg of [1,1,1-¹³C] tripalmitin for long-chain
109 saturated FA, 210 mg of [1,1,1-¹³C] triolein for unsaturated FA and 90 mg of [1,1,1-¹³C]
110 trioctanoin for short- and medium-chain FA (99 atom% ¹³C, Eurisotop, Saint-Aubin, France).
111 For the emulsion test, melted labeled milk fat was coarsely pre-mixed in skimmed milk
112 (ProScientific Inc., Oxford, USA) and then further finely emulsified (4x 1 min, Vibra-cell™
113 Ultrasonic Processor, Sonics, Newtown, USA) (Supplemental Figure 2). Test products were
114 then kept at 4°C overnight.

115 A second meal was served 5 hours after breakfast, containing pasta (200 g), turkey (100
116 g), butter (10 g), olive oil (10 g), bread (50 g), stewed fruit (100 g) which provided 713 kcal
117 (2985 kJ) with 29%, 51% and 20% of energy as lipids (22.7 g), carbohydrates (91.5 g) and
118 proteins (35.7 g) respectively. All subjects were given 10 minutes to eat breakfast and 30
119 minutes for lunch. During the test, participants were allowed to drink 200 mL of water.

120

121 **Test fat characterization**

122 Emulsion droplet size was measured by Dynamic Light Scattering (Zetasizer Nano S,
123 Malvern, France). Specific surface area of emulsion droplets was calculated using Laser Light
124 Scattering (Mastersizer 2000, Malvern, France). The melting temperature and crystalline state
125 of the fat was characterized by Differential Scanning Calorimetry (DSC) using a Q1000 DSC

126 (TA Instruments, New Castle, USA) and by powder X-ray Diffraction (XRD) using a D8
127 Advance diffractometer (Bruker, Germany).

128

129 **Hunger assessment**

130 Subjective assessment of hunger was measured on a 10-cm visual analogue scale 2 min
131 before breakfast and 2 min before lunch. Specific question to assess hunger was “How hungry
132 do you feel?”.

133

134 **Metabolic explorations**

135 Blood samples were obtained at baseline and at regular intervals after the meal, from an
136 antecubital arm vein through a catheter and collected in vacutainer sterile tubes (with EDTA
137 when necessary). Plasma was separated by centrifugation (1500 g, 10 min, 4°C) and stored at
138 -20°C until analysis or at 4°C for separation of the chylomicron-rich fraction (CMRF).

139 Metabolic tests were divided into postprandial phases including a first period of 5 hours (0
140 to 300 min) post-breakfast in the morning, a second period of 3 hours post-lunch (300 to 480
141 min) and the entire exploration day (0 to 480 min).

142 Indirect calorimetry was performed during metabolic testing using a Deltatrac II™
143 calorimeter (Sensormedics, Yorba Linda, CA, USA). Respiratory exchanges (VO_2 and VCO_2)
144 were recorded for periods of 30 or 60 minutes during the 8-h test period. Substrate oxidations
145 were calculated using Ferrannini's equations (19). Urine was collected at 0, 300 and 480 min
146 to determine nitrogen excretion for oxidation calculations. For breath test, expired gas
147 samples were obtained at baseline, each 30 min for 8-h and then at 720 and 1440 min to check
148 return to baseline. Subjects had to collect and freeze their stools individually over 72 h after
149 the test day.

150

151 **Plasma metabolite and hormone measurements**

152 Non esterified fatty acid (NEFA) concentrations were determined by an enzymatic
153 method Wako® (Neuss, Deutschland). ApoB48 was measured by ELISA (Gentaur, France).
154 Insulin concentration was determined by RIA (CISBIO Bi insuline IRMA, France).

155

156 **Isolation and analysis of chylomicron-rich fractions**

157 To collect the CMRF, containing chylomicrons and their large remnants, 250 µL of
158 plasma was deposited below a layer of 850 µL of distilled/deionized water and centrifuged at
159 80 000 rpm for 30 min using a Sorvall Kendro ultracentrifuge (Asheville, NC, USA). The
160 floating layer was collected and stored at -80°C. TAG and cholesterol concentrations of
161 CMRF were measured with a lipase glycerokinase and a cholesterol esterase/oxidase method,
162 respectively, on a AU 2700 Beckman Coulter® (O' Cllagan's mils, Ireland) and expressed as
163 differences in concentration over baseline. Hydrodynamic diameter of CMRF was measured
164 by dynamic light scattering at 25°C using a ZetaSizer NanoS (Malvern, UK) using 1.0658 cP
165 and 1.33 as viscosity and refractive index of the aqueous phase, respectively. Gravity-
166 separated fraction containing chylomicrons and their large remnants are reported to size in the
167 range 70-450 nm (20-22), and postprandial increase in cholesterol content of such fraction is
168 about <0.2 mM (23). Our fraction collected by ultracentrifugation contained particles of mean
169 diameter up to 200-300 nm and with postprandial increase in cholesterol of <0.15 mM in NW
170 and <0.25 mM in obese subjects. Our CMRF is thus typical of fractions that are rich in
171 chylomicrons and that also contain large remnants.

172

173 **¹³C-fatty acids in plasma lipids, NEFA, CMRF and stools**

174 *Sample preparation.* Internal standards were added according to the fraction analyzed
175 (heptadecanoic acid or glycerol triheptadecanoate).

176 *Plasma processing.* Plasma samples were submitted to direct methylation as described
177 previously (24).

178 *NEFA processing.* Total lipids were extracted from plasma aliquots at 120 min after
179 breakfast consumption (700 μ L) with 3 mL of a mixture of chloroform/methanol (2:1 v/v)
180 according to Folch method (25). NEFA fractions were obtained therefrom by TLC on silica-
181 gel plates with a mobile phase of hexane/diethyl ether/acetic acid (80:20:1 v/v/v). NEFA were
182 derivatized to Fatty Acid Methyl Esters (FAMES) (24).

183 *CMRF processing.* Lipids were extracted from CMRF at 120 min after breakfast
184 according to Folch method (25). TAG fractions were then processed as the NEFA fractions
185 above, to obtain the FAMES from CMRF.

186 *Stool processing.* Fecal collections were weighed, homogenized and a precisely weighed
187 aliquot was collected. Total lipids were extracted according to a modification of the Folch
188 method and derivatized to obtain FAMES (24).

189 *Sample analysis.* The amounts of FA in stools, plasma and NEFA were assessed by
190 GC/MS using a quadrupole mass spectrometer connected to a gas chromatograph (MS 5975
191 and GC6890, Agilent Technologies, Massy, France). The isotopic enrichment of palmitic and
192 oleic acids was determined using GC/C/IRMS (Isoprime, GV Instruments, Manchester,
193 UK)(24). The ^{13}C enrichments were expressed as atom percent excess (APE). The plasma
194 concentrations of non-esterified labelled palmitic and oleic acids (called [^{13}C -NEFA]) and
195 non-esterified unlabelled palmitic and oleic acids (called [^{12}C -NEFA]) were also obtained
196 from these analyses.

197

198 **Calculations associated with apparent dietary fatty acid “spillover”**

199 NEFA analysis at 120 min after breakfast was used to calculate ^{13}C enrichment in plasma
200 NEFA as: $[\text{C-NEFA}] / ([\text{C-NEFA}] + [\text{C-NEFA}])$ (expressed in % enrichment). The
201 proportion of exogenous NEFA in total plasma NEFA, expressed in %, was estimated by the

202 ratio of the ^{13}C enrichment in plasma NEFA to the ^{13}C enrichment of corresponding FA in the
 203 ingested milk fat. The proportion of exogenous fatty acids in plasma that was present in non-
 204 esterified form in the sum of pools NEFA+CMRF was calculated as: $[\text{}^{13}\text{C-NEFA}]_{\text{plasma}} /$
 205 $([\text{}^{13}\text{C-NEFA}]_{\text{plasma}} + [\text{}^{13}\text{C-FA}_{\text{CMRF}}]_{\text{plasma}})$, where $[\text{}^{13}\text{C-FA}_{\text{CMRF}}]_{\text{plasma}}$ is the plasma concentration
 206 of $^{13}\text{C-FA}$ esterified in CMRF-TAG = $[\text{}^{13}\text{C-FA}]_{\text{CMRF}} / ([\text{}^{13}\text{C-FA}]_{\text{CMRF}} + [\text{}^{12}\text{C-FA}]_{\text{CMRF}}) \times 3 \times$
 207 $[\text{CMRF-TAG}]_{\text{plasma}}$.

208

209 **Calculations of exogenous lipid oxidation from indirect calorimetry & breath tests**

210 Exogenous lipid oxidation was calculated according to Binnert *et al.* (18) from data of
 211 indirect calorimetry and breath tests. Here the formula was adapted to our use of 3 labeled
 212 triglycerides as follows:

213 Exogenous lipid oxidation (% of ingested fat) =

$$214 \frac{\{[\{[AP \text{CO}_2(t) + AP \text{CO}_2(t_{-30})] / 2\} - AP \text{CO}_2(t_0)] / 100\} \times \dot{V} \text{CO}_2}{\{(A) + (B) + (C)\} \times 22.4 \times dARF} \times 100$$

215

$$216 \text{ With: } (A) = \left\{ \left[AP \text{}^{13}\text{TG C8:0} \right] / 100 \right\} \times \left[(0.09 / 473.66) \times 27 \right]$$

$$217 (B) = \left\{ \left[AP \text{}^{13}\text{TG C16:0} \right] / 100 \right\} \times \left[(0.30 / 810.30) \times 51 \right]$$

$$218 (C) = \left\{ \left[AP \text{}^{13}\text{TG C18:1} \right] / 100 \right\} \times \left[(0.21 / 888.40) \times 57 \right]$$

219 Where AP $\text{CO}_2(t)$ is the AP value of the expired CO_2 at time t, AP $\text{CO}_2(t_0)$ is the AP value of
 220 the expired CO_2 at time t_0 , AP tracers is the calculated AP value of the labeled mixture of
 221 TAG (tracers) and $\dot{V} \text{CO}_2$ is the production rate of expired CO_2 (indirect calorimetry). Mean
 222 molecular weights of trioctanoin, tripalmitin and triolein are 473.66 g/mol, 810.30 g/mol and
 223 888.40 g/mol, respectively. Mean number of carbons in trioctanoin, tripalmitin and triolein
 224 are 27, 51 and 57, respectively. dARF (Acetate Recovery Factor) is the correction factor for

225 incomplete recovery of ^{13}C bicarbonate (0.505 for NW; 0.453 for obese (26)) and 22.4 is the
226 molar volume (L) of CO_2 .

227

228 **Kinetic parameters**

229 We calculated the incremental area under curve (iAUC); maximum postprandial
230 concentration, delta and diameter (C_{max} , Δ_{max} , d_{max}); time for appearance of these maximum
231 parameters (t_{max}) and appearance/enlargement-rates between 0 and 60 min.

232

233 **Statistical analysis**

234 Each subject served as his own control. All data are presented as means \pm SEM ($n=9$ per
235 group) and were analyzed with Statview 5.0 software (Abacus Concept, Berkeley, CA).
236 Postprandial data were compared by analysis of variance (ANOVA) for repeated measures
237 followed by post-hoc test (Fisher PLSD) for statistical effects of (i) time alone (P_{time}) over the
238 first postprandial period (0-300min), (ii) meal alone (P_{meal}) independently of the time in the
239 postprandial period and (iii) interaction of both factors, time and meal ($P_{\text{time}\times\text{meal}}$). Kinetic
240 parameters were compared by two-way ANOVA followed by Fisher PLSD according to meal
241 and BMI (P_{meal} , P_{BMI} , $P_{\text{meal}\times\text{BMI}}$) and time period before/after lunch ($P_{\text{meal}\times\text{BMI}\times\text{time}}$). Multiple
242 comparisons regarding tracer excretion in feces were performed using ANOVA followed by
243 Bonferroni post-hoc test. Comparisons between meals within subject groups were performed
244 using a paired Student's t -test and comparisons between subject groups within meals with an
245 unpaired Student's t -test. Differences were considered significant at the $P < 0.05$ level.

246

247 **RESULTS**

248 **Properties of emulsion vs spread fat**

249 The emulsion droplet size (Table 3; Supplemental Figure 3) indicates the homogenization
250 was effective in producing the emulsions. The emulsion had an approximately ~ 70000 -fold

251 greater surface area than the spread fat. To control for the possibility that the different
252 metabolic effects could be attributed to the fat melting properties, we measured melting
253 profiles and crystalline structures in all conditions (with or without tracers, emulsified or not,
254 Supplemental Figure 4). According to these analyses, the test fat differed only by their
255 structure (Table 3).

256

257 **Hunger feeling**

258 At the end of the first postprandial period (0-300 min) just before lunch, normal-weight
259 subjects felt similarly hungry regardless of breakfast type (Supplemental figure 5). In contrast,
260 obese subjects felt hungrier after emulsion than spread fat ($P < 0.05$; Supplemental figure 5).
261 Of note, before breakfasts, all subjects felt equally hungry (not shown).

262

263 **Postprandial concentration profile and size of chylomicron-rich fraction**

264 **Figure 1A-B** shows that in both groups, CMRF-TAG rapidly increased (60 min) after
265 ingestion of emulsified fat and peaked at 3-4 h (t_{\max} in **Table 4**). The emulsion induced a
266 significantly earlier and sharper increase in CMRF-TAG than the spread fat (Table 4: t_{\max} and
267 appearance-rate_{0-60min}; $P < 0.001$). These differences were dramatically marked in the obese
268 subjects, with a significant delay in absorption of the spread fat from 0 to 300 min compared
269 to NW subjects ($P < 0.01$, Table 4). At the end of the test, CMRF-TAG of NW subjects
270 returned to lower values regardless of fat structure. The obese subjects showed different
271 profiles, with CMRF-TAG remaining elevated at the end of the spread fat test: e.g. at 480
272 min, 0.61 ± 0.15 mmol/L for spread vs 0.27 ± 0.06 mmol/L for emulsion above fasting
273 baseline ($P < 0.05$). These differences in profile before and after lunch according to obese state
274 and meal type are supported by different BMI x meal x time interactions for the Δ_{\max} and
275 iAUC of CMRF-TAG (Table 4).

276 Mean CMRF size sharply increased in both groups from the first hour after emulsion
277 (Figure 1 C-D; $P_{\text{meal}} < 0.05$ for enlargement-rate_{0-60min}, Table 4). For NW subjects, CMRF
278 diameters were similar for spread fat and emulsion all along the test. In obese subjects, CMRF
279 diameters became equal for both meals at 300 min. We can note that in the period from 0 to
280 240 min, CMRF diameter in obese subjects was higher after emulsion vs spread fat from 0 to
281 240 min ($P_{\text{meal}} < 0.05$ and $P_{\text{time}} < 0.001$). Altogether, obese subjects presented larger CMRF
282 than NW subjects ($P < 0.01$ for d_{max} 0-480 min, Table 4) with persistence of large CMRF after
283 the second meal for spread fat.

284

285 **Plasma concentration profile of ApoB48**

286 Figure 1E-F shows that plasma ApoB48 changed over time in both groups after both
287 breakfasts ($P_{\text{time}} < 0.0001$) and differently according to the type of breakfast for NW subjects
288 ($P_{\text{time} \times \text{meal}} = 0.001$). At 120 min, NW subjects accumulated more ApoB48 after consumption of
289 emulsion than spread fat (7.08 ± 0.86 mg/L for spread fat vs 9.73 ± 0.69 mg/L for emulsion, $P <$
290 0.05) and compared with obese subjects for emulsion (7.47 ± 0.78 mg/L, $P < 0.05$).

291

292 **Plasma concentration profile of ¹³C-fatty acid tracers and fecal loss**

293 **Figure 2A-B-C-D** shows a change in plasma ¹³C-palmitic and ¹³C-oleic acids over time in
294 both groups after both breakfasts ($P_{\text{time}} < 0.0001$). ¹³C-palmitic acid appeared earlier and
295 sharper in plasma when it was in emulsion, differences between breakfasts being greater for
296 obese ($P_{\text{meal}} = 0.007$). Plasma concentrations of ¹³C-oleic acid were higher during 5h of
297 emulsion digestion, especially for obese subjects ($P_{\text{meal}} = 0.018$ and $P_{\text{time} \times \text{meal}} = 0.0002$). For
298 both tracers, a second peak was observed at 360 min, after ingestion of the second meal.

299 During the first 300 min for obese subjects, the iAUC for plasma ¹³C-FA were
300 significantly higher after consumption of emulsion vs spread fat ($P < 0.05$, Figure 2B-D).
301 iAUC after spread fat were lower for obese vs NW subjects ($P < 0.05$, **Figure 2B-D**).

302 Fecal excretion of ^{13}C -palmitic acid was higher than that of ^{13}C -oleic acid (Figure 2E).
303 There was no effect of breakfast type on fecal excretion of ^{13}C -palmitic acid or ^{13}C -oleic acid
304 in the two groups.

305

306 **Plasma concentration profile of insulin and NEFA and apparent dietary FA spillover**

307 **Figure 3A-B-C-D** shows a significant change in plasma insulin and NEFA over time after
308 the two breakfasts in both groups ($P_{\text{time}} < 0.0001$). Over the first 300 min, the NEFA profile
309 indicated a meal type x time interaction in both groups, with the decrease in plasma NEFA at
310 120 min being lower for emulsion than spread (Figure 3C-D). Therefore, we measured ^{13}C
311 enrichment in plasma NEFA at 120 minutes (Figure 3E) to estimate whether this would result
312 from the contribution of exogenous FA, so-called apparent fatty acid “spillover”. We
313 observed higher apparent “spillover” during the postprandial phase of emulsion vs spread fat,
314 in both groups ($P < 0.05$ for NW and $P < 0.01$ for obese subjects, Figure 3E). The contribution
315 of exogenous FA to total NEFA in NW subjects was 42% for spread vs 79% for emulsion,
316 and was lower in obese, 4% for spread vs 50% for emulsion (meal effect, $P < 0.01$; BMI
317 effect, $P < 0.01$; no meal x BMI interaction). Moreover, in NW subjects, the proportion of
318 exogenous FA being in non-esterified form in plasma at 120 min was 10.6 % for spread vs
319 18.7 % for emulsion, i.e. 1.8-fold increase, and in obese 10.1 % for spread vs 15.0% for
320 emulsion, i.e. 1.5-fold increase ($P < 0.05$ for spread vs emulsion; no significant effect of BMI
321 nor meal x BMI interaction). In the same time emulsification increased total plasma NEFA by
322 2.2-fold in NW and by 1.6-fold in obese subjects (Figure 3C-D). Altogether, this means that
323 more than 80 % of the increase of plasma NEFA due to emulsification may be explained by
324 an increased amount of exogenous fatty acids being released non-esterified in plasma
325 (spillover).

326

327 **Postprandial appearance of label in expired CO₂**

328 The ¹³C appearance in expired CO₂ represents the final product of FA β-oxidation. A
329 significant change in APE occurred over time in both groups ($P_{\text{time}} < 0.0001$; **Figure 4A-B**). In
330 both groups, APE was higher after consumption of emulsion *vs* spread fat over the first 300
331 min, indicating improvement of dietary fat β-oxidation using emulsified form. APE returned
332 to baseline after 720 min. An effect of BMI was also observed with higher appearance-
333 rate_{0-60min} in NW than obese subjects ($P < 0.01$; Table 4).

334 Inserts in Figure 4A-B show that AUC of expired ¹³CO₂ after 300 min for the emulsion
335 was significantly higher than for spread fat in both groups ($P < 0.01$ for NW subjects and $P <$
336 0.001 for obese subjects). Besides, over 0 to 720 min, obese subjects presented higher AUC
337 of expired ¹³CO₂ after consuming emulsion *vs* spread fat ($P < 0.05$, Figure 4B insert).
338 Altogether, the structuring of fat in the meal significantly affected the kinetic parameters of
339 ¹³CO₂ air enrichment regardless of BMI while BMI affected AUC and appearance-rate_{0-60min}
340 (Table 4).

341

342 **Exogenous lipid fate**

343 We studied the metabolic handling of exogenous lipids by evaluating the fractions of
344 ingested lipids that have been either oxidized or lost in feces, and so estimated the remaining
345 fraction stored in body pools. For the same quantity and composition of ingested fat, all
346 subjects β-oxidized FA better when fat was emulsified (Figure 4C). In turn, the calculated
347 fraction of ingested lipids oriented towards storage in body pools was lower after emulsion *vs*
348 spread consumption. After accounting for the part of exogenous lipids lost in feces (Figure
349 4C), the percentage of exogenous lipid oxidation according to the fraction that has been
350 intestinally absorbed was higher for NW and obese subjects after emulsion *vs* spread fat ($P <$
351 0.05 and $P < 0.01$, respectively, Figure 4D). Total lipid oxidation was unchanged with the two

352 test meals in both groups but with a greater relative contribution of exogenous lipids ingested
353 at breakfast as emulsion vs spread fat (Figure 4E).

354

355 **DISCUSSION**

356 Postprandial triglyceridemia is the first step in the metabolization of dietary lipids.
357 Ingested FA are firstly present in plasma TAG in the form of intestinally secreted
358 chylomicrons, which further lead to large remnants after hydrolysis by lipoprotein lipase (10,
359 22). The next step concerns trafficking of FA towards β -oxidation or storage that is of utmost
360 importance regarding the metabolic impact of these dietary FA. We therefore investigated
361 whether structuring fat in the meal could modify postprandial lipid metabolism, from the
362 amount and size of chylomicrons to β -oxidation, including fecal loss. To this aim, labeled
363 breakfasts containing either spread or emulsified fat were fed to NW and obese subjects. Test
364 meals were designed to be of equal composition. Thus, factors like FA composition or protein
365 content cannot be involved in the presently observed differences in lipid metabolism, which
366 can be uniquely attributed to the fat physico-chemical structure in the meal. The postprandial
367 chylomicron TAG profile after emulsion consumption differed from that of the spread fat,
368 with the peak being more rapidly achieved, more pronounced and more quickly cleared,
369 especially in obese subjects. This is consistent with reports of enhanced FA absorption when a
370 simple bolus of vegetable oil was emulsified in humans (27) and rodents(14, 15, 28). One
371 explanation is that our emulsion had ~70000-times greater surface area available for lipases
372 than spread fat, which is reported to enhance lipolysis and absorption (29). Enteral emulsions
373 of different droplet sizes, ~1 vs ~10 μm (14.5-fold difference in fat surface area), were shown
374 to result in small differences only in postprandial lipemia in humans (17). The dramatic
375 differences observed in the present work are due to the greatest differences in fat structure. Of
376 note, postprandial lipid metabolization was previously found faster using unemulsified than
377 emulsified milk fat in rats (14). Differences with the present results can be explained by (i)

378 rodent physiology of bile flow that is different from humans (30), and (ii) unemulsified
379 melted milk fat being force-fed intragastrically, prior to the proteinaceous phase. This could
380 have favored lipid emptying in the upper intestine and a rapid rise of plasma TAG in rats.
381 Our study also provides a proof of concept that effects of fat structure in the meal can occur in
382 a real mixed meal, while previous studies used oil or emulsion bolus fed orally or
383 intragastrically (17, 29). Regarding emulsifier type, our fat was emulsified by the proteins
384 naturally present in skim milk. Emulsions stabilized with caseins and monoacylglycerols were
385 recently reported to result in lower postprandial plasma TAG than those formulated with
386 lecithin in non-obese humans (31), which can be explained by lower *in vitro* digestive
387 lipolysis (32). Because obese men were the most affected by emulsification, further work
388 should test the effect of emulsifiers on postprandial lipid metabolism. The importance of
389 sensory exposure to lipids on postprandial metabolism in humans was also recently revealed
390 (5). Therefore, we cannot exclude a contribution of oral fat perception in our results.

391 TAG-rich lipoproteins remaining elevated all along the postprandial phase are an
392 independent CVD risk factor (3, 4), which can be mechanistically related to the atherogenic
393 potential of small chylomicron remnants (21, 22). In this study, we collected CMRF fractions
394 containing chylomicrons and their large remnants. The latter do not contribute significantly to
395 the formation of small atherogenic remnants because of direct hepatic clearance (33).
396 However, the role of TAG-rich particles is still debated, underlining that this is still an open
397 field needing further studies (1, 2). In this context, our study shows for the first time in NW
398 and obese men that postprandial lipemia profile can be modulated by structuring the fat in a
399 mixed meal. Therefore, the study of atherogenic small particles in the postprandial phase after
400 consumption of differently structured lipids in the meal should now be performed.

401 Obese men presented a delayed increase of CMRF-TAG after spread fat. Overweight men
402 were also reported to present delayed TAG-rich lipoprotein metabolism after a high-fat load
403 (34). Our observed differences between the two breakfasts in obese men could be explained

404 by their lower pancreatic secretion and lower levels of gallbladder emptying compared with
405 lean men (35). Therefore, obese subjects can better hydrolyze fat when it is pre-emulsified.
406 Moreover, fine stable emulsions were reported to be emptied faster and to cause greater
407 release of cholecystokinin than those that broke and layer in the stomach (36, 37). We can
408 thus suggest that the fat absorption delay observed with spread fat can be due to layering in
409 the stomach and thus delayed emptying. Of note, immediately after lunch, a peak of ^{13}C -FA
410 appeared in plasma. This so-called “second-meal effect” is known as the contribution of lipids
411 from a meal to lipemia after the next meal (38). For obese men, the marked delay in lipemia
412 appearance after spread fat cumulated with the second meal effect, causing high lipemia until
413 the test ended. In contrast, obese subjects (without fasting hyperlipidemia herein) did not have
414 difficulties in absorbing the emulsion, with a final return of lipemia to baseline.

415 In NW subjects, higher CMRF-TAG after the emulsion corresponded transiently to an
416 increased number of particles, as shown by the similar CMRF-size with an increased ApoB48
417 level at 120 min. In obese subjects however, ApoB48 levels remained similar, i.e., the
418 increase in lipemia after emulsion was due to an increased CMRF size. High particle numbers
419 estimated by ApoB48 level are reported to lead to increased chylomicron remnant numbers,
420 hence potentially increased atherosclerotic risk (10, 21). It would now be useful to explore the
421 chronic metabolic impact of fat structure, especially regarding ApoB48-containing particles.

422 The few reports about metabolic effects of emulsions have solely studied lipemia or
423 plasma FA concentrations as endpoints. For the first time to our knowledge, our study shows
424 that fat emulsification further affects the metabolic handling of exogenous FA, including β -
425 oxidation. Early appearance of $^{13}\text{CO}_2$ was due to the rapid β -oxidation of short-chain FA that
426 are directly absorbed in the portal vein and oxidized by the liver (39). Obesity is associated
427 with a defect in the β -oxidation of dietary FA (18, 40-42). Hodson *et al.* recently challenged
428 this idea by showing greater FA β -oxidation in obese men and attributed this to specific FA
429 acid partitioning (43). We highlight that exogenous FA oxidation can be enhanced in obese

430 men by emulsifying fat. Discrepancies between reports can thus be explained by the present
431 “fast vs slow lipid” notion. Indeed, lower β -oxidation in obese vs lean subjects was observed
432 using a single oil bolus (18) whereas higher β -oxidation in obese was observed when the
433 tracer was dispersed into an emulsion (43). This aspect had not been taken into account by
434 previous authors. Moreover, emulsification is now advised to enhance the intestinal
435 absorption of essential fatty acids (27). However, our results highlight the risk that such
436 essential FA quickly absorbed can be lost in the β -oxidation process rather than being
437 bioavailable for cell membrane turnover. Therefore, further studies on the structuring of oils
438 rich in essential PUFA should now investigate their final postprandial metabolic fate.

439 The effect of emulsification on exogenous lipid oxidation cannot be due to differences in
440 intestinal absorption because of similar fecal excretion. Total lipid oxidation during the test
441 day was unchanged by fat structuring, as well as total energy expenditure and diet-induced
442 thermogenesis. However, the source of β -oxidized FA was different: using emulsion,
443 exogenous FA ingested at breakfast were shunted towards β -oxidation pathways. Using
444 spread fat, more endogenous FA and/or exogenous FA ingested at lunch were oxidized so that
445 exogenous FA ingested at breakfast were more oriented towards storage. Therefore,
446 regardless of energy balance, FA metabolism is changed by lipid structure. This is consistent
447 with the greater FA spillover after emulsion. It can be explained by the faster intestinal
448 absorption, resulting in enhanced lipolysis of chylomicrons that generates exogenous NEFA
449 (44). Their early influx can serve as fuel for tissues and explain their higher contribution to
450 total FA oxidation with emulsion. However, high NEFA can also constitute a risk for ectopic
451 fat accumulation (44). Another aspect in obesity research concerns energy balance and satiety
452 regulation (45, 46). Just before lunch, our obese subjects felt hungrier after emulsion than
453 spread fat (visual analog scales, Supplemental figure 5). Further trials could test the impact of
454 fat structuring at breakfast on satiety regulation at lunch and energy balance.

455 In summary, we demonstrate that the postprandial metabolic handling of dietary FA can be
456 significantly modified by emulsifying the fat in the meal, especially in obese subjects. The
457 clinical perspectives of this first study should thus not be underrated. This study supports the
458 further exploration of a possible dietary concept of “fast vs slow lipid” for the nutritional
459 management of metabolic diseases through food formulation. Our results in the postprandial
460 phase raise the questions of whether (i) daily ingestion of “fast vs slow fat” would result in
461 different lipid metabolisms, adiposity and/or cardiovascular risk markers in the long term and
462 (ii) the composition and structuring of dietary lipids could be optimized to this aim.

463

464 **Acknowledgements**

465 We thank volunteers for their involvement. J Peyrat, C Maitrepierre, N Torche are
466 acknowledged for technical assistance, M Sothier for dietary survey analysis, and C Louche-
467 Pelissier, A Faure, M Cervantes, C Norbert and S Gonin for management of lipid analyses. B
468 Morio, JM Antoine, C Boyer and C Marmonier are acknowledged for useful discussions. We
469 thank RJ Ward for editing the English language. F Bellisle is acknowledged for advice on
470 visual analog scales for hunger responses. Corman SA (N Vaghefi) is acknowledged for
471 kindly providing AMF, Lactalis R&D (P Le Ruyet) for providing skimmed milk powder, and
472 Sodiaal-Candia R&D (O Cousin, G Vialaret) for preparing and processing bottled flavoured
473 skimmed milk.

474

475 The authors' responsibilities were as follows — CV: designed research, conducted research,
476 analyzed data, wrote paper, statistical analysis; GP: conducted research, technical support;
477 LG: conducted research, technical support, analyzed data; JD: conducted research, technical
478 and material support, analyzed data; CLP: conducted research, technical support; CD:
479 acquisition of data, technical support; DL: critical revision of the manuscript for important
480 intellectual content; MD: conducted research, analyzed data; SD: technical support, analyzed

481 data; SLP: designed research, administrative support; HV: critical revision of the manuscript
482 for important intellectual content; ML: critical revision of the manuscript for important
483 intellectual content; MCM: designed research, study supervision, analyzed data, wrote paper,
484 had primary responsibility for final content. All authors read and approved the final
485 manuscript. No conflict of interest exists for this manuscript.

REFERENCES

1. Lopez-Miranda J, Williams C, Lairon D. Dietary, physiological, genetic and pathological influences on postprandial lipid metabolism. *Br J Nutr* 2007;98(3):458-73. doi: 10.1017/S000711450774268X.
2. Nordestgaard BG, Benn M, Schnohr P, Tybjaerg-Hansen A. Nonfasting triglycerides and risk of myocardial infarction, ischemic heart disease, and death in men and women. *JAMA* 2007;298(3):299-308. doi: 10.1001/jama.298.3.299.
3. Chapman MJ, Ginsberg HN, Amarenco P, Andreotti F, Boren J, Catapano AL, Descamps OS, Fisher E, Kovanen PT, Kuivenhoven JA, et al. Triglyceride-rich lipoproteins and high-density lipoprotein cholesterol in patients at high risk of cardiovascular disease: evidence and guidance for management. *Eur Heart J* 2011;32(11):1345-61. doi: 10.1093/eurheartj/ehr112.
4. Maggi FM, Raselli S, Grigore L, Redaelli L, Fantappie S, Catapano AL. Lipoprotein remnants and endothelial dysfunction in the postprandial phase. *J Clin Endocrinol Metab* 2004;89(6):2946-50. doi: 10.1210/jc.2003-031977.
5. Chavez-Jauregui RN, Mattes RD, Parks EJ. Dynamics of fat absorption and effect of sham feeding on postprandial lipema. *Gastroenterology* 2010;139(5):1538-48. doi: 10.1053/j.gastro.2010.05.002.
6. Sakr SW, Attia N, Haourigui M, Paul JL, Soni T, Vacher D, Girard-Globa A. Fatty acid composition of an oral load affects chylomicron size in human subjects. *Br J Nutr* 1997;77(1):19-31.
7. Lai HC, Ney DM. Gastric digestion modifies absorption of butterfat into lymph chylomicrons in rats. *J Nutr* 1998;128(12):2403-10.
8. Renner F, Samuelson A, Rogers M, Glickman RM. Effect of saturated and unsaturated lipid on the composition of mesenteric triglyceride-rich lipoproteins in the rat. *J Lipid Res* 1986;27(1):72-81.
9. Feldman EB, Russell BS, Hawkins CB, Forte T. Intestinal lymph lipoproteins in rats fed diets enriched in specific fatty acids. *J Nutr* 1983;113(11):2323-34.
10. Xiao C, Lewis GF. Regulation of chylomicron production in humans. *Biochim Biophys Acta* 2012;1821(5):736-46. doi: 10.1016/j.bbalip.2011.09.019.
11. Willett W, Manson J, Liu S. Glycemic index, glycemic load, and risk of type 2 diabetes. *Am J Clin Nutr* 2002;76(1):274S-80S.
12. Hur SJ, Decker A, McClements DJ. Influence of initial emulsifier type on microstructural changes occurring in emulsified lipids during in vitro digestion. *Food Chemistry* 2009;114:253-62.
13. Michalski MC, Soares AF, Lopez C, Leconte N, Briard V, Geloën A. The supramolecular structure of milk fat influences plasma triacylglycerols and fatty acid profile in the rat. *Eur J Nutr* 2006;45(4):215-24. doi: 10.1007/s00394-006-0588-9.
14. Michalski MC, Briard V, Desage M, Geloën A. The dispersion state of milk fat influences triglyceride metabolism in the rat--a ^{13}C breath test study. *Eur J Nutr* 2005;44(7):436-44. doi: 10.1007/s00394-005-0551-1.
15. Laugerette F, Vors C, Geloën A, Chauvin MA, Soulage C, Lambert-Porcheron S, Peretti N, Alligier M, Burcelin R, Laville M, et al. Emulsified lipids increase endotoxemia: possible role in early postprandial low-grade inflammation. *J Nutr Biochem* 2011.

16. McQuaid SE, Hodson L, Neville MJ, Dennis AL, Cheeseman J, Humphreys SM, Ruge T, Gilbert M, Fielding BA, Frayn KN, et al. Downregulation of adipose tissue fatty acid trafficking in obesity: a driver for ectopic fat deposition? *Diabetes* 2011;60(1):47-55. doi: 10.2337/db10-0867.
17. Armand M, Pasquier B, Andre M, Borel P, Senft M, Peyrot J, Salducci J, Portugal H, Jaussan V, Lairon D. Digestion and absorption of 2 fat emulsions with different droplet sizes in the human digestive tract. *Am J Clin Nutr* 1999;70(6):1096-106.
18. Binnert C, Pachiaudi C, Beylot M, Hans D, Vandermander J, Chantre P, Riou JP, Laville M. Influence of human obesity on the metabolic fate of dietary long- and medium-chain triacylglycerols. *Am J Clin Nutr* 1998;67(4):595-601.
19. Ferrannini E. The theoretical bases of indirect calorimetry: a review. *Metabolism* 1988;37(3):287-301.
20. Hussain MM, Kedeas MH, Singh K, Athar H, Jamali NZ. Signposts in the assembly of chylomicrons. *Front Biosci* 2001;6:D320-31.
21. Pang J, Chan DC, Barrett PH, Watts GF. Postprandial dyslipidaemia and diabetes: mechanistic and therapeutic aspects. *Curr Opin Lipidol* 2012;23(4):303-9. doi: 10.1097/MOL.0b013e328354c790.
22. Su JW, Nzekwu MM, Cabezas MC, Redgrave T, Proctor SD. Methods to assess impaired post-prandial metabolism and the impact for early detection of cardiovascular disease risk. *Eur J Clin Invest* 2009;39(9):741-54. doi: 10.1111/j.1365-2362.2009.02179.x.
23. Dubois C, Beaumier G, Juhel C, Armand M, Portugal H, Pauli AM, Borel P, Latge C, Lairon D. Effects of graded amounts (0-50 g) of dietary fat on postprandial lipemia and lipoproteins in normolipidemic adults. *Am J Clin Nutr* 1998;67(1):31-8.
24. Gabert L, Vors C, Louche-Pelissier C, Sauvinet V, Lambert-Porcheron S, Draï J, Laville M, Desage M, Michalski MC. ¹³C tracer recovery in human stools after digestion of a fat-rich meal labelled with [1,1,1-¹³C₃]tripalmitin and [1,1,1-¹³C₃]triolein. *Rapid Commun Mass Spectrom* 2011;25(19):2697-703. doi: 10.1002/rcm.5067.
25. Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem* 1957;226(1):497-509.
26. Antoun E, Momken I, Bergouignan A, Villars C, Platat C, Schoeller DA, Blanc S, Simon C. The [1-¹³C]acetate recovery factor to correct tracer-derived dietary fat oxidation is lower in overweight insulin-resistant subjects. *The European e-Journal of Clinical Nutrition and Metabolism* 2010;5(4):173-9.
27. Garaiova I, Guschina IA, Plummer SF, Tang J, Wang D, Plummer NT. A randomised cross-over trial in healthy adults indicating improved absorption of omega-3 fatty acids by pre-emulsification. *Nutr J* 2007;6:4. doi: 10.1186/1475-2891-6-4.
28. Couedelo L, Boue-Vaysse C, Fonseca L, Montesinos E, Djoukitch S, Combe N, Cansell M. Lymphatic absorption of alpha-linolenic acid in rats fed flaxseed oil-based emulsion. *Br J Nutr* 2011;105(7):1026-35. doi: 10.1017/S000711451000454X.
29. Delorme V, Dhouib R, Canaan S, Fotiadu F, Carriere F, Cavalier JF. Effects of surfactants on lipase structure, activity, and inhibition. *Pharm Res* 2011;28(8):1831-42. doi: 10.1007/s11095-010-0362-9.
30. Martins PN, Neuhaus P. Surgical anatomy of the liver, hepatic vasculature and bile ducts in the rat. *Liver Int* 2007;27(3):384-92. doi: 10.1111/j.1478-3231.2006.01414.x.

31. Keogh JB, Wooster TJ, Golding M, Day L, Otto B, Clifton PM. Slowly and rapidly digested fat emulsions are equally satiating but their triglycerides are differentially absorbed and metabolized in humans. *J Nutr* 2011;141(5):809-15. doi: 10.3945/jn.110.131110.
32. Vors C, Capolino P, Guerin C, Meugnier E, Pesenti S, Chauvin MA, Monteil J, Peretti N, Cansell M, Carriere F, et al. Coupling in vitro gastrointestinal lipolysis and Caco-2 cell cultures for testing the absorption of different food emulsions. *Food Funct* 2012;3(5):537-46. doi: 10.1039/c2fo10248j.
33. Karpe F, Olivecrona T, Hamsten A, Hultin M. Chylomicron/chylomicron remnant turnover in humans: evidence for margination of chylomicrons and poor conversion of larger to smaller chylomicron remnants. *J Lipid Res* 1997;38(5):949-61.
34. Nabeno-Kaeriyama Y, Fukuchi Y, Hayashi S, Kimura T, Tanaka A, Naito M. Delayed postprandial metabolism of triglyceride-rich lipoproteins in obese young men compared to lean young men. *Clin Chim Acta* 2010;411(21-22):1694-9. doi: 10.1016/j.cca.2010.07.004.
35. Wisen O, Johansson C. Gastrointestinal function in obesity: motility, secretion, and absorption following a liquid test meal. *Metabolism* 1992;41(4):390-5.
36. Marciani L, Wickham M, Singh G, Bush D, Pick B, Cox E, Fillery-Travis A, Faulks R, Marsden C, Gowland PA, et al. Enhancement of intragastric acid stability of a fat emulsion meal delays gastric emptying and increases cholecystokinin release and gallbladder contraction. *Am J Physiol Gastrointest Liver Physiol* 2007;292(6):G1607-13. doi: 10.1152/ajpgi.00452.2006.
37. Marciani L, Faulks R, Wickham MS, Bush D, Pick B, Wright J, Cox EF, Fillery-Travis A, Gowland PA, Spiller RC. Effect of intragastric acid stability of fat emulsions on gastric emptying, plasma lipid profile and postprandial satiety. *Br J Nutr* 2009;101(6):919-28. doi: 10.1017/S0007114508039986.
38. Silva KD, Wright JW, Williams CM, Lovegrove JA. Meal ingestion provokes entry of lipoproteins containing fat from the previous meal: possible metabolic implications. *Eur J Nutr* 2005;44(6):377-83. doi: 10.1007/s00394-004-0538-3.
39. Michalski MC. Specific molecular and colloidal structures of milk fat affecting lipolysis, absorption and postprandial lipemia. *European Journal of Lipid Science and Technology* 2009;111(5):413-31. doi: DOI 10.1002/ejlt.200800254.
40. Schutz Y, Flatt JP, Jequier E. Failure of dietary fat intake to promote fat oxidation: a factor favoring the development of obesity. *Am J Clin Nutr* 1989;50(2):307-14.
41. Giacco R, Clemente G, Busiello L, Lasorella G, Riviaccio AM, Rivellese AA, Riccardi G. Insulin sensitivity is increased and fat oxidation after a high-fat meal is reduced in normal-weight healthy men with strong familial predisposition to overweight. *Int J Obes Relat Metab Disord* 2004;28(2):342-8. doi: 10.1038/sj.ijo.0802589.
42. Bessesen DH, Rupp CL, Eckel RH. Dietary fat is shunted away from oxidation, toward storage in obese Zucker rats. *Obes Res* 1995;3(2):179-89.
43. Hodson L, McQuaid SE, Humphreys SM, Milne R, Fielding BA, Frayn KN, Karpe F. Greater dietary fat oxidation in obese compared with lean men: an adaptive mechanism to prevent liver fat accumulation? *Am J Physiol Endocrinol Metab* 2010;299(4):E584-92. doi: 10.1152/ajpendo.00272.2010.

44. Fielding B. Tracing the fate of dietary fatty acids: metabolic studies of postprandial lipaemia in human subjects. *Proc Nutr Soc* 2011;70(3):342-50. doi: 10.1017/S002966511100084X.
45. Halford JC, Harrold JA. Satiety-enhancing products for appetite control: science and regulation of functional foods for weight management. *Proc Nutr Soc* 2012;71(2):350-62. doi: 10.1017/S0029665112000134.
46. Simpson K, Parker J, Plumer J, Bloom S. CCK, PYY and PP: the control of energy balance. *Handb Exp Pharmacol* 2012(209):209-30. doi: 10.1007/978-3-642-24716-3_9.

Table 1. Anthropometric and fasting metabolic subject parameters.

	Normal weight (n=9)	Obese (n=9)	P value
Anthropometric parameters			
Age (years)	28.3 ± 1.4	30.2 ± 2.2	<i>ns</i>
Body weight (kg)	72.0 ± 2.1	101.2 ± 1.9	< 0.0001
BMI (kg.m ²)	22.3 ± 0.5	31.7 ± 0.3	< 0.0001
Waist circumference (cm)	83.3 ± 1.6	105.9 ± 0.8	< 0.0001
Fasting metabolic parameters			
Glucose (mM)	4.94 ± 0.16	5.19 ± 0.15	<i>ns</i>
Insulin (mIU/L)	3.75 ± 0.59	7.14 ± 0.95	0.008
HOMA	0.85 ± 0.14	1.69 ± 0.25	0.009
Total cholesterol (mM)	4.85 ± 0.22	4.89 ± 0.24	<i>ns</i>
HDL cholesterol (mM)	1.51 ± 0.10	1.09 ± 0.06	0.004
LDL cholesterol (mM)	3.03 ± 0.27	3.11 ± 0.21	<i>ns</i>
Triacylglycerols (mM)	0.85 ± 0.06	1.39 ± 0.18	0.017

Data are means ± SEM. Groups are compared using unpaired Student's *t*-test.

Table 2. Nutritional composition of the test breakfasts containing either spread or emulsified fat enriched with ^{13}C -labelled triglycerides.

Breakfast composition¹				
	Quantity (g or mL)	Carbohydrates (g)	Proteins (g)	Lipids (g)
Anhydrous Milk Fat ²	40	-	-	40
Skimmed milk ³	160	7.5	5.3	0.3
Bread	50	28	4	0.5
[1,1,1- $^{13}\text{C}_3$] triolein	0.09	-	-	0.09
[1,1,1- $^{13}\text{C}_3$] tripalmitin	0.30	-	-	0.30
[1,1,1- $^{13}\text{C}_3$] triolein	0.21	-	-	0.21
Total (g)	250.6	35.5	9.3	41.4
% caloric intake		26	7	67

¹ Identical nutrient composition for both spread and emulsion breakfasts.

² Fatty acid profile of TAG includes 68.6% SFA, 28.1% MUFA and 3.3% PUFA.

³ Natural vanillia-flavoured.

Table 3. Physico-chemical properties of fat used in the formulation of test breakfasts.

Structure (type of breakfast)	Fat properties ¹			
	Droplet size ² (μm)	d_{32} ³ (μm)	Fat surface area in meal ⁴ (m^2)	Melting temperature ⁵ ($^{\circ}\text{C}$)
Spread	-	-	0.006	42
Emulsion	1.04	0.63	410	40

¹Mixture of milk fat + ¹³C-TAG tracers.

²Diameter of the peak of maximum intensity measured by DLS.

³Surface averaged diameter measured by LLS.

⁴For Spread fat: calculated as the surface of an equivalent sphere of 40 g. For Emulsion: calculated from the specific surface area (m^2 per g fat) calculated by the software, further multiplied by fat content in the meal.

⁵Temperature at which the entire fat amount is in liquid form.

Table 4. Kinetic parameters after digestion of test breakfasts in NW vs obese subjects.

Parameter	Spread fat 40 g		Emulsified fat 40 g		P value			
	Normal-weight	Obese	Normal-weight	Obese	P_{meal}^1	P_{BMI}^1	$P_{meal \times BMI}^1$	$P_{meal \times BMI \times time}^2$
Δ CMRF TAG								
0-480 min:								
Δ_{max} (mmol/L)	0.63 ± 0.13	0.80 ± 0.12	0.75 ± 0.09	0.94 ± 0.25	<i>ns</i>	<i>ns</i>	<i>ns</i>	
iAUC (mmol·min/L)	132.8 ± 29.1	165.6 ± 23.1	180.4 ± 28.2	218.1 ± 53.2	<i>ns</i>	<i>ns</i>	<i>ns</i>	
t_{max} (min)	293 ± 23	367 ± 33	220 ± 42	207 ± 25	< 0.001	<i>ns</i>	<i>ns</i>	
Appearance-rate _{0-60 min} (μmol/L/min)	0.32 ± 0.27	0.14 ± 0.32	2.85 ± 0.72	2.93 ± 0.6	<0.0001	<i>ns</i>	<i>ns</i>	
0-300 min:								
Δ_{max} (mmol/L)	0.56 ± 0.10	0.53 ± 0.09	0.74 ± 0.09	0.94 ± 0.25	0.05	<i>ns</i>	<i>ns</i>	
iAUC (mmol·min/L)	81.4 ± 18.2	57.6 ± 10.7	127.7 ± 17.4	159.3 ± 39.6	<i>ns</i>	< 0.01	<i>ns</i>	
t_{max} (min)	233 ± 16	267 ± 15	167 ± 17	193 ± 17	<0.0001	< 0.1	<i>ns</i>	
300-480 min:								
Δ_{max} (mmol/L)	0.54 ± 0.13	0.80 ± 0.12	0.54 ± 0.11	0.56 ± 0.14	<i>ns</i>	<i>ns</i>	<i>ns</i>	< 0.05
iAUC (mmol·min/L)	51.4 ± 12.9	107.9 ± 17.7	52.7 ± 11.2	58.8 ± 14.7	< 0.05	<i>ns</i>	<i>ns</i>	< 0.01
t_{max} (min)	347 ± 5	390 ± 23	397 ± 19	420 ± 24	< 0.05	< 0.1	<i>ns</i>	< 0.1
CMRF Size								
0-480 min:								
d_{max} (nm)	253 ± 34	494 ± 93	262 ± 20	344 ± 58	<i>ns</i>	< 0.01	<i>ns</i>	
t_{max} (min)	243 ± 25	307 ± 23	207 ± 30	237 ± 33	< 0.1	0.1	<i>ns</i>	
Enlargement-rate _{0-60min} (μmol/L/min)	-0.06 ± 0.07	-0.02 ± 0.11	0.65 ± 0.11	1.26 ± 0.78	< 0.05	<i>ns</i>	<i>ns</i>	
0-300 min:								
d_{max} (nm)	246 ± 35	296 ± 70	239 ± 12	336 ± 58	<i>ns</i>	< 0.1	<i>ns</i>	
t_{max} (min)	180 ± 17	200 ± 41	200 ± 26	193 ± 28	<i>ns</i>	<i>ns</i>	<i>ns</i>	
300-480 min:								
d_{max} (nm)	195 ± 10	451 ± 96	207 ± 28	236 ± 46	< 0.1	< 0.05	< 0.05	0.055
t_{max} (min)	337 ± 4	367 ± 17	340 ± 5	360 ± 16	<i>ns</i>	< 0.05	<i>ns</i>	<i>ns</i>
¹³CO₂ enrichment								
0-720 min:								
C_{max} (%)	0.019 ± 0.001	0.013 ± 0.001	0.019 ± 0.001	0.016 ± 0.001	< 0.01	<i>ns</i>	<i>ns</i>	
AUC (%.min)	6.9 ± 0.5	4.8 ± 0.8	7.7 ± 0.2	6.4 ± 0.5	< 0.01	< 0.001	<i>ns</i>	
t_{max} (min)	310 ± 21	347 ± 23	267 ± 26	267 ± 17	< 0.01	<i>ns</i>	<i>ns</i>	
Appearance-rate _{0-60 min} (%/min)	6.0 · 10 ⁻⁵ ± 1.7 · 10 ⁻⁵	2.0 · 10 ⁻⁵ ± 0.5 · 10 ⁻⁵	13.6 · 10 ⁻⁵ ± 1.9 · 10 ⁻⁵	9.8 · 10 ⁻⁵ ± 0.8 · 10 ⁻⁵	< 0.0001	< 0.01	<i>ns</i>	
0-300 min:								
C_{max} (mmol/L)	0.018 ± 0.001	0.011 ± 0.001	0.019 ± 0.001	0.016 ± 0.001	< 0.05	< 0.0001	< 0.05	
AUC (%.min)	2.9 ± 0.3	1.6 ± 0.4	3.9 ± 0.2	3.2 ± 0.2	< 0.0001	< 0.0001	<i>ns</i>	
t_{max} (min)	283 ± 17	293 ± 7	257 ± 23	260 ± 14	< 0.1	<i>ns</i>	<i>ns</i>	
300-720 min:								
C_{max} (mmol/L)	0.019 ± 0.001	0.013 ± 0.001	0.017 ± 0.001	0.016 ± 0.001	<i>ns</i>	0.0001	< 0.05	<i>ns</i>
AUC (%.min)	3.9 ± 0.4	3.2 ± 0.5	3.8 ± 0.3	3.1 ± 0.4	<i>ns</i>	< 0.1	<i>ns</i>	<i>ns</i>
t_{max} (min)	333 ± 3	360 ± 18	330 ± 0	330 ± 0	< 0.1	<i>ns</i>	<i>ns</i>	<i>ns</i>

Data are means ± SEM, $n=9$ per group. Parameters calculated over the indicated time period: C_{max} indicates maximum concentration; d_{max} indicates maximum diameter; Δ_{max} indicates maximum concentration delta; iAUC, incremental area under the curve; AUC, area under the curve. P values (P_{meal} , P_{BMI} and $P_{meal \times BMI}$) obtained by ANOVA followed by post hoc Fisher PLSD.

¹ P values of two-way ANOVA for meal and BMI effects and their interactions.

² P values of two-way ANOVA for repeated measures regarding both time periods (before and after 300 min), for meal x BMI x time period interactions.

Figure Caption

Figure 1. Postprandial profile after consuming spread fat (□, ■) or emulsion (○, ●): CMRF-TAG (mM) in NW (A) and obese subjects (B) and corresponding iAUC; CMRF size (nm) in NW (C) and obese subjects (D); ApoB48 (mg/L) in NW (E) and obese subjects (F). Data are means \pm SEM, $n=9$ per group; P_{time} , P_{meal} and $P_{\text{time}\times\text{meal}}$ for postprandial period from 0 to 300 min (repeated measures-ANOVA followed by post-hoc Fisher PLSD).

(B) ** $P < 0.01$ for time 420 min emulsion vs spread fat (paired Student's t -test); * $P < 0.05$ for emulsion vs spread fat at time 480 min and for iAUC₀₋₃₀₀ (paired Student's t -test); § $P < 0.05$ for obese vs NW regarding spread fat iAUC₃₀₀₋₄₈₀ (unpaired Student's t -test).

(D) * $P < 0.05$ for time 120 min emulsion vs spread fat (paired Student's t -test)

(C-D) §§ $P < 0.01$ for time 120 min obese vs NW subjects (unpaired Student's t -test);

(E-F) * $P < 0.05$ for time 120 min emulsion vs spread fat (paired Student's t -test), § $P < 0.05$ for time 120 min obese vs NW subjects (unpaired Student's t -test).

Figure 2. Postprandial concentration profile and iAUC of total plasma lipids of ¹³C-palmitic acid (mM) and ¹³C-oleic acid (mM) in NW (A & C, respectively) and obese subjects (B & D, respectively) consuming spread fat (□, ■) or emulsion (○, ●). (E) Fecal excretion of ¹³C-palmitic acid and ¹³C-oleic acid in NW (white bars) and obese subjects (black bars) consuming spread fat (plain bars) or emulsion (droplet pattern bars).

Data are means \pm SEM, $n=9$ per group.

(A-B-C-D) P_{time} , P_{meal} and $P_{\text{time} \times \text{meal}}$ for postprandial period from 0 to 300 min (repeated measures-ANOVA followed by post-hoc Fisher PLSD).

(A-B) * $P < 0.05$ for obese iAUC 0-300 min emulsion vs spread fat (paired Student's t -test), § $P < 0.05$ for spread fat iAUC 0-300 min obese vs NW subjects (unpaired Student's t -test), § $P < 0.1$ for obese iAUC 0-480 min emulsion vs spread fat (paired Student's t -test); (C-D) * $P < 0.05$ for obese iAUC 0-300 min emulsion vs spread fat (paired Student's t -test), § $P < 0.05$ for spread fat iAUC 0-300 min obese vs NW subjects (unpaired Student's t -test); (E) no common letter with another bar indicates a statistical difference, $P < 0.001$ (ANOVA followed by post-hoc Bonferroni).

Figure 3. Postprandial concentration profile of insulin (mUI/L) and NEFA (μM) in NW (A & C, respectively) and obese subjects (B & D, respectively) consuming spread fat (\square , \blacksquare) or emulsion (\circ , \bullet). (E) ^{13}C -enrichment of plasma NEFA at 120 min, estimating so-called apparent fatty acid “spillover” in NW (white bars) and obese subjects (black bars) consuming spread fat (plain bars) or emulsion (droplet pattern bars).

Data are means \pm SEM, $n=9$ per group; P_{time} , P_{meal} and $P_{\text{time} \times \text{meal}}$ for postprandial period from 0 to 300 min (repeated measures-ANOVA followed by post-hoc Fisher PLSD).

(C) ** $P < 0.01$ for NW subjects at 120 min emulsion vs spread fat (paired Student's t -test);

(D) * $P < 0.05$ for obese subjects at 120 min emulsion vs spread fat (paired Student's t -test);

(E) * $P < 0.05$ for NW subjects at 120 min emulsion vs spread fat (paired Student's t -test);

** $P < 0.01$ for obese subjects at 120 min emulsion vs spread fat (paired Student's t -test);

ANOVA analysis revealed meal effect ($P < 0.01$) and BMI effect ($P < 0.01$) but no significant meal x BMI interaction.

Figure 4. Postprandial profile of ^{13}C -appearance in breath in NW subjects (A) and obese subjects (B) consuming spread fat (\square , \blacksquare) or emulsion (\circ , \bullet); (C) Exogenous lipid fate, either oxidized (white), lost in feces (black) or stored (grey, calculated as “total – (lost+oxidized)”) in NW and obese subjects over 480 min; (D) Oxidation of intestinally absorbed lipids over 480 min in NW (white bars) and obese subjects (black bars) consuming spread fat (plain bars) or emulsion (droplet pattern bars); (E) Total lipid oxidation (total bar) and fraction of cumulative exogenous lipid oxidation (dashed part) over 480 min after consuming test breakfasts in NW vs obese subjects.

Data are means \pm SEM, $n=9$ per group.

(C-D-E) * $P < 0.05$ for NW subjects and ** $P < 0.01$ for obese subjects emulsion vs spread fat (paired Student's t -test). ANOVA analysis revealed no significant meal x BMI interaction ($P=0.087$).