



Contrasted styles of rifting in the eastern Gulf of Aden: A combined wide-angle, multichannel seismic, and heat flow survey

Sylvie Leroy, F. Lucazeau, Elia D'Acremont, Louise Watremez, Julia Autin,
Stéphane Rouzo, Nicolas Bellahsen, Christel Tiberi, C. Ebinger, Marie-Odile
Beslier, et al.

► **To cite this version:**

Sylvie Leroy, F. Lucazeau, Elia D'Acremont, Louise Watremez, Julia Autin, et al.. Contrasted styles of rifting in the eastern Gulf of Aden: A combined wide-angle, multichannel seismic, and heat flow survey. *Geochemistry, Geophysics, Geosystems*, AGU and the Geochemical Society, 2010, 11 (7), pp.Q07004. <10.1029/2009GC002963>. <hal-00504866>

HAL Id: hal-00504866

<https://hal.archives-ouvertes.fr/hal-00504866>

Submitted on 2 May 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Contrasted styles of rifting in the eastern Gulf of Aden: A combined wide-angle, multichannel seismic, and heat flow survey

Sylvie Leroy

ISTeP, UMR 7193, UPMC, Université Paris 06, CNRS, F-75005 Paris, France (sylvie.leroy@upmc.fr)

Francis Lucazeau

Géosciences Marines, UMR 7097, IPGP, Case 89, 4 place Jussieu, F-75252 Paris CEDEX 05, France

**Elia d'Acremont, Louise Watremez, Julia Autin, Stéphane Rouzo, Nicolas Bellahsen, and
Christel Tiberi**

ISTeP, UMR 7193, UPMC, Université Paris 06, CNRS, F-75005 Paris, France

Cynthia Ebinger

Department of Earth and Environmental Sciences, University of Rochester, Rochester, New York 14627, USA

Marie-Odile Beslier

Géosciences Azur, UMR 6526, BP 48, F-06235 Villefranche sur Mer CEDEX, France

Julie Perrot

Domaines Océaniques, IUEM, CNRS, Place Nicolas Copernic, F-29280 Plouzané, France

Philippe Razin

EGID, University of Bordeaux III, 1 allée Daguin, F-33607 Pessac CEDEX, France

Frédérique Rolandone

ISTeP, UMR 7193, UPMC, Université Paris 06, CNRS, F-75005 Paris, France

Heather Sloan

Department of Environmental, Geographic, and Geological Sciences, Lehman College, City University of New York, Bronx, New York 10468, USA

Graham Stuart

School of Earth and Environment, University of Leeds, 15-19 Hyde Terrace, Leeds LS2 9JT, UK

Ali Al Lazki and Khalfan Al-Toubi

College of Science and Earthquake Monitoring Center, Sultan Qaboos University, PC 120 Al Khod, Sultanate of Oman

François Bache

ISTeP, UMR 7193, UPMC, Université Paris 06, CNRS, F-75005 Paris, France

Alain Bonneville and Bruno Goutorbe

Géosciences Marines, UMR 7097, IPGP, Case 89, 4 place Jussieu, F-75252 Paris CEDEX 05, France

Philippe Huchon*ISTeP, UMR 7193, UPMC, Université Paris 06, CNRS, F-75005 Paris, France***Patrick Unternehr***TOTAL, Tour Coupole, 2, Place de la Coupole, F-92078 Paris La Défense CEDEX, France***Khaled Khanbari***Department of Environmental and Earth Sciences, Sana'a University, Sana'a, Republic of Yemen*

[1] Continental rifts and passive continental margins show fundamental along-axis segmentation patterns that have been attributed to one or a number of different processes: extensional fault geometry, variable stretching along strike, preexisting lithospheric compositional and structural heterogeneities, oblique rifting, and the presence or absence of eruptive volcanic centers. The length and width scales of the rift stage fault-bounded basin systems change during the late evolution of the new plate boundary, and the role of magmatism may increase as rifting progresses to continental rupture. Along obliquely spreading ridges, first-order mid-ocean ridge geometries originate during the synrift stage, indicating an intimate relationship between magma production and transform fault spacing and location. The Gulf of Aden rift is a young ocean basin in which the earliest synrift to breakup structures are well exposed onshore and covered by thin sediment layers offshore. This obliquely spreading rift is considered magma-poor and has several large-offset transforms that originated during late stage rifting and control the first-order axial segmentation of the spreading ridge. Widely spaced geophysical transects of passive margins that produce only isolated 2-D images of crust and uppermost mantle structure are inadequate for evaluation of competing rift evolution models. Using closely spaced new geophysical and geological observations from the Gulf of Aden we show that rift sectors between transforms have a large internal variability over short distances (~10 km): the ocean-continent transition (OCT) evolves from a narrow magmatic transition to wider zones where continental mantle is probably exhumed. We suggest that this small-scale variability may be explained (1) by the distribution of volcanism and (2) by the along-strike differences in time-averaged extension rate of the oblique rift system. The volcanism may be associated with (1) the long-offset Alula-Fartak Fracture Zone, which may enhance magma production on its younger side, or (2) channeled flow from the Afar plume material along the newly formed OCT and the spreading ridge. Oblique extension and/or hot spot interactions may thereby have a significant control on the styles of rifting and continental breakup and on the evolution of many magma-poor margins.

Components: 7800 words, 5 figures, 1 table.

Keywords: magma-poor rifted margins; oblique rifting; mantle exhumation; Gulf of Aden.

Index Terms: 8105 Tectonophysics: Continental margins: divergent (1212); 8106 Tectonophysics: Continental margins: transform (8031); 8159 Tectonophysics: Rheology: crust and lithosphere (8031).

Received 17 November 2009; **Revised** 19 April 2010; **Accepted** 19 May 2010; **Published** 8 July 2010.

Leroy, S., et al. (2010), Contrasted styles of rifting in the eastern Gulf of Aden: A combined wide-angle, multichannel seismic, and heat flow survey, *Geochem. Geophys. Geosyst.*, 11, Q07004, doi:10.1029/2009GC002963.

1. Introduction

[2] Continental rifting leading to plate rupture is achieved through lithospheric crust and mantle thinning in a direction parallel to the plate opening; the resulting margin architecture constrains crust and mantle processes [e.g., Buck, 2004]. Variability of crust and mantle structures along the strike of

the rift also influence processes, further restricting acceptable models: where the production of melt is large (volcanic margins), the transition from continent to ocean is narrow and substantial amounts of magma are added to the lower crust; where melt production is small (magma-poor margins), the transition can extend over hundreds of kilometers with possible exhumation of the continental mantle

along low-angle detachment faults [Hopper and Buck, 1996; Buck *et al.*, 1999]. Along-axis mid-ocean ridge segmentation may be related to one or several processes such as extensional fault geometry, variable rate of extension along strike, structural inheritance and presence or absence of volcanism [e.g., Ebinger *et al.*, 1987; Dunbar and Sawyer, 1989; Hopper *et al.*, 1992; Brun, 1999; Buck *et al.*, 1999; Lizarralde *et al.*, 2007; Péron-Pinvidic and Manatschal, 2009]. The geometry of the fault-bounded systems may change during rifting, and the influence of magmatism may differ when it occurs before or after the continental breakup [White, 1992; Yamasaki and Gernigon, 2009]. Zones that accommodate deformation between extensional basins of obliquely spreading rift systems during late synrift stages could be precursors of the first-order transform faults that appear along the OCT during or soon after continental rupture [Corti *et al.*, 2003; d'Acremont *et al.*, 2005]. It has also been proposed that transform fault configuration may not be inherited from earlier structures at all [Taylor *et al.* 2009]. Finally, long-offset transform fault and fracture zone geometry at obliquely spreading ridges may affect melt supply to axial segments from the onset of seafloor spreading following rifting [Hoofft *et al.*, 2000] and possibly control magma flow in plume-ridge interactions [Georgen *et al.*, 2001].

[3] Lateral structure of rifted margins can be highly variable, but is poorly constrained due to the wide spacing between geophysical transects (more than 100 km). This was the case for the Gulf of Aden where previously sparse studies consisted of seafloor bathymetry, limited gravity transects and seismic profiles that gave the impression that there was a lack of east-west structural variation. In this paper, we focus on the structure of the northern margin of the Gulf of Aden between the Alula-Fartak Fracture Zone (AFFZ) and the Socotra Hadbeen Fracture Zone (SHFZ). Each of these sections of the margin corresponds to a first-order ridge segment. This study of this first-order segment of the rifted margin is based on new detailed geological and geophysical observations and analysis, including seismic velocity models from wide-angle data, collocated with multichannel seismic (MCS) profiles and high-resolution heat flow measurements. Because of its recent formation and the absence of salt deposits, the Gulf of Aden provides ideal conditions to study the processes of rifting and continental breakup with seismic imaging techniques. This part of the margin reveals a complexity unexpected at this scale. Our analysis reveals significant change in the width and character of the OCT over a lateral dis-

tance of only 10 km. From 15 km wide with postbreakup magmatism, the OCT broadens to 50 km with low-angle detachment faults and little or no magmatism.

2. Tectonic Setting

[4] The Gulf of Aden is an active oblique rift system with young margins [d'Acremont *et al.*, 2005] along the Arabia-Somalia plate boundary connects to the Indian Ocean to the east and to the Red Sea to the west (Figure 1). Extension began at 35 Ma [Roger *et al.*, 1989; Watchorn *et al.*, 1998] following the closure of the Neo-Tethys Ocean and the development of the Afar plume [Bellahsen *et al.*, 2003]. Earliest rift structures strike N110°E, ca. 40° from the strike of late synrift structures and from the strike of the Gulf [d'Acremont *et al.*, 2005; Bellahsen *et al.*, 2006] (Figure 1). Seafloor spreading followed at 17.6 Ma in the eastern part of the Gulf [Leroy *et al.*, 2004; d'Acremont *et al.*, 2006] as far west as the Shukra-El Sheik transform fault (SSTf) [Leroy *et al.*, 2010] (Figure 1). Finally, the rift propagated toward the Afar plume [Audin *et al.*, 2001]. The along-axis segmentation, controlled by continental margin structure during the early stage of rifting, was replaced during the late rifting stage by a new geometry of shorter axial segments [d'Acremont *et al.*, 2010]. This late stage segmentation persisted through the onset of seafloor spreading and continues to evolve along the present-day spreading ridge [Leroy *et al.*, 2004; d'Acremont *et al.*, 2006, 2010]. The high degree of margin segmentation is attributable to oblique rifting. As a consequence of margin segmentation, the nascent OCT of the study area is located along the eastern side of the Mesozoic Jeza-Qamar basin, which was reactivated during the opening of the Gulf of Aden (Figure 1 [Bellahsen *et al.*, 2006; d'Acremont *et al.*, 2006]).

3. Data Acquisition

[5] A combined onshore-offshore survey was carried out in 2006 along the southern part of Oman in order to image the deep structures of the eastern Gulf of Aden margin (Figure 1). We have collected a large amount of geological and geophysical data in the study area, including multibeam bathymetry, gravity, magnetism and heat flow in addition to the multichannel and wide-angle seismic studies. Thirty-five Ocean Bottom Seismometers (OBS) deployed at 10–15 km intervals and 19 broadband seismometers onshore in Oman recorded the air gun shots




Figure 1. Map of the Encens experiment in the northeastern Gulf of Aden. (top) Gray shading denotes the Afar-related volcanism. SSTf, Shukra el Sheik transform fault; AFTf, Alula Fartak transform fault; Aden R, Aden ridge; ShR, Sheba ridge; Jeza Qamar b., inherited Jeza Qamar basin. The red line indicates the active oceanic ridge, and blue indicates the spreading axis at Chron 5c (16 Ma) and in the east of AFFZ; the dark blue line indicates Chron 5d (17.6 Ma) [Leroy *et al.*, 2010]. The black arrows show relative plate motions. (bottom) Shaded bathymetric map from multibeam soundings recorded during the Encens cruise. The Ashawq-Salalah, Taqah, and Mirbat segments are separated by dashed lines (accommodation zones). Purple dots and red triangles are seismic instrument locations of the overall experiment, and the three yellow and black lines are locations of the three seismic transects presented here. Salmon-colored shading indicates the extent of the ocean-continent transition (OCT) data coverage. AFFZ, Alula Fartak fracture zone; SHFZ, Socotra Hadbeen fracture zone.

up to 150 km inland (Figure 1). Source and receiver placement allowed for acquisition of excellent quality data along the margin between the AFFZ and SHFZ. We focus in this paper on one first-order segment of the margin.

4. Margin Structure

[6] Our results reveal fundamental differences in the crust and upper mantle structure of the second-order Ashawq-Salalah, Taqah, and Mirbat segments over short distance (Figure 1 and Table 1). We identified three second-order segments within the first-order AFFZ-SHFZ segment. The structure of these

second-order segments is interpreted in the following subsections from the P wave velocity, magnetic, gravity, MCS and heat flow data (Figures 2 to 4 and auxiliary material).¹ Based upon these interpretations, we define the location of the OCT as the domain in the distal margin where (1) the crust has neither geophysical nor geological characteristics of classical continental crust or of classical oceanic crust, (2) syn-OCT sediments are observed rather than typical synrift wedges, and (3) no oceanic magnetic anomalies are clearly identifiable. The tran-

¹Auxiliary materials are available in the HTML. doi:10.1029/2009GC002963.




Figure 2

sition from continental domain also corresponds to an increase in the seismic velocities from 6 to 7.6 km s⁻¹ for the lower crust, in the surface heat flow from 40 to 120 mWm⁻², and in the free-air gravity anomaly from -40 to 40 mGals (Figures 2–4 and auxiliary material). The OCT, which includes a zone of exhumed continental mantle (ZECM [Péron-Pinvidic and Manatschal, 2009]) is considered as the locus of change in crustal character between classical continental and oceanic. The formation of the OCT occurred when the continental crust broke up leaving behind no lower and/or upper continental crust. It extends from this point to the location where oceanic crust formed at the onset of seafloor spreading.

4.1. Western Segment: Ashawq-Salalah

[7] The Ashawq-Salalah subsegment has a length of 94 km (along margin, Figure 1). The OCT in this second-order segment is very narrow (15 km across margin); its 13 km thick crust is composed of two main parts separated by an internal reflector at ~7.5 km depth (Figure 2 and auxiliary material). The upper part is 5.5 km thick (P wave velocity of 4 to 7.2 km s⁻¹) and the lower, 7.5 km thick, has high P wave velocity ranging from 7.6 to 7.8 km s⁻¹. The mantle beneath the crust of the OCT and oceanic crust has slow P wave velocity (7.8 to 7.9 km s⁻¹; Figure 2). The oceanic crust has a total thickness of 9 km, with an internal reflector at 4 km. Although a clearly identifiable oceanic magnetic anomaly exists to the south of the OCT (the 5d anomaly [d'Acremont et al., 2006, 2010]), the oceanic crust does not have a typical low-velocity gradient (Figure 2 and auxiliary material). There is evidence of volcanic activity after continental

breakup. A volcano has been identified on the seafloor within the OCT based on morphology, thick seismic reflection patterns typical of volcanic flows [Autin et al., 2010] with the associated crustal velocities ranging from 5 to 6 km s⁻¹, and by an extremely high heat flow (~900 mWm⁻²). These observations suggest the persistence of activity until recent times (~100,000 yr) [Lucazeau et al., 2009] (Figure 2). Dipping thick deep reflectors are also observed in the oceanic crust (Figure 2) and have been related to magmatism [Autin et al., 2010]. Deformation of this margin segment is characterized in the north by large-offset normal faults dipping seaward that define two rifted basins and in the south by closely spaced normal faults (5 km) that thinned the continental crust during late stage rifting (Figures 2 and 5) [Autin et al., 2010].

4.2. Central Segment: Taqah

[8] In the 36 km long Taqah subsegment, the characteristic of the margin changes significantly: there is no evidence of postbreakup extrusive magmatism, the OCT is wider (35 km) with thinner crust (8 km) where the internal reflector observed in the other two segments is not present (Figures 1 and 3). Lower crustal velocities contrast with the high velocities observed in the Ashawq-Salalah adjoining segment. A mussel-shaped topographic high emerges from the morphology in the OCT (Figure 1). P wave velocities increase within this high with a continuous gradient from 5 km s⁻¹ at the seafloor to 7.9 km s⁻¹ at the base of the crust, a characteristic typical of serpentinized rocks [Christensen, 1972; Dean et al., 2000; Minshull, 2009]. Similar features at other margins have been related to

Figure 2. The Ashawq-Salalah segment of the Encens experiment. (top) Model of seismic velocity structure. Velocity contours are color coded and labeled in units of km s⁻¹. Red circles are the instrument locations. OCT indicates the interpreted ocean-continent transition located in between the vertical dashed lines. Models are determined using a combined forward/inverse travelttime modeling approach (see auxiliary material). Examples of data are presented in the auxiliary material (Chi² = 1.358; Nrays = 5101; thick solid lines are the reflection points). The crustal seismic models are shown with marine magnetic anomalies (blue line) from shipboard data with interpreted chrons and the free-air gravity anomaly (red line), both acquired jointly during the experiment. (middle) Deep seismic reflection and heat flow profiles and interpretation. Red arrows show the locations of the heat flow measurements in the seismic profile. The black circles are uncorrected heat flow, and red circles are corrected for sedimentation, topography, and refraction. Error bars represent a 68% confidence interval. The seismic profile is presented with interpretation of faults (faults active during early synrift are shown in red, faults active during late synrift are shown in pink [Autin et al., 2010], and faults associated with active landslides are shown in purple) and of the main sedimentary units. Blue arrow indicates the location of the major fault imaged and reported on the model in Figure 2 (top). The colored dotted lines are derived from the velocity models converted in two-way travelttime seconds corresponding to main interfaces below the acoustic basement. (bottom) Schematic representation of the segment reconstructed with the southern conjugate margin [d'Acremont et al., 2006], showing the margin style prior to seafloor spreading along the Ashawq-Salalah profile near the Alula Fartak Fracture zone (AFFZ). The 3-D block diagram shows the location of the AFFZ in relation to the segment and the distribution of the faults close to the cross section. Sedimentary series are not shown.

exhumed continental mantle that was serpentinized to within a few km of the surface rather than continental or basaltic crust [Boillot *et al.*, 1989; Brun and Beslier, 1996; Whitmarsh *et al.*, 1998; Dean *et al.*, 2000; Lavier and Manatschal, 2006; Minshull, 2009]. The mantle here also has low P wave velocities (7.8 to 7.9 km s⁻¹; Figure 3) as observed in margins in Newfoundland [Reid, 1994], Nova Scotia [Funck *et al.*, 2004] and Goban spur [Bullock and Minshull, 2005]. In addition, in the northern part of the mussel-shaped high, low-amplitude magnetic anomalies are present, which is also a characteristic of serpentinized rocks. These anomalies are interpreted as Chron 6 (20 Ma) in the northern part of the OCT domain (Figures 1 and 3). The oceanic domain displays typical crustal thickness (7.5 km) for an oceanic crust aged 18 Ma (Chron 5d), a flat Moho, a very steep P wave velocity gradient, high heat flow values and identifiable oceanic magnetic anomalies (Chron 5d; Figures 1 and 3). As in the previous subsegment, normal faults dipping seaward are closely spaced near the OCT and within the OCT domain. The mussel-shaped bathymetric high has been recently faulted as evidenced by many submarine landslides observed on the seafloor morphology (Figure 3). Other sedimentary instabilities are located in the continental domain (Figures 1 and 3 and auxiliary material), revealing active uplift of this margin segment.

4.3. Eastern Segment: Mirbat

[9] In the 65 km long Mirbat subsegment, the characteristics of the margin is again different from the two others. The OCT is wider (50 km) and its crustal thickness decreases from more than 8 km near the continental domain to 5.5 km near the oceanic domain with notable necking (Figure 4). A spur-shaped basement structure is observed in the OCT at its boundary with continental domain (Figure 1) where P wave velocities in the crust increase from 6 to 7.3 km s⁻¹ and the surface heat flow increases abruptly to a value of 120 mWm⁻², close to that expected for seafloor (Figure 4) [Lucazeau *et al.*, 2008]. As in the Taqah subsegment, there is no evidence of volcanism at the surface or at depth and the mantle here also has low P wave velocities (7.8 to 7.9 km s⁻¹; Figure 4). The continuous gradient of P wave velocity in the OCT from 5 to 7.8 km s⁻¹ from the surface to the base of the crust, and especially for the spur-shaped basement high, as well as low-amplitude magnetic anomaly (possibly Chron 6) (Figures 1 and 4) suggest the presence of serpentinized continental

mantle which may have been exhumed by one or several faults [Dean *et al.*, 2000; Funck *et al.*, 2003; Bullock and Minshull, 2005; Minshull, 2009]. Further south in the OCT domain, a buried isolated basement structure displays velocities that could be attributable to the presence of continental crust material (between 5 and 5.4 km s⁻¹ abscissa 40–55 km; Figure 4). The deformation of this segment is characterized (1) by two major seaward normal faults that bound blocks of thinned continental crust (one is imaged by the seismic profile and the other one crops out onshore; Figure 5), (2) possibly by exhumed serpentinized mantle, and (3) possibly by an extensional allochthon of continental crust (Figure 4). In the continental slope, active sedimentary instabilities also reveal persistent uplift of the margin during postrift (Figures 1 and 4 and auxiliary material).

5. Interpretations and Discussion

[10] Our primary observations are variations in margins geometry, OCT width and magmatism over small spatial scales. We describe a wide OCT for the magma-poor rifted margin segment (two eastern segments) and a narrow OCT associated with postbreakup magma-rich segment (western segment). Why do such extreme differences exist between these three segments of the eastern Gulf of Aden margin? We consider possible interpretations based on the P wave velocities, MCS, gravity, heat flow and magnetism such as postbreakup volcanism, exhumation of subcontinental mantle and widening of the OCT in relation to the increase of average rate of extension.

5.1. Postbreakup and Postrift Volcanism

[11] Based on the presence of internal reflectors and unusually high P wave velocities at large depth, we propose that the high-velocity body existing exclusively at the base of the crust in the Ashawq-Salalah segment is related to an underplated magmatic body. Its origin may not be related to the other volcanic activity along the margin. Its seismic velocities are much higher than those reported for underplated material along the North Atlantic margins [Raum *et al.*, 2002]. Also, the volume of magmatism is smaller in the eastern Gulf of Aden than for other volcanic margins where underplating has been interpreted. Moreover, in a magma-poor margin setting where no synrift volcanism has been documented [d'Acremont *et al.*, 2005; Autin *et al.*, 2010], it is unlikely that igneous underplated body




Figure 3

has an origin similar to that of the volcanic rifted margins [e.g., *Hirsch et al.*, 2009]. In Atlantic margins, high-velocity bodies are usually associated with flood basalt volcanism [*Meyer et al.*, 2007]. Here, at the top of the crust, volcanism is observed in the OCT and occurs soon after the breakup, as the OCT formed, and persists during the onset of seafloor spreading [*Lucazeau et al.*, 2009; *Autin et al.*, 2010; *Leroy et al.*, 2010]. If the origin of this high-velocity body in the lower crust corresponds to magmatic sills accreted after the breakup and/or early during the seafloor spreading phase, we would expect dikes to appear within the crust. We do not observe any dikes, but they are probably below the resolution of our seismic profiles. Volcanism is observed within the OCT of the longest margin segment and close to the long-offset AFFZ. At the ridge segment that corresponds to this section of the OCT, oceanic crust production is greater than the global average [*d'Acremont et al.*, 2010]. The origin of this postrift volcanism within the OCT may be related either to crustal thickening and magma production at the edge of the long-offset transform [*Hooff et al.*, 2000; *Gregg et al.*, 2007] as proposed by *d'Acremont et al.* [2010] for the 180 km long AFFZ offset, which has been active at least since the time of OCT formation [*d'Acremont et al.*, 2010]. Alternatively, the volcanism and associated underplated body may be related to the anomalous presence of melt beneath the Gulf of Aden fed by channeled flow of plume material from the Afar hot spot along the nascent ridge in a context of plume-ridge interaction as suggested by *Leroy et al.* [2010]. In other segmented margins with long-offset fracture zones, the oceanic transforms may act as long-lived magmatic pathways for melt in the lithosphere [*Gernigon et al.*, 2009]. Thus, the relatively sharp along-strike change in the OCT structure between the Ashawq-Salalah segment and the two others that are not associated large-offset transform faults may be related to the postbreakup volcanism that controlled the geometry of the OCT and that of the nascent spreading ridge. Volcanism is indeed considered to be capable of reducing fault spacing

and heave and as a result decreasing the width of the axial valley [*Behn and Ito*, 2008].

5.2. Exhumation of Subcontinental Mantle

[12] Based on the observations of a continuous gradient of P wave velocity from surface to depth on the two basement highs (mussel-shaped and spur-shaped) and the presence of low-amplitude magnetic anomaly in the OCT of the Taqah and Mirbat segments, we propose that continental crust breakup first began by the exhumation of the subcontinental mantle by faults that flatten near the seafloor [*Brun and Beslier*, 1996; *Lavier and Manatschal*, 2006] (Figure 3–5). In the Ashawq-Salalah segment, exhumation of the continental mantle may also have occurred [*Autin et al.*, 2010], but evidence for this may have been overprinted by the postrift volcanic activity (Figures 2 and 5).

[13] Examples of fault systems that exhume continental mantle have been imaged by seismic studies [*Dean et al.*, 2000] and drilled [*Boillot et al.*, 1988; *Whitmarsh et al.*, 1998] in the Iberia abyssal plain or exposed in the Alps [*Lavier and Manatschal*, 2006] and Pyrenees [*Lagabrielle and Bodinier*, 2008]. In these locations, exhumed mantle in the OCT is associated with shallow crustal detachment systems, and is serpentinized at temperatures under 600°C at depths shallower than 10 km where hydrothermal circulation is supposed to have occurred [*Lavier and Manatschal*, 2006; *Cannat et al.*, 2009]. We believe that the very low-amplitude Chron 6 magnetic anomaly could be associated with serpentinization of the exhumed mantle and not formed as a typical oceanic magnetic anomaly would be. The contribution of peridotites to magnetic anomalies becomes significant only where they are affected by a high degree of serpentinization and associated presence of magnetite [*Oufi et al.*, 2002]. Here the low-amplitude anomalies, which are not centered over the basement highs, may be associated with low degrees of serpentinization. In the Ashawq-Salalah segment, the recent volcano lying on the OCT [*Autin et al.*, 2010] may have modified the magnetic signal. This may be why Chron 6 was not

Figure 3. The Taqah segment of the Encens study. (top) Model of seismic velocity structure ($\text{Chi}2 = 1.919$; $N_{\text{rays}} = 2689$; thick solid lines are the reflection points), marine magnetic anomalies (blue line) from shipboard data with interpreted chrons, and the free-air gravity anomaly (red line). (middle) Deep seismic reflection and joint heat flow profiles and interpretation. The seismic profile is presented with interpretation of faults (faults active during early syn-rift are shown in red, faults active during exhumation phase are shown in yellow, faults active during postrift are shown in orange, and faults associated with active submarine landslides are shown in purple). (bottom) Schematic 3-D representation of the spatial evolution of the segment reconstructed with the southern conjugate margin.




Figure 4




Figure 5. Modes of extension leading to continental breakup. Conceptual crustal sections without vertical exaggeration of the three different segments based on observations and data interpretation. The thickness of the crust in the proximal margin is from receiver function study [Tiberi et al., 2007]. The crust is thinned by seaward dipping normal faults that slipped during different stages of rifting. The earliest faults form the main grabens, and the latest localize deformation at the axis of paleorifts or are responsible for exhumation of the subcontinental mantle.

recorded in the western segment. The P wave velocity versus serpentinization rate [Horen et al., 1996] however suggests 100% and 60% of serpentinization rate for mussel-shaped and spur-shaped highs, respectively. These basement highs are controlled by recently active faults that reach the surface and may be acting as recharge zones for fluid circulation within the crust. This notion is supported by the decreased heat flow over these two highs (Figure 3). The recently active faults uplifted the ridges of serpentinized peridotites to their present-day positions. Mantle with a velocity of 7.8 km/s underlies both the thinned continental and oceanic crust. A gradual downward increase to normal mantle velocities has been interpreted to reflect decreasing degree of serpentinization with depth [Funck et al., 2003]. Normal mantle velocities of 8.0 km/s are observed ~6 km below the base of the

crust as it is the case in several others margins such as Newfoundland [Reid, 1994].

5.3. Widening of the OCT

[14] We propose that the widening of the OCT toward the east between AFFZ and SHFZ is related to the oblique opening of the Gulf of Aden and could be explained either by an eastward increase in the amount of extension or by a westward propagating rift system. In the absence of well data to constrain ages, we rely on magnetic chron patterns and analogue modeling (J. Autin et al., Analogue models of oblique rifting in a cold lithosphere, submitted to *Tectonics*, 2010). In the highly segmented margins, distal grabens present a morphology related to an opening system, with deepening toward the wider eastern side caused by rotation of

Figure 4. The Mirbat segment of the Encens experiment. (top) Model of seismic velocity structure ($\chi^2 = 2.407$; $N_{rays} = 3546$; thick solid lines are the reflection points), marine magnetic anomalies (blue line) from shipboard data with interpreted chrons, and the free-air gravity anomaly (red line). (middle) Deep seismic reflection and joint heat flow profiles and interpretation. The seismic profile is presented with interpretation of faults (faults active during early synrift are shown in red, faults active during exhumation phase are shown in yellow, faults active during postrift are shown in orange, and faults associated with active submarine landslides are shown in purple). (bottom) Schematic 3-D representation of the spatial evolution of the segment reconstructed with the southern conjugate margin.

Table 1. Summary of Across-Margin Characteristics of Segments on the Profiles

	Ashawq-Salalah	Taqah	Mirbat
Segment length (km)	94	36	65
OCT width (km)	15	35	50
OCT crust thickness (km)	13	8	7–8
Oceanic crust thickness (km)	9	7.5	5.5

tilted blocks at the end of rifting [Autin, 2008]. Evidence for such block rotations in the young obliquely opening Main Ethiopian Rift can be seen in paleomagnetic analysis [Kidane *et al.*, 2006]. In a V-shaped basin such as the Woodlark basin or West Iberia margin for instance, widening of the rift is often related to the propagation of the rift system [Taylor *et al.*, 1999; Péron-Pinvidic and Manatschal, 2009]. In such propagating systems, different phases of extension must be occurring coincidentally [Lavie and Manatschal, 2006]. In our study area, the grabens deepen and widen toward the east and no propagation is observed (Figure 1). We, therefore, prefer a rifting model without propagation in an oblique rifting context added to the effects of the volcanism in the western segment of the OCT. We believe that the OCT formed with a higher extension rate where there is exhumed continental mantle and no magmatism. Where the OCT is wide, allochthonous relics of continental crust lying on the exhumed mantle are observed (Figure 4) as in the West Iberia margin [Péron-Pinvidic and Manatschal, 2009].

6. Concluding Remarks

[15] The rifted margin sectors between large offset transforms in the eastern Gulf of Aden reveal fundamental along-strike variations in distribution and in style of lithospheric strain over distances < 50 km. Our primary observations indicate the presence of a narrow segment of OCT within an otherwise wide magma-poor OCT rift sector that may have been produced by enhanced melting or by a supply of Afar plume material channeled by the ridge to the large-offset transform fault zone that bounds the segment. We believe the two rifting mechanisms led to the observed structure of the margin and present-day seafloor spreading system: oblique rifting near evolving large-offset transform faults, and lithospheric weakening by magma and localized mantle exhumation associated with margin segmentation. Studies of other passive margins with along-strike changes show similarities with our interpretations although pub-

lished data sets from these other locations cannot constrain the lateral evolution over such a short distance (Figure 4, middle). Studies in the Gulf of California and northwestern Australia show that synrift and early postrift magmatism varies between rift segments [Hopper *et al.*, 1992; Lizarralde *et al.*, 2007], but the short lateral changes, which may be related to the geological and tectonic setting, are unknown. The results presented in the Gulf of California highlight the importance of inherited mantle fertility/hydration resulting from the presence of prerift ignimbrites of the former arc system [Lizarralde *et al.*, 2007]. The structural inheritance of the Gulf of Aden consists of previous episodes of rifting with some associated rifted basins [d'Acremont *et al.*, 2005]. In the case of the Gulf of Aden, the proximity of the Afar plume may have had an influence on the rift system since its inception [Bellahsen *et al.*, 2003]. Therefore, unlike the interpretation proposed for the Gulf of California [Lizarralde *et al.*, 2007], the along-axis variations in the Gulf of Aden margin do not seem to be related only to mantle depletion versus mantle fertility. Instead, we believe that the genetic link between the widening of the OCT toward the east, the exhumation of subcontinental mantle, and the postrift volcanism is due to the obliquity of the rifting as well as in the influence of the Afar plume. Plume-ridge interaction may have affected the rift system and its margins as early as immediately after OCT emplacement [Leroy *et al.*, 2010].

[16] Similar processes have affected other margins either in a context of oblique opening with strike-slip components as has been proposed for the Woodlark basin [Taylor *et al.*, 1999] or for some segments of Atlantic margins where influence of plume material during the OCT formation may have occurred [e.g., Moulin *et al.*, 2005]. Thus it is important to consider these profound along-axis variations in lithospheric strain accommodation and OCT structure, which can affect sedimentary and thermal structure of continental margins as well as evolution of petroleum systems located there. Our observations and interpretations demonstrate the need for a fully 3-D approach to rifted margin studies.

Acknowledgments

[17] We thank the captains and crews of the R/V *L'Atalante* (IFREMER-GENAVIR) and *Dhelkoot*; the OBS teams; Y. Hello, A. Anglade, O. Desprez, and P. Charvis of IRD-Geosciences Azur; L. Beguery, O. Aouji, R. Daniel, and S. Singh of INSU-IPGP for technical support; Hilal Al-Azri, Salim Al

Busaidi, Harub Al Hashmi, Sami Zubedi, Said Abahalaf, Sheikh A. Haroon, and C. Deplus for logistical support; and GDR Marges, Actions Marges, CNRS-INSU, TOTAL, NERC grant NE/C514031/1, UMR7072, ANR YOCMAL, and MOCI for financial support.

References

- Audin, L., I. Manighetti, P. Tapponnier, F. Metivier, E. Jacques, and P. Huchon (2001), Fault propagation and climatic control of sedimentation on the Ghoubbet Rift floor: Insights from the Tadjouraden cruise in the western Gulf of Aden, *Geophys. J. Int.*, *144*, 391–413, doi:10.1046/j.0956-540x.2000.01322.x.
- Autin, J. (2008), Déchirure continentale et segmentation du golfe d'Aden oriental en contexte de rifting oblique, Ph.D. thesis, 302 pp., Univ. Paris 06, Paris.
- Autin, J., S. Leroy, M. O. Beslier, E. d'Acremont, P. Razin, A. Ribodetti, N. Bellahsen, C. Robin, and K. Al-Toubi (2010), Continental break-up history of a deep magma-poor margin from seismic reflection data (northeastern Gulf of Aden margin, offshore Oman), *Geophys. J. Int.*, *180*, 501–519, doi:10.1111/j.1365-246X.2009.04424.x.
- Behn, M. D., and G. Ito (2008), Magmatic and tectonic extension at mid-ocean ridges: 1. Controls on fault characteristics, *Geochem. Geophys. Geosyst.*, *9*, Q08O10, doi:10.1029/2008GC001965.
- Bellahsen, N., C. Faccenna, F. Funiciello, J.-M. Daniel, and L. Jolivet (2003), Why did Arabia separate from Africa? Insights from 3-D laboratory experiments, *Earth Planet. Sci. Lett.*, *216*, 365–381, doi:10.1016/S0012-821X(03)00516-8.
- Bellahsen, N., M. Fournier, E. d'Acremont, S. Leroy, and J.-M. Daniel (2006), Fault reactivation and rift localization: Northeastern Gulf of Aden margin, *Tectonics*, *25*, TC1007, doi:10.1029/2004TC001626.
- Boillot, G., J. Girardeau, and J. Kornprobst (1988), Rifting of the Galicia margin: Crustal thinning and emplacement of mantle rocks on the seafloor, *Proc. Ocean Drill. Program Sci. Results*, *103*, 741–756.
- Boillot, G., G. Féraud, M. Recq, and J. Girardeau (1989), Undercrusting by serpentinite beneath rifted margins, *Nature*, *341*, 523–525, doi:10.1038/341523a0.
- Brun, J.-P. (1999), Narrow rifts versus wide rifts: Inferences for the mechanics of rifting from laboratory experiments, *Philos. Trans. R. Soc. London, Ser. A*, *357*, 695–712, doi:10.1098/rsta.1999.0349.
- Brun, J.-P., and M.-O. Beslier (1996), Mantle exhumation at passive margins, *Earth Planet. Sci. Lett.*, *142*, 161–173, doi:10.1016/0012-821X(96)00080-5.
- Buck, W. R. (2004), Consequences of asthenospheric variability on continental rifting, in *Rheology and Deformation of the Lithosphere at Continental Margins*, edited by G. Karner et al., pp. 1–30, Columbia Univ. Press, New York.
- Buck, W. R., L. L. Lavier, and A. N. B. Poliakov (1999), How to make a rift wide, *Philos. Trans. R. Soc. London, Ser. A*, *357*, 671–693, doi:10.1098/rsta.1999.0348.
- Bullock, A. D., and T. A. Minshull (2005), From continental extension to seafloor spreading: Crustal structure of the Goban Spur rifted margin, southwest of the UK, *Geophys. J. Int.*, *163*, 527–546, doi:10.1111/j.1365-246X.2005.02726.x.
- Cannat, M., G. Manatschal, D. Sauter, and G. Péron-Pinvidic (2009), Assessing the conditions of continental breakup at magma-poor rifted margins: What can we learn from slow spreading mid-ocean ridges?, *C. R. Geosci.*, *341*, 406–427, doi:10.1016/j.crte.2009.01.005.
- Christensen, N. I. (1972), The abundance of serpentinites in the oceanic crust, *J. Geol.*, *80*, 709–719, doi:10.1086/627796.
- Corti, G., V. Wijk, M. Bonini, D. Sokoutis, S. Cloetingh, F. Innocenti, and P. Manetti (2003), Transition from continental break-up to punctiform seafloor spreading: How fast, symmetric and magmatic, *Geophys. Res. Lett.*, *30*(12), 1604, doi:10.1029/2003GL017374.
- d'Acremont, E., S. Leroy, M. O. Beslier, N. Bellahsen, M. Fournier, C. Robin, M. Maia, and P. Gente (2005), Structure and evolution of the eastern Gulf of Aden conjugate margins from seismic reflection data, *Geophys. J. Int.*, *160*, 869–890, doi:10.1111/j.1365-246X.2005.02524.x.
- d'Acremont, E., S. Leroy, M. Maia, P. Patriat, M. O. Beslier, N. Bellahsen, M. Fournier, and P. Gente (2006), Structure and evolution of the eastern Gulf of Aden: Insights from magnetic and gravity data (Encens-Sheba/MD117 cruise), *Geophys. J. Int.*, *165*, 786–803, doi:10.1111/j.1365-246X.2006.02950.x.
- d'Acremont, E., S. Leroy, M. Maia, P. Gente, and J. Autin (2010), Volcanism, jump and propagation on the Sheba Ridge, eastern Gulf of Aden: Segmentation evolution and implications for accretion processes, *Geophys. J. Int.*, *180*, 535–551, doi:10.1111/j.1365-246X.2009.04448.x.
- Dean, S. M., T. A. Minshull, R. B. Whitmarsh, and K. E. Loudon (2000), Deep structure of the ocean-continent transition in the southern Iberia Abyssal Plain from seismic refraction profiles: The IAM-9 transect at 40°20'N, *J. Geophys. Res.*, *105*, 5859–5885, doi:10.1029/1999JB900301.
- Dunbar, J., and D. S. Sawyer (1989), How preexisting weaknesses control the style of continental breakup, *J. Geophys. Res.*, *94*, 7278–7292, doi:10.1029/JB094iB06p07278.
- Ebinger, C., B. Rosendahl, and D. Reynolds (1987), Tectonic model of the Malaŵi rift, Africa, *Tectonophysics*, *141*, 215–235, doi:10.1016/0040-1951(87)90187-9.
- Funck, T., J. R. Hopper, H. C. Larsen, K. E. Loudon, B. E. Tucholke, and W. S. Holbrook (2003), Crustal structure of the ocean-continent transition at Flemish Cap: Seismic refraction results, *J. Geophys. Res.*, *108*(B11), 2531, doi:10.1029/2003JB002434.
- Funck, T., H. R. Jackson, K. E. Loudon, S. A. Dehler, and Y. Wu (2004), Crustal structure of the northern Nova Scotia rifted continental margin (eastern Canada), *J. Geophys. Res.*, *109*, B09102, doi:10.1029/2004JB003008.
- Georgen, J. E., J. Lin, and H. J. B. Dick (2001), Evidence from gravity anomalies for interactions of the Marion and Bouvet hotspots with the southwest Indian ridge: Effects of transform offsets, *Earth Planet. Sci. Lett.*, *187*, 283–300, doi:10.1016/S0012-821X(01)00293-X.
- Gernigon, L., O. Olesen, J. Ebbing, S. Wienecke, C. Gaina, J. O. Mogaard, M. Sand, and R. Myklebust (2009), Geophysical insights and early spreading history in the vicinity of the Jan Mayen Fracture Zone, Norwegian-Greenland Sea, *Tectonophysics*, *468*, 185–205, doi:10.1016/j.tecto.2008.04.025.
- Gregg, P. M., J. Lin, M. D. Behn, and L. G. J. Montesi (2007), Spreading rate dependence of gravity anomalies along oceanic transform faults, *Nature*, *448*, 183–187, doi:10.1038/nature05962.
- Hirsch, K., K. Bauer, and M. Scheck-Wenderoth (2009), Deep structure of the western South African passive margin—Results of a combined approach of seismic, gravity and isostatic investigations, *Tectonophysics*, *470*, 57–70, doi:10.1016/j.tecto.2008.04.028.

- Hooft, E. E., R. S. Detrick, D. R. Toomey, J. A. Collins, and J. Lin (2000), Crustal thickness and structure along three contrasting spreading segments of the Mid-Atlantic Ridge, 33.5°–35°N, *J. Geophys. Res.*, *105*, 8205–8226, doi:10.1029/1999JB900442.
- Hopper, J. R., and W. R. Buck (1996), The effect of lower crustal flow on continental extension and passive margin formation, *J. Geophys. Res.*, *101*, 20,175–20,194, doi:10.1029/96JB01644.
- Hopper, J. R., et al. (1992), Magmatism and rift margin evolution: Evidence from northwest Australia, *Geology*, *20*, 853–857, doi:10.1130/0091-7613(1992)020<0853:MARMEE>2.3.CO;2.
- Horen, H., M. Zamora, and G. Dubuisson (1996), Seismic waves velocities and anisotropy in serpentinitized peridotites from xigaze ophiolite: Abundance of serpentinite in slow spreading ridge, *Geophys. Res. Lett.*, *23*, 9–12, doi:10.1029/95GL03594.
- Kidane, T., E. Platzman, C. J. Ebinger, B. Abebe, and P. Rochette (2006), Palaeomagnetic constraints on continental break-up processes: Observations from the Main Ethiopian Rift, in *The Afar Volcanic Province Within the East African Rift System*, *Geol. Soc. Spec. Publ.*, *259*, 165–183, doi:10.1144/GSL.SP.2006.259.01.14.
- Lagabrielle, Y., and J.-L. Bodinier (2008), Submarine reworking of exhumed subcontinental mantle rocks: Field evidence from the Lherz peridotites, French Pyrenees, *Terra Nova*, *20*, 11–21.
- Lavier, L. L., and G. Manatschal (2006), A mechanism to thin the continental lithosphere at magma-poor margins, *Nature*, *440*, 324–328, doi:10.1038/nature04608.
- Leroy, S., et al. (2004), From rifting to spreading in the eastern Gulf of Aden: A geophysical survey of a young oceanic basin from margin to margin, *Terra Nova*, *16*, 185–192, doi:10.1111/j.1365-3121.2004.00550.x.
- Leroy, S., E. d'Acremont, C. Tiberi, C. Basuyau, J. Autin, F. Lucazeau, and H. Sloan (2010), Recent off-axis volcanism in the eastern Gulf of Aden: Implications for plume-ridge interaction, *Earth Planet. Sci. Lett.*, *293*, 140–153, doi:10.1016/j.epsl.2010.02.036.
- Lizarralde, D., G. J. Axen, H. E. Brown, J. M. Fletcher, A. Gonzalez-Fernandez, A. J. Harding, W. S. Holbrook, G. M. Kent, P. Paramo, and F. Sutherland (2007), Variation in styles of rifting in the Gulf of California, *Nature*, *448*, 466–469, doi:10.1038/nature06035.
- Lucazeau, F., et al. (2008), Persistent thermal activity at the Eastern Gulf of Aden after continental break-up, *Nat. Geosci.*, *1*, 854–858, doi:10.1038/ngeo359.
- Lucazeau, F., S. Leroy, J. Autin, A. Bonneville, B. Goutorbe, F. Rolandone, E. d'Acremont, L. Watremez, D. Düsünür, and P. Huchon (2009), Post-Rift volcanism and high heat-flow at the ocean-continent transition of the eastern Gulf of Aden, *Terra Nova*, *21*, 285–292, doi:10.1111/j.1365-3121.2009.00883.x.
- Meyer, R., J. Van Wijk, and L. Gernigon (2007), The North Atlantic Igneous Province: A review of models for its formation, *Spec. Pap. Geol. Soc. Am.*, *430*, 525–552.
- Minshull, T. A. (2009), Geophysical characterisation of the ocean-continent transition at magma-poor rifted margins, *C. R. Geosci.*, *341*, 382–393, doi:10.1016/j.crte.2008.09.003.
- Moulin, M., D. Aslanian, J.-L. Olivet, I. Contrucci, L. Matias, L. Géli, F. Klingelhoefer, H. Nouzé, J.-P. Réhault, and P. Unternehr (2005), Geological constraints on the evolution of the Angolan margin based on reflection and refraction seismic data (ZaiAngo project), *Geophys. J. Int.*, *162*, 793–810, doi:10.1111/j.1365-246X.2005.02668.x.
- Oufi, O., M. Cannat, and H. Horen (2002), Magnetic properties of variably serpentinitized abyssal peridotites, *J. Geophys. Res.*, *107*(B5), 2095, doi:10.1029/2001JB000549.
- Péron-Pinvidic, G., and G. Manatschal (2009), The final rifting evolution at deep magma-poor passive margins from Iberia-Newfoundland: A new point of view, *Int. J. Earth Sci.*, *98*, 1581–1597, doi:10.1007/s00531-008-0337-9.
- Raum, T., R. Mjelde, P. Digranes, H. Shimamura, H. Shiobara, S. Kodaira, G. Haatvedt, N. Serenes, and T. Thorbjurnsen (2002), Crustal structure of the southern part of the Voring Basin, mid-Norway margin, from wide-angle seismic and gravity data, *Tectonophysics*, *355*, 99–126, doi:10.1016/S0040-1951(02)00136-1.
- Reid, I. D. (1994), Crustal structure of a nonvolcanic rifted margin east of Newfoundland, *J. Geophys. Res.*, *99*, 15,161–15,180.
- Roger, J., J. P. Platel, C. Cavelier, and C. Bourdillon-de-Grisac (1989), Données nouvelles sur la stratigraphie et l'histoire géologique du Dhofar (Sultanat d'Oman), *Bull. Soc. Geol. Fr.*, *2*, 265–277.
- Taylor, B., A. Goodliffe, and F. Martinez (1999), How continents break up: Insights for Papua New Guinea, *J. Geophys. Res.*, *104*, 7497–7512, doi:10.1029/1998JB900115.
- Taylor, B., A. Goodliffe, and F. Martinez (2009), Initiation of transform faults at rifted continental margins, *C. R. Geosci.*, *341*, 428–438, doi:10.1016/j.crte.2008.08.010.
- Tiberi, C., S. Leroy, E. d'Acremont, N. Bellahsen, C. Ebinger, A. Al Lazki, and A. Pointu (2007), Crustal geometry of the northeastern Gulf of Aden passive margin; Localization of the deformation inferred from receiver function analysis, *Geophys. J. Int.*, *168*, 1247–1260, doi:10.1111/j.1365-246X.2006.03294.x.
- Watchorn, F., G. J. Nichols, and D. W. J. Bosence (1998), Rift-related sedimentation and stratigraphy, southern Yemen (Gulf of Aden), in *Sedimentation and Tectonics in the Rift Basins: Red Sea-Gulf of Aden*, edited by B. Purser and D. Bosence, pp. 165–189, Chapman and Hall, London.
- White, R. S. (1992), Magmatism during and after continental break-up, *Geol. Soc. London Spec. Pap.*, *68*, 1–16, doi:10.1144/GSL.SP.1992.068.01.01.
- Whitmarsh, R. B., et al. (1998), *Proceedings of the Ocean Drilling Program, Initial Reports*, vol. 173, Ocean Drill. Program, College Station, Tex.
- Yamasaki, T., and L. Gernigon (2009), Styles of lithospheric extension controlled by underplated mafic bodies, *Tectonophysics*, *468*, 169–184, doi:10.1016/j.tecto.2008.04.024.