



Assessment of seven culture media for the growth and isolation of *Capnocytophaga* spp.

Elodie Ehrmann, Martine Bonnaure-Mallet, Thierry Fosse

► To cite this version:

Elodie Ehrmann, Martine Bonnaure-Mallet, Thierry Fosse. Assessment of seven culture media for the growth and isolation of *Capnocytophaga* spp.. Bulletin- Groupement International pour la Recherche Scientifique en Stomatologie et Odontologie, Groupement International pour la Recherche Stomatologie et Odontologie, 2011, 50 (2), pp.9-10. <inserm-00826672>

HAL Id: inserm-00826672

<http://www.hal.inserm.fr/inserm-00826672>

Submitted on 3 Jun 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SHORT COMMUNICATION

O-4. ASSESSMENT OF SEVEN CULTURE MEDIA FOR THE GROWTH AND ISOLATION OF *CAPNOCYTOPHAGA SPP*

E. Ehrmann^{1,2,3}, M. Bonnaure-Mallet³, T. Fosse².

¹ Faculté d'odontologie, Université de Nice Sophia-Antipolis, 24 avenue des Diablos Bleus, 06357 Nice, France ² Laboratoire de bactériologie et d'épidémiologie moléculaire, Centre Hospitalier Universitaire de Nice, 151 Route de St Antoine de Ginestière 06200 Nice, France. ³ Equipe de Microbiologie, EA 1254, Université de Rennes 1, Rennes, France.

Introduction

The genus *Capnocytophaga* comprises a group of capnophilic/facultatively anaerobic Gram-negative bacilli and consists of eight species, namely *C. ochracea*, *C. sputigena*, *C. gingivalis*, *C. granulosa*, *C. leadbetteri*, *C. haemolytica*, *C. canimorsus* and *C. cynodegmi* 1,2. The ecologic niche for the first six species is the human oral cavity. They are associated with varying degrees of periodontal health and disease and are often reported to cause bacteremia and various severe general infections in compromised as well as in noncompromised patients 3,4. Both last ones species are a member of the oral cavity of dogs and cats and are associated most commonly with dog-bite infections. Data regarding the isolation frequency of *Capnocytophaga* spp. from clinical samples are conflicting, primarily due to the different microbial procedure employed, but also due to the difficult growth of this bacterial genus flooded in the wealth of the oral microbiome⁵. Isolation of these organisms is important for proper diagnosis and treatment of the systemic infections which they cause but also for its epidemiological study⁶. The aim of this investigation was to perform a quantitative and qualitative comparison of several media for the culture of the different *Capnocytophaga* species.

Materials and Methods

Ten clinical *Capnocytophaga* spp. isolated from oral human cavity and five ATCC reference strains were included. All clinical strains were identified by 16s RNA genes se-

quencing. Five selective culture media: VCAT (vancomycin, colistin, amphotericin B and colistin), bacitracin, Fuso, VK (vancomycin and kanamycin), ANC (nalidixic acid and colistin) were compared with two nonselective media: a blood agar (SBA) and a chocolate agar. The numbers of CFU/mL for each strain on the different media were calculated from the average of two replicate plates. MICs of antibiotics incorporated in various selective media (vancomycin, colistin, trimethoprim, bacitracin, gentamycin and nalidixic acid) were determined by Etest or diffusion method.

Results

Nonselective medium and Fuso medium were equivalent and able of supporting growth of all the different species of *Capnocytophaga*. They also yielded the maximum number of CFU/mL for all type strains even if *C. canimorsus* preferred Fuso medium and SBA than chocolate agar. Otherwise, ANC, VK, VCAT, and bacitracin media were able of supporting the growth of respectively 90%, 70%, 50%, and 40% of tested strains. All strains were highly resistant to trimethoprim, colistin and gentamycin while they were variably resistant or sensitive to bacitracin, vancomycin and nalidixic acid.

Discussion

Selective media used in this study like Mashimo⁷ and Cap medium⁸ were formulated to select for *Capnocytophaga* spp. from polymicrobial clinical specimens. But we show that susceptibility of certain strains to present anti-

biotics in media prevents the bacterial growth. Nonselective media is successful for growth of all species but often can't show Capnocytophaga strains from a dental plaque sample containing numerous other fast growth microorganisms.

Conclusions

The culture medium is a determining factor in isolating the full range of Capnocytophaga spp. from clinical isolates. It would be necessary to formulate a new medium considering these observations.

Acknowledgment:

This work was supported by IFRO.

References

1. Leadbetter ER, et al. (1979) Arch Microbiol 122 : 9-33.
2. Frandsen EV et al (2008) Int J Syst Evol Microbiol 58 :324-36.
3. Wang HK et al (2007) Journal of clinical Microbiology 45: 645-7
4. Martino R. et al Clin Infect Dis. (2001) Aug 15;33(4):E20-2.
5. Ciantar M. et al (2001) Clinical Microbiology and infectious diseases 7 : 158-60
6. Rummens JL. et al (1986) Antimicrob Agents Chemother 30 :739-42
7. Mashimo PA. et al (1983) Journal of Clinical Microbiology 17 :187-91
8. Rummens JL. et al (1985) Journal of Clinical Microbiology 22 :375-8