



# Galilean Covariance versus Gauge Invariance

Germain Rousseaux

► **To cite this version:**

Germain Rousseaux. Galilean Covariance versus Gauge Invariance. 4 pages. 2009. <hal-00440826>

**HAL Id: hal-00440826**

**<https://hal.archives-ouvertes.fr/hal-00440826>**

Submitted on 11 Dec 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Galilean Covariance versus Gauge Invariance

Germain Rousseaux

*Université de Nice-Sophia Antipolis, Laboratoire J.-A. Dieudonné,  
UMR CNRS-UNS 6621, Parc Valrose, 06108 Nice Cedex 02, France, European Union.*

(Dated: December 11, 2009)

We demonstrate for the first time and unexpectedly that the Principle of Relativity dictates the choice of the "gauge conditions" in the canonical example of a Gauge Theory namely Classical Electromagnetism. All the known "gauge conditions" of the literature are interpreted physically as electromagnetic continuity equations hence the "gauge fields". The existence of a Galilean Electromagnetism with TWO dual limits ("electric" and "magnetic") is the crux of the problem [1]. A phase-space with the domains of validity of the various "gauge conditions" is provided and is shown to depend on three characteristic times : the magnetic diffusion time, the charge relaxation time and the transit time of electromagnetic waves in a continuous medium [2].

The Standard Model of Physics is based on the assumed existence of a superior principle called Gauge Symmetry which would rule all the laws of Physics: *Physical theories of fundamental significance tend to be gauge theories. These are theories in which the physical system being dealt with is described by more variables than there are physically independent degree of freedom. The physically meaningful degrees of freedom then reemerge as being those invariant under a transformation connecting the variables (gauge transformation). Thus, one introduces extra variables to make the description more transparent and brings in at the same time a gauge symmetry to extract the physically relevant content. It is a remarkable occurrence that the road to progress has invariably been towards enlarging the number of variables and introducing a more powerful symmetry rather than conversely aiming at reducing the number of variables and eliminating the symmetry* [3]. Wolfgang Pauli was used to ask at the end of tiresome seminars he attended loosely if the principal result presented by the speaker was "gauge invariant" [4]. Hence, the concept of Gauge Theory has emerged progressively in Physics such that the equations feature variables ("gauge fields") which are underdetermined and in order to remove this degree of liberty ("gauge transformations") a closure assumption ("gauge condition") is formulated [5]. Similarly, the Principle of Relativity is known to be a robust safeguard when scaffolding a new theory since the proposed new laws must be covariant with respect to the transformations of space-time.

The goal of this paper is to remove the Gauge symmetry in the most famous example of a supposed Gauge Theory namely Classical Electromagnetism by revealing a conflict with another symmetry that is the Principle of Relativity. To do so, we first emphasize the Riemann-Lorenz approach to Electromagnetism. Therein the central role is played by the vector and scalar potentials  $\mathbf{A}$  and  $V$ , unlike the Heaviside-Hertz approach, which rather relies on the fields  $\mathbf{B}$  and  $\mathbf{E}$  themselves (for a justification, see [6] and [7]). In this formulation, the fields are defined as a function of the potentials (and not the reverse) according to  $\mathbf{B} = \nabla \times \mathbf{A}$  and  $\mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}$ . As a consequence of these definitions and using obvious vec-

torial identities, the fields obey the following constraints  $\nabla \cdot \mathbf{B} = 0$  and  $\frac{\partial \mathbf{B}}{\partial t} = -\nabla \times \mathbf{E}$ . But how are defined the potentials themselves ? They are the mathematical solutions of the Maxwell-Minkowski equations written for the excitations:

$$\nabla \cdot \mathbf{D} = \rho \quad \text{and} \quad \nabla \times \mathbf{H} = \frac{\partial \mathbf{D}}{\partial t} + \mathbf{J}. \quad (1)$$

We have to relate the excitations to the fields thanks to the constitutive relations for media at rest and then the fields to the potentials thanks to their definitions above. The current density features two terms  $\mathbf{J} = \mathbf{J}_{\text{constitutive}} + \mathbf{J}_{\text{external}}$ . The constitutive current which expresses the matter response to the fields depends on the medium. For example, in Ohmic conductors, we have  $\mathbf{J}_{\text{Ohm}} = \sigma \mathbf{E} = \sigma (-\nabla V - \frac{\partial \mathbf{A}}{\partial t})$  whereas in a Superconductor [8], the constitutive relation becomes  $\mathbf{J}_{\text{Supra}} = \frac{\hbar n e^*}{m} (\nabla \phi - \frac{e^*}{\hbar} \mathbf{A})$ . For continuous media at rest the excitations are related to the fields according to  $\mathbf{D} = \epsilon \mathbf{E}$  and  $\mathbf{B} = \mu \mathbf{H}$ . We get a system of equations where the unknowns are the potentials  $\mathbf{S}(\mathbf{A}, V; \epsilon, \mu, \rho, \mathbf{J}) = 0$  provided the sources are given or expressed in function of the potentials which vanish far from the latter or take prescribed values on given boundaries. However, the system  $\mathbf{S} = 0$  cannot be solved unless another equation is added. This closure assumption is usually known as the "gauge condition" in the Heaviside-Hertz formulation since the potentials are *de facto* underdetermined (by the "gauge transformations"  $\mathbf{A}' = \mathbf{A} + \nabla f$  and  $V' = V - \frac{\partial f}{\partial t}$  [5]) if and only if they are defined in function of the fields and not the reverse as in the Riemann-Lorenz formulation.

In the following, we will show that the closure assumption is a consequence of the Relativistic or Galilean nature of the problem under study. For that purpose, we will recall the Stratton "gauge condition" which is, according to us, the most general physical constraint which can be used all the times. Then, thanks to the Galilean limits of Classical Electromagnetism [1, 9, 10, 11, 12], we will approximate the Stratton "gauge condition" depending on the context and we will recover the other "gauge conditions" introduced in the literature by pointing out their domain of validity.

The Stratton "gauge condition" was introduced in Physics at M.I.T. in 1941 by Julius Adams Stratton [13] to cope with the propagation of electromagnetic waves in Ohmic conductors such that the sources are given by  $\rho = 0$  and  $\mathbf{J}_{\text{constitutive}} = \mathbf{J}_{\text{Ohm}}$ . Its temporal Fourier transformation was known as early as 1928 by communication engineers like John Renshaw Carson from Bell System [14]. Indeed, from the temporal Fourier transformation of the Maxwell-Ampère equation  $\nabla \times \hat{\mathbf{H}} = i\omega\epsilon\hat{\mathbf{E}} + \sigma\hat{\mathbf{E}}$ , Carson introduced a complex permittivity  $\underline{\epsilon} = \epsilon - i\frac{\sigma}{\omega}$  into the temporal Fourier transformation of the Lorenz "gauge condition"  $\nabla \cdot \mathbf{A} + \mu\epsilon\frac{\partial V}{\partial t} = 0$  [5, 16] to obtain the temporal Fourier transformation of the Stratton "gauge condition"  $\nabla \cdot \hat{\mathbf{A}} + (i\omega\mu\epsilon + \mu\sigma)\hat{V} = 0$ .

According to Stratton's alternative procedure, Gauss' law  $\nabla \cdot \mathbf{E} = 0$  implies immediately:

$$\nabla^2 V + \frac{\partial}{\partial t} (\nabla \cdot \mathbf{A}) = 0 \quad (2)$$

which can be solved if and only if the potentials are constrained by the Stratton "gauge condition":

$$\nabla \cdot \mathbf{A} + \mu\epsilon\frac{\partial V}{\partial t} = -\mu\sigma V. \quad (3)$$

In the simple case of constant permeability  $\mu$  and permittivity  $\epsilon$ , Stratton deduced from the Maxwell-Minkowski's set the following equations ( $\mathbf{S}_{\text{Stratton}} = 0$ ):

$$\nabla^2 V - \mu\epsilon\frac{\partial^2 V}{\partial t^2} - \mu\sigma\frac{\partial V}{\partial t} = 0 \quad (4)$$

$$\nabla^2 \mathbf{A} - \mu\epsilon\frac{\partial^2 \mathbf{A}}{\partial t^2} - \mu\sigma\frac{\partial \mathbf{A}}{\partial t} = -\mu\mathbf{J}_{\text{external}} \quad (5)$$

which are the well-known "telegrapher's equations". They were derived previously for the tension and the current by Vaschy and Heaviside starting from the global electrical equations of Kirchhoff for circuitry and not directly from the local Maxwell-Minkowski equations for the fields. As an example, they described the propagation of waves in a coaxial cable with Ohmic dissipation. Later, Paul Poincelot derived its tensorial expression since the Stratton "gauge condition" is not manifestly Relativistic covariant under the Lorentz transformations of space-time [15]. The more famous Lorenz "gauge condition" [5, 16] is the dissipation-free version of the Stratton's constraint ( $\sigma = 0$ ). As a partial conclusion, it is very surprising to notice that the Stratton "gauge condition" is completely absent from modern textbooks and is not even mentioned in the benchmark review paper on the history of Gauge Invariance [5].

Now, we recall the reader of the physical meaning of the potentials [7] and their constraints. As for the Stratton "gauge condition", the following interpretations of the "gauge conditions" are nowhere in modern treatments of Classical Electromagnetism. The Lorenz "gauge condition" for vacuum  $\nabla \cdot \mathbf{A} + \frac{1}{c^2}\frac{\partial V}{\partial t} = 0$  is analogous to

the mass continuity equation for compressible flows in the particular case of the linearized acoustic perturbations. As a matter of fact, the mass conservation of a flowing fluid is encoded in the following law [17]:

$$\nabla \cdot (\rho\mathbf{u}) + \frac{\partial \rho}{\partial t} = 0. \quad (6)$$

If we perturb the density, pressure and velocity around a basic state at rest:  $\rho = \rho_0 + \delta\rho$ ,  $p = p_0 + \delta p$  and  $\mathbf{u} = \mathbf{0} + \delta\mathbf{u}$ , the continuity equation can be recast in a Lorenz "gauge condition" form:

$$\nabla \cdot (\delta\mathbf{u}) + \frac{1}{c_s^2}\frac{\partial}{\partial t} \left( \frac{\delta p}{\rho_0} \right) = 0 \quad (7)$$

where  $c_s = \frac{1}{\sqrt{\rho\kappa}} = \sqrt{\frac{\partial p}{\partial \rho}} \simeq \sqrt{\frac{\delta p}{\delta \rho}}$  is the speed of sound analogous to the speed of light in vacuum  $c = \frac{1}{\sqrt{\mu_0\epsilon_0}}$ .

The Coulomb "gauge condition"  $\nabla \cdot \mathbf{A} = 0$  is analogous to the mass continuity equation for incompressible flows  $\nabla \cdot \mathbf{u} = 0$  [17] provided that the compressibility (permittivity) vanishes i.e.  $\kappa \rightarrow 0$  at constant density  $\rho_0$  (permeability). As we will see later on, this approximation corresponds to the Galilean (magnetic) limit of the Lorenz "gauge condition" [9, 10, 11].

The Stratton "gauge condition" is a generalized continuity equation for the vector potential :

$$\nabla \cdot \mathbf{A} + \mu\epsilon\frac{\partial V}{\partial t} = -\mu\sigma V. \quad (8)$$

The right-hand side is a sink term. The vector potential is dissipated by Ohmic conduction. Loci of high scalar potential are sinks for the vector potential whose flux is directed towards them. The Stratton "gauge condition" is analogous to the mass continuity equation with nuclear reactions acting as a sink.

Thanks to the above analogy with Fluid Mechanics, it is now obvious to the reader that the vector (scalar) potential is a kind of electromagnetic momentum (energy) per unit charge [7]. Once again, modern Physics has almost completely forgotten the physical meaning of the potentials as it was formulated by James Clerk Maxwell in the nineteenth century and part of his results are rediscovered from time to time either by historians of science or Physics teachers [7].

We have just recalled three examples of "gauge conditions". It is clear that the analogy with Fluid Mechanics advocates for different domains of validity depending on the underlying Physics. Here, we will discuss how to choose a "gauge condition" depending on the context. Our method will be dimensional analysis as often in Fluid Mechanics. Our guide will be Relativistic or Galilean Covariance. That is why we start by a recap on Galilean Electromagnetism as described by Physicists following Lévy-Leblond and Le Bellac [1, 9, 10, 11, 12] and Engineers following another M.I.T. researcher James Melcher [2].

We list first the dimensional quantities. An electromagnetic phenomenon happens in a spatial arena of extension  $L$  in a duration  $\tau$ . The arena is a continuous medium with constitutive properties  $\epsilon, \mu$  and  $\sigma$  taken as constant for simplicity (otherwise they are tensors with time and space dependence). Applying the Vaschy-Buckingham theorem of dimensional analysis [17], we can construct dimensionless parameters which would characterize the electromagnetic response of the continuous medium. As we will deal with Galilean approximations, we introduce  $v \approx \frac{L}{\tau}$  the typical velocity of the system and we compare it with  $c_m = \frac{1}{\sqrt{\mu\epsilon}}$  the light celerity in the continuous medium. The Galilean limit (quasi-static approximation) corresponds to  $v \ll c_m$ . If we neglect time dependence in the Stratton system  $\mathbf{S}_{Stratton} = 0$  ( $\partial/\partial t = 0$  or  $\partial/\partial t \simeq 0$ ), we get  $\nabla^2 \mathbf{A} = -\mu \mathbf{J}$  and  $\nabla^2 V = -\frac{\rho}{\epsilon}$ . In terms of orders of magnitude [1, 9, 10, 11, 12] (the tilde means order of magnitude), we deduce  $\tilde{A} \approx L^2 \mu \tilde{J}$  and  $\tilde{V} \approx \frac{L^2}{\epsilon} \tilde{\rho}$ . Hence, we construct by hand the dimensionless parameter:

$$\frac{\frac{1}{\sqrt{\epsilon\mu}} \tilde{A}}{\tilde{V}} \approx \frac{\tilde{J}}{\tilde{\rho} \frac{1}{\sqrt{\epsilon\mu}}} \quad (9)$$

which characterizes the type of regime [1, 9, 10, 11, 12]: (i)  $v \simeq c_m$  and  $c_m \tilde{A} \simeq \tilde{V} \rightarrow$  Relativistic regime; (ii)  $v \ll c_m$  and  $c_m \tilde{A} \gg \tilde{V}$  ( $\tilde{V} \simeq v \tilde{A}$ )  $\rightarrow$  Galilean magnetic limit (magnetoquasi-statics or MQS); (iii)  $v \ll c_m$  and  $c_m \tilde{A} \ll \tilde{V}$  ( $\tilde{A} \simeq \frac{v}{c_m} \tilde{V}$ )  $\rightarrow$  Galilean electric limit (electroquasi-statics or EQS).

The Stratton's continuity equation becomes (the bar denotes a dimensionless quantity):

$$\frac{\tilde{A}}{L} \nabla \cdot \bar{\mathbf{A}} + \mu \epsilon \frac{\tilde{V}}{\tau} \frac{\partial \bar{V}}{\partial \bar{t}} = -\mu \sigma \tilde{V} \bar{V} \quad (10)$$

that is:

$$\nabla \cdot \bar{\mathbf{A}} + \frac{\frac{L}{\tau}}{\frac{1}{\sqrt{\epsilon\mu}}} \frac{\tilde{V}}{\frac{1}{\sqrt{\epsilon\mu}} \tilde{A}} \frac{\partial \bar{V}}{\partial \bar{t}} = -\frac{L}{l_*} \frac{\tilde{V}}{\frac{1}{\sqrt{\epsilon\mu}} \tilde{A}} \bar{V} \quad (11)$$

whose mathematical form is simply  $I + II = III$  with the following dimensionless ratios:

$$\frac{II}{I} \approx \frac{v}{c_m} \frac{\tilde{V}}{c_m \tilde{A}} = \frac{\tau_{em}}{\tau} \frac{\tilde{V}}{c_m \tilde{A}} \quad (12)$$

$$\frac{III}{I} \approx \frac{L}{l_*} \frac{\tilde{V}}{c_m \tilde{A}} = \frac{\tau_m}{\tau_{em}} \frac{\tilde{V}}{c_m \tilde{A}} = \frac{\tau_{em}}{\tau_e} \frac{\tilde{V}}{c_m \tilde{A}} \quad (13)$$

$$\frac{III}{II} \approx \frac{\tau}{\tau_e} \quad (14)$$

and where we introduced the following parameters [2]:

$l_* = \frac{1}{\sigma} \sqrt{\frac{\epsilon}{\mu}}$  the constitutive length,  $\tau_{em} = \frac{L}{c_m}$  the light transit time,  $\tau_e = \sqrt{\frac{\epsilon}{\sigma}}$  the charge relaxation time and  $\tau_m = \mu \sigma L^2$  the magnetic diffusion time such that  $\tau_{em} = \sqrt{\tau_e \tau_m}$ .

The Figure 1 displays the different approximations of the Stratton's constraint depending on the Relativistic or Galilean (Magnetic, Electric or Statics) regime for a given problem. In practice, we compare the magnitude of the three terms  $I, II$  and  $III$  in the Stratton's constraint using the scaling laws (i), (ii) or (iii).




FIG. 1: Domains of validity of (a) the Galilean limits and (b) the various "gauge conditions" in a log-log plot inspired by Melcher [2] with dimensionless times as variables.

Hence, the "gauge conditions" are continuity equations whose domains of validity depend on the Relativistic or Galilean nature of the underlying phenomenon and have nothing to do with mathematical closure assumptions taken without physical motivations.

According to our results, Gauge Invariance is NOT a fundamental symmetry of Physics since (1) the "gauge transformations" can be avoided by a direct definition of the potentials as mathematical solutions of the Maxwell-Minkowski equations; (2) the "gauge conditions" are interpreted physically as electromagnetic continuity equations; (3) the "gauge fields" are interpreted phys-

ically as electromagnetic energy and momentum per unit charge; (4) the "gauge conditions" have domains of validity derived from Relativistic or Galilean Covariance.

The author would like to thank Francesca Rapetti for playing the role of a sounding board.

- 
- [1] M. Le Bellac and J.-M. Lévy-Leblond, Galilean Electromagnetism, *Il Nuovo Cimento*, **14**, p. 217-233, 1973.
- [2] H. H. Woodson and J. R. Melcher, *Electromechanical Dynamics*, Wiley, New York (1968).  
J. R. Melcher, *Continuum Electromechanics*, The M.I.T. Press (1981).  
M. Zahn and H. A. Haus, Contributions of Prof. James R. Melcher to Engineering-Education, *Journal of Electrostatics*, **34**, p. 109-162, 1995.  
J. R. Melcher and H. A. Haus, *Electromagnetic Fields and Energy*, Hypermedia Teaching Facility, M.I.T. (1998).  
(Book Available at:  
[http://web.mit.edu/6.013\\_book/www/](http://web.mit.edu/6.013_book/www/)).
- [3] M. Henneaux and C. Teitelboim, *Quantization of gauge systems*, Princeton University Press (1992).
- [4] C. P. Enz, *No time to be brief: A scientific biography of Wolfgang Pauli*, Oxford University Press (2002).
- [5] J. D. Jackson and L. B. Okun, Historical roots of gauge invariance, *Reviews of Modern Physics*, Vol. 73, p. 663-680, 2001.
- [6] T. Levi-Civita, Sulla riducibilità delle equazioni elettrodinamiche di Helmholtz alla forma hertziana, *Il Nuovo Cimento*, VI (4), p. 93-108, 1897.  
T. Levi-Civita, Sur le champ électromagnétique engendré par la translation uniforme d'une charge électrique parallèlement à un plan conducteur indéfini, *Annales de la faculté des sciences de Toulouse*, Sér. 2, 4, p. 5-44, 1902.  
(Article available at: [http://archive.numdam.org/article/AFST\\_1902\\_2\\_4\\_5\\_0.pdf](http://archive.numdam.org/article/AFST_1902_2_4_5_0.pdf)).  
A. O'Rahilly, *Electromagnetic Theory, a Critical Examination of Fundamentals*, New York: Dover (1965).  
C. C. Su, Explicit definitions of electric and magnetic fields in potentials and derivation of Maxwell's equations, *European Journal of Physics*, **22**, L5-L8, 2001.  
C. Mead, *Collective Electrodynamics : Quantum Foundations of Electromagnetism*, The M.I.T. Press (2002).  
O. D. Jefimenko, Presenting electromagnetic theory in accordance with the principle of causality, *European Journal of Physics*, **25**, p. 287-296, 2004.
- [7] A. Bork, Maxwell and the Vector Potential, *Isis*, Vol. 58, p. 210-222, 1967.  
E. J. Konopinski, What the electromagnetic vector potential describes, *American Journal of Physics*, **46** (5), p. 499-502, 1978.  
W. Gough and J. P. G. Richards, Electromagnetic or electromagnetic induction?, *European Journal of Physics*, **7**, p. 195-197, 1986.  
J. Roche, Explaining electromagnetic induction: a critical re-examination. The clinical value of history in physics, *Physics Education*, **22**, p. 91-99, 1987.  
J. Roche, A critical study of the vector potential, in *Physicists Look Back* edited by John Roche, Adam Hilger, Chap. 9, p. 144-168, 1990.
- R. Anderson, On an Early Application of the Concept of Momentum to Electromagnetic Phenomena: The Whewell-Faraday Interchange, *Studies in the History and Philosophy of Science*, **25**, p. 577-594, 1994.
- M. D. Semon and J. R. Taylor, Thoughts on the magnetic vector potential, *American Journal of Physics*, **64** (11), p. 1361-1369, 1996.
- A. Tonomura, *The quantum world unveiled by electron waves*, World Scientific (1998).
- A. C. T. Wu and C. N. Yang, Evolution of the concept of the vector potential in the description of fundamental interactions, *International Journal of Modern Physics A*, Vol. 21, No. 16, p. 3235-3277, 2006.
- G. Rousseaux, R. Kofman and O. Minazzoli, The Maxwell-Lodge effect : significance of electromagnetic potentials in the classical theory, *The European Physical Journal D*, Volume 42, Number 2, p. 249-256, 2008.
- [8] M. Thinkham, *Introduction to Superconductivity* (second ed.), McGraw-Hill, New York (1996).
- [9] M. de Montigny, F. C. Khanna and A. E. Santana, Non-relativistic wave equations with gauge fields, *International Journal of Theoretical Physics*, **42**, p. 649-71, 2003.
- [10] G. Rousseaux, Lorenz or Coulomb in Galilean electromagnetism?, *EuroPhysics Letters*, **71**, p. 15-20, 2005.
- [11] M. de Montigny and G. Rousseaux, On some applications of Galilean electrodynamics of moving bodies, *American Journal of Physics*, **75**, p. 984-992, 2007.
- [12] G. Rousseaux, On the electrodynamics of Minkowski at low velocities, *EuroPhysics Letters*, **84**, p. 20002, 2008.
- [13] J. A. Stratton, *Electromagnetic Theory*, McGraw-Hill, New York (1941).
- [14] J. R. Carson, The Rigorous and Approximate Theories of Electrical Transmission Along Wires, *Bell System Technical Journal*, **7**, p. 11-25, 1928.  
(Article available at:  
[http://puhep1.princeton.edu/~mcdonald/examples/EM/carson\\_bstj\\_7\\_11\\_28](http://puhep1.princeton.edu/~mcdonald/examples/EM/carson_bstj_7_11_28)).
- [15] P. Poincelot, Généralisation de la condition de Lorentz, *Annales des Télécommunications*, Tome 18, Numéros 9-10, p. 174-176, 1963.
- [16] Dutch physicist Hendrik Anton Lorentz is often credited for the gauge condition, whereas it is actually due to Danish physicist Ludvig Valentin Lorenz. For a justification, see:  
O. Keller, *Optical works of L.V. Lorenz* in *Progress in Optics XXXVII*, edited by E. Wolf (Amsterdam: North-Holland) 257-343, 1997.
- [17] E. Guyon, J.-P. Hulin, L. Petit and C. D. Mitescu, *Physical Hydrodynamics*, Oxford University Press (2001).