



## New values of gravitational moments $J_2$ and $J_4$ deduced from helioseismology

Redouane Mecheri, Toufik Abdelatif, Abdanour Irbah, Janine Provost,  
Gabrielle Berthomieu

### ► To cite this version:

Redouane Mecheri, Toufik Abdelatif, Abdanour Irbah, Janine Provost, Gabrielle Berthomieu. New values of gravitational moments  $J_2$  and  $J_4$  deduced from helioseismology. Solar Physics, Springer Verlag, 2004, v. 222 (Issue 2), pp.191-197. <10.1023/B:SOLA.0000043563.96766.21>. <hal-00433235>

**HAL Id: hal-00433235**

**<https://hal.archives-ouvertes.fr/hal-00433235>**

Submitted on 20 Nov 2009

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# New values of gravitational moments $J_2$ and $J_4$ deduced from helioseismology

R. Mecheri, T. Abdelatif and A. Irbah

*C.R.A.A.G - Observatoire d'Alger BP 63 Bouzareah, Alger, Algeria.*

J. Provost and G. Berthomieu

*Departement Cassini, UMR CNRS 6529 - Observatoire de la Côte d'Azur, BP 4229, 06304 Nice CEDEX4, France.*

**Abstract.** By applying the theory of slowly rotating stars to the Sun, the solar quadrupole and octopole moments  $J_2$  and  $J_4$  were computed using a solar model obtained from *CESAM* stellar evolution code (Morel (1997)) combined with a recent model of solar differential rotation deduced from helioseismology (Corbard *et al.* (2002)). This model takes into account a near-surface radial gradient of rotation which was inferred and quantified from *MDI f-modes* observations by Corbard and Thompson (2002). The effect of this observational near-surface gradient on the theoretical values of the surface parameters  $J_2$ ,  $J_4$  is investigated. The results show that the octopole moment  $J_4$  is much more sensitive than the quadrupole moment  $J_2$  to the subsurface radial gradient of rotation.

## 1. Introduction

Several theoretical determinations of the  $J_2$  and  $J_4$  gravitational moments have been undertaken in case of different solar differential rotation laws : (i) only radius dependence (Goldreich and Schubert, 1968; Paternò, Sofia, and Di Mauro, 1996), (ii) quadratic expansion in colatitude cosine terms (Ulrich and Hawkins (1981a and 1981b)), (iii) angular velocity distribution with a slowly latitude variation determined by mean of helioseimology technics (Gough (1982)). More recent determinations are those performed by : (i) Armstrong and Kuhn (1999) using a quadratic expansion rotation law with coefficients obtained by fitting higher resolution helioseismic interior rotation data from MDI (Scherrer *et al.* (1995)), (ii) Godier and Rozelot (1999) and (iii) Roxburgh (2001) using the differential rotation model given by Kosovichev (1996) from *BBSO p-modes* observations. This model takes into account the presence of a constant near-surface radial gradient based on the assumption that the angular momentum is preserved in the supergranulation layer. The aim of the present work is a contribution to  $J_2$  and  $J_4$  determinations using a new analytical model of solar differential rotation provided by Corbard *et al.* (2002) which has a latitudinal dependent profile of the near-surface radial gradient of rotation.

If we consider the Sun as an axial symmetry distribution of matter in



© 2009 Kluwer Academic Publishers. Printed in the Netherlands.

rotation, the outer gravitational field  $\phi_{out}$  is expressed as :

$$\phi_{out}(r, \theta) = -\frac{GM_{\odot}}{r} \left[ 1 - \sum_{n=1}^{\infty} \left( \frac{R_{\odot}}{r} \right)^{2n} J_{2n} P_{2n}(\cos \theta) \right] \quad (1)$$

where  $J_{2n}$  are the gravitational moments,  $P_{2n}$  the Legendre polynomials and  $r, \theta$  respectively the distance from the Sun centre and the angle to the symmetry axis (colatitude).

Since the solar rotation is slow, it induces small perturbations around the spherical equilibrium. These perturbations can be expanded on Legendre polynomial basis. The distribution of the gravitational potential in the Sun can be written :

$$\phi(r, \theta) = \phi_0(r) + \phi_1(r, \theta) = \phi_0(r) + \sum_{n=1}^{\infty} \phi_{12n}(r) P_{2n}(u) \quad (2)$$

where  $u = \cos \theta$ .

The gravitational moments  $J_{2n}$  are obtained assuming the continuity of the gravitational potential at the surface :

$$J_{2n} = \frac{R_{\odot}}{GM_{\odot}} \phi_{12n}(R_{\odot}) \quad (3)$$

The perturbed potential is obtained by linearization of the equation of hydrostatic equilibrium and the Poisson equation, leading to :

$$\begin{aligned} \frac{d^2 \phi_{12n}}{dr^2} + \frac{2}{r} \frac{d\phi_{12n}}{dr} - (2n(2n+1) + UV) \frac{\phi_{12n}}{r^2} = U[(V+2)B_{2n} + \\ + r \frac{dB_{2n}}{dr} + \frac{4n+1}{2} \int_{-1}^{+1} (1-u^2) P_{2n}(u) \Omega(r, u)^2 du] \end{aligned} \quad (4)$$

where  $U = 4\pi\rho_0 r^3/M_r$ ,  $V = d \ln \rho_0 / d \ln r$ ,  $M_r$  is the mass contained in a sphere of radius  $r$  and  $\Omega(r, u)$  the angular velocity.  $B_{2n}$  is given by:

$$B_{2n}(r) = -\frac{1}{2n!} \frac{4n+1}{2^{2n+1}} \int_{-1}^{+1} u \Omega(r, u)^2 \frac{d^{2n-1}}{du^{2n-1}} ((u^2-1)^{2n}) du \quad (5)$$

Equation (3) is integrated with the usual boundary conditions, using  $U$  and  $V$  provided by a solar model obtained from the *CESAM* stellar evolution code (Morel (1997)) and a rotation law derived from helioseismology.

## 2. Analytical model of solar differential rotation

We consider the recent Corbard *et al.* (2002) model of solar differential rotation and, for comparison, the Kosovichev one (1996) already used by Roxburgh (2001) and Godier and Rozelot (1999). The Corbard model contains a near-surface radial gradient of rotation inferred from the radial dependence of the *MDI f-modes* observations (Corbard and Thompson (2002)). Two estimations of this gradient have been derived from different sets of modes leading to two rotation models denoted afterward by (a) and (b). As for Kosovichev's model, the surface rotation is forced to surface plasma observations (Snodgrass, 1992).

Figure 1 shows the solar rotation profiles corresponding to these models computed for different latitudes. Kosovichev's model presents a negative constant subsurface radial gradient. The Corbard models have a negative value of the radial gradient at low latitude which is twice smaller than Kosovichev ones. At high latitude, the gradients are positive with larger magnitudes for the Corbard model (b).

We use an analytical expression derived respectively by Corbard *et al.* (2002) and Dikpati *et al.* (2002) for the Corbard rotation laws and for the Kosovichev one. We recall hereafter the full set of equations and parameters they give for these rotation laws :

$$\Omega(r, u) = A_1(r, u) + \Psi_{tac}(r) (\Omega_{cz} - \Omega_0 + a_2 u^2 + a_4 u^4) \quad (6)$$

where

$$A_1(r, u) = \Omega_0 + \Psi_{cz}(r) \{\alpha(u)(r - r_{cz})\} + \Psi_s(r) \{\Omega_{eq} - \Omega_{cz} - \beta(u)(r - R_\odot) - \alpha(u)(r - r_{cz})\} \quad (7)$$




Figure 1. Profiles of the solar rotation from  $0.55R_\odot$  to the surface for different latitudes computed each  $10^\circ$  from the Equator (top) to the Pole (bottom). (1) Model of Kosovichev. (2) Model of Corbard (a). (3) Model of Corbard (b).

with

$$\alpha(u) = \frac{\Omega_{eq} - \Omega_{cz} + \beta(u)(R_{\odot} - r_s)}{r_s - r_{cz}}$$

$\Omega_0$ ,  $\Omega_{eq}$  and  $\Omega_{cz}$  are respectively the constant rotation of the radiative interior zone, the equatorial rate at the surface and at the top  $r_{cz}$  of the tachocline localized at  $r_{tac}$ . The  $a_2$  and  $a_4$  constants describe the latitudinal differential rotation.  $\beta(u)$  represents the latitudinal dependence of the rotation radial gradient below the surface down to a radius  $r_s$ . This gradient depends on the latitude through the  $\beta_0$ ,  $\beta_3$  and  $\beta_6$  constants by the Equation :  $\beta(u) = \beta_0 + \beta_3 u^3 + \beta_6 u^6$ .

The  $\Psi_x$  function where  $x$  stands for  $tac$ ,  $cz$  or  $s$ , models the transition between different gradients. An error function centered at  $r_x$  with width  $\omega_x$  is used for this goal :  $\Psi_x(r) = 0.5 (1 + erf [2(r - r_x)/\omega_x])$ .

All the rotation laws have the following common parameters :  $\Omega_0 = 435nHz$ ,  $\Omega_{eq} = 452.5nHz$ ,  $\Omega_{cz} = 453.5nHz$ ,  $r_{tac} = 0.69R_{\odot}$ ,  $r_{cz} = 0.71R_{\odot}$ ,  $a_2 = -61nHz$ ,  $a_4 = -73.5nHz$ . The parameters which are different for the three laws are given in Table I.

Table I. The non common parameter values between the rotation models. The  $\beta_0$ ,  $\beta_3$  and  $\beta_6$  parameters are given in  $nHz/R_{\odot}$ .

model	$\omega_{tac}/R_{\odot}$	$\omega_{cz}/R_{\odot}$	$\omega_s/R_{\odot}$	$r_s/R_{\odot}$	$\beta_0$	$\beta_3$	$\beta_6$
Kosovichev (1996)	0.1	0	0	0.983	891.5	0	0
Corbard (a) (2002)	0.05	0.05	0.05	0.97	437	-214	-503
Corbard (b) (2002)	0.05	0.05	0.05	0.97	437	0	-1445

### 3. Results and discussion

We present in Table II the computed values of  $J_2$  and  $J_4$  obtained with different solar rotation models described in Section 2 and with an uniform rotation equal to the rotation rate of the solar radiative zone  $\Omega_0$ , for comparison. Table III gives also some values of  $J_2$  and  $J_4$  presented by other authors.

Our results show that the differential rotation in the convective zone reduces  $J_2$  value of about 0.8% when the Corbard models are considered and about 0.5% in the case of Kosovichev's model. For this last case, our  $J_2$  determination is larger than the value found by Godier and Rozelot (1999) but in agreement with the one obtained by Roxburgh (2001), both using Kosovichev's model. Our values are also in agreement with

Table II. The  $J_2$  and  $J_4$  values corresponding to the different rotation models.  $\Omega_0 = 2.733 \mu rd/s$  is the rotation in the radiative zone.

Model of rotation	$J_2(\times 10^{-7})$	$J_4(\times 10^{-9})$
Uniform rotation ( $\Omega_0$ )	2.217	0
Kosovichev (1996)	2.205	-4.455
Corbard (a) (2002)	2.201	-5.601
Corbard (b) (2002)	2.198	-4.805

those given by Paternò *et al.* (1996), Pijpers (1998) and Armstrong and Kuhn (1999) (see Table III). All these values deviate from the range obtained by Ulrich and Hawkins (1981a and 1981b) in the case of the rotation law defined as a simple quadratic expansion. The difference between the subsurface radial gradients induces only a small reduction on  $J_2$  values. It is about 0.25% between Kosovichev's model and Corbard's ones. This difference is however about 0.1% between the two Corbard models.

As expected, the effect of the subsurface radial gradient is more important on  $J_4$  gravitational moment.  $J_4$  absolute values obtained using the models of Corbard (a) and (b) are respectively about 20% and 7% larger than the one obtained with Kosovichev's model. The (a) Corbard model increases the  $J_4$  absolute value of about 14% compared to the one obtained from the (b) Corbard model. Our  $|J_4|$  value corresponding to Kosovichev's model is in agreement with the one given by Roxburgh (2001) using the same model. However, those obtained from Corbard's models are larger than other values given in Table III. All these values

Table III. Some computed values of  $J_2$  and  $J_4$  obtained by other authors. The large value of Gough (1982) is due to an estimation of the internal rotation deduced from earlier helioseismic observations

Authors	$J_2(\times 10^{-7})$	$J_4(\times 10^{-9})$
Ulrich and Hawkins (1981)	$1.0 < J_2 < 1.5$	$2.0 <  J_4  < 5.0$
Gough (1982)	36	-
Paternò <i>et al.</i> (1996)	2.22	-
Pijpers (1998)	2.18	-
Godier and Rozelot (1999)	1.6	-
Armstrong and Kuhn (1999)	2.22	-3.84
Roxburgh (2001)	2.206	-4.45

are consistent with the range given by Ulrich and Hawkins (1981a and 1981b) except for the (a) Corbard model.

Rotation induces a distortion of the solar surface which can be roughly related to  $J_2$  through the following quantity often called oblateness :

$$\frac{R_e - R_p}{R_\odot} \approx \frac{3}{2}J_2 + \frac{\Omega_s^2 R_\odot^3}{2GM_\odot} \quad (8)$$

where  $\Omega_s$  is an effective rotation rate.  $R_e$ ,  $R_p$  and  $R_\odot$  are respectively the equatorial, polar and mean solar radius. This formula is strictly valid for an uniform rotation or for a rotation constant on cylinders. For a solar rotation which presents a complex profile not constant on cylinders, Paternò, Sofia, and Di Mauro (1996) proposed an expression of  $\Omega_s$  derived from the surface rotation  $\Omega(R_\odot, u)$ . In our case,  $\Omega_s$  will be the same for the three models since they are built so as to have the same surface rotation. Thus, in this rough description, the effect of different subsurface radial gradients of rotation on the oblateness appears through the modification of the  $J_2$  gravitational moment. It is negligible since the main term in Equation (8) is the surface rotation term. The value of the oblateness found is  $9.1 \times 10^{-6}$ . It is in agreement with observations of Lydon and Sofia (1996) and Rozelot and Rösch (1997) but slightly larger than those of Kuhn *et al.* (1998) and Armstrong and Kuhn (1999). Rozelot, Godier, and Lefebvre (2001) presented new developments taking into account the surface latitudinal differential rotation to link  $J_2$  and  $J_4$  to the solar equatorial and polar radius. To the lowest order, their formula reduces to Equation (8) with another definition of  $\Omega_s$ . For our surface rotation, their formula leads to an oblateness value equal to  $11.4 \times 10^{-6}$ . We have estimated that the effects of the second terms are of the order of (4/1000) in the case of Kosovichev's model. New constraints on the oblateness and the shape of the solar surface will hopefully be provided by the future *PICARD* microsatellite *CNES-mission* (Thuillier *et al.* (2003)).

In conclusion, the octopole moment  $J_4$  is much more sensitive than the quadrupole moment  $J_2$  to the inclusion of the latitudinal and radial differential rotation in the convective zone and particularly to the subsurface radial gradient of rotation. Indeed, an important subsurface radial gradient at high latitude decreases significantly the value of  $|J_4|$  while it does not affect significantly the  $J_2$  value.

### Acknowledgements

We thank Thierry Corbard for providing his model of rotation and the referee for his constructive remarks and comments. This work has

been performed with support of the French Foreign Affaire Ministry in the framework of scientific cooperation between France and Algeria (contract 00MDU501)

## References

- Armstrong, J. and Kuhn, J. R.: 1999, *Astrophys. J.* **525**, 533.
- Corbard, T. and Thompson, M. J.: 2002, *Solar Phys.* **205**, 211.
- Corbard, T., Dikpati, M., Gilman, P.A. and Thompson, M.J.: 2002, in A. Wilson (ed.), *The SOHO 11 Symposium on From Solar Min to Max: Half a Solar Cycle with SOHO*, Davos, Switzerland, ESA SP-508, Noordwijk: ESA Publications Division, ISBN 92-9092-818-2, p. 75.
- Dikpati, M., Corbard, T., Thompson, M. J. and Gilman, P. A.: 2002, *Astrophys. J.* **575**, 41.
- Godier, S. and Rozelot, M. J.: 2000, *Astron. Astrophys.* **355**, 365.
- Godier, S. and Rozelot, M. J.: 1999, *Astron. Astrophys.* **350**, 310.
- Gough, D. O.: 1982, *Nature.* **298**, 334.
- Kosovichev, A. G.: 1996, *Astrophys. J.* **469**, 61.
- Kuhn, J. R., Bush, R. I., Scheick, X. and Scherrer, P.: 1998, *Nature.* **392**, 155.
- Lydon, T. J. and Sofia, S.: 1996, *Phys. Rev. Letters.* **76**, 177.
- Morel, P.: 1997, *Astron. Astrophys.* **124**, 597.
- Paternò, L., Sofia, S. and Di Mauro, M. P.: 1996, *Astron. Astrophys.* **314**, 940.
- Pijpers, F. P.: 1998, *Monthly Notices Royal Astron. Soc.* **297**, 76.
- Roxburgh, I. W.: 2001, *Astron. Astrophys.* **377**, 688.
- Rozelot, J. P., Godier, S. and Lefebvre, S.: 2001, *Solar Phys.* **198**, 223.
- Rozelot, J. P. and Rösch, J.: 1997, *Solar Phys.* **192**, 11.
- Scherrer, P. H., et al.: 1995, *Solar Phys.* **162**, 129.
- Snodgrass, H. B.: 1992, in ASP Conf. Serf. 27, *The Solar Cycle*, ed. K. L. Harvey (San Francisco : ASP), 205.
- Thuillier, G., Joukoff, A. and Schmutz, W.: 2003, *Proc. ISCS 2003 Symposium, Solar Variability as an Input to the Earth's Environment*, Tatransk Lomnica, Slovakia, 23-28 june 2003 (ESA SP-535, September 2003)
- Ulrich, R. K. and Hawkins, G. W.: 1981a, *Astrophys. J.* **246**, 985.
- Ulrich, R. K. and Hawkins, G. W. "Erratum": 1981b, *Astrophys. J.* **249**, 831.



