



Non trivial generalizations of the Schwinger pair production result II

Herbert M. Fried, Yves Gabellini

► To cite this version:

Herbert M. Fried, Yves Gabellini. Non trivial generalizations of the Schwinger pair production result II. *Physical Review D*, American Physical Society, 2006, 73, pp.011901. <10.1103/PhysRevD.73.011901>. <hal-00012766>

HAL Id: hal-00012766

<https://hal.archives-ouvertes.fr/hal-00012766>

Submitted on 27 Oct 2005

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Non trivial generalizations of the Schwinger pair production result II

H. M. FRIED

*Department of Physics
Brown University
Providence R.I. 02912 USA*

Y. GABELLINI

*Institut Non Linéaire de Nice
UMR 6618 CNRS
1361 Route des Lucioles
06560 Valbonne France*

Abstract

It is suggested that Schwinger's (1951) vacuum persistence probability against pair production by an intense but constant electric field is a very good approximation to the corresponding quantity if the field does not vary appreciably over distances less than $m/e|E|$.

The object of this brief letter is to point out an argument previously overlooked, which suggests that the vacuum persistence probability for the production of charged lepton pairs in Schwinger's seminal 1951 paper [1], calculated for constant electromagnetic fields, is, in fact, a very good approximation to the same quantity in the limit of sufficiently intense fields. Previous approximations [2] have given qualitative results for "slowly-varying" fields; but the relevant observation here is that, unless those fields vary appreciably over space-time distances on the order of $m^{-1}(m^2/e|F|)$, which for intense fields can be far smaller than the Compton wavelength of each member of the produced pair, a respectable approximation to Schwinger's formula is obtained by merely replacing his constant $F_{\mu\nu}$ by $F_{\mu\nu}(x)$.

For equivalently small distances, we also note that the vacuum persistence probability for the production of high mass charged tachyon pairs, as suggested in a recent QTD model [3], can be given an expression quite similar to that of Schwinger; and that the resulting approximation for intense, external fields is essentially the same. Estimates of lepton pair production by crossed laser beams of high intensity [4] are also simplified.

We follow the notation and functional techniques previously used [2, 5] for this problem, in terms of a convenient Fradkin representation [6], where the vacuum persistence probability P is given by $P = \exp[2\text{Re}L[A]]$, with :

$$\begin{aligned}
L[A] = & -\frac{1}{2} \int_0^\infty \frac{ds}{s} e^{-ism^2} \exp \left[i \int_0^s ds' \sum_\mu \frac{\delta^2}{\delta v_\mu^2(s')} \right] \\
& \times \int \frac{d^4p}{(2\pi)^4} e^{ip \cdot \int_0^s ds' v(s')} \int d^4x e^{-ie \int_0^s ds' v_\mu(s') A_\mu(x - \int_0^{s'} v)} \\
& \times \text{tr} \left(e^{\int_0^s ds' \sigma \cdot F(x - \int_0^{s'} v)} \right) \Big|_{v_\mu \rightarrow 0}
\end{aligned} \tag{1}$$

In fact, $L[A]$ is a completely gauge invariant functional of the $F_{\mu\nu}$; this may be seen by following the little known functional argument presented in the book of ref[6], wherein the $A_\mu(x - \int_0^{s'} v)$ of (1) may, for this closed-loop calculation, be replaced by :

$$A_\mu(s') \equiv A_\mu(x, u(s')) \equiv \int_0^1 \lambda d\lambda F_{\mu\nu}(x - \lambda u(s')) u_\nu(s') \tag{2}$$

with $u_\nu(s') = \int_0^{s'} ds'' v_\nu(s'')$; and the $L[A]$ of (1) is then clearly independent of gauge : $L[A] = L[F]$.

As in [2], it is convenient to formally introduce $u_\mu(s')$ as an independent variable, by means of the functional identity :

$$1 = \int d[u] \delta[u(s') - \int_0^{s'} v] = N' \int d[u] \int d[\Omega] e^{i \int_0^s ds' \Omega_\mu(s') [u(s') - \int_0^{s'} v]}$$

where N' is the appropriate normalization constant. With the aid of (2), (1) can be rewritten as :

$$L[F] = -\frac{1}{2} \int_0^\infty \frac{ds}{s} e^{-ism^2} N' \int d[u] \int d[\Omega] e^{i \int \Omega u} \int \frac{d^4p}{(2\pi)^4} \int d^4x \text{tr} \left(e^{\int_0^s ds' \sigma \cdot F(x - u(s'))} \right) \Big|_+$$

$$\times e \int_0^s \frac{\delta^2}{\delta v^2} e \int_0^s ds' v_\mu(s') [p_\mu - eA_\mu(s') - \int_{s'}^s \Omega(s'')] \Big|_{v_\mu \rightarrow 0} \quad (3)$$

where Abel's trick has been used in rewriting $\int_0^s ds' \Omega_\mu(s') \int_0^{s'} ds'' v_\mu(s'')$.

The functional linkage operation of (3) is immediate, and one finds for the second line of (3) :

$$\begin{aligned} \exp\left[-i \int_0^s ds' [p - eA(s') - \int_{s'}^s \Omega(s'')]^2\right] &= \exp\left[-isp^2 - ie^2 \int_0^s A^2 - i \int_0^s \int_0^s ds_1 ds_2 \Omega_\mu(s_1) h(s_1, s_2) \Omega_\mu(s_2) \right. \\ &\quad \left. + 2iep \cdot \int_0^s A + 2ip \cdot \int_0^s ds' s' \Omega(s') - 2ie \int_0^s ds' \Omega_\mu(s') \int_0^{s'} ds'' A_\mu(s'')\right] \end{aligned} \quad (4)$$

where $h(s_1, s_2) = \int_0^s ds' \theta(s' - s_1) \theta(s' - s_2) = \theta(s_1 - s_2) s_2 + \theta(s_2 - s_1) s_1$. The p -dependent terms of (4) then yield :

$$\int \frac{d^4 p}{(2\pi)^4} e^{-isp^2 + 2ip \cdot \int_0^s ds' [eA(s') + s' \Omega(s')]} = \frac{-i}{16\pi^2 s^2} \exp\left\{ \frac{i}{s} \left[\int_0^s ds' (eA(s') + s' \Omega(s')) \right]^2 \right\} \quad (5)$$

and, combining (5) with the remaining of (3), one finds :

$$\begin{aligned} L[F] &= \frac{i}{32\pi^2} \int_0^\infty \frac{ds}{s^3} e^{-ism^2} \int d^4 x N' \int d[u] \int d[\Omega] \text{tr} \left(e \int_0^s ds' \sigma \cdot F(x - u(s')) \right)_+ \\ &\quad \times e^{-ie^2 \left[\int_0^s ds' A^2(s') - \frac{1}{s} \left(\int_0^s ds' A(s') \right)^2 \right]} \\ &\quad \times e^{-i \int_0^s \int_0^s ds_1 ds_2 \Omega_\mu(s_1) J(s_1, s_2) \Omega_\mu(s_2) + i \int_0^s ds' f_\mu(s') \Omega_\mu(s')} \end{aligned} \quad (6)$$

where $J(s_1, s_2) = h(s_1, s_2) - s_1 s_2 / s$, and $f_\mu(s') = u_\mu(s') - 2e \int_0^s ds'' A_\mu(s'') [\theta(s' - s'') - s' / s]$.

The observation we wish to make follows upon performing the Gaussian functional integral :

$$\sqrt{N'} \int d[\Omega] e^{-i \int_0^s \int_0^s \Omega_\mu J \Omega_\mu + i \int_0^s f_\mu \Omega_\mu} = e^{-\frac{1}{2} \text{Tr} \ln(2J)} e^{+\frac{i}{4} \int_0^s \int_0^s f_\mu J^{-1} f_\mu} \quad (7)$$

where $J^{-1}(s_1, s_2)$ satisfies $\delta(s_1 - s_2) = \int_0^s ds' J^{-1}(s_1, s') J(s', s_2) = \int_0^s ds' J(s_1, s') J^{-1}(s', s_2)$

The functional integral (FI) that results from the combination of (7) and (6) cannot, in general, be evaluated exactly; but, because it can be represented as a gaussian-weighted FI over $u_\mu(s')$, proportional to :

$$\begin{aligned} \sqrt{N'} \int d[u] e^{\frac{i}{4} \int_0^s \int_0^s f_\mu J^{-1} f_\mu} \text{tr} \left(e \int_0^s ds' \sigma \cdot F(x - u(s')) \right)_+ \\ \times e^{-ie^2 \left[\int_0^s ds' A^2(s') - \frac{1}{s} \left(\int_0^s ds' A(s') \right)^2 \right]} \end{aligned} \quad (8)$$

and contains non-gaussian corrections of u -dependence buried inside fields of form $F_{\mu\nu}(x - u(s'))$ and $\int_0^1 \lambda d\lambda F_{\mu\nu}(x - \lambda u(s')) u_\nu(s')$, certain approximations are indeed possible.

Firstly, let us assume that $e|F|/m^2 \leq 1$, $f_\mu(s') \sim u_\mu(s')$, i.e., the fields are weak. Then (8) becomes essentially a gaussian FI, with the natural constraints on the size of $|u(s')|$ given by $|\iint u J^{-1} u| \leq 1$. Since $O(J^{-1}) \sim 1/s^3$, and because the scale of s is set by the factor $\exp[-ism^2]$, $O(sm^2) \leq 1$, it follows that $O(|u|) \leq \sqrt{s} \leq 1/m$. Hence, if the fields $F_{\mu\nu}(x - \lambda u)$ do not vary appreciably over distances of a Compton wavelength, the $u(s')$ dependence inside the $F_{\mu\nu}$ can be neglected; one then finds that the term of (8) proportional to $\exp[-ie^2 s < (A - < A >)^2 >]$ becomes unity¹, and one is left with the Schwinger, constant-field solution, although written in an unconventional way, where $F_{\mu\nu} \rightarrow F_{\mu\nu}(x)$. In comparison with Schwinger's constant electric field result for the vacuum persistence probability :

$$P = \exp\left\{-\frac{\alpha}{\pi^2} \int d^4x E^2 \sum_{n=1}^{\infty} \frac{1}{n^2} e^{-n\pi m^2/e|E|}\right\} \quad (9)$$

The restriction $e|E|/m^2 \leq 1$ means that each term in the expansion of $\ln P$ is exponentially small, P remains close to unity, and the probability of pair production is very small.

In contrast, let us now assume that $e|E|/m^2 > 1$. Then the dominant term of $f_\mu(s')$ is not $u_\mu(s')$ but, rather, $-2e \int_0^s ds'' [\theta(s' - s'') - s'/s] \int_0^1 \lambda d\lambda F_{\mu\nu}(x - \lambda u(s')) u_\nu(s')$, which produces a quadratic gaussian weighting in (8) of order $e^2 |F|^2 \iint u J^{-1} u$, which quantity is now restricted to be < 1 . This, in turn, leads to the condition that $O(|u|) < \frac{1}{m} (m^2/e|F|)$, which can be much smaller than the bound of the weak field case. Therefore, unless the intense fields vary significantly over distances less than $(m/e|F|)$, one is justified in dropping the $u(s')$ dependence inside the $F(x - \lambda u(s'))$ and $F(x - u(s'))$ of (8); when this is done, there is no difference between Schwinger's constant field solution for P and the expression generated by (8), except that $F_{\mu\nu} \rightarrow F_{\mu\nu}(x)$. And, in this case, the exponential terms in the expansion of $\ln P$ of (9) can take on significant values, at least until $n = n_{max} \sim e|E|/\pi m^2$.

For realistic external fields $F_{\mu\nu}(x)$, it is just that simple. This observation should have been made by the present authors two years ago, in ref[2].

References

- [1] J. Schwinger, Phys. Rev. **82**, 664 (1951).
- [2] J. Avan, H.M. Fried and Y. Gabellini, Phys. Rev. **D67**, 16003 (2003). We take this opportunity to correct the misprint which occurred in eq.(18) of this reference, where the exponential factors of $\delta/\delta u_\mu$ were incorrectly written in place of the correct $\delta/\delta f_\mu$, so that $\delta/\delta u_\mu$ should be replaced by $\int (\delta u_\nu/\delta f_\mu)(\delta/\delta u_\nu)$.

¹This statement is literally true if only an electric field $\mathcal{E}_3(x_\pm)$ in one spatial direction is considered, where $x_\pm = x_3 \pm x_0$; but the inclusion of this exponential factor cannot change our order-of-magnitude estimates, unless exceptional cancellations were to occur.

- [3] H.M. Fried and Y. Gabellini, arXiv: hep-th/0505272. ;
- [4] H.M. Fried, Y. Gabellini, B.H.J. McKellar and J. Avan, Phys. Rev. D**63**, 125001 (2001).
- [5] H.M. Fried and R.P. Woodard, Phys. Lett. B**524**, 233 (2002).
- [6] E.S. Fradkin, Nucl. Phys. **76**, 588 (1966); a detailed explanation and derivation of the Fradkin representation can be found in the book by H.M. Fried, “ Functional Methods and Eikonal Models ” , Editions Frontières, Gif sur Yvette, France, 1990.