



The Permeability Transition in Plant Mitochondria: The Missing Link

Marco Zancani*, Valentino Casolo, Elisa Petrusa, Carlo Peresson, Sonia Patui, Alberto Bertolini, Valentina De Col, Enrico Braidot, Francesco Boscutti and Angelo Vianello

Plant Biology Unit, Department of Agricultural and Environmental Sciences, University of Udine, Udine, Italy

The synthesis of ATP in mitochondria is dependent on a low permeability of the inner membrane. Nevertheless, mitochondria can undergo an increased permeability to solutes, named permeability transition (PT) that is mediated by a permeability transition pore (PTP). PTP opening requires matrix Ca^{2+} and leads to mitochondrial swelling and release of intramembrane space proteins (e.g., cytochrome c). This feature has been initially observed in mammalian mitochondria and tentatively attributed to some components present either in the outer or inner membrane. Recent works on mammalian mitochondria point to mitochondrial ATP synthase dimers as physical basis for PT, a finding that has been substantiated in yeast and *Drosophila* mitochondria. In plant mitochondria, swelling and release of proteins have been linked to programmed cell death, but in isolated mitochondria PT has been observed in only a few cases and in plant cell cultures only indirect evidence is available. The possibility that mitochondrial ATP synthase dimers could function as PTP also in plants is discussed here on the basis of the current evidence. Finally, a hypothetical explanation for the origin of PTP is provided in the framework of molecular exaptation.

Keywords: permeability transition, plant mitochondria, ATP synthase, exaptation, environmental stress

OPEN ACCESS

Edited by:

Ildikó Szabó,
University of Padova, Italy

Reviewed by:

Hao Peng,
Washington State University, USA
Paolo Pinton,
University of Ferrara, Italy

*Correspondence:

Marco Zancani
marco.zancani@uniud.it

Specialty section:

This article was submitted to
Plant Physiology,
a section of the journal
Frontiers in Plant Science

Received: 30 September 2015

Accepted: 26 November 2015

Published: 15 December 2015

Citation:

Zancani M, Casolo V, Petrusa E,
Peresson C, Patui S, Bertolini A,
De Col V, Braidot E, Boscutti F
and Vianello A (2015)
The Permeability Transition in Plant
Mitochondria: The Missing Link.
Front. Plant Sci. 6:1120.
doi: 10.3389/fpls.2015.01120

THE PERMEABILITY TRANSITION

ATP synthesis in mitochondria occurs by a chemiosmotic coupling of substrate oxidation and phosphorylation (Mitchell, 1961). This explanation is based on the highly selective permeability of the inner mitochondrial membrane (IMM) and on utilization of protonmotive force by the F_1F_0 ATP synthase (F-ATPase) for the synthesis of ATP. Nevertheless, a sudden increase in permeability of the IMM has been described in the 1950s (Raaflaub, 1953a,b) and characterized in the late 1970s (Haworth and Hunter, 1979; Hunter and Haworth, 1979a,b). Initially considered an artifact, later it has been named Permeability Transition (PT) and associated to a pore, the Permeability Transition Pore (PTP). The appreciation of its relevance has increased since it has been related to many diseases in mammals, including reperfusion injury of the heart and muscular dystrophy (Bernardi, 2013a). This mitochondrial PT requires matrix Ca^{2+} and is favored by matrix P_i , as well as benzodiazepine Bz-423 and thiol oxidants, while it can be inhibited by Mg^{2+} , thiol reductants, ADP and ATP (Bernardi, 2013b). Cyclosporin A (CsA) acts as inhibitor of PT (Crompton et al., 1988) by binding with the peptidyl-prolyl isomerase Cyclophilin D (CyPD) (Halestrap and Davidson, 1990). The features of PTP (e.g., pore diameter of ~ 2.8 nm and size exclusion of about 1500 Da) are consistent with those described for the Mitochondrial Mega-Channel (MMC), a high-conductance channel, which is considered to be its electrophysiological equivalent (Szabó and Zoratti, 1992).

THE PT IN PLANTS

The first evidence of a Ca^{2+} -induced and CsA-delayed collapse of transmembrane electrical potential difference ($\Delta\Psi$) in pea stem mitochondria dates back to 1995 (Vianello et al., 1995). PT has been then observed in different plant species, although the features of this phenomenon cannot be summarized in a straightforward model (Table 1). Potato tuber mitochondria exhibit a typical $\text{Ca}^{2+}/\text{P}_i$ -induced PT, inhibited (Arpagaus et al., 2002) or not (Fortes et al., 2001) by CsA. These mitochondria do not show any Ca^{2+} uptake, suggesting an external effect of Ca^{2+} on PT (Fortes et al., 2001), which is not consistent with the observations in mammals (Bernardi et al., 2015). The PT described in oat leaves (Curtis and Wolpert, 2002) and wheat roots (Virolainen et al., 2002) shows a $\text{Ca}^{2+}/\text{P}_i$ -induced $\Delta\Psi$ collapse and matrix swelling, which are CsA-insensitive. Calcium uptake by isolated plant mitochondria occurs spontaneously in wheat, but requires the addition of the $\text{Ca}^{2+}/\text{H}^+$ ionophore A23187 in oat.

Indirect evidence of PT in plants has been also based on the CsA-induced inhibition of programmed cell death (PCD), reviewed by Vianello et al. (2007, 2012). However, the prevention of PCD might depend on CsA binding to cytosolic Cyclophilin A (a ubiquitous enzyme) that drives enzymatic cascades (Lu et al., 2007), linked to oxidative stress (Nigro et al., 2013).

THE MITOCHONDRIAL Ca^{2+} ACCUMULATION IN PLANTS

The PT requires Ca^{2+} accumulation into the mitochondrial matrix (i.e., matrix Ca^{2+} is a permissive factor, although it may not be sufficient *per se*). Calcium transport in isolated plant mitochondria exhibits distinct features. The uptake could be mediated by a low-affinity electrophoretic P_i -dependent symport, with low or no sensitivity to ruthenium red and lanthanides (Dieter and Marme, 1980; Akerman and Moore, 1983; Silva et al., 1992), but also by a uniport mechanism (Zottini and Zannoni, 1993). CsA inhibits mitochondrial Ca^{2+} transport in *Citrus* (de Oliveira et al., 2007), suggesting its synergic effect with PT. A low concentration of matrix free Ca^{2+} (~100 nM) is maintained under steady state, where influx is balanced by an efflux through a yet speculative Na^+ -independent $\text{Ca}^{2+}/\text{H}^+$ antiport mechanism (Nomura and Shiina, 2014). The influx of Ca^{2+} in plant mitochondria is highly variable, depending on species and tissues, or might be even completely absent (Martins and Vercesi, 1985). *In vivo*

Ca^{2+} dynamics have been monitored by fluorescent probes targeted to plant mitochondria (Manzoor et al., 2012; Loro and Costa, 2013). Matrix Ca^{2+} uptake can be induced by abiotic stresses such as heat, oxidative stress, or anoxia, and follows the cytosolic Ca^{2+} pattern (Subbaiah et al., 1998; Logan and Knight, 2003; Schwarzländer et al., 2012; Rikhvanov et al., 2014).

Homologue genes of mammalian mitochondrial Ca^{2+} uniporter (MCU) and its regulatory protein MICU1 have been found in plants (Bick et al., 2012; Stael et al., 2012; Rikhvanov et al., 2014). The MICU1 homologue in *Arabidopsis* (AtMICU) is a negative regulator of mitochondrial Ca^{2+} uptake in root tips, providing strong evidence for the operation of a mitochondrial Ca^{2+} uniporter in plants (Wagner et al., 2015).

THE INVOLVEMENT OF PT/PCD IN PLANT DEVELOPMENT AND STRESS RESPONSES

The physiological role of mitochondrial PT in plants is often related to developmental processes (Reape et al., 2015) and mild environmental stresses, which involve also PCD in many cases. However, the mechanistic link between PT and PCD remains still speculative.

Permeability transition/programmed cell death are fundamental in the selection of damaged cells and in sculpturing new anatomical and morphological structures (Van Hautegeem et al., 2015). Morphological modifications are also needed for adaptive responses to environment (e.g., climate changes) and, more in general, for fitness increase. In particular, *Aponogeton madagascariensis* forms lacunae on its leaves by executing PCD, which is inhibited by CsA, suggesting the involvement of PT (Lord et al., 2013). In aerenchyma formation, lack of oxygen induces stress characterized by mitochondrial PT, ATP depletion, and PCD induction (Yamauchi et al., 2013). Consistently, stressed pea plants show cytochrome *c* release, followed by DNA fragmentation (Sarkar and Gladish, 2012).

Programmed cell death is a common response in plants subjected to abiotic and biotic stresses, which may be linked to the sessile lifestyle, providing a survival strategy for the whole organism. Excess of UV-C stimulates reactive oxygen species (ROS) formation and collapse of $\Delta\Psi$ in *Arabidopsis* mitochondria (Gao et al., 2008). The role of PT has also been described in case of extreme temperatures. In *Arabidopsis* protoplasts, heat stress induces mitochondrial swelling, and $\Delta\Psi$ loss, but these damages are counteracted by a heat shock

TABLE 1 | Characteristics of permeability transition (PT) in plant mitochondria.

Plant material	Ca^{2+} stimulation	CsA inhibition	Sucrose swelling	Cytochrome <i>c</i> release	Reference
Etiolated pea stem	Yes	Yes	No	Not detected	Vianello et al., 1995
Potato tuber	Yes (external)	No	Yes	Yes	Fortes et al., 2001
Potato tuber	Yes	Yes	Yes	Yes	Arpagaus et al., 2002
Oat leaves	Yes (with A23187)	No	Yes	Yes	Curtis and Wolpert, 2002
Wheat roots	Yes	No	Yes	Yes	Virolainen et al., 2002

transcription factor (Zhang et al., 2009). Similarly, ROS and mild heat shock induce mitochondrial PT and the subsequent induction of cell death in *Arabidopsis* protoplasts, which are prevented by the superoxide dismutase analog TEMPOL, by the Ca²⁺ channel-blocker lanthanum chloride, and by CsA (Scott and Logan, 2008). The role of mitochondria in PCD is confirmed in heat-stressed rice protoplasts, where mHSP70 overexpression maintains mitochondrial $\Delta\Psi$, partially inhibits cytochrome *c* release and suppresses PCD by lowering ROS formation (Qi et al., 2011). In wheat cells subjected to freezing, ROS-dependent PCD is associated to $\Delta\Psi$ collapse and cytochrome *c* release (Lyubushkina et al., 2014). In salt-stressed tobacco protoplasts, PCD is triggered by ROS produced by mitochondria, through a process controlled by a CsA-sensitive PT (Lin et al., 2006).

The response to heavy metals requires the participation of mitochondrial PT. In particular, aluminum triggers a high ROS production in peanut, by plasmalemma NADPH oxidases, which induce mitochondrial mediated-PCD (Huang et al., 2014). Consistently, metal phytotoxicity appears to be also mediated by PT in aluminum-treated *Arabidopsis* protoplasts (Li and Xing, 2011) and in cadmium-treated rice roots (Yeh et al., 2007).

Biotic stress, such as pathogen attack, may lead to protoplast shrinkage, mitochondria swelling and cytochrome *c* release. These responses appear to be associated to PCD involvement during the hypersensitive response, a strategy to counteract biotrophic pathogens. The generation of a defensive layer, promoted by PT-induced PCD, has been shown in *Arabidopsis*. In particular, PCD is mediated by a rapid decrease in mitochondrial $\Delta\Psi$, which is partially counteracted by CsA (Yao et al., 2004). Finally, there is evidence on the release of cytochrome *c* induced by elicitors such as harpin or victorin (Curtis and Wolpert, 2002; Krause and Durner, 2004).

THE MOLECULAR STRUCTURE OF PTP

The components involved in PTP formation initially included the voltage-dependent anion channel, the benzodiazepine receptor, the adenine nucleotide translocase and the phosphate carrier. This model has been questioned, since isolated mitochondria from organisms where the expression of each of these proteins has been suppressed still exhibit a PT (Kokoszka et al., 2004; Krauskopf et al., 2006; Baines et al., 2007; Gutiérrez-Aguilar et al., 2014; Šileikytė et al., 2014).

Recent evidence shows that F-ATPase is involved in PTP formation in different species and *taxa* (Bernardi, 2013b; Bonora et al., 2013; Alavian et al., 2014). This enzyme is highly conserved in both prokaryotes and eukaryotes (Hamasur and Glaser, 1992; Heazlewood et al., 2003), consisting in the hydrophilic F₁ and the hydrophobic F₀ sectors, which operate in concert to carry out distinct functions (Antonieli et al., 2014).

The F₁ contains five subunits: α and β forming the catalytic region, while γ , δ , and ϵ are organized in the central stalk. In all eukaryotes these subunits show a high degree of similarity in the sequences (Hamasur and Glaser, 1992; Antonieli et al., 2014; Jiko et al., 2015), while the subunit composition of the F₀ varies

among different *taxa* and species (Hamasur and Glaser, 1992). For details about F-ATPase components in mammals, fungi and algae, see Vázquez-Acevedo et al. (2006), van Lis et al. (2007), Dabbeni-Sala et al. (2012), Antonieli et al. (2014), Lee et al. (2015) and Liu et al. (2015). Specific subunits have been characterized in plants such as sweet potato (Morikami et al., 1992), potato (Dell'Orto et al., 1993; Polgreen et al., 1995) and soybean (Smith et al., 1994).

Plant F₁ includes the classical five-subunit structure (Hamasur and Glaser, 1990, 1992), and also a 24 kDa protein (Li et al., 2012), but the picture of F₀ components remains still incomplete. Several proteins belonging to F₀ have been identified in spinach (Hamasur and Glaser, 1992), potato (Jänsch et al., 1996), rice (Heazlewood et al., 2003), and *Arabidopsis* (Heazlewood et al., 2003; Meyer et al., 2008; Klodmann et al., 2011). As shown by Klodmann et al. (2011) and by Li et al. (2012), F₀ includes subunits a, c, d, 4 (corresponding to subunit b or orf25, Heazlewood et al., 2003), a 6 kDa protein (plant specific), subunit 8 (also called AL6 or orfB, Heazlewood et al., 2003), ATP17 (plant specific) and Oligomycin Sensitivity-Confering Protein (OSCP), sometimes referred to as δ' in plants (Morikami et al., 1992), for some authors belonging to F₁ (Jänsch et al., 1996). Subunit g was found detached from F-ATPase monomer, suggesting that it could represent a dimer-specific protein (Meyer et al., 2008; Klodmann et al., 2011). Plant subunit e sequences have been identified so far only in protein databases for few species (e.g., rice and *Medicago truncatula*).

Multimeric structures of F-ATPase are present in animal, fungi (Davies et al., 2011; Seelert and Dencher, 2011; Liu et al., 2015) and plant mitochondria (Eubel et al., 2003, 2004; Krause et al., 2004; Bultema et al., 2009). Eubel et al. (2003) highlighted the presence of F-ATPase dimers in *Arabidopsis*, potato, bean, and barley. The relative abundance of dimers in plants is low, with respect to the total F-ATPase, and even lower when comparing different organisms (Eubel et al., 2003, 2004).

Rows of F-ATPase dimers in *cristae* seem to be a universal feature of all mitochondria (Davies et al., 2011) that enable the formation of highly curved ridges in *cristae* (Davies et al., 2012). The Inhibitory factor 1 (IF₁) that binds F-ATPase at low pH (Campanella et al., 2008) could favor dimer formation even if it is not clear how it improves dimer stability. The arrangement of F-ATPase in mammals and fungi is different from that of potato, being the angle between monomers in the latter larger ($\sim 115^\circ$) than in the former ($\sim 80^\circ$) (Davies et al., 2011). Interestingly, this correlates with *cristae* morphology observed for many plant mitochondria, where irregular saccular structures with a less convex curvature appear particularly prevalent (Douce, 1985). In aging *Podospora anserina* (Ascomycetes) mitochondria, the IMM is progressively vesiculated, the *cristae* collapse and the F-ATPase dimers are disassembled (Daum et al., 2013). The impairment of ATP synthesis, and the outer membrane rupture by swelling, lead to the release of pro-apoptotic factors and, finally, to cell death.

Animal mitochondria F-ATPase dimers have been shown to act as pores with properties of the PTP (Giorgio et al., 2013). CyPD modulates F-ATPase activity by binding OSCP (Giorgio et al., 2009) and this interaction is favored by P_i, while CsA


displaces CyPD from the enzyme. F-ATPase is inhibited by Bz-423, which binds to OSCP (Cleary et al., 2007). These features are consistent with those observed for PT regulation. Magnesium, Ca^{2+} , adenine nucleotides, membrane potential and matrix pH are also key modulators of both F-ATPase activity and PTP. Electrophysiological experiments, after isolation and insertion of F-ATPase dimers in artificial phospholipid bilayers, showed that the pore activity matches that of PTP-MMC (Giorgio et al., 2013).

The involvement of F-ATPase dimers in PTP formation has been extended and confirmed in yeast and *Drosophila*, even if these organisms show specific differences. In yeast mitochondria the ionophore ETH129 is needed for Ca^{2+} uptake in the matrix and the PT displays a low conductance (around 300 pS). Phosphate acts as an inhibitor of PT, while CsA does not interfere with PTP. Yeast mutants lacking of subunits e and g, which are involved in dimerization, display a striking resistance to PTP opening (Carraro et al., 2014). In *Drosophila* (von Stockum et al., 2015), PTP has been initially identified as mitochondrial Ca^{2+} -induced Ca^{2+} release channel (mCrC). The main differences

between mCrC and mammalian PTP are: (i) absence of swelling; (ii) absence of CsA effect, since no CyPD is present in this species; (iii) sensitivity to rotenone, an inhibitor of Complex I; (iv) inhibition of mCrC by P_i ; (v) low conductance (around 53 pS) of the F-ATPase dimers in artificial bilayer.

Other research groups have also suggested that F-ATPase is involved in pore formation by the channel activity within the c-ring formed by c subunits of F_0 (Bonora et al., 2013; Alavian et al., 2014). Nevertheless, this hypothesis is still under debate, since it does not justify the different pore size observed in bovine, yeast, and *Drosophila*, where similar c-rings are present (Bernardi et al., 2015). Finally, the possible involvement of IF_1 in modulation of PTP through F-ATPase dimerization needs further investigations (Faccenda et al., 2013; Bernardi et al., 2015).

The presence in plants of many common components and features of F-ATPase lead us to speculate that, similarly to mammals, yeast, and *Drosophila*, PT function could be exerted by F-ATPase dimers also in such organisms.



THE EMERGENCE OF PT DURING EVOLUTION

Evolution does not always proceed by adaptations. It may also develop a non-adaptive exaptation/cooptation (pre-adaptation), where the term exaptation/cooptation means a trait evolved to accomplish a specific function (or even no function), which may be then exapted/coopted to perform a novel function (or to acquire a function) (Gould and Vrba, 1982).

It has been suggested that the structure of PTP (as a multicomponent complex, Bernardi, 2013a) may have arisen by a mechanism of molecular exaptation, a phenomenon largely recognized at different levels of complexity (genes, proteins, organs), during evolution (Vianello et al., 2012; Barve and Wagner, 2013). The new model, involving F-ATPase dimer in PTP formation, does not contradict our previous interpretation on its origin, but rather appears to support it further. The dimer appears to be the result of a molecular exaptation/cooptation, where two monomers are assembled to perform an additional function (Figure 1A). In other words, F-ATPase seems to have a “Janus double face”, catalyzing the synthesis of ATP, but in some circumstances preventing such a synthesis (Bernardi et al., 2015). This dimer could even possess a “triple face”, because the dimerization induces also the curvature of the IMM.

The F-ATPase dimer is present in eukaryotes, but not in prokaryotes, because the F-ATPase of the latter is lacking of some crucial subunits (e and g) involved in dimer formation (Antonieli et al., 2014). It is thus reasonable to assume that the dimer/PTP may be arisen after the endosymbiosis between an alpha-proteobacterium and an archaeon (Martin and Müller, 1998). At the beginning, these dimers could have transferred ATP from the endosymbiont to the cytoplasm of the host cell, because the former presumably did not have ATP/ADP transporters. PTP was then maintained to dissipate the protonmotive force, thus regulating both ATP synthesis and exchanges of solutes between the cytoplasm and the mitochondrial matrix.

The presence of F-ATPase dimer has been assessed in different evolutionary divergent eukaryotes, some of which exhibit mitochondrial PT, such as ‘*Unikonts*’ (*Opisthokonts*) and *Plantae* (Arpagaus et al., 2002; Giorgio et al., 2013; Carraro et al., 2014; von Stockum et al., 2015). To understand the phylogenesis of this structure/function, a cladogram has been generated by comparing the ancestral sequences of F_O subunit g from bovine and *Drosophila* (animals), yeast (fungi, *Ascomycetes*), and *Arabidopsis* (*Plantae*) (Figure 1B). The tree suggests an early differentiation at higher taxonomical levels (supergroups): *Plantae* show the highest phylogenetic distance and within the

Opisthokonts, mammals, and insects exhibit similar distances, whereas yeast shows a higher distance. These phylogenetic patterns are consistent with the main evolutionary life tree (e.g., Keeling et al., 2005).

It has been suggested that F-ATPase shows a progressive differentiation along the main steps of evolution. In turn, some features of PTP seem to be occurred independently from changes in ATP synthase. As an example, swelling of mitochondria occurs only in bovine (Bernardi et al., 2015) and in some plants (see Table 1), suggesting that PTP has been differently shaped by exaptation during the evolution. Hence, exaptation leading to PT seems to have occurred in diverse contexts during life history, depending on the molecular characteristics of F-ATPase structure and the specific requirements of the respective *taxa*.

FUTURE DIRECTIONS

The molecular nature of PTP in plants is still elusive. Further structural and functional studies are required to verify if F-ATPase dimers represent the channel associated to PT also in plants. This is needed to understand better the relationship between mitochondrial PT and PCD in plants.

AUTHOR CONTRIBUTIONS

MZ and AV co-supervised the manuscript and co-wrote the article. VC, EP, CP, SP, AB, and EB co-wrote the article. VDC and FB performed the phylogenetic analyses and co-wrote the article.

FUNDING

This work was supported by the Italian Ministry of Education, University and Research (National Research Program PRIN2010CSJX4F) and the European Social Fund (Program 2007/2013).

ACKNOWLEDGMENTS

We thank Paolo Bernardi (University of Padova), Markus Schwarzländer (University of Bonn), and Giovanna Lippe (University of Udine) for their critical reading of the manuscript and Manuela Antoniel (University of Padova) for help in drawing F-ATPase.

REFERENCES

- Akerman, K. E., and Moore, A. L. (1983). Phosphate dependent, ruthenium red insensitive Ca²⁺ uptake in mung bean mitochondria. *Biochem. Biophys. Res. Commun.* 114, 1176–1181. doi: 10.1016/0006-291X(83)90686-1
- Alavian, K. N., Beutner, G., Lazrove, E., Sacchetti, S., Park, H.-A., Licznarski, P., et al. (2014). An uncoupling channel within the c-subunit ring of the F₁F₀ ATP synthase is the mitochondrial permeability transition pore. *Proc. Natl. Acad. Sci. U.S.A.* 111, 10580–10585. doi: 10.1073/pnas.1401591111
- Antonieli, M., Giorgio, V., Fogolari, F., Glick, G. D., Bernardi, P., and Lippe, G. (2014). The oligomycin-sensitivity conferring protein of mitochondrial ATP synthase: emerging new roles in mitochondrial pathophysiology. *Int. J. Mol. Sci.* 15, 7513–7536. doi: 10.3390/ijms15057513
- Arpagaus, S., Rawlyer, A., and Braendle, R. (2002). Occurrence and characteristics of the mitochondrial permeability transition in plants. *J. Biol. Chem.* 277, 1780–1787. doi: 10.1074/jbc.M109416200
- Baines, C. P., Kaiser, R. A., Shelko, T., Craigen, W. J., and Molkenin, D. (2007). Voltage-dependent anion channels are dispensable for mitochondrial-dependent cell death. *Nat. Cell Biol.* 9, 550–555. doi: 10.1038/ncb1575

- Barve, A., and Wagner, A. (2013). A latent capacity for evolutionary innovation through exaptation in metabolic systems. *Nature* 500, 203–206. doi: 10.1038/nature12301
- Bernardi, P. (2013a). “Mitochondrial permeability transition pore,” in *Encyclopedia of Biological Chemistry*, ed. W. J. L. D. Lane (Waltham: Academic Press), 162–167.
- Bernardi, P. (2013b). The mitochondrial permeability transition pore: a mystery solved? *Front. Physiol.* 4:95. doi: 10.3389/fphys.2013.00095
- Bernardi, P., Lisa, F. D., Fogolari, F., and Lippe, G. (2015). From ATP to PTP and back: a dual function for the mitochondrial ATP synthase. *Circ. Res.* 116, 1850–1862. doi: 10.1161/CIRCRESAHA.115.306557
- Bick, A. G., Calvo, S. E., and Mootha, V. K. (2012). Evolutionary diversity of the mitochondrial calcium uniporter. *Science* 336, 886–886. doi: 10.1126/science.1214977
- Bonora, M., Bononi, A., De Marchi, E., Giorgi, C., Lebedzinska, M., Marchi, S., et al. (2013). Role of the c subunit of the FO ATP synthase in mitochondrial permeability transition. *Cell Cycle* 12, 674–683. doi: 10.4161/cc.23599
- Bultema, J. B., Braun, H.-P., Boekema, E. J., and Kouril, R. (2009). Megacomplex organization of the oxidative phosphorylation system by structural analysis of respiratory supercomplexes from potato. *Biochim. Biophys. Acta* 1787, 60–67. doi: 10.1016/j.bbabi.2008.10.010
- Campanella, M., Casswell, E., Chong, S., Farah, Z., Wieckowski, M. R., Abramov, A. Y., et al. (2008). Regulation of mitochondrial structure and function by the F₁F₀-ATPase inhibitor protein, IF1. *Cell Metab.* 8, 13–25. doi: 10.1016/j.cmet.2008.06.001
- Carraro, M., Giorgio, V., Šileikytė, J., Sartori, G., Forte, M., Lippe, G., et al. (2014). Channel formation by yeast F-ATP synthase and the role of dimerization in the mitochondrial permeability transition. *J. Biol. Chem.* 289, 15980–15985. doi: 10.1074/jbc.C114.559633
- Cleary, J., Johnson, K. M., Opipari, A. W. Jr., and Glick, G. D. (2007). Inhibition of the mitochondrial F₁F₀-ATPase by ligands of the peripheral benzodiazepine receptor. *Bioorg. Med. Chem. Lett.* 17, 1667–1670. doi: 10.1016/j.bmlc.2006.12.102
- Crompton, M., Ellinger, H., and Costi, A. (1988). Inhibition by cyclosporin A of a Ca²⁺-dependent pore in heart mitochondria activated by inorganic phosphate and oxidative stress. *Biochem. J.* 255, 357–360.
- Curtis, M. J., and Wolpert, T. J. (2002). The oat mitochondrial permeability transition and its implication in victorin binding and induced cell death. *Plant J.* 29, 295–312. doi: 10.1046/j.0960-7412.2001.01213.x
- Dabbeni-Sala, F., Rai, A. K., and Lippe, G. (2012). “F₁F₀ ATP Synthase: a fascinating challenge for proteomics,” in *Proteomics: Human Diseases and Protein Functions*, eds T.-K. Man and R. J. Flores (Rijeka: InTech), 161–188.
- Daum, B., Walter, A., Horst, A., Osiewacz, H. D., and Kühlbrandt, W. (2013). Age-dependent dissociation of ATP synthase dimers and loss of inner-membrane cristae in mitochondria. *Proc. Natl. Acad. Sci. U.S.A.* 110, 15301–15306. doi: 10.1073/pnas.1305462110
- Davies, K. M., Anselmi, C., Wittig, I., Faraldo-Gómez, J. D., and Kühlbrandt, W. (2012). Structure of the yeast F₁F₀-ATP synthase dimer and its role in shaping the mitochondrial cristae. *Proc. Natl. Acad. Sci. U.S.A.* 109, 13602–13607. doi: 10.1073/pnas.1204593109
- Davies, K. M., Strauss, M., Daum, B., Kief, J. H., Osiewacz, H. D., Rycovska, A., et al. (2011). Macromolecular organization of ATP synthase and complex I in whole mitochondria. *Proc. Natl. Acad. Sci. U.S.A.* 108, 14121–14126. doi: 10.1073/pnas.1103621108
- Dell’Orto, P., Moenne, A., Vincent Graves, P., and Jordana, X. (1993). The potato mitochondrial ATP synthase subunit 9: gene structure, RNA editing and partial protein sequence. *Plant Sci.* 88, 45–53. doi: 10.1016/0168-9452(93)90108-C
- de Oliveira, H. C., Saviani, E. E., de Oliveira, J. F. P., and Salgado, I. (2007). Cyclosporin A inhibits calcium uptake by *Citrus sinensis* mitochondria. *Plant Sci.* 172, 665–670. doi: 10.1016/j.plantsci.2006.12.002
- Dieter, P., and Marme, D. (1980). Ca²⁺ transport in mitochondrial and microsomal fractions from higher plants. *Planta* 150, 1–8. doi: 10.1007/BF00385606
- Douce, R. (1985). *Mitochondria in Higher Plants: Structure, Function, and Biogenesis*. New York, NY: Elsevier.
- Edgar, R. C. (2004). MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* 5:113. doi: 10.1186/1471-2105-5-113
- Eubel, H., Heinemeyer, J., Sunderhaus, S., and Braun, H.-P. (2004). Respiratory chain supercomplexes in plant mitochondria. *Plant Physiol. Biochem.* 42, 937–942. doi: 10.1016/j.plaphy.2004.09.010
- Eubel, H., Jänsch, L., and Braun, H.-P. (2003). New insights into the respiratory chain of plant mitochondria. Supercomplexes and a unique composition of complex II. *Plant Physiol.* 133, 274–286. doi: 10.1104/pp.103.024620
- Faccenda, D., Tan, C. H., Seraphim, A., Duchon, M. R., and Campanella, M. (2013). IF1 limits the apoptotic-signalling cascade by preventing mitochondrial remodeling. *Cell Death Differ.* 20, 686–697. doi: 10.1038/cdd.2012.163
- Fortes, F., Castilho, R. F., Catisti, R., Carnieri, E. G. S., and Vercesi, A. E. (2001). Ca²⁺ induces a cyclosporin A-insensitive permeability transition pore in isolated potato tuber mitochondria mediated by reactive oxygen species. *J. Bioenerg. Biomembr.* 33, 43–51. doi: 10.1023/A:1005672623709
- Gao, C., Xing, D., Li, L., and Zhang, L. (2008). Implication of reactive oxygen species and mitochondrial dysfunction in the early stages of plant programmed cell death induced by ultraviolet-C overexposure. *Planta* 227, 755–767. doi: 10.1007/s00425-007-0654-4
- Giorgio, V., Bisetto, E., Soriano, M. E., Dabbeni-Sala, F., Basso, E., Petronilli, V., et al. (2009). Cyclophilin D modulates mitochondrial F₀F₁-ATP synthase by interacting with the lateral stalk of the complex. *J. Biol. Chem.* 284, 33982–33988. doi: 10.1074/jbc.M109.020115
- Giorgio, V., von Stockum, S., Antoniel, M., Fabbro, A., Fogolari, F., Forte, M., et al. (2013). Dimers of mitochondrial ATP synthase form the permeability transition pore. *Proc. Natl. Acad. Sci. U.S.A.* 110, 5887–5892. doi: 10.1073/pnas.1217823110
- Gould, S. J., and Vrba, E. S. (1982). Exaptation – A missing term in the science of form. *Paleobiology* 8, 4–15.
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321. doi: 10.1093/sysbio/syq010
- Gutiérrez-Aguilar, M., Douglas, D. L., Gibson, A. K., Domeier, T. L., Molkenin, J. D., and Baines, C. P. (2014). Genetic manipulation of the cardiac mitochondrial phosphate carrier does not affect permeability transition. *J. Mol. Cell. Cardiol.* 72, 316–325. doi: 10.1016/j.yjmcc.2014.04.008
- Halestrap, A. P., and Davidson, A. M. (1990). Inhibition of Ca²⁺-induced large-amplitude swelling of liver and heart mitochondria by cyclosporin is probably caused by the inhibitor binding to mitochondrial-matrix peptidyl-prolyl *cis-trans* isomerase and preventing it interacting with the adenine nucleotide translocase. *Biochem. J.* 268, 153–160. doi: 10.1042/bj2680153
- Hamasur, B., and Glaser, E. (1990). F₀F₁-ATPase of plant mitochondria: isolation and polypeptide composition. *Biochem. Biophys. Res. Commun.* 170, 1352–1358. doi: 10.1016/0006-291X(90)90543-V
- Hamasur, B., and Glaser, E. (1992). Plant mitochondrial F₀F₁ ATP synthase. Identification of the individual subunits and properties of the purified spinach leaf mitochondrial ATP synthase. *Eur. J. Biochem.* 205, 409–416. doi: 10.1111/j.1432-1033.1992.tb16794.x
- Haworth, R. A., and Hunter, D. R. (1979). The Ca²⁺-induced membrane transition in mitochondria: II. Nature of the Ca²⁺ trigger site. *Arch. Biochem. Biophys.* 195, 460–467. doi: 10.1016/0003-9861(79)90372-2
- Heazlewood, J. L., Whelan, J., and Millar, A. H. (2003). The products of the mitochondrial *orf25* and *orfB* genes are F₀ components in the plant F₁F₀ ATP synthase. *FEBS Lett.* 540, 201–205. doi: 10.1016/S0014-5793(03)00264-3
- Huang, W., Yang, X., Yao, S., LwinOo, T., He, H., Wang, A., et al. (2014). Reactive oxygen species burst induced by aluminum stress triggers mitochondria-dependent programmed cell death in peanut root tip cells. *Plant Physiol. Biochem.* 82, 76–84. doi: 10.1016/j.plaphy.2014.03.037
- Hunter, D. R., and Haworth, R. A. (1979a). The Ca²⁺-induced membrane transition in mitochondria: I. The protective mechanisms. *Arch. Biochem. Biophys.* 195, 453–459. doi: 10.1016/0003-9861(79)90371-0
- Hunter, D. R., and Haworth, R. A. (1979b). The Ca²⁺-induced membrane transition in mitochondria: III. Transitional Ca²⁺ release. *Arch. Biochem. Biophys.* 195, 468–477. doi: 10.1016/0003-9861(79)90373-4
- Jänsch, L., Kruff, V., Schmitz, U. K., and Braun, H.-P. (1996). New insights into the composition, molecular mass and stoichiometry of the protein complexes of plant mitochondria. *Plant J.* 9, 357–368. doi: 10.1046/j.1365-313X.1996.09030357.x

- Jiko, C., Davies, K. M., Shinzawa-Ittoh, K., Tani, K., Maeda, S., Mills, D. J., et al. (2015). Bovine F₁F₀ ATP synthase monomers bend the lipid bilayer in 2D membrane crystals. *Elife* 4, e06119. doi: 10.7554/eLife.06119
- Keeling, P. J., Burger, G., Durnford, D. G., Lang, B. F., Lee, R. W., Pearlman, R. E., et al. (2005). The tree of eukaryotes. *Trends Ecol. Evol.* 20, 670–676. doi: 10.1016/j.tree.2005.09.005
- Klodmann, J., Senkler, M., Rode, C., and Braun, H.-P. (2011). Defining the protein complex proteome of plant mitochondria. *Plant Physiol.* 157, 587–598. doi: 10.1104/pp.111.182352
- Kokoszka, J. E., Waymire, K. G., Levy, S. E., Sligh, J. E., Cai, J., Jones, D. P., et al. (2004). The ADP/ATP translocator is not essential for the mitochondrial permeability transition pore. *Nature* 427, 461–465. doi: 10.1038/nature02229
- Krause, F., Reifschneider, N. H., Vocke, D., Seelert, H., Rexroth, S., and Dencher, N. A. (2004). “Respirasome”-like supercomplexes in green leaf mitochondria of spinach. *J. Biol. Chem.* 279, 48369–48375. doi: 10.1074/jbc.M406085200
- Krause, M., and Durner, J. (2004). Harpin inactivates mitochondria in *Arabidopsis* suspension cells. *Mol. Plant Microbe Interact.* 17, 131–139. doi: 10.1094/MPMI.2004.17.2.131
- Krauskopf, A., Eriksson, O., Craigen, W. J., Forte, M. A., and Bernardi, P. (2006). Properties of the permeability transition in VDAC1^{-/-} mitochondria. *Biochim. Biophys. Acta* 1757, 590–595. doi: 10.1016/j.bbabi.2006.02.007
- Lee, J., Ding, S., Walpole, T. B., Holding, A. N., Montgomery, M. G., Fearnley, I. M., et al. (2015). Organization of subunits in the membrane domain of the bovine F₁-ATPase revealed by covalent cross-linking. *J. Biol. Chem.* 290, 13308–13320. doi: 10.1074/jbc.M115.645283
- Li, L., Carrie, C., Nelson, C., Whelan, J., and Millar, A. H. (2012). Accumulation of newly synthesized F₁ in vivo in *Arabidopsis* mitochondria provides evidence for modular assembly of the plant F₁F₀ ATP synthase. *J. Biol. Chem.* 287, 25749–25757. doi: 10.1074/jbc.M112.373506
- Li, Z., and Xing, D. (2011). Mechanistic study of mitochondria-dependent programmed cell death induced by aluminium phytotoxicity using fluorescence techniques. *J. Exp. Bot.* 62, 331–343. doi: 10.1093/jxb/Erq279
- Lin, J., Wang, Y., and Wang, G. (2006). Salt stress-induced programmed cell death in tobacco protoplasts is mediated by reactive oxygen species and mitochondrial permeability transition pore status. *J. Plant Physiol.* 163, 731–739. doi: 10.1016/j.jplph.2005.06.016
- Liu, S., Charlesworth, T. J., Bason, J. V., Montgomery, M. G., Harbour, M. E., Fearnley, I. M., et al. (2015). The purification and characterization of ATP synthase complexes from the mitochondria of four fungal species. *Biochem. J.* 468, 167–175. doi: 10.1042/BJ20150197
- Logan, D. C., and Knight, M. R. (2003). Mitochondrial and cytosolic calcium dynamics are differentially regulated in plants. *Plant Physiol.* 133, 21–24. doi: 10.1104/pp.103.026047
- Lord, C. E. N., Dauphinee, A. N., Watts, R. L., and Gunawardena, A. H. L. A. N. (2013). Unveiling interactions among mitochondria, caspase-like proteases, and the actin cytoskeleton during plant programmed cell death (PCD). *PLoS ONE* 8:e57110. doi: 10.1371/journal.pone.0057110
- Loro, G., and Costa, A. (2013). Imaging of mitochondrial and nuclear Ca²⁺ dynamics in *Arabidopsis* roots. *Cold Spring Harb. Protoc.* 8, 781–785. doi: 10.1101/pdb.prot073049
- Lu, K. P., Finn, G., Lee, T. H., and Nicholson, L. K. (2007). Prolyl cis-trans isomerization as a molecular timer. *Nat. Chem. Biol.* 3, 619–629. doi: 10.1038/nchembio.2007.35
- Lyubushkina, I. V., Grabelnykh, O. I., Pobezhimova, T. P., Stepanov, A. V., Fedyaeva, A. V., Fedoseeva, I. V., et al. (2014). Winter wheat cells subjected to freezing temperature undergo death process with features of programmed cell death. *Protoplasma* 251, 615–623. doi: 10.1007/s00709-013-0562-3
- Manzoor, H., Chiltz, A., Madani, S., Vatsa, P., Schoefs, B., Pugin, A., et al. (2012). Calcium signatures and signaling in cytosol and organelles of tobacco cells induced by plant defense elicitors. *Cell Calcium* 51, 434–444. doi: 10.1016/j.ceca.2012.02.006
- Martin, W., and Müller, M. (1998). The hydrogen hypothesis for the first eukaryote. *Nature* 392, 37–41. doi: 10.1038/32096
- Martins, I. S., and Vercesi, A. E. (1985). Some characteristics of Ca²⁺ transport in plant mitochondria. *Biochem. Biophys. Res. Commun.* 129, 943–948. doi: 10.1016/0006-291x(85)91982-5
- Meyer, E. H., Taylor, N. L., and Millar, A. H. (2008). Resolving and identifying protein components of plant mitochondrial respiratory complexes using three dimensions of gel electrophoresis. *J. Proteome Res.* 7, 786–794. doi: 10.1021/pr700595p
- Mitchell, P. (1961). Coupling of phosphorylation to electron and hydrogen transfer by a chemi-osmotic type of mechanism. *Nature* 191, 144–148. doi: 10.1038/191144a0
- Morikami, A., Aiso, K., Asahi, T., and Nakamura, K. (1992). The delta-subunit of higher plant six-subunit mitochondrial F₁-ATPase is homologous to the delta-subunit of animal mitochondrial F₁-ATPase. *J. Biol. Chem.* 267, 72–76.
- Nigro, P., Pompilio, G., and Capogrossi, M. C. (2013). Cyclophilin A: a key player for human disease. *Cell Death Dis.* 4, e888. doi: 10.1038/cddis.2013.410
- Nomura, H., and Shiina, T. (2014). Calcium signaling in plant endosymbiotic organelles: mechanism and role in physiology. *Mol. Plant* 7, 1094–1104. doi: 10.1093/Mp/Ssu020
- Polgreen, K. E., Featherstone, J., Willis, A. C., and Harris, D. A. (1995). Primary structure and properties of the inhibitory protein of the mitochondrial ATPase (H⁺-ATP synthase) from potato. *Biochim. Biophys. Acta* 1229, 175–180. doi: 10.1016/0005-2728(94)00193-9
- Qi, Y., Wang, H., Zou, Y., Liu, C., Liu, Y., Wang, Y., et al. (2011). Over-expression of mitochondrial heat shock protein 70 suppresses programmed cell death in rice. *FEBS Lett.* 585, 231–239. doi: 10.1016/j.febslet.2010.11.051
- Raaflaub, J. (1953a). Mechanism of adenosinetriphosphate as cofactor of isolated mitochondria. *Helv. Physiol. Pharmacol. Acta* 11, 157–165.
- Raaflaub, J. (1953b). Swelling of isolated mitochondria of the liver and their susceptibility to physicochemical influences. *Helv. Physiol. Pharmacol. Acta* 11, 142–156.
- Reape, T. J., Kacprzyk, J., Brogan, N., Sweetlove, L., and McCabe, P. F. (2015). Mitochondrial markers of programmed cell death in *Arabidopsis thaliana*. *Methods Mol. Biol.* 1305, 211–221. doi: 10.1007/978-1-4939-2639-8_15
- Rikhvanov, E. G., Fedoseeva, I. V., Pyatrikas, D. V., Borovskii, G. B., and Voinikov, V. K. (2014). Role of mitochondria in the operation of calcium signaling system in heat-stressed plants. *Russ. J. Plant Physiol.* 61, 141–153. doi: 10.1134/S1021443714020125
- Sarkar, P., and Gladish, D. K. (2012). Hypoxic stress triggers a programmed cell death pathway to induce vascular cavity formation in *Pisum sativum* roots. *Physiol. Plant.* 146, 413–426. doi: 10.1111/j.1399-3054.2012.01632.x
- Schwarzländer, M., Logan, D. C., Johnston, I. G., Jones, N. S., Meyer, A. J., Fricker, M. D., et al. (2012). Pulsing of membrane potential in individual mitochondria: a stress-induced mechanism to regulate respiratory bioenergetics in *Arabidopsis*. *Plant Cell* 24, 1188–1201. doi: 10.1105/tpc.112.096438
- Scott, I., and Logan, D. C. (2008). Mitochondrial morphology transition is an early indicator of subsequent cell death in *Arabidopsis*. *New Phytol.* 177, 90–101. doi: 10.1111/j.1469-8137.2007.02255.x
- Seelert, H., and Dencher, N. A. (2011). ATP synthase superassemblies in animals and plants: two or more are better. *Biochim. Biophys. Acta* 1807, 1185–1197. doi: 10.1016/j.bbabi.2011.05.023
- Šileikytė, J., Blachly-Dyson, E., Sewell, R., Carpi, A., Menabò, R., Lisa, F. D., et al. (2014). Regulation of the mitochondrial permeability transition pore by the outer membrane does not involve the peripheral benzodiazepine receptor (Translocator Protein of 18 kDa (TSPO)). *J. Biol. Chem.* 289, 13769–13781. doi: 10.1074/jbc.M114.549634
- Silva, M. A. P., Carnieri, E. G. S., and Vercesi, A. E. (1992). Calcium-transport by corn mitochondria – evaluation of the role of phosphate. *Plant Physiol.* 98, 452–457. doi: 10.1104/Pp.98.2.452
- Smith, M. K., Day, D. A., and Whelan, J. (1994). Isolation of a novel soybean gene encoding a mitochondrial ATP synthase subunit. *Arch. Biochem. Biophys.* 313, 235–240. doi: 10.1006/abbi.1994.1382
- Stael, S., Wurzing, B., Mair, A., Mehlmer, N., Vothknecht, U. C., and Teige, M. (2012). Plant organellar calcium signalling: an emerging field. *J. Exp. Bot.* 63, 1525–1542. doi: 10.1093/jxb/err394
- Subbiah, C. C., Bush, D. S., and Sachs, M. M. (1998). Mitochondrial contribution to the anoxic Ca²⁺ signal in maize suspension-cultured cells. *Plant Physiol.* 118, 759–771. doi: 10.1104/Pp.118.3.759
- Szabó, I., and Zoratti, M. (1992). The mitochondrial megachannel is the permeability transition pore. *J. Bioenerg. Biomembr.* 24, 111–117. doi: 10.1007/BF00769537

- Van Hautegeem, T., Waters, A. J., Goodrich, J., and Nowack, M. K. (2015). Only in dying life: programmed cell death during plant development. *Trends Plant Sci.* 20, 102–113. doi: 10.1016/j.tplants.2014.10.003
- van Lis, R., Mendoza-Hernández, G., Groth, G., and Atteia, A. (2007). New insights into the unique structure of the F₀F₁-ATP synthase from the chlamydomonad algae *Polytomella* sp. and *Chlamydomonas reinhardtii*. *Plant Physiol.* 144, 1190–1199. doi: 10.1104/pp.106.094060
- Vázquez-Acevedo, M., Cardol, P., Cano-Estrada, A., Lapaille, M., Remacle, C., and González-Halphen, D. (2006). The mitochondrial ATP synthase of chlorophycean algae contains eight subunits of unknown origin involved in the formation of an atypical stator-stalk and in the dimerization of the complex. *J. Bioenerg. Biomembr.* 38, 271–282. doi: 10.1007/s10863-006-9046-x
- Vianello, A., Casolo, V., Petrusa, E., Peresson, C., Patui, S., Bertolini, A., et al. (2012). The mitochondrial permeability transition pore (PTP) — An example of multiple molecular exaptation? *Biochim. Biophys. Acta* 1817, 2072–2086. doi: 10.1016/j.bbabi.2012.06.620
- Vianello, A., Macri, F., Braidot, E., and Mokhova, E. (1995). Effect of cyclosporin A on energy coupling in pea stem mitochondria. *FEBS Lett.* 371, 258–260. doi: 10.1016/0014-5793(95)00897-1
- Vianello, A., Zancani, M., Peresson, C., Petrusa, E., Casolo, V., Krajinakova, J., et al. (2007). Plant mitochondrial pathway leading to programmed cell death. *Physiol. Plant.* 129, 242–252. doi: 10.1111/j.1399-3054.2006.00767.x
- Virolainen, E., Blokhina, O., and Fagerstedt, K. (2002). Ca²⁺-induced high amplitude swelling and cytochrome c release from wheat (*Triticum aestivum* L.) mitochondria under anoxic stress. *Ann. Bot.* 90, 509–516. doi: 10.1093/Aob/Mcf221
- von Stockum, S., Giorgio, V., Trevisan, E., Lippe, G., Glick, G. D., Forte, M. A., et al. (2015). F-ATPase of *Drosophila melanogaster* forms 53-pS channels responsible for mitochondrial Ca²⁺-induced Ca²⁺ release. *J. Biol. Chem.* 290, 4537–4544. doi: 10.1074/jbc.C114.629766
- Wagner, S., Behera, S., De Bortoli, S., Logan, D. C., Fuchs, P., Carraretto, L., et al. (2015). The EF-hand Ca²⁺ binding protein MICU choreographs mitochondrial Ca²⁺ dynamics in *Arabidopsis*. *Plant Cell* (in press). doi: 10.1105/tpc.15.00509
- Yamauchi, T., Shimamura, S., Nakazono, M., and Mochizuki, T. (2013). Aerenchyma formation in crop species: a review. *Field Crops Res.* 152, 8–16. doi: 10.1016/j.fcr.2012.12.008
- Yao, N., Eisfelder, B. J., Marvin, J., and Greenberg, J. T. (2004). The mitochondrion – an organelle commonly involved in programmed cell death in *Arabidopsis thaliana*. *Plant J.* 40, 596–610. doi: 10.1111/j.1365-313X.2004.02239.x
- Yeh, C.-M., Chien, P.-S., and Huang, H.-J. (2007). Distinct signalling pathways for induction of MAP kinase activities by cadmium and copper in rice roots. *J. Exp. Bot.* 58, 659–671. doi: 10.1093/jxb/erl240
- Zhang, L., Li, Y., Xing, D., and Gao, C. (2009). Characterization of mitochondrial dynamics and subcellular localization of ROS reveal that HsfA2 alleviates oxidative damage caused by heat stress in *Arabidopsis*. *J. Exp. Bot.* 60, 2073–2091. doi: 10.1093/jxb/erp078
- Zottini, M., and Zannoni, D. (1993). The use of fura-2 fluorescence to monitor the movement of free calcium ions into the matrix of plant mitochondria (*Pisum sativum* and *Helianthus tuberosus*). *Plant Physiol.* 102, 573–578.

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Zancani, Casolo, Petrusa, Peresson, Patui, Bertolini, De Col, Braidot, Boscutti and Vianello. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.