



# Phytobase: a tool for the integrated synusial approach of vegetation classification at regional to national scale

François Gillet

## ► To cite this version:

François Gillet. Phytobase: a tool for the integrated synusial approach of vegetation classification at regional to national scale. Chytry M; Zeleny D; Hettengergerova E. Understanding broad-scale vegetation patterns, Jul 2015, Brno, Czech Republic. 58th Symposium of the International Association for Vegetation Science (IAVS) – Understanding broad-scale vegetation patterns, Brno, République Tchèque, pp.136, 2015. <hal-01211283>

**HAL Id: hal-01211283**

**<https://hal.archives-ouvertes.fr/hal-01211283>**

Submitted on 4 Oct 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Phytobase: a tool for the integrated synusial approach of vegetation classification at regional to national scale

François Gillet<sup>1,2,3</sup>

<sup>1</sup> UMR 6249 Chrono-environnement, Université de Bourgogne Franche-Comté – CNRS, Besançon, France, [francois.gillet@univ-fcomte.fr](mailto:francois.gillet@univ-fcomte.fr)

<sup>2</sup> Ecological Systems laboratory, Ecole Polytechnique Fédérale de Lausanne, Ecublens, Switzerland

<sup>3</sup> Soil and Vegetation laboratory, University of Neuchâtel, Neuchâtel, Switzerland

## Integrated Synusial Phytosociology



**Integrated Synusial Phytosociology (ISP)** considers vegetation as organized in nested hierarchical levels, each of them being potentially described in a classification system (1, 2, 3).

The first, finest, level is the **synusia**, a concrete plant community composed of plant species co-occurring in a local similar environment, with comparable size and habitat use. *Phytobase* distinguishes three categories of synusiae, i.e. tree, shrub, herb and moss synusiae, classified into **elementary syntaxa**.


Synusiae are integrated into **phytocoenoses**, the second level of vegetation description, based on a list of synusial elementary syntaxa co-occurring in a given phytocoenosis.

Phytocoenoses are classified into **elementary coenotaxa** according to their synusial composition (elementary syntaxa or alliances).


- (1) Barkman J.J. 1980. Synusial approaches to classification. In: Whittaker R.H. (ed.), *Classification of plant communities*, pp. 111–165. W. Junk, The Hague, NL.
- (2) Gillet F., de Foucault B. & Julve P. 1991. La phytosociologie synusiale intégrée : objets et concepts. *Candollea* 46: 315–340.
- (3) Gillet F. & Galland J.D. 1996. Integrated synusial phytosociology: some notes on a new, multiscalar approach to vegetation analysis. *Journal of Vegetation Science* 7: 13–18.




## Synusial plot record



## Phytosociological diagnosis



## Ecological diagnosis



Phytosociological and ecological diagnostic of a synusial relevé in *Phytobase* 8.3.

## Main applications


*Phytobase* has been applied to various ecosystems (wood-pastures, forests, floodplains, wetlands and grasslands) in many regions, mostly in Switzerland and in France. ISP and *Phytobase* have proved particularly useful when studying structurally and dynamically complex vegetation, such as wooded pastures and alluvial forests, which are poorly described by the Braun-Blanquet approach.

## Phytobase: a relational database for ISP

*Phytobase* is a relational database management system (RDBMS) devoted to the management of vegetation survey data, following the concepts of the ISP. This application has been developed in the 4D environment (4) from the 1990s at the University of Neuchâtel, at EPFL and now at the University of Besançon.

*Phytobase* is freely available from the *Tela-botanica* website (5), including a sample of relevés and a user guide (6). It is used by more than 200 registered users in Europe and registered in the GIVD metadatabase (7).

The current standalone application (version 8.3 for Windows or Mac OS X, built with 4D v13) has a user interface in French language but an English version is planned to allow a broader diffusion to the international community of vegetation scientists.




- (4) <http://www.4d.com>
- (5) [http://www.tela-botanica.org/page\\_liste\\_projets?id\\_projets=18&act=documents&id\\_repertoire=16428](http://www.tela-botanica.org/page_liste_projets?id_projets=18&act=documents&id_repertoire=16428)
- (6) Gillet F. 2014. *Guide d'utilisation de Phytobase (version 8.3), base de données phytosociologiques*. Université de Franche-Comté, Besançon.
- (7) <http://www.givd.info/ID/EU-00-015>

## Phytosociological and ecological diagnostic tools

A comprehensive diagnostic of synusial plot records is implemented in *Phytobase*, including:

- **Synsystematic diagnosis:** assignment to a class, an order and an alliance based on the relative number or cover of characteristic and differential species according to Julve's *baseveg* classification system (8, 9);
- **Syntaxonomic diagnosis:** assignment to an elementary syntaxon stored in the database, based on the combination of various indices (10);
- **Ecological diagnosis:** ecological indicator values (according to Landolt (11), Ellenberg or Julve), life forms, CSR strategies, taxonomic diversity indices, pastoral value, etc.



Centroid relevé and homogeneity assessment of a synusial elementary syntaxon in *Phytobase* 8.3.

Each elementary syntaxon or coenotaxon is described by the centroid of all plot records that were assigned to it. This centroid relevé includes fidelity indices (e.g., Phi (12), IndVal (13)) for species or elementary syntaxa, used in phytosociological diagnoses. An assessment of compositional homogeneity, taking or not into account dominance, is also implemented.

- (8) Julve P. 1993. Synopsis phytosociologique de la France (communautés de plantes vasculaires). *Lejeunia* N.S. 140: 1-160.
- (9) Julve P. 1998-2015. *Baseveg - Répertoire synonymique des unités phytosociologiques de France*. Available online: <http://perso.wanadoo.fr/philippe.julve/catminat.htm>
- (10) Tichý L. 2005. New similarity indices for the assignment of relevés to the vegetation units of an existing phytosociological classification. *Plant Ecology* 179: 67-72.
- (11) Landolt E. et al. 2010. *Flora indicativa - Ecological indicator values and biological attributes of the flora of Switzerland and the Alps*. Haupt, Bern.
- (12) Chytrý M., Tichý L., Holt J. & Botta-Dukát Z. 2002. Determination of diagnostic species with statistical fidelity measures. *Journal of Vegetation Science* 13: 79-90.
- (13) Dufrene M. & Legendre P. 1997. Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological Monographs* 67: 345-366.