


Structural characterization of α -amino acid complexes of molybdates: a spectroscopic and DFT

View metadata, citation and similar papers at core.ac.uk

brought to you by  CORE

provided by Archivio della Ricerca - Università di Pisa

Lorenzo Biancalana,^a Marco Bortoluzzi,^b Claudia Forte,^c Fabio Marchetti^a and Guido Pampaloni^{*a}

Received 10th November 2014
Accepted 24th December 2014

DOI: 10.1039/c4ra14236e

www.rsc.org/advances

The reactions of L-enantiopure α -amino acids (aaH) with Na_2MoO_4 led to the high-yield isolation of Mo(VI) complexes of general formula $\text{Mo}_2\text{O}_4(\text{OH})_4(\text{aaH})$. A similar outcome was achieved by using $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$ in the place of Na_2MoO_4 . Solid-state IR and NMR spectroscopy indicated the presence of *cis*- MoO_2 units linked by a zwitterionic amino acid ligand, *via* a bidentate bridging coordination through the carboxylate group. Thus possible dinuclear and polynuclear structures are proposed on the basis of DFT calculations.

Introduction

Molybdenum is found in living organisms at the active site of several enzymes involved in redox reactions, whose clear understanding is still a big challenge for science.¹ The main bio-available form of molybdenum is the tetrahedral molybdate ion, *i.e.* $[\text{MoO}_4]^{2-}$, which is stable and inert in non-acidic solutions,² and presumed to have played a crucial role in biological evolution.³

The study of the interaction between molybdenum compounds and naturally-occurring molecules in aqueous solution is intended to contribute to the coordination chemistry of Mo(VI) in water and the related concerns, including the biological ones.⁴

On the other hand, α -amino acids are good candidates as potential ligands, indeed their coordination to a variety of metal ions has been extensively investigated.⁵ It is noteworthy that the association of natural, chiral α -amino acids with water soluble, biocompatible metal compounds may be a privileged way to obtain potential catalysts for environmentally friendly asymmetric synthesis.⁶

In this framework, the chemistry of a variety of Mo(VI) species with α -amino acids in aqueous medium has aroused huge interest.⁷ In particular, a series of studies were carried out on complexation equilibria involving $[\text{MoO}_4]^{2-}$.⁸ Working

in mildly acidic conditions ($4.5 < \text{pH} < 7$), complex formation was observed for those α -amino acids with an ionisable side-chain, however isolation in the solid state was limited to a few cases.^{9,10} On the other hand, the products of the molybdate/ α -amino acid interaction could be isolated more favourably at lower pH values ($\text{pH} = 0-2$).^{5a,9,11}

The exploration of this piece of chemistry bears an important drawback: the products are insoluble materials whose crystallographic characterization still remains a hard, not achieved task. As a matter of fact, the reported characterisations relied on limited analytical and spectroscopic (IR) data, and for this reason the structure of the Mo-oxido backbone and the amino acid coordination mode have not been clearly elucidated hitherto.^{5a,9,11}

In the light of this preface, we decided to deal with the reactions of molybdates, *i.e.* Na_2MoO_4 and $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$, with a selection of α -amino acids (glycine, L-phenylalanine, L-leucine, L-methionine, L-proline and *N,N*-dimethyl-L-phenylalanine), in acidic aqueous solution. We present herein the results of our investigation: a detailed structural characterization of the products has been possible for the first time, based on solid-state spectroscopic techniques (IR and NMR) and DFT calculations.

Results and discussion

Molybdenum(VI)/amino acid complexes (1–6, see Scheme 1) were prepared from aqueous solutions of α -amino acids (aaH)

^aUniversity of Pisa, Dipartimento di Chimica e Chimica Industriale, Via Moruzzi 3, I-56124 Pisa, Italy. E-mail: guido.pampaloni@unipi.it; Web: <http://www.dcci.unipi.it/guido-pampaloni.html>; fabmar@dcci.unipi.it; Tel: +39-050-2219219

^bCa'Foscari University of Venice, Dipartimento di Scienze Molecolari e Nanosistemi, Dorsoduro 2137, I-30123 Venezia, Italy


^cIstituto di Chimica dei Composti Organometallici, CNR, Area della Ricerca di Pisa, Via Moruzzi 1, I-56124 Pisa, Italy

† Electronic supplementary information (ESI) available: Experimental details; ¹³C CP-MAS NMR and IR spectra of 1–6 and related α -amino acids and α -ammonium acid nitrates. The Cartesian coordinates of optimized geometries are collected in a separated .xyz file. See DOI: 10.1039/c4ra14236e

α -amino acid (aaH)	complex $\text{Mo}_2\text{O}_4(\text{OH})_4(\text{aaH})$
Glycine (GlyH)	1
L-Phenylalanine (PheH)	2
L-Leucine (LeuH)	3
L-Methionine (MetH)	4
L-Proline (ProH)	5
<i>N,N</i> -Dimethyl-L-phenylalanine (dmPheH)	6


Scheme 1 Synthesis of compounds 1–6.

and a molybdate salt, either Na_2MoO_4 or $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24}$, with $\text{Mo}/\text{aaH} = 2 : 1$ (eqn (1)). The resulting mixtures were acidified to $\text{pH} = 2$, then the precipitation of neutral colourless solids **1–6** followed in variable times (73–92% yields).¹²



The reactions were investigated on varying the reaction conditions (pH , Mo/aaH ratio, temperature, nature of molybdate salt and acid; see ESI[†]). As a matter of fact, the formation of **1–6** was generally found to be selective and high yielded. Compounds **1–6** were isolated in good to high yields even by using a large excess of aaH. The synthesis of **1** was effectively performed also at $\text{pH} \text{ ca. } 0$ (HNO_3 , $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, room temperature) and at $\text{pH} = 1$ (HCl , $(\text{NH}_4)_2\text{MoO}_4$, *ca.* 100 °C), in spite of the pH -dependent equilibria affecting the molybdate ion.² In other terms, the same products were isolated from distinct solutions containing different cations (Na^+ or NH_4^+) and anions (Cl^- or NO_3^-); in addition, chlorine was not incorporated in the isolated materials when using HCl . All of these features agree with the idea that **1–6** are neutral species.

In the case of the reaction between Na_2MoO_4 and glycine, we carefully monitored the pH of the reaction solution under different initial conditions (see ESI[†]); our data indicated a stoichiometric H^+/Mo ratio of 4 (eqn (2)).



Complexes **1–6**, once isolated in the solid state, resulted almost insoluble, both in H_2O and in 1 M HNO_3 or HCl , while a stronger acidic medium (4 M HCl or HNO_3) was generally needed to obtain colourless solutions. Concerning organic solvents, complexes **1–6** did not dissolve in MeOH , EtOH , Me_2CO and MeCN ; instead they dissolved into yellow solutions when treated with DMSO .

The generally observed insolubility of **1–6** prompted us to their characterization by means of solid-state techniques.

As described in the following sections, spectroscopic (IR, NMR) and analytical data indicated the general basic structure $[(\text{MoO}_2(\text{OH}))_2(\mu\text{-OH})_2(\mu\text{-aaH-}\kappa\text{O},\kappa\text{O}')]]$. This outcome has been supported also by DFT calculations and appears independent on the nature of the ammonium group [*i.e.* primary (GlyH, PheH, LeuH, MetH), secondary cyclic (ProH) or tertiary (dmPheH)]. The general coordination mode proposed for aaH is shared also by **4** despite the additional, potential donor atom (S) contained in MetH.

Solid-state NMR characterisation

In general, ^{13}C chemical shifts of α -amino acids in D_2O vary with pH .¹³ More precisely, an upfield shift of ^{13}C resonances has been observed on going from the amino acidate anion (aa^-) to the zwitterion (aaH) and then to the ammonium acid (aaH_2^+). This effect^{13d} is significant for the carboxylic unit and the adjacent carbons; conversely the remaining nuclei, included those belonging to the side-chain and the N-substituents,

undergo negligible shift. As an example, Fig. 1 reports the case of L-leucine.^{13f}

The solid-state ^{13}C cross polarization with magic angle spinning (CP-MAS) spectra of **1–6** showed one, two or three sets of isotropic resonances attributed to the aaH ligand (see ESI, Table S2[†]). The appearance of more than one signal for a given carbon is presumably due to the presence of different molecular or crystallographic environments.¹⁴

In addition, we recorded ^{13}C CP-MAS spectra of the α -amino acids employed in this work and of their nitrate salts, $[\text{aaH}_2][\text{NO}_3]$, $\text{aaH} = \text{GlyH}$, PheH, LeuH, MetH.¹⁵ The latter compounds were obtained as crystalline materials by slow evaporation of the solvent from aqueous solutions containing aaH and HNO_3 . A comparison of the isotropic ^{13}C chemical shifts related to aaH with the corresponding ones detected for the respective $[\text{aaH}_2]^+$ species evidenced variations that are analogous to those occurring in D_2O (see above).

As reported in Table S2,[†] chemical shifts for the carboxylic-, α - and β -carbons of **1–4** are very close to those found for the corresponding ammonium acid nitrate (*i.e.*, they are deshielded if compared to aaH). The spectrum of **5** shows three groups of resonances: one of these follows the general trend observed for **1–4**, while the other two are only slightly shifted with respect to ProH. Chemical shift values for **6** are deshielded if compared to dmPheH, with the exception of the α -carbon.

It may be concluded that, in general, the coordination of the zwitterionic α -amino acid to the metal centre, $\text{aaH} \rightarrow \text{Mo}_2\text{O}_4(\text{OH})_4(\text{aaH})$, mimics the electronic effects of protonation, $\text{aaH} \rightarrow [\text{aaH}_2]^+$. This feature supports the hypothesis that, in **1–6**, the α -amino acid actually acts as an α -ammonium-carboxylate ligand.

Solid-state IR characterisation

The oxido-molybdenum backbone. The stretching vibrations of $\text{Mo}=\text{O}$ bonds usually give rise to strong and characteristic infrared bands.¹⁶ In fact the IR spectra of **1–6** showed strong absorptions around 944, 915 and 902 cm^{-1} (see ESI, Table S3[†]), which are diagnostic for the *cis*- MoO_2 group.¹⁷ The presence of three stretching absorptions instead of two [*i.e.*, $\nu_{\text{sym}}(\text{MoO}_2)$ and $\nu_{\text{asym}}(\text{MoO}_2)$] might be ascribed to a solid-state effect^{17a,g} or to the fact that more than one $\{\text{MoO}_2\}$ unit is present within a polynuclear complex.¹⁸ A medium-intensity band around 760 cm^{-1} has been assigned to the antisymmetric stretching of a bent $\text{Mo}-\text{O}-\text{Mo}$ bridge,^{16,17g,18,19} while the absorption around 470 cm^{-1} is presumably due to the corresponding $\nu_{\text{sym}}(\text{Mo}_2\text{O})$.^{19a,b,20}




Fig. 1 ^{13}C chemical shifts at different pH values for L-leucine in D_2O .^{13f}

Moreover, the strong and broad band around 500 cm^{-1} may be related to other types of bridging oxido groups (e.g. $\text{Mo}_2(\mu\text{-O})_2$),^{21,22} however $\text{Mo-O}_{\text{ligand}}$ stretching vibrations may also contribute.^{17g,22,23}

The carboxylate group. The wavenumber difference between the anti-symmetric and the symmetric stretching vibration of a carboxylate ligand ($\Delta\nu_{\text{a-s}}$) has proven to be a useful parameter in order to discriminate between monodentate, chelating and bridging-bidentate coordination modes.²⁴ More specifically, the separation between carboxylate stretching wavenumbers relative to a reference ionic carboxylate compound; hence $\Delta\nu_{\text{a-s}}^{\text{rel}} = \Delta\nu_{\text{a-s}}(\text{metal complex}) - \Delta\nu_{\text{a-s}}(\text{ionic reference})$.

We found a good correlation between $\Delta\nu_{\text{a-s}}^{\text{rel}}$ and the coordination mode of a carboxylate ligand by analyzing the IR data reported for a variety of crystallographically-characterized dioxidomolybdenum(vi) complexes.²⁵ Thus $\Delta\nu_{\text{a-s}}^{\text{rel}}$ fall within the range $+140\text{ cm}^{-1}$ to $+60\text{ cm}^{-1}$ in the case of monodentate ligands,²⁶ -40 cm^{-1} to -60 cm^{-1} for chelating ligands,²⁷ and $+30\text{ cm}^{-1}$ to -30 cm^{-1} for bidentate bridging ligands.^{28,7d,7e,21a} As shown in Table S3,† $\Delta\nu_{\text{a-s}}^{\text{rel}}$ values calculated for **1-6** are distributed within the range $+33\text{ cm}^{-1}$ to -38 cm^{-1} (zwitterionic amino acids in the solid state have been taken as ionic references). According to the aforementioned empirical rule, this evidence supports the bidentate-bridging coordination of the carboxylate moiety in **1-6**.

The ammonium group and other considerations. In the $1400\text{-}1600\text{ cm}^{-1}$ spectral region, the antisymmetric $\delta_{\text{asym}}(\text{NH}_3^+)$ and symmetric $\delta_{\text{sym}}(\text{NH}_3^+)$ bands of the ammonium group are recognizable for **1-4**.²⁹ One deformation band is observed for **5**, as expected for a secondary ammonium ion.³⁰ In the spectrum of **6**, the $\nu(\text{N-CH}_3)$ has been recognized.³¹ All of these absorptions are slightly shifted with respect to the corresponding, uncoordinated zwitterionic amino acid (see Table S3†). In addition, the IR spectra of **1-4** resemble the spectra of the corresponding $[\text{aaH}_2]\text{NO}_3$ species, for what regards the C-H and N-H stretching region ($2800\text{-}3300\text{ cm}^{-1}$),³² see Fig. S7.† All of these observations agree with the presence of a protonated ammonium group in compounds **1-6**.

The deformation of the $-\text{SCH}_3$ group gives rise to another characteristic and intense band,³³ falling at 1316 cm^{-1} in the spectrum of MetH and at 1334 cm^{-1} in the spectrum of its molybdenum complex, **4**. A medium intensity absorption at 1333 cm^{-1} was found in the IR spectra of two structurally determined complexes of Nb(v) and Ta(v) bearing a *L*-methionine ligand bonded to the metal centre through the amino and the carboxylate groups.³⁴ This suggests that the sulphur atom in **4** is not involved in coordination.

Theoretical calculations

In summary, analytical and spectroscopic characterization of **1-6** highlighted: (i) an elemental composition coherent with the empirical formula $\text{Mo}_2\text{H}_4\text{O}_8(\text{aaH})$, (ii) the presence of terminal (*cis*- MoO_2) and bridging oxido groups, (iii) a bidentate bridging coordination, *via* the carboxylate group, of the zwitterionic amino acid ligand.

The computational investigation was initially aimed to ascertain possible dinuclear structures for **1** and **6**. The calculated structures were subjected to DFT optimization both in the gas-phase (EDF2 functional) and in the presence of implicit solvation for water (PBE functional with Grimme's corrections for dispersion), using valence-double- ζ quality basis sets. The lowest energy structures are shown in Fig. 2, while selected bond distances and angles are reported in Table 1. The Cartesian coordinates are given as ESI† as a separated .xyz file.

Both structures are built up of dinuclear molecules $[(\text{MoO}_2(\text{OH}))_2(\mu\text{-OH})_2(\mu\text{-O}_2\text{CCHRNR}'_2-\kappa\text{O},\kappa\text{O}')] (R = \text{H}, R' = \text{H}$ for **1**, $R = \text{CH}_2\text{Ph}, R' = \text{Me}$ for **6**). Each molybdenum is bonded to two terminal oxido and one terminal hydroxido ligands. The two $\text{MoO}_2(\text{OH})$ units are bridged by two hydroxido ligands and the carboxylate group of the α -amino acid. Moreover, one bridging OH ligand and one carboxylic oxygen are *trans* to a terminal oxido ligand. The coordination spheres around the Mo centres can be described as distorted octahedra, as observable from the $\text{O}_{\text{terminal}}\text{-Mo-O}_{\text{terminal}}$ ($<85^\circ$) and $\text{O}_{\text{bridging}}\text{-Mo-O}_{\text{bridging}}$ (105°) angles. Both features (*trans* influence of oxido ligands and distortion from octahedral coordination) are commonly observed in dioxidomolybdenum(vi) complexes.³⁵ An intramolecular hydrogen bond involving the ammonium and the nearest $\text{Mo}=\text{O}$ group is present in the dinuclear model for **1** while is missing in the analogous for **6**: this seems to indicate that intermolecular $\text{N-H}\cdots\text{O}$ interactions are not essential for the stability of this type of structures.

As discussed above, the DFT-optimised structures of **1** and **6** are similar to each other: this agrees with the homogeneous spectroscopic data collected, as discussed in the previous sections. Concerning other possible structures, the formal replacement of the bridging hydroxido groups with oxido ligands caused, on theoretical grounds, the dissociation of the carboxylate moiety and/or the migration of one H^+ from the ammonium to one oxido. Also the presence of a single bridging ligand, besides the amino acid, was ruled out by DFT calculations, because geometry optimizations of several Mo-O-Mo and Mo-OH-Mo starting structures gave stationary points with two bridging OH ligands or structures having very




Fig. 2 Lowest-energy DFT optimized dinuclear structures of **1** and **6** (gas-phase EDF2 calculations). White = hydrogen, grey = carbon, blue = nitrogen, red = oxygen, light blue = molybdenum. Carbon-bonded hydrogen atoms have been omitted for clarity.

Table 1 Selected bond distances (Å) and angles (°) for the dinuclear structures **1** and **6**. Data computed using the Grimme-corrected PBE functional in the presence of COSMO solvation model for water. Gas-phase EDF2 data in parentheses

Atoms	1	6
Mo(1)–O(1)	1.735 (1.699)	1.742 (1.710)
Mo(2)–O(7)	1.731 (1.691)	1.740 (1.707)
Mo(1)–O(2)	1.743 (1.708)	1.740 (1.710)
Mo(2)–O(6)	1.760 (1.765)	1.741 (1.713)
Mo(1)–O(4)	2.221 (2.263)	2.312 (2.323)
Mo(1)–O(5)	2.096 (2.089)	2.072 (2.041)
Mo(2)–O(5)	2.217 (2.130)	2.304 (2.323)
Mo(2)–O(4)	2.067 (2.050)	2.085 (2.098)
Mo(1)–O(3)	1.942 (1.904)	1.954 (1.960)
Mo(2)–O(8)	1.936 (1.899)	1.949 (1.919)
Mo(1)–O(9)	2.422 (2.572)	2.269 (2.310)
Mo(2)–O(10)	2.385 (2.582)	2.298 (2.351)
O(1)–Mo(1)–O(2)	105.5 (105.5)	106.3 (105.8)
O(5)–Mo(1)–O(9)	85.9 (83.1)	77.8 (76.5)
O(4)–Mo(1)–O(5)	68.1 (64.8)	75.5 (69.8)
O(4)–Mo(2)–O(5)	68.6 (67.9)	75.4 (70.0)

high relative energies. Thus the $\text{Mo}_2(\mu\text{-OH})_2$ core is strongly suggested by theoretical calculations. Moreover, the coordination of water molecules to the Mo(vi) centres was very weak in all of the cases considered here. The position of the terminal OH ligands *cis* to the carboxylate groups resulted favoured with respect to the *trans* configuration ($\Delta G = -10.4$ kcal mol⁻¹ for **1**, -4.3 kcal mol⁻¹ for **6**; see the ESI†.xyz file for the **1-trans** and **6-trans** optimized geometries).

It should be mentioned that a series of X-ray determined molybdates present a structural motif analogous to that proposed for **1-6**: *cis*-MoO₂ groups bridged by two oxido and one carboxylato unit derived from a zwitterionic ligand LH (α -, β - or γ -amino acids, or nicotinic acid).^{7d,7e,21,28} In such cases, the resulting dinuclear frames, *i.e.* $[\text{Mo}_2\text{O}_4(\mu\text{-O})_2(\text{LH})]$, may condense to three-units rings – in the presence of a central heteroatom such as V(v), Se(IV) or As(III)^{7d,7e,28} – or to infinite oxido-bridged rail-like chains.²¹

Therefore, a theoretical approach was attempted by considering the polymerization of the dinuclear structures (Fig. 2). Preliminary DFT calculations indicated the condensation of two terminal OH groups as the only way to the formation of polymer chains maintaining the dinuclear $\text{Mo}_2\text{O}_4(\mu\text{-OH})_2(\mu\text{-aaH-}\kappa\text{O},\kappa\text{O}')$ core.³⁶ In order to restrict the influence of N–H⋯O hydrogen bonds during the research of the most stable minima, the dinuclear structure optimized for **6** was selected as the starting point. Fig. 3 shows the lowest-energy structures for condensation products containing four (**6T**), six (**6H**) and eight (**6O**) metal centres. Selected ranges of bond distances and angles are reported in Table 2, while the Cartesian coordinates are collected in a separated .xyz file.†

Mo₂ subunits in the polynuclear structures are held together not only by a covalent oxygen bridge, but also by an intramolecular hydrogen bond between two OH groups, acting as bridging ligands within distinct Mo₂ frames. These interactions are sketched in Fig. 4 for clarity. The presence of a single bent




Fig. 3 Lowest-energy DFT optimized polynuclear structures of **6** (**6T**, **6H** and **6O**) (PBE/Grimme/COSMO, in the presence of water as implicit solvent). White = hydrogen, grey = carbon, blue = nitrogen, red = oxygen, light blue = molybdenum. The substituents on the carboxylate-bonded carbons have been omitted for clarity.

Mo–O–Mo bridge is supported also by IR characterization (see above).

On theoretical grounds, the condensation reactions leading to polynuclear structures (see eqn (3) for aaH = dmPheH and $n = 1-3$) appear thermodynamically favoured, as observable from the ΔG variations (including solvation) reported in Table 3 for each condensation. This point suggests that **1-6** probably possess, to some extent, a polymeric structure, in accordance with their insolubility in water and in organic solvents. By assuming further condensation reactions between Mo₂ subunits ($n > 3$), an oxido-bridged 2D coordination polymer of general formula $\{\text{Mo}_2\text{O}_4(\text{OH})_2(\text{aaH})(\mu\text{-O})\}_n \cdot \text{H}_2\text{O}$ would be obtained.




Table 2 Selected ranges of bond distances (Å) and angles (°) for the computed polynuclear structures of **6**. PBE/COSMO calculations, EDF2 data in parentheses^a

Atoms	6T	6H	6O
Mo-O(a)	1.737–1.747 (1.708–1.723)	1.738–1.764 (1.703–1.741)	1.738–1.763
Mo-O(b)	1.942–1.959 (1.923–1.950)	1.941–1.952 (1.923–1.956)	1.949–1.951
Mo-O(c)	2.236–2.339 (2.237–2.401)	2.257–2.347 (2.268–2.427)	2.250–2.358
	2.028–2.104 (2.013–2.099)	2.017–2.107 (1.979–2.099)	2.014–2.125
Mo-O(d)	1.920–1.925 (1.887–1.915)	1.921–1.937 (1.883–1.955)	1.912–1.953
Mo-O(e)	2.247–2.329 (2.311–2.358)	2.228–2.327 (2.285–2.345)	2.243–2.321
O(c)···O(c) (H-bond)	2.571 (2.601)	2.548–2.828 (2.598–2.724)	2.579–2.719
O(a)–Mo–O(a)	103.5–106.5 (104.6–105.5)	103.5–105.5 (104.1–105.6)	103.5–106.7
O(c)–Mo–O(c)	73.1–74.0 (69.8–71.3)	71.6–74.4 (69.9–71.2)	71.4–74.1
Mo–O(d)–Mo	151.6 (157.8)	138.6–152.8 (140.9–160.6)	136.4–153.9

^a Mo–O(a) = Mo=O(terminal); Mo–O(b) = Mo–OH(terminal); Mo–O(c) = Mo–OH(bridging); Mo–O(d) = Mo–O(bridging); Mo–O(e) = Mo–O(COO).




Fig. 4 Sketch of the covalent and hydrogen-bond interactions between Mo₂ units.

Table 3 Computed ΔG variations (including implicit solvation, water as solvent, 298 K) for the condensation steps leading from **6** to **6O** (PBE/Grimme/COSMO calculations)

Step	ΔG , kcal mol ⁻¹
6 + 6 → 6T + H ₂ O	-10.0
6T + 6 → 6H + H ₂ O	-7.1
6H + 6 → 6O + H ₂ O	-4.1

Conclusions

We have carried out a detailed solid state characterization of Mo(vi) compounds isolated by reactions of molybdate salts with α -amino acids in acidic aqueous solutions. The

combination of experimental results with DFT calculations has allowed the unprecedented structural identification of the dinuclear core and the coordination mode of the α -amino acid. Moreover, the presumable formation of coordination polymers by condensation reactions has been discussed. This paper contributes to clarify the structural features of a series of compounds obtained by a simple reaction, arousing great interest for the implications associated with the nature of the reactants and the use of water as medium (see Introduction).

Experimental

General

All manipulations were performed in air with common laboratory glassware. Reactions were carried out using deionised water. The following reagents and solvents were used as received: Na₂MoO₄·2H₂O, (NH₄)₆Mo₇O₂₄·4H₂O, CaH₂ (Alfa Aesar), glycine (GlyH), L-phenylalanine (PheH), L-leucine (LeuH), L-proline (ProH), L-methionine (MetH), *N,N*-dimethyl-L-phenylalanine (dmPheH) (Apollo Scientific), NaOH, DMSO (Carlo Erba), potassium hydrogen phthalate (Fluka), MoO₂(acac)₂, 65% HNO₃, 37% HCl, methanol, 96% ethanol, acetone, diethyl ether (Sigma-Aldrich). (NH₄)₂MoO₄ was prepared from (NH₄)₆Mo₇O₂₄·4H₂O according to the literature.³⁷ An Orion pH-meter equipped with a Hamilton glass pH-electrode was used for pH measurements. The instrument was routinely calibrated with standard pH = 4.0 and pH = 7.0 buffer solutions (Fluka). Infrared spectra were recorded on a Perkin Elmer FT-IR spectrometer. The UATR sampling accessory was used in the 650–4000 cm⁻¹ range. Spectra in the 360–650 cm⁻¹ region were recorded in the transmission mode on Nujol mulls. NMR solution spectra were recorded on a Bruker Avance DRX-400 instrument equipped with a BBFO broadband probe. Chemical shifts for ¹H and ¹³C were referenced to tetramethylsilane. CP-MAS NMR spectra were recorded on a Bruker AMX300WB spectrometer working at 75.47 MHz for carbon, using a CP pulse sequence with a 90° pulse of 3.4 μ s, a contact time of 2 ms and a recycle delay of 4 s. 4 mm zirconia rotors were used. A 9 kHz spinning speed was used except when, due to the consistency of the sample, homogeneous packing within the rotor was difficult and a stable rotation was achieved at a lower spinning speed (5 kHz). Carbonyl and aromatic spinning sidebands were detected at \pm 119 ppm and \pm 66 ppm for 9 kHz and 5 kHz rotation, respectively. Spectral editing techniques³⁸ were used to aid the spectral assignment in the case of dmPheH and its molybdenum complex. Carbon, hydrogen and nitrogen analyses were performed on a Carlo Erba mod. 1106 instrument. Molybdenum content was determined with the spectrophotometric method proposed by Crouthamel e Johnson³⁹ after dissolution of a weighted amount of sample (30–60 mg) in 100 mL of 4 M HCl. A calibration curve was obtained using (NH₄)₆Mo₇O₂₄·4H₂O as standard ($R^2 = 0.999$), while MoO₂(acac)₂ was used as a reference compound (anal. found: Mo 30.2%. C₁₀H₁₄MoO₄ requires Mo 29.8%).

Synthesis of $\text{Mo}_2\text{O}_4(\text{OH})_4(\text{aaH})$, 1–6

Compound $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ or $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$ (4.00 mmol of Mo) and the α -amino acid (aaH, 2.00 mmol) were dissolved in water (30 mL), then 1 M HNO_3 was added until $\text{pH} = 1.6$ – 2.0 . The resulting pale-yellow solution ($C_{\text{Mo}} = 0.11$ M) was stirred at room temperature. A colourless solid (1–6) was obtained in a variable time, depending on the amino acid employed. Then the reaction mixture was filtered (filtrated solution $\text{pH} = 2.1$ – 2.5). The solid was washed with acetone, diethyl ether and then dried *in vacuo* over CaH_2 . Details for single preparations with $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ are given below; analogous outcomes were obtained by using $(\text{NH}_4)_6\text{Mo}_7\text{O}_{24} \cdot 4\text{H}_2\text{O}$.

$\text{Mo}_2\text{O}_4(\text{OH})_4(\text{GlyH})$, 1. A precipitate formed in 25 minutes (715 mg, 90%). Anal. found: C, 6.03; H, 2.13; Mo, 49.1; N, 3.67%. $\text{C}_2\text{H}_9\text{Mo}_2\text{NO}_{10}$ requires C, 6.02; H, 2.27; Mo, 48.1; N, 3.51%.⁴⁰ IR (ATR and Nujol): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3565w, 3424m, 3362m, 3225m, 3045m, 3011m, 2926m, 2757m, 2650m, 1672w, 1623s, 1590s, 1512s, 1456s, 1412s, 1340s, 1148w, 1118w, 1100w-sh, 946vs, 918vs, 903vs, 761m, 732w-sh, 531vs-br, 473m-sh, 384w. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 172.0 (CO), 41.3 (CH_2).

$\text{Mo}_2\text{O}_4(\text{OH})_4(\text{PheH})$, 2. A precipitate formed in 2.5 hours (805 mg, 90%). Anal. found: C, 22.05; H, 2.85; Mo, 40.1; N, 2.75%. $\text{C}_9\text{H}_{15}\text{Mo}_2\text{NO}_{10}$ requires C, 22.10; H, 3.09; Mo, 39.2; N, 2.86%.⁴⁰ IR (ATR and Nujol): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3605w-sh, 3561w, 3136w-br, 3033w, 2945w, 2908w, 2831w, 2764w, 1607s, 1525m, 1498m-sh, 1454m, 1426s, 1346m, 1300w, 1248w, 1143w, 1105w, 1084w, 942vs, 913vs-sh, 904vs, 856w-sh, 756m, 738w-sh, 705w, 534vs-br. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 172.6 (CO), 134.5 (Ph), 133.6 (Ph), 132.5 (Ph), 129.6 (Ph), 57.3 (CH), 35.6 (CH_2).

$\text{Mo}_2\text{O}_4(\text{OH})_4(\text{LeuH})$, 3. No precipitate was observed after 18 hours. Thus the solvent was completely removed *in vacuo* at 50 °C affording a pale-yellow residue. This residue was washed with water, acetone, diethyl ether and dried *in vacuo* over CaH_2 ; a colourless solid was obtained (704 mg, 77%). Anal. found: C, 16.15; H, 3.70; Mo, 42.7; N, 3.01%. $\text{C}_6\text{H}_{17}\text{Mo}_2\text{NO}_{10}$ requires C, 15.83; H, 3.76; Mo, 42.17; N, 3.08%.⁴⁰ IR (ATR and Nujol): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3487m-br, 3133m-sh, 3055m, 2956m, 2931m-sh, 2874m, 1620s-sh, 1599s, 1513m, 1506m, 1494m, 1468m, 1428s, 1389m, 1369m, 1350m, 1338m, 1318m, 1293w, 1172w, 1133w, 1123w, 939vs, 909vs-sh, 896vs, 768w, 750w-sh, 537vs-br, 479s-sh, 380w. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 173.2 (CO), 53.2 (CHN), 39.2 (CH_2), 25.1 (CHMe_2), 21.3 (CH_3), 20.1 (CH_3).

$\text{Mo}_2\text{O}_4(\text{OH})_4(\text{MetH})$, 4. A precipitate formed in 40 minutes (781 mg, 82%). Anal. found: C, 12.38; H, 2.99; Mo, 41.5; N, 3.10%. $\text{C}_5\text{H}_{15}\text{Mo}_2\text{NO}_{10}\text{S}$ requires C 12.69; H 3.20; Mo 40.56; N 2.96%.⁴⁰ IR (ATR and Nujol): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3600w-sh, 3457w-br, 3365w-br, 3143m, 3066m, 2992w, 2922m, 2766–2443w, 1603vs, 1575s, 1558m-sh, 1504s, 1440m-sh, 1427s, 1358m, 1334s, 1317m-sh, 1280w, 1252w, 1195w, 1148m, 1103w, 1073w, 968w-sh, 942vs, 913vs, 894vs, 762m, 748w-sh, 540vs-br, 486m-sh. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 174.4 (CO), 171.8 (CO), 54.7 (CH), 31.5 (CH_2CH), 30.4 (CH_2CH), 28.7 (SCH_2), 15.3 (SCH_3), 14.3 (SCH_3), 12.4 (SCH_3).

$\text{Mo}_2\text{O}_4(\text{OH})_4(\text{ProH})$, 5. A precipitate formed in 25 minutes (642 mg, 73%). Anal. found: C, 13.83; H, 2.85; Mo, 44.8; N,

3.25%. $\text{C}_5\text{H}_{13}\text{Mo}_2\text{NO}_{10}$ requires C, 13.68; H, 2.98; Mo, 43.71; N, 3.19%.⁴⁰ IR (ATR and Nujol): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3545w-br, 3157w-br, 3040w-br, 2776w, 1603s, 1558m-sh, 1472m, 1432s, 1394m, 1372m, 1338m, 1301w, 1258w, 1175w, 944vs, 914vs, 901vs, 767m, 543vs-br, 463m-sh, 389w. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 176.8 (CO), 175.2 (CO), 173.5 (CO), 65.0 (CH), 61.0 (CH), 49.7 (CH_2N), 47.3 (CH_2N), 46.9 (CH_2N), 30.2 (CH_2CH), 29.4 (CH_2CH), 26.2 ($\text{CH}_2(\text{CH}_2)_2$), 25.8 ($\text{CH}_2(\text{CH}_2)_2$), 23.7 ($\text{CH}_2(\text{CH}_2)_2$).

$\text{Mo}_2\text{O}_4(\text{OH})_4(\text{dmPheH})$, 6. A precipitate formed immediately (862 mg, 83%). Anal. found: C, 26.05; H, 3.55; Mo, 37.5; N, 2.73%. $\text{C}_{11}\text{H}_{19}\text{Mo}_2\text{NO}_{10}$ requires C, 25.55; H, 3.70; Mo, 37.11; N, 2.71%.⁴⁰ IR (ATR and Nujol): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3563w-br, 3410w, 3055w, 3040w, 3025w, 2965w, 2943w, 2734w, 1626s, 1471m, 1455m, 1432m-sh, 1495m, 1412s, 1358m, 1260w, 1185w, 1170w, 1081w, 1046w, 946vs, 916vs, 901vs-sh, 769m, 742m, 705m, 547vs-br, 484m-sh, 438w, 385w. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 171.6 (CO), 140.9 (Ph), 130.8 (Ph), 128.8 (Ph), 127.8 (Ph), 69.9 (CH), 42.0 (NCH_3), 36.6 (NCH_3), 30.4 (CH_2).

Preparation of $[\text{aaH}_2]\text{NO}_3$, aaH = GlyH, PheH, LeuH, MetH

The α -amino acid (2–4 mmol) was dissolved in 1 M HNO_3 (10 mL for GlyH, PheH and LeuH; a stoichiometric amount of HNO_3 for MetH) and the resulting colourless solution was allowed to evaporate slowly at room temperature. Colourless crystals of $[\text{aaH}_2]\text{NO}_3$ were formed in a period of 5–12 days. These were collected and dried *in vacuo* over CaH_2 .

$[\text{GlyH}_2]\text{NO}_3$. Anal. found: C, 17.23; H, 4.11; N, 20.01%. $\text{C}_2\text{H}_6\text{N}_2\text{O}_5$ requires C, 17.40; H, 4.38; N, 20.29%. IR (ATR): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3205m, 3150w, 3032m, 2966m, 2909m, 2837m, 2748w, 2719w, 2640m, 2541w, 2442w, 2332w, 2262w, 2236w, 2221w, 1986w, 1724s, 1627w, 1596w, 1517m, 1442m, 1416s, 1355s, 1331s, 1215s, 1124s, 1043s, 914s, 890s, 868s, 813s, 738m, 712w, 662w. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 171.7 (CO), 41.1 (CH_2).

$[\text{PheH}_2]\text{NO}_3$. Anal. found: C, 47.01; H, 5.11; N, 12.01%. $\text{C}_9\text{H}_{12}\text{N}_2\text{O}_5$ requires C, 47.37; H, 5.30; N, 12.28%. IR (ATR): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3148m-br, 3026m-br, 2972m-br, 1717s, 1667m, 1606w, 1584w, 1505m, 1496m, 1471w, 1453m, 1445m, 1408m, 1357s, 1335s, 1304m, 1200s, 1150m, 1119m, 1082w, 1045w, 1033w, 986w, 972w, 943w, 922w, 904w, 873w, 851w, 814m, 796w, 759w, 743w-sh, 737m, 693s. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 173.9 (CO), 173.2 (CO), 135.2 (Ph), 132.8 (Ph), 132.3 (Ph), 129.8 (Ph), 129.3 (Ph), 127.6 (Ph), 127.3 (Ph), 56.3 (CH), 55.5 (CH), 37.0 (CH_2), 36.3 (CH_2).

$[\text{LeuH}_2]\text{NO}_3$. Anal. found: C, 37.21; H, 6.99; N, 14.25%. $\text{C}_6\text{H}_{14}\text{N}_2\text{O}_5$ requires C, 37.11; H, 7.27; N, 14.43%. IR (ATR): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3221w, 3197w, 3149w, 2973m, 2958m, 2918m, 2875m, 2768w, 2701w, 2649w, 2583w, 2537w, 1717s, 1624m, 1579w, 1507s, 1473w, 1454m, 1413s, 1386s, 1344s, 1323s, 1297m, 1212s, 1171s, 1142s, 1075m, 1035s, 995m, 939m, 912s, 807s, 752w, 730w, 711m. ¹³C NMR (75.47 MHz, CP-MAS): δ/ppm 173.9 (CO), 173.1 (CO), 52.9 (CHN), 39.2 (CH_2), 24.7 (CHMe_2), 23.3 (CH_3), 22.3 (CH_3), 21.7 (CH_3), 20.0 (CH_3).

$[\text{MetH}_2]\text{NO}_3$. Anal. found: C, 27.99; H, 5.46; N, 13.01%. $\text{C}_5\text{H}_{12}\text{N}_2\text{O}_5\text{S}$ requires C, 28.30; H, 5.70; N, 13.20%. IR (ATR): $\tilde{\nu}_{\text{max}}/\text{cm}^{-1}$ 3182m, 3044m, 2964m, 2921m, 2853m, 2713w,

1 2640w, 2604w, 1724s, 1620w, 1573w, 1506m, 1407s, 1381s,
1340s, 1287s, 1238m, 1225m, 1181s, 1135s, 1100m, 1073m,
1049m, 1035m, 984m, 948m, 913m, 864s, 817m, 773m, 756m,
5 731m, 700m, 655w.¹³C NMR (75.47 MHz, CP-MAS): δ /ppm 173.7
(CO), 172.2 (CO), 54.6 (CH), 54.4 (CH), 32.9 (CH₂CH), 30.2
(SCH₂), 16.0 (SCH₃), 15.5 (SCH₃).

Computational details

10 The geometry optimization of the complexes was carried out
without symmetry constrains, using the hybrid DFT functional
EDF2,⁴¹ in combination with the split-valence double- ζ polar-
ized basis set LACVP** (6-31G** on the light atoms, LANL2DZ
15 effective core potential on molybdenum centres),⁴² and with the
GGA PBE functional combined with an atom-centred doubly
numerical polarized basis set.⁴³ Dispersion correction was
added through the Grimme's approach⁴⁴ and the COSMO
implicit solvation model for water was added.⁴⁵ On the basis of
20 the experimentally observed diamagnetism, in all cases the
molecules were considered in the singlet state and the
"restricted" approach was adopted. IR simulations on the
equilibrium geometries were carried out using the harmonic
approximation, from which zero-point vibrational energies and
25 thermal corrections to enthalpy and entropy were obtained
using standard formulae.⁴⁶ The Cartesian coordinates of the
most stable geometries are collected in a separated .xyz file.†

Acknowledgements

30 CINECA is gratefully acknowledged for computational support.

References and Notes

- 35 1 (a) B. M. Hoffman, D. Lukoyanov, Z.-Y. Yang, D. R. Dean and
L. C. Seefeldt, *Chem. Rev.*, 2014, **114**, 4041–4062; (b)
M. V. Ribbe, Y. Hu, K. O. Hodgson and B. Hedman, *Chem.*
Rev., 2014, **114**, 4063–4080; (c) B. M. Hoffman,
D. Lukoyanov, D. R. Dean and L. C. Seefeldt, *Acc. Chem.*
40 *Res.*, 2013, **46**, 587–595; (d) T. Spatzal, M. Aksoyoglu,
L. Zhang, S. L. A. Andrade, E. Schleicher, S. Weber,
D. C. Rees and O. Einsle, *Science*, 2011, **334**, 940; (e)
K. M. Lancaster, M. Roemelt, P. Ettenhuber, Y. Hu,
M. W. Ribbe, F. Neese, U. Bergmann and S. DeBeer,
45 *Science*, 2011, **334**, 974–977; (f) J. Emsley, in *Nature's*
Building Blocks: An A-Z Guide to the Elements, Oxford
University Press, Oxford, 2001; (g) G. C. Tucci,
J. P. Donhaue and R. H. Holm, *Inorg. Chem.*, 1998, **37**,
1602–1608; (h) R. Hille and H. Sprecher, *J. Biol. Chem.*,
50 1987, **262**, 10914–10917; (i) R. H. Holm and J. M. Berg, *Acc.*
Chem. Res., 1986, **19**, 363–370.
- 2 (a) J. J. Cruywagen, *Adv. Inorg. Chem.*, 2000, **49**, 127–182; (b)
J. J. Cruywagen, *Inorg. Chem.*, 1980, **19**, 552–554.
- 55 3 (a) D. Wang, *Front. Microbiol.*, 2012, **3**, 1–7; (b) G. Rayner-
Canham and J. Grandy, *Educ. Chem.*, 2011, 144–147.
- 4 (a) D. Collison, C. D. Garner and J. A. Joule, *Chem. Soc. Rev.*,
1996, 25–32; (b) R. Hille, *Chem. Rev.*, 1996, **96**, 2757–2816; (c)
T. J. Huang and G. P. Haight Jr, *J. Am. Chem. Soc.*, 1970, **92**,

2336–2342; (d) J. T. Spence, *Coord. Chem. Rev.*, 1969, **4**,
475–498.

- 5 (a) K. Vassilev, M. Dimitrova, S. Turmanova and R. Milina,
Synth. React. Inorg. Met.-Org. Chem., 2013, **43**, 243–249; (b)
5 W. J. Shaw, *Catal. Rev.: Sci. Eng.*, 2012, **54**, 489–550; (c)
D. Desbouis, I. P. Troitsky, M. J. Belousoff, L. Spiccia and
B. Graham, *Coord. Chem. Rev.*, 2012, **256**, 897–937; (d)
K. Micskei, T. Patonay, L. Caglioti and G. Palyi, *Chem.*
Biodiversity, 2010, **7**, 1660–1669; (e) K. Severin, R. Bergs and
W. Beck, *Angew. Chem., Int. Ed.*, 1998, **37**, 1635–1654.
- 10 6 J. Paradowska, M. Stodulski and J. Mlynarski, *Angew. Chem.,*
Int. Ed., 2009, **48**, 4288–4297.
- 7 See for instance: (a) W. S. Zhu, H. Li, Q. Q. Gu, P. Wu, G. Zhu,
Y. Yan and G. J. Chen, *J. Mol. Catal. A: Chem.*, 2011, **336**, 16–
22; (b) M. L. Ramos, L. L. G. Justino and H. D. Barroiwis,
15 *Dalton Trans.*, 2011, 4374–4383; (c) E. Cartuyvels, K. Van
Hecke, L. Van Meervelt, C. Görller-Walrand and
T. N. Parac-Vogt, *J. Inorg. Biochem.*, 2008, **102**, 1589–1598;
(d) U. Kortz, J. Vaissermann, R. Thouvenot and P. Gouzerh,
20 *Inorg. Chem.*, 2003, **42**, 1135–1139; (e) U. Kortz,
M. G. Savelieff, F. Y. A. Ghali, L. M. Khalil, S. A. Maalouf
and D. I. Sinno, *Angew. Chem., Int. Ed.*, 2002, **41**, 4070–
4073; (f) J. Lu and Y. Xu, *Chem. Mater.*, 1998, **10**, 4141–
4147; (g) C. Djordjevic, N. Vuletic, B. A. Jacobs, M. Lee-
25 Renslo and E. Sinn, *Inorg. Chem.*, 1997, **36**, 1798–1805; (h)
B. Kamenar, B. Korpar-Colig, M. Penavic and M. Cindric, *J.*
Chem. Soc., Dalton Trans., 1990, 1125–1130.
- 8 Some selected references are: (a) J. J. Cruywagen,
J. B. B. Heyns and E. A. Rohwer, *J. Chem. Soc., Dalton*
30 *Trans.*, 1993, 1713–1717; (b) A. Domènech, E. Llopis,
E. Garcia-España and A. Cervilla, *Transition Met. Chem.*,
1990, **15**, 425–428; (c) D. O. Martire, M. R. Feliz and
A. L. Capparelli, *Polyhedron*, 1989, **8**, 2225–2232; (d)
M. A. Freeman, F. A. Schultz and C. N. Reilley, *Inorg.*
35 *Chem.*, 1982, **21**, 567–576; (e) A. M. V. S. V. Cavaleiro,
J. D. Pedrosa de Jesus, V. M. S. Gil, R. D. Gillard and
P. A. Williams, *Transition Met. Chem.*, 1982, **7**, 75–79.
- 9 A. Kay and P. C. Mitchell, *J. Chem. Soc. A*, 1970, 2421–2428.
- 10 R. J. Butcher, H. K. J. Powell, C. J. Wilkins and S. H. Yong, *J.*
Chem. Soc., Dalton Trans., 1976, 356–359.
- 11 (a) M. Castillo and P. Palma, *Synth. React. Inorg. Met.-Org.*
Chem., 1984, **14**, 1173–1182; (b) K. S. Nagaraja and
M. R. Udupa, *Indian J. Chem., Sect. A: Inorg., Phys., Theor.*
45 *Anal.*, 1983, **22**, 531–532; (c) H. Eguchi, T. Takeuchi,
A. Ouchi and A. Furuhashi, *Bull. Chem. Soc. Jpn.*, 1969, **42**,
3585–3588.
- 12 As explained later in the text, the proposed structure for 1–6
corresponds to the dinuclear complex [(MoO₂(OH))₂(μ -
50 OH)₂(μ -aaH- κ O, κ O')], which may give rise to an oxido-
bridged polymer though the condensation of terminal –OH
ligands. The shortened formula Mo₂O₄(OH)₄(aaH) will be
used throughout this paper and the ESI.†
- 13 (a) H. L. Surprenant, J. E. Sarneski, R. R. Key, J. T. Byrd and
55 C. N. Reilley, *J. Magn. Reson.*, 1980, **40**, 231–243; (b)
J. G. Batchelor, J. Feeney and G. C. K. Roberts, *J. Magn.*
Reson., 1975, **20**, 19–38; (c) A. R. Quirt, J. R. Lyerla,
I. R. Peat Jr, J. S. Cohen, W. F. Reynolds and

- 1 M. H. Freedman, *J. Am. Chem. Soc.*, 1974, **96**, 570–574; (d) 20 R. M. Wing and K. P. Callahan, *Inorg. Chem.*, 1969, **8**, 871–
1 M. Christl and J. D. Roberts, *J. Am. Chem. Soc.*, 1972, **94**, 874.
- 5 F. R. Gurd, P. J. Lawson, D. W. Cochran and E. Wenkert, *J. Biol. Chem.*, 1971, **246**, 3725–3730. 21 (a) S.-M. Chen, C.-Z. Lu, Y.-Q. Yu, Q.-Z. Zhang and X. He, *Acta Crystallogr., Sect. C: Cryst. Struct. Commun.*, 2004, **60**, m437–
5 m439; (b) T. Yamase, M. Inoue, H. Naruke and K. Fukaya, *Chem. Lett.*, 1999, **7**, 563–564.
- 14 G. R. Hays, *J. Chem. Soc., Perkin Trans. 2*, 1983, 1049–1052. 22 K. Wieghardt, W. Holzbach and J. Weiss, *Inorg. Chem.*, 1981, **20**, 3436–3439.
- 15 (a) S. Asath Bahadur, R. K. Rajaram, M. Nethaji and S. Z. Natarajan, *Z. Kristallogr.*, 2010, **203**, 93–100; (b) 23 C. Knobler, B. R. Penfold, W. T. Robinson, C. J. Wilkins and S. H. Yong, *J. Chem. Soc., Dalton Trans.*, 1980, 248–252. 10
10 F. Berrah, N. Benali-Cherif and H. Lamraoui, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2005, **61**, o1517–
10 o1519; (c) S. Pandiarajan, B. Sridhar and R. K. Rajaram, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2002, **58**, o882–
15 o884; (d) B. Sridhar, N. Srinivasan and R. K. Rajaram, *Acta Crystallogr., Sect. E: Struct. Rep. Online*, 2002, **58**, o1103–
15 o1105; (e) P. Narayanan and S. Venkataraman, *J. Cryst. Mol. Struct.*, 1975, **5**, 15–26.
- 16 (a) L. Favero, F. Marchetti, G. Pampaloni and S. Zacchini, *Dalton Trans.*, 2014, **43**, 495–504; (b) F. Marchetti, 20
20 G. Pampaloni and S. Zacchini, *Dalton Trans.*, 2013, **42**, 15226–15234; (c) F. Marchetti, G. Pampaloni and S. Zacchini, *Dalton Trans.*, 2013, **42**, 2477–2487; (d) S. Dolci, F. Marchetti, G. Pampaloni and S. Zacchini, *Eur. J. Inorg. Chem.*, 2013, **42**, 1371–1380; (e) W. P. Griffith, *J. Chem. Soc. A*, 1969, 211–218. 25
25
- 17 (a) X. Ma, K. Starke, C. Schulzke, H.-G. Schmidt and M. Noltemeyer, *Eur. J. Inorg. Chem.*, 2006, 628–637; (b) M. J. Taylor, C. E. F. Rickard and L. A. Kloo, *J. Chem. Soc., Dalton Trans.*, 1998, 3195–3198; (c) B. Kojic-Prodic, Z. Ruzic-Toros, D. Grdenic and L. Golic, *Acta Crystallogr., Sect. B: Struct. Crystallogr. Cryst. Chem.*, 1974, **30**, 300–305; (d) B. Kamenar, B. M. Penavic and C. K. Prout, *Cryst. Struct. Commun.*, 1973, **2**, 41–44; (e) L. O. Atovmyan and Y. A. Sokolova, *Zh. Strukt. Khim.*, 1971, **12**, 851–858; (f) B. M. Craven, K. C. Ramey and W. B. Wise, *Inorg. Chem.*, 1971, **10**, 2626–2628; (g) R. H. Fenn, *J. Chem. Soc. A*, 1969, 1764–1769; (h) W. P. Griffith and T. D. Wickins, *J. Chem. Soc. A*, 1968, 400–404; (i) L. R. Florian and E. R. Corey, *Inorg. Chem.*, 1968, **7**, 722–725; (j) F. W. Moore and R. E. Rice, *Inorg. Chem.*, 1968, **7**, 2510–2514; (k) M. Cousins and M. L. H. Green, *J. Chem. Soc.*, 1964, 1567–1572. 30
30
- 18 For example, $K_2[HM_o_6VO_{22}(NH_3CH_2COO)_3] \cdot 8H_2O$ is a structurally determined compound with dinuclear $(MoO_2)_2(\mu-O)_2$ units and three absorptions at 940, 926 and 891 cm^{-1} are reported for MoO_2 stretchings in M. Cindric, N. Strukan, T. Kajfez, G. Giester and B. Kamenar, *Inorg. Chem. Commun.*, 2000, **3**, 281–284. 35
35
- 19 (a) J. Zhang, Y. Guo, W. Lu, X. Guo, M. Xu and H. Guo, *J. Ind. Eng. Chem.*, 2012, **18**, 1824–1827; (b) A. C. Coelho, M. Nolasco, S. S. Balula, M. M. Antunes, C. C. L. Pereira, F. A. Almeida Paz, A. A. Valente, M. Pillinger, P. Ribeiro-Claro, J. Klinowski and L. S. Gonçalves, *Inorg. Chem.*, 2011, **50**, 525–538; (c) R. Aguado, J. Escribano, M. R. Pedrosa, A. De Cian, R. Sanz and F. J. Arnáiz, *Polyhedron*, 2007, **26**, 3842–3848; (d) B.-Z. Lin and P.-D. Liu, *J. Mol. Struct.*, 2003, **654**, 55–60; (e) D. J. Hewkin and W. P. Griffith, *J. Chem. Soc. A*, 1966, 472–475. 40
40
- 20 (a) B. Modec, D. Dolenc and M. Kasunic, *Inorg. Chem.*, 2008, **47**, 3625–3633; (b) E. Cartuyvels, K. Van Hecke, L. Van Meervelt, C. Görller-Walrand and T. N. Parac-Vogt, *J. Inorg. Biochem.*, 2008, **102**, 1589–1598; (c) M. Cindric, T. K. Novak, S. Kraljevic, M. Kralj and B. Kamenar, *Inorg. Chim. Acta*, 2006, **359**, 1673–1680; (d) Z.-H. Zhou, S.-Y. Hou, Z.-X. Cao, H.-L. Wan and S.-W. Ng, *J. Inorg. Biochem.*, 2004, **98**, 1037–1044; (e) Z.-H. Zhou, H. Zhao and K.-R. Tsai, *J. Inorg. Biochem.*, 2004, **98**, 1787–1794; (f) C. F. Edwards, W. P. Griffith, A. J. P. White and D. J. Williams, *J. Chem. Soc., Dalton Trans.*, 1993, 3813–3819. 45
45
- 21 (a) N. Gharah, B. Chattopadhyay, S. K. Maiti and M. Mukherjee, *Transition Met. Chem.*, 2010, **35**, 531–539; (b) S. K. Maiti, K. M. A. Malik, S. Gupta, S. Chakraborty, A. K. Ganguli, A. K. Mukherjee and R. Bhattacharyya, *Inorg. Chem.*, 2006, **45**, 9843–9857; (c) B. Korpar-Colig, M. Cindric, D. Matkovic-Calogovic, V. Vrdoljak and B. Kamenar, *Polyhedron*, 2002, **21**, 147–153. 50
50
- 22 M. Cindric, N. Strukan, M. Devcic and B. Kamenar, *Inorg. Chem. Commun.*, 1999, **2**, 558–560. 55
55
- 23 P. J. Stone, J. C. Craig and H. W. Thompson, *J. Chem. Soc.*, 1958, 52–54. 50
50
- 24 R. A. Heacock and L. Marion, *Can. J. Chem.*, 1956, **34**, 1782–1795. 55
55
- 25 C. C. Watson, *Spectrochim. Acta*, 1960, **16**, 1322–1327. 50
50
- 26 T. Shimanouchi, *Table of Molecular Vibrational Frequencies, Vol. 1*, U.S. National Standards Reference Data System, 1972. 55
55

- 1 33 B. B. Koleva, *Vib. Spectrosc.*, 2007, **44**, 30–35. 1
- 34 M. Hayatifar, F. Marchetti, G. Pampaloni and S. Zacchini, *Inorg. Chem.*, 2013, **52**, 4017–4025.
- 5 35 (a) R. J. Butcher, B. R. Penfold and E. Sinn, *J. Chem. Soc., Dalton Trans.*, 1979, 668–675; (b) E. I. Stiefel, *Prog. Inorg. Chem.*, 1977, **22**, 1–223; (c) R. H. Fenn, *J. Chem. Soc. A*, 1969, 1764–1769.
- 10 36 The condensation of two Mo–OH groups leads to the formation of a Mo–O–Mo bridge and does not involve a pH change: $2\text{Mo-OH} \rightarrow \text{Mo-O-Mo} + \text{H}_2\text{O}$, hence the stoichiometric ratio $\text{H}^+/\text{Mo} = 4$ required for the formation of $\text{Mo}_2\text{O}_4(\text{OH})_4(\text{aaH})$ (see above) is not affected by successive condensation reactions leading to $\{\text{Mo}_2\text{O}_4(\text{OH})_2(\mu\text{-O})(\text{aaH})\}_n$.
- 15 37 M. Dittmann and E. Schweda, *Z. Anorg. Allg. Chem.*, 1998, **624**, 2033–2037.
- 38 (a) J. Z. Hu, J. K. Harper, C. Taylor, R. J. Pugmire and D. M. Grant, *J. Magn. Reson.*, 2000, **142**, 326–330; (b) X. Wu, S. T. Burns and K. W. Zilm, *J. Magn. Reson., Ser. A*, 1994, **111**, 29–36.
- 20 39 C. E. Crouthamel and C. E. Johnson, *Anal. Chem.*, 1954, **26**, 1284–1291.
- 25 40 Calculated values refer to the formula $\text{Mo}_2\text{O}_4(\text{OH})_4(\text{aaH})$, where aaH = amino acid. The corresponding 25
- experimental results are generally lower for H and higher for Mo, N and C, respectively. This can be explained by the formation of $\{\text{Mo}_2\text{O}_4(\text{OH})_2(\text{aaH})(\mu\text{-O})\}_n$ species *via* water loss (see eqn (3)).
- 41 C. Y. Lin, M. W. George and P. M. W. Gill, *Aust. J. Chem.*, 2004, **57**, 365–370. 5
- 42 (a) M. Dolg, in *Modern Methods and Algorithms of Quantum Chemistry*, ed. J. Grotendorst, John Neumann Institute for Computing, NIC series, Jülich, 2000, pp. 479–508; (b) P. J. Hay and W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 299–310; (c) P. J. Hay and W. R. Wadt, *J. Chem. Phys.*, 1985, **82**, 270–283; (d) W. J. Hehre, R. Ditchfield and J. A. Pople, *J. Chem. Phys.*, 1972, **56**, 2257–2261. 10
- 43 J. P. Pedrew, K. Bulke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865–3868. 15
- 44 S. Grimme, *J. Comput. Chem.*, 2006, **27**, 1787–1789.
- 45 A. Klamt and G. Schüürmann, *J. Chem. Soc., Perkin Trans. 2*, 1993, 799–805.
- 46 (a) F. Jensen, in *Introduction to Computational Chemistry*, Wiley, Chichester, 2nd edn, 2007; (b) C. J. Cramer, in *Essentials of Computational Chemistry*, Wiley, Chichester, 2nd edn, 2004. 20
- 30 30
- 35 35
- 40 40
- 45 45
- 50 50
- 55 55