

Heart Institute
University of São Paulo Medical
School
Sao Paulo, SP Brazil

References

- Hannan EL, Wu C, Smith CR, Higgins RS, Carlson RE, Culliford AT, et al. Off-pump versus on-pump coronary artery bypass graft surgery: differences in short-term outcomes and in long-term mortality and need for subsequent revascularization. *Circulation*. 2007;116:1145-52.
- Polomsky M, He X, O'Brien SM, Puskas JD. Outcomes of off-pump versus on-pump coronary artery bypass grafting: Impact of preoperative risk. *J Thorac Cardiovasc Surg*. 2013;145:1193-8.
- Mejia OA, Lisboa LA, Puig LB, Moreira LF, Dallan LA, Jatene FB. On-pump or off-pump? Impact of risk scores in coronary artery bypass surgery. *Rev Bras Cir Cardiovasc*. 2012;27:503-11.
- Puskas JD, Thourani VH, Kilgo P, Cooper W, Vassiliades T, Vega JD, et al. Off-pump coronary artery bypass disproportionately benefits high-risk patients. *Ann Thorac Surg*. 2009;88:1142-7.
- Tatoulis J. Giant leaps in surgical myocardial revascularisation. *Heart Lung Circ*. 2011;20:149-56.

<http://dx.doi.org/10.1016/j.jtcvs.2013.07.085>

B-TYPE NATRIURETIC PEPTIDE IN CHILDREN UNDERGOING PEDIATRIC CARDIAC SURGERY: JUST A MARKER OF DISEASE SEVERITY STRONGLY RELATED TO AGE OR MUCH MORE?

To the Editor:

We read with interest the article of Radman and colleagues¹ entitled, "The effect of preoperative nutritional status on postoperative outcomes in children undergoing surgery for congenital heart defects in San Francisco (UCSF) and Guatemala City (UNICAR)," recently published in the Journal. We agree with Radman and colleagues¹ that body fat mass and acute and chronic malnourishment are important determinants of worse outcome in children undergoing surgery¹⁻⁴; the relationship of these factors to preoperative B-type natriuretic peptide (BNP) values, however, merits further consideration.

Actually, Radman and colleagues¹ included in their study a wide

spectrum of ages (interquartile range, 3.2-47.1 months) and congenital heart diseases (from simple septal defects to univentricular heart). It has been widely demonstrated that preoperative BNP values as well as postoperative BNP variations should be interpreted as the consequences of 2 major factors: disease severity and age.^{2,3} The influence of these variables is particularly remarkable in the neonatal and infant setting, because neonatal cardiac surgery usually carries a higher surgical risk as a result of the severity of the disease²⁻⁴ and many maturational variations in endocrine function within the first month of life.^{2,3} The cardiac natriuretic peptide system is a relevant component of a complex and integrated network that includes the endocrine, nervous, and immune systems.⁵ For the same age and disease severity, a huge number of biologic substances, environmental factors, and physiologic parameters contribute to BNP response⁵; body fat mass and malnutrition are only 2 factors among these. Unfortunately, the small sample size (only 71 patients) of the study by Radman and colleagues¹ does not allow a risk stratification analysis that includes all these parameters. In particular, low body fat mass and malnutrition, frequently observed in children with more severe cardiac defects,^{1,4} are important factors that contribute to the severity of the disease state and in turn to the rise in BNP values.

In conclusion, recent data support the use of BNP as a useful adjunct prognostic and disease severity marker in children undergoing cardiac surgery. Both preoperative values and postsurgical variations in BNP should be interpreted first according to age and disease severity^{2,3} and only secondarily as the consequence of additional factors, including malnutrition and body fat mass.

Massimiliano Cantinotti, MD^a
Giorgio Iervasi, MD^{a,b}

Aldo Clerico, MD^{a,c}
^aFondazione Toscana G. Monasterio
Massa, Italy
^bInstitute of Clinical Physiology
Pisa, Italy
^cScuola Superiore Sant'Anna
Pisa, Italy

References

- Radman M, Mack R, Barnoya J, Castanada A, Rosales M, Azakie A, et al. The effect of preoperative nutritional status on postoperative outcomes in children undergoing surgery for congenital heart defects in San Francisco (UCSF) and Guatemala City (UNICAR). *J Thorac Cardiovasc Surg*. April 9, 2013 [Epub ahead of print].
- Cantinotti M, Lorenzoni V, Storti S, Moschetti R, Murzi B, Crocetti M, et al. Thyroid and brain natriuretic peptide response in children undergoing cardiac surgery for congenital heart disease—age-related variations and prognostic value. *Circ J*. 2012;77:188-97.
- Amimov R, Keller RL, Herrera C, Hsu JH, Datar S, Karl TR, et al. B-type natriuretic peptide levels predict outcomes in infants undergoing cardiac surgery in a lesion-dependent fashion. *J Thorac Cardiovasc Surg*. 2013;145:1279-87.
- Lacour-Gayet F, Clarke D, Jacobs J, Comas J, Daebritz S, Daenen W, et al. The Aristotle score: a complexity-adjusted method to evaluate surgical results. *Eur J Cardiothorac Surg*. 2004;25:911-24.
- Clerico A, Giannoni A, Vittorini S, Passino C. Thirty years of the heart as an endocrine organ: physiological role and clinical utility of cardiac natriuretic hormones. *Am J Physiol Heart Circ Physiol*. 2011; 301:H12-20.

<http://dx.doi.org/10.1016/j.jtcvs.2013.07.086>

TAVI WITHOUT SURGICAL STANDBY: IS HISTORY REPEATING ITSELF? A WORD OF CAUTION

To the Editor:

Current practice guidelines require that transcatheter aortic valve implantation (TAVI) be performed by a heart team composed of interventional cardiologists and surgeons in a hybrid operating room.¹ The occurrence of a case, treated in a hospital without cardiac surgery on site, in which the implanted prosthesis migrated into the left ventricle and required surgical removal, prompted this report.

A 69-year-old woman underwent TAVI in a facility without on-site cardiac surgery. Shortly after transfemoral implantation, the prosthesis