

Environ Sci Pollut Res (2015) 22:5711–5741
DOI 10.1007/s11356-014-3974-5

REVIEW ARTICLE

Bisphenol A, nonylphenols, benzophenones, and benzotriazoles in soils, groundwater, surface water, sediments, and food: a review

Alessandro Careghini · Andrea Filippo Mastorgio ·
Sabrina Saponaro · Elena Sezenna

Received: 6 October 2014 / Accepted: 8 December 2014 / Published online: 30 December 2014
© The Author(s) 2014. This article is published with open access at Springerlink.com

Abstract Contaminants of emerging concern (CECs) are not commonly monitored in the environment, but they can enter the environment from a variety of sources. The most worrying consequence of their wide use and environmental diffusion is the increase in the possible exposure pathways for humans. Moreover, knowledge of their behavior in the environment, toxicity, and biological effects is limited or not available for most CECs. The aim of this work is to edit the state of the art on few selected CECs having the potential to enter the soil and aquatic systems and cause adverse effects in humans, wildlife, and the environment: bisphenol A (BPA), nonylphenol (NP), benzophenones (BPs), and benzotriazole (BT). Some reviews are already available on BPA and NP, reporting about their behavior in surface water and sediments, but scarce and scattered information is available about their presence in soil and groundwater. Only a few studies are available about BPs and BT in the environment, in particular in soil and groundwater. This work summarizes the information available in the literature about the incidence and behavior of these compounds in the different environmental matrices and food. In particular, the review focuses on the physical-chemical properties, the environmental fate, the major degradation byproducts, and the environmental evidence of the selected CECs.

Keywords Contaminants of emerging concern · Soil · Groundwater · Surface water · Sediments · Food

Responsible editor: Leif Kronberg

A. Careghini · A. F. Mastorgio · S. Saponaro (✉) · E. Sezenna
DICA - Sezione Ambientale, Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133 Milan, Italy
e-mail: sabrina.saponaro@polimi.it

Introduction

Contaminants of emerging concern (CECs) are defined as any synthetic or naturally occurring chemical that is not commonly monitored in the environment, though having the potential to enter soil and aquatic ecosystems and cause adverse effects in humans, wildlife, and the environment. CECs include synthesized and commercialized chemicals that have just gained entry into the environment and a range of chemicals that have been produced and released into the environment for long, for which new concerns (occurrence, fate, adverse effects on human health and the environment) have recently raised (Focazio et al. 2008). CECs include industrial, agricultural, and household chemicals, such as flame retardants and plasticizers, pesticides, gasoline additives, fluorinated compounds and nanomaterials, as well as pharmaceuticals and personal care products, such as human and veterinary antibiotics and drugs, fragrances, sunscreen agents, antimicrobial cleaning agents, and excipients (Daughton and Ternes 1999; Thomaidis et al. 2012; EUGRIS 2013; Molnar et al. 2013; NORMAN 2013).

CECs can enter the environment from a variety of sources, such as industrial wastes, drain from urbanized areas and transportation systems, sewage treatment plants, atmospheric deposition, etc. (Kolpin et al. 2002; Anderson et al. 2012). On agricultural land, field application of biosolids (manure or sludge) and polymers for modern intensive agriculture (i.e., mulch films, drip irrigation tubes, string, clips, pots, etc.) and irrigation with reclaimed water can be significant sources of CECs (Kolpin et al. 2002; Christian et al. 2003; Kumar et al. 2005). Moreover, due to their continuous release into the environment, these contaminants can accumulate and cause adverse effects in ecosystems, as their transformation/removal rate can be exceeded by their high loading rate (Anderson et al. 2012; USEPA 2014a).

The most worrying consequence of wide use and environmental diffusion of CECs is the increase in the possible exposure pathways for humans, such as ingestion of food plants cultivated on contaminated land or irrigated with reclaimed water, ingestion of meat/animal products from pasture on contaminated land, and consumption of tap water from polluted groundwater or surface water (Weber et al. 2005; Molnar et al. 2013). However, knowledge about the behavior in soil-water systems, toxicity, and biological effects is limited or not available for most CECs; thus, human exposure and related health effects and potential toxicological significance in terrestrial and aquatic ecosystems are mostly unknown and caution is advised. Accordingly, regulatory concentration limits or sound guidance and standard or trigger values for the environmental media have not been established yet (Molnar et al. 2013).

In soil, the behavior of organic contaminants is governed by a variety of complex dynamic physical, chemical, and biological processes, including sorption/desorption, volatilization, leaching, chemical and biological degradation, plant uptake, and runoff (Arias-Estévez et al. 2008). These processes directly control contaminant mobility and fate through the soil and their transfer from soil to water, air, or food. The rate and relative importance of these processes vary with the chemical nature of the contaminant and the chemical, biological, and hydraulic properties of soil (Kibbey et al. 2007). Some compounds, though at trace levels in the sources, accumulate in soils (Kinney et al. 2006; Ternes et al. 2007; Xu et al. 2009), whereas others easily runoff from soil into surface waters or leach to groundwater affecting water reservoirs (Koschorreck et al. 2002). Transport mechanisms related to colloidal material have been underlined for some CECs (Yamamoto and Liljestrand 2003; Zhou et al. 2007).

Most degradation studies were carried out in the aqueous environment (Richardson and Bowron 1985; Buser et al. 1998; Zwiener and Frimmel 2003; Lin and Reinhard 2005; Yu et al. 2006), sewage sludge (Kimura et al. 2007; Zhao et al. 2008), or sediments (Ying and Kookana 2005). Only few studies investigated CEC degradation in soil (Tolls 2001; Gao and Pedersen 2005; Ying and Kookana 2005; Williams and Adamsen 2006; Chefetz et al. 2008; Xuan et al. 2008), showing that some organic contaminants are biodegradable at a certain extent, whereas others exhibit very slow biodegradation rates or are sequestered within soil particles, being inaccessible for microbial degradation. Moreover, most of the previous studies on soil focused on sorption/desorption at equilibrium conditions and degradation under optimal conditions in batch tests, instead of assessing transport dynamics under field conditions (Wehrhan et al. 2007).

The aim of this work is to edit the state of the art on few selected contaminants of emerging concern (bisphenol A, nonylphenol, benzophenones, and benzotriazole) that have

been already measured in many European environmental samples and belong to different classes of widely used emerging substances (plasticizers, surfactants, personal care products, and industrial chemicals) (EUGRIS 2013; Molnar et al. 2013; NORMAN 2013). In particular, the review focuses on their physical-chemical properties in relation with their environmental fate and transport, major degradation byproducts, and environmental evidence.

Bisphenol A (BPA)

Bisphenol A (2,2-bis(4-hydroxyphenyl)propane) is an organic compound composed of two phenol molecules bonded by a methyl bridge and two methyl groups (Table 1).






BPA is used as an intermediate (binding, plasticizing, and hardening) in plastics, paints/lacquers, binding materials, and filling materials. Furthermore, it is used as an additive for flame-retardants, brake fluids, and thermal papers. About 95 % of BPA produced in industry is used to make plastics, in particular polycarbonate resins (71 %) and epoxy resins (29 %) (RIKZ 2001; Huang et al. 2012). Due to the increasing demand for polycarbonates and epoxy resins, BPA production has constantly grown in the last years: the global demand was 3.2, 3.9, and 5.0 million tons in 2003, 2006, and 2010, respectively (Flint et al. 2012; Huang et al. 2012).

Toxicity

BPA is listed as an endocrine disrupter. It has been proven to have estrogenic activity even at concentrations below $1 \mu\text{g}/\text{m}^3$ (Rykowska and Wasiak 2006). Estrogenic compounds can have deleterious effects on living organisms because they can disrupt natural hormone balance in both men and women. The effects of exposure to BPA can be particularly harmful to fetus, infants, and young children, because of lack of feedback regulating the activity, synthesis, and elimination of hormones (RIKZ 2001; Rykowska and Wasiak 2006). The acute toxicity of BPA is relatively low. In subacute toxicity studies, a marked reduction in the rate of body weight increase was observed in treated animals (RIKZ 2001). There is limited evidence for carcinogenicity in animals; according to the IARC classification, BPA belongs to group 3 (“not classifiable as to its carcinogenicity to humans”) (IARC 1999).

Ingestion through contaminated food is the major exposure pathway for humans; inhalation and dermal contact are significant exposure pathways for workers involved in the manufacture of BPA. Rykowska and Wasiak (2006) recommended a reference dose (RfD) for oral exposure of $0.01 \text{ mg}/\text{kg}/\text{day}$. USEPA estimated a reference dose of $50 \mu\text{g}$ of BPA/kg of body weight/day (USEPA-IRIS 2014).

Table 1 List of chemical compounds studied in the paper

Compound	CAS number	Formula	Chemical Structure
Bisphenol A (BPA)	80-05-7	C ₁₅ H ₁₆ O ₂	
Nonylphenol (NP)	104-40-5	C ₁₅ H ₂₄ O	
Benzophenone (BP)	119-61-9	C ₁₃ H ₁₀ O	
Benzophenone-3 (BP-3)	131-57-7	C ₁₄ H ₁₂ O ₃	
Benzotriazole (BT)	95-14-7	C ₆ H ₅ N ₃	

Environmental fate and transport

BPA is not produced naturally; it can be released into the environment during production and transport operations, from many products during their use or after their disposal in landfill, through effluent from wastewater treatment plants and from sewage sludge used in agriculture (Huang et al. 2012).

BPA is a moderately water-soluble compound at ambient temperature (Table 2). It has low vapor pressure and does not tend to volatilize significantly from water or dry soil surfaces (Flint et al. 2012). Based on the organic carbon/water partition coefficient (*K*_{OC}) value, significant sorption of BPA on soil and sediments is expected. Based on the octanol/water partition coefficient *K*_{OW}, BPA has modest capacity for bioaccumulation, which occurs only at high doses (Fent et al. 2003; Flint et al. 2012; Roberts et al. 2014).

Hydrolysis is expected to be negligible under environmental conditions due to the absence of hydrolysable groups, but BPA not tied to organic matter undergoes photolysis in water at wavelengths above 290 nm (RIKZ 2001). Pseudo-first-order degradation constants between 0.0053 and 0.008 l/min were observed in aqueous solutions with 10 g/m³ of humic substances and BPA between 1 and 20 g/m³; no photodegradation occurred in pure water (Zhan et al. 2006).

BPA can be readily biodegraded in soil and sediments under aerobic conditions, with estimated half-life values in

soils between 3 and 37.5 days. No degradation was observed in anaerobic soils during 70 day experiments or in anoxic estuarine sediments during 120 day experiments (RIKZ 2001; Fent et al. 2003; Flint et al. 2012; Yu et al. 2013; Chang et al. 2014; Yang et al. 2014). BPA is not expected to be persistent in the environment (USEPA 2010a; Michałowicz 2014).

Degradation byproducts

Many bacterial strains capable of growing on BPA as a sole source of carbon and energy were isolated from different environmental matrices; they included both gram-negative and gram-positive strains (Zhang et al. 2013a). Different degradation pathways for BPA have been proposed in the literature. In particular, biodegradation of BPA proceeds via complicated metabolic routes that leads to formation of several kinds of byproducts (Spivack et al. 1994; Ike et al. 2002; Zhan et al. 2006; Ye et al. 2011).

Zhan et al. (2006) proposed a photodegradation pathway in aqueous solution with humic substances based on the results of structural analyses for intermediate photoproducts. Mono-hydroxylated BPA, glycerol, 2-hydroxy-propanoic acid, and *p*-hydroquinone were identified as degradation products.

Zhang et al. (2007) studied BPA degradation by a microbial strain isolated from the compost leachate of a municipal solid waste; 4-hydroxybenzaldehyde, 4-hydroxybenzoic acid, and

Table 2 Physical-chemical properties of BPA, 4-NP, BP, BP-3, and BT

Properties	Units	BPA	4-NP	BP	BP-3	BT
Molar weight	g/mol	228.3	220.3	182.2	228.2	119.1
Melting point	°C	158 (RIKZ 2001)	-8 (EC 2002)	49 (Rajendra 2000; Jeon et al. 2006; IARC 2010) Insoluble (Rajendra 2000; IARC 2010)	68 (Jeon et al. 2006)	99 (DECOS 2000)
Water solubility (20 °C)	g/m ³	120–300 (RIKZ 2001; Kalmykova et al. 2013)	6.0 (EC 2002)		28.6 (Rodil et al. 2009)	20,000–28,000 (DECOS 2000; Harris et al. 2007)
Henry's law constant	Pa m ³ /mol	10 ⁻⁵ –10 ⁻⁶ (RIKZ 2001; Fent et al. 2003)	0.84–11.02 (CCME 2002; EC 2002; Soares et al. 2008)	0.19 (IARC 2010)	1.5 × 10 ⁻³ (Liu et al. 2012a)	1.49 × 10 ⁻² (Liu et al. 2011a)
Vapor pressure (25 °C)	Pa	1.1 × 10 ⁻⁷ –5.3 × 10 ⁻⁶ (RIKZ 2001; Kalmykova et al. 2013)	2.07 × 10 ⁻² –0.3 (EC 2002; Soares et al. 2008)	0.26 (IARC 2010)	7.01 × 10 ⁻⁴ (Guidechem 2013)	5.3 (at 20 °C) (TOXNET 2013a)
Octanol-water partition coefficient log(<i>K</i> _{ow})	–	2.2–3.4 (RIKZ 2001; Kalmykova et al. 2013)	4.48–5.76 (EC 2002; Soares et al. 2008)	3.18–3.38 (Rajendra 2000; IARC 2010)	3.52–3.82 (Jeon et al. 2006; Brunete et al. 2011; Liu et al. 2012a)	1.23–1.44 (Cornell et al. 2000; Bi et al. 2007; Zhang et al. 2011)
Organic carbon-water partition coefficient log(<i>K</i> _{oc})	–	2.5–4.5 (RIKZ 2001; Fent et al. 2003; Flint et al. 2012)	3.4–5.6 (Sekela et al. 1999; Hou et al. 2006)	2.6–2.7 (TOXNET 2013b)	3.43 (Guidechem 2013)	1.27–1.97 (Yu et al. 2009; Liu et al. 2011a)

p-hydroquinone were the observed metabolic intermediates. Also, Dodgen et al. (2014) detected 4-hydroxybenzaldehyde and 4-hydroxybenzoic acid, together with 4-hydroxyacetophenone, as transformation products of BPA in degradation tests using different artificially contaminated soils.

Spivack et al. (1994) studied the degradation pathway for a gram-negative aerobic bacterium (pure culture). The major route (80 %) was BPA cleavage to *p*-hydroxyacetophenone (*p*-HAP) and *p*-hydroxybenzaldehyde (*p*-HBAL), followed by further degradation via *p*-hydroxybenzoic acid (*p*-HBA). The remaining 20 % of BPA was converted into 2,3-bis-(4-hydroxyphenyl)-1,2-propanediol (tetraol-IV) via bis-(4-hydroxyphenyl)-1-propanol. Although tetraol-IV can be slowly degraded to *p*-hydroxyphenacyl alcohol (*p*-HPOH), both these byproducts accumulated in the medium. In Ike et al. (2002), BPA degradation by mixed microbial consortia from activated sludge or river water also led in most cases to the accumulation of the minor route byproducts tetraol-IV and *p*-HPOH, generally identified as the dead-end compounds. Based on the studies reported above, BPA photodegradation and biodegradation do not seem to mineralize the compound.

Ye et al. (2011) investigated BPA metabolism in rat and human liver microsomes. The oxidative metabolism of BPA to BPA catechol was a major pathway when using male rat microsomes, but only a minor pathway (less than 10 % BPA catechol was formed) when using human liver microsomes.

Environmental evidence

The major studies reported in the literature about BPA presence in environmental matrices and food are reported in Table 3. Industrial activities (mainly chemical plants) and wastewater treatment plants were the major sources of BPA in the surface waters and sediments. High concentrations in soil and groundwater were detected especially for agricultural fields irrigated with treated wastewater and/or amended with biosolids or near landfills (Heemken et al. 2001; Kawahata et al. 2004; Cespedes et al. 2005; Vethaak et al. 2005; Loos et al. 2007; Yoon et al. 2010; Félix-Cañedo et al. 2013; Wu et al. 2013; Gorga et al. 2014; Michałowicz 2014).

BPA in soil and sediments

BPA concentrations in soils span between 0.55 and 147 µg/kg on dry weight basis (d.w.), with higher values generally found in agricultural fields amended with biosolids or irrigated with wastewater. Kinney et al. (2008) observed higher maximum concentrations of BPA in soils at a not-amended site (147 µg/kg d.w.) than at a site amended with biosolids (81 µg/kg d.w.); no detectable concentration of BPA was observed in site receiving liquid swine manure. Low values were observed by Xu et al. (2008) in soils from a golf course in southern

Table 3 BPA concentrations in various environmental matrices and in food (percentages between brackets represent the detection frequency)

Reference	Location	Units	Value
Soils			
Kinney et al. (2008)	Agricultural fields, USA	µg/kg d.w.	<32–147 Mean 59
Xu et al. (2008)	Golf course irrigated with reclaimed wastewater, California, USA	µg/kg d.w.	0.55–2
Gibson et al. (2010)	Agricultural fields irrigated with wastewater, Tula Valley, Mexico	µg/kg d.w.	1.6–30.2 Mean 8.3
Staples et al. (2010)	Soils amended with biosolids, North America (data collected in the period 1990–2006)	µg/kg d.w.	Median 1.15 95th percentile 21
Staples et al. (2010)	Soils amended with biosolids, Europe (data collected in the period 1990–2006)	µg/kg d.w.	Median 0.24 95th percentile 140
USEPA (2010a)	Range of values in USA	µg/kg d.w.	4–14 Mean 6–7
Sediments			
Heemken et al. (2001)	Elbe River and some of its tributaries, Germany	µg/kg d.w.	66–343 Mean 163
Kawahata et al. (2004)	Estuarine and marine sediments from Okinawa and Ishigaki Islands, Japan	µg/kg d.w.	<0.5–13 Mean 3.2
Vethaak et al. (2005)	Fresh, marine, and estuarine sediments, The Netherlands	µg/kg d.w.	<1.1–43 Median 3.2 (78 %)
Fu et al. (2007)	Estuarine and marine sediments from Jiaozhou Bay and surrounding rivers, China	µg/kg d.w.	0.7–27.3
Pojana et al. (2007)	Sediments from Venice Lagoon, Italy	µg/kg d.w.	<2.0–118 Mean 36
USEPA (2010a)	Fresh sediments, USA	µg/kg d.w.	1.4–140
USEPA (2010a)	Marine sediments, USA	µg/kg d.w.	1.5–5.0
Gorga et al. (2014)	Ebro River basin, Spain	µg/kg d.w.	<0.24–100
Michalowicz (2014)	Elba River sediments, Germany	µg/kg d.w.	10–380
Michalowicz (2014)	16 major rivers' sediments, Taiwan	µg/kg d.w.	0.37–492
Stewart et al. (2014)	Estuarine sediments from Auckland, New Zealand	µg/kg d.w.	<50–145 Mean 57
Wu et al. (2013)	Huangpu River and its tributaries, China	µg/kg d.w.	0.96–14.44 Mean 7.22
Gorga et al. (2015)	Different rivers, Spain	µg/kg d.w.	<0.24–117
Groundwater			
Lacorte et al. (2002)	Agricultural area in Catalonia, Spain	mg/m ³	<0.01–0.35
Latorre et al. (2003)	Agricultural areas in northern Spain	mg/m ³	0.05–0.18
Godejohann et al. (2009)	Ammunition disposal site, Switzerland	mg/m ³	12–13
Loos et al. (2010)	Survey on European groundwaters	mg/m ³	<0.001–2.299 Mean 0.079 90th percentile 0.073
USEPA (2010a)	Range of mean values in USA	mg/m ³	0.004–1.9
Stuart et al. (2011)	Groundwater, England	mg/m ³	up to 20
Félix-Cañedo et al. (2013)	Groundwater in Mexico City, Mexico	mg/m ³	<0.0005–0.010 (63 %)
Luo et al. (2014)	Groundwater in Europe	mg/m ³	Mean 0.079, maximum 2.299
Luo et al. (2014)	Groundwater in USA	mg/m ³	Mean 2.550
Michalowicz (2014)	Groundwater contaminated with leachate from refuse dump in Osaka, Japan	mg/m ³	740
Surface water			
Azevedo et al. (2001)	River and coastal waters, Portugal	mg/m ³	0.07–4.0 Mean 1.0
Heemken et al. (2001)	Elbe River and some of its tributaries, Germany	mg/m ³	0.017–0.776 Mean 0.105

Table 3 (continued)

Reference	Location	Units	Value
Basheer et al. (2004)	Surface coastal water, Singapore	mg/m ³	<0.002–2.47 Mean 0.40
Kawahata et al. (2004)	Estuarine and marine waters from Okinawa, and Ishigaki Islands, Japan	mg/m ³	<0.005–0.08 Mean 0.02
Cespedes et al. (2005)	Llobregat River basin, Spain	mg/m ³	<0.09–2.97 Mean 0.44
Vethaak et al. (2005)	Fresh, marine and estuarine water, The Netherlands	mg/m ³	<0.009–1.0 Median 0.045 (52 %)
Patrolecco et al. (2006)	Tiber River, Italy	mg/m ³	<0.03–0.14 Mean 0.07
Vousta et al. (2006)	Glatt River, Switzerland	mg/m ³	0.009–0.076
Fu et al. (2007)	Estuarine and marine water from Jiaozhou Bay, China	mg/m ³	0.0015–0.262
Loos et al. (2007)	River water, Belgium	mg/m ³	0.003–0.055 Mean 0.031
Loos et al. (2007)	River water, Italy	mg/m ³	<0.002–0.175 Mean 0.065
Pojana et al. (2007)	Venice Lagoon, Italy	mg/m ³	<0.001–0.145 Mean 0.014
Yoon et al. (2010)	Han River, South Korea	mg/m ³	0.0069–0.059 Mean 0.027
Yoon et al. (2010)	Effluent-dominated creeks discharging into Han River, South Korea	mg/m ³	0.011–0.120 Mean 0.062
USEPA (2010a)	Range of mean values in USA	mg/m ³	0.012–0.14
Félix-Cañedo et al. (2013)	Surface water (dams) in Mexico City, Mexico	mg/m ³	<0.0005–0.007 (52 %)
Esteban et al. (2014)	Manzanares and Jarama rivers, Spain	mg/m ³	0.006–0.126
Luo et al. (2014)	Canada	mg/m ³	Mean 0.0021 Maximum 0.087
Luo et al. (2014)	China	mg/m ³	0.006–0.881
Luo et al. (2014)	Germany	mg/m ³	0.192–0.215
Luo et al. (2014)	Greece	mg/m ³	0.055–0.152
Luo et al. (2014)	Korea	mg/m ³	0.0075–0.334
Luo et al. (2014)	UK	mg/m ³	0.006–0.068
Michalowicz (2014)	Range of concentrations in rivers, Portugal	mg/m ³	0.029–0.098
Melo and Brito (2014)	Rivers crossing Sao Luis island, Brazil	mg/m ³	<0.46
Michalowicz (2014)	Elba River, Germany	mg/m ³	4–92
Michalowicz (2014)	16 major rivers, Taiwan	mg/m ³	0.01–45
Wu et al. (2014)	Huangpu River and its tributaries, China	mg/m ³	0.0071–0.1115 Mean 0.0276
Xu et al. (2014)	Cape D' Aguilar Marine Reserve, Hong Kong, wet season	mg/m ³	0.011–0.41 Mean 0.0645
Xu et al. (2014)	Cape D' Aguilar Marine Reserve, Hong Kong, dry season	mg/m ³	0.025–0.24 Mean 0.0695
Zhang et al. (2014)	North Tai Lake Basin, Eastern China	mg/m ³	0.024–1.175 Mean 0.270
Gorga et al. (2015)	Iberian rivers (Ebro, Llobregat, Júcar and Guadalquivir)	mg/m ³	0.00011–0.649
	Food		
Basheer et al. (2004)	Seafood from supermarkets, Singapore	µg/kg f.w.	13.3–213.1 Mean 82.5
Sun et al. (2006)	Canned vegetables, fruits, and meats from local supermarkets, Singapore	µg/kg f.w.	32.8–164.5 Mean 72.5
Isobe et al. (2007)	Green mussel from India, Indonesia, Singapore, Malaysia, Thailand, Cambodia, Vietnam, and the Philippines during 1994–1999	µg/kg d.w.	1.1–13.7
Isobe et al. (2007)	Tokyo Bay	µg/kg d.w.	0.54–13.4

Table 3 (continued)

Reference	Location	Units	Value
Shao et al. (2007)	Meat/seafood from supermarkets in Beijing, China	µg/kg f.w.	<0.30–7.08 Mean 0.71
Cao et al. (2011)	Different foods from stores in Quebec City, Canada	µg/kg f.w.	0.2–106 Mean 7.7
Noonan et al. (2011)	Canned food from local supermarkets in Washington and Maryland, USA	µg/kg f.w.	<2–790 Mean 509
Gyllenhammar et al. (2012)	Fruits, meats, and vegetables commercially available, Sweden	µg/kg f.w.	<2.0–29.0 Mean 3.8
Dodgen et al. (2013)	Lettuce and collards, steam and leaves	µg/kg f.w.	0.22–3.05
Dodgen et al. (2013)	Lettuce and collards, roots	µg/kg f.w.	199.6–441.7
Li et al. (2013c)	Soft commercial drinks	mg/m ³	<0.02–0.86 Mean 0.31
Lu et al. (2013)	Vegetables and fruits in Florida, USA	µg/kg f.w.	0.2–9.0 Mean 4.2
Maggioni et al. (2013)	Drinking water from public drinking fountains, Italy	mg/m ³	<0.00073–0.102
Maggioni et al. (2013)	Bottled mineral water, Italy	mg/m ³	<0.00073–0.00113
Michałowicz (2014)	Meat products, worldwide	µg/kg f.w.	0.49–56
Michałowicz (2014)	Fish, worldwide	µg/kg f.w.	7.1–103
Michałowicz (2014)	Vegetables and fruits, worldwide	µg/kg f.w.	11–95
Michałowicz (2014)	Cereals, worldwide	µg/kg f.w.	1.0–3.8
Michałowicz (2014)	Various tinned products, including vegetables, fruits, and seafood, worldwide	µg/kg f.w.	0.1–267

California irrigated with reclaimed wastewater, but the authors pointed out the potential exposition of groundwater to contamination due to the accumulation of BPA over time. Gibson et al. (2010) investigated agricultural fields irrigated with wastewater for many years (up to 90 years) at different horizons; concentrations of BPA up to 30.2 µg/kg d.w. were measured, suggesting little evidence of BPA accumulation in soil and no evidence of transport through the different horizons.

Concentrations of BPA in sediments range between <0.24 and 492 µg/kg d.w. Values between 66 and 343 µg/kg d.w. were measured by Heemken et al. (2001) in the Elbe River; the highest concentration was observed downstream to a chemical factory. The authors monitored also the changes during 21 months of the BPA concentration in one of the most impacted sampling point, observing concentrations ranging between 127 and 322 µg/kg d.w. In Kawahata et al. (2004), the highest values were observed in areas affected by high pollution and/or downstream of commercial and industrial sites. Fu et al. (2007) studied the distribution of BPA in sediments collected in the Jiaozhou Bay (China) and at five monitoring stations located in the surrounding rivers; BPA was detected in all samples from the bay at concentrations between 0.7 and 20.3 µg/kg d.w. and at concentrations between 2.4 and 27.3 µg/kg d.w. in samples from the river sediments, resulting in similar ranges of concentrations. Pojana et al. (2007) investigated the occurrence of BPA in

sediment samples collected at four stations in the Venice Lagoon close to municipal wastewater and industrial discharges; BPA was detected in seven out of eight samples at values up to 118 µg/kg d.w. in the sampling station nearest to the plant discharge.

BPA in groundwater and surface water

Reported concentrations of BPA in groundwater vary between 0.001 and 20 mg/m³. Lacorte et al. (2002) analyzed groundwater collected at an agriculture area of Catalonia (Spain) treated with a pesticide containing traces of BPA (1.5 mg/m³ in the pesticide), resulting in concentrations between <0.01 and 0.35 mg/m³. Latorre et al. (2003) performed measurements in groundwater collected from agriculture sites in northern Spain; the highest level (>1.5 mg/m³) was found next to grape cultivations. Godejohann et al. (2009) performed systematic analyses of groundwater near a former ammunition destruction site in Switzerland; BPA concentrations were about 12–13 mg/m³. In USEPA (2010a), the average concentrations of BPA in groundwater in the USA vary between 0.0041 and 1.9 mg/m³, with a range of values of 0.006–2.55 mg/m³. Stuart et al. (2011) reported about a survey on micropollutants in groundwater in England, with BPA concentrations up to 20 mg/m³; the distribution of detections was clearly split into two areas, southern England and Midlands.

BPA in surface water is reported between <0.001 and 92 mg/m^3 . Azevedo et al. (2001) detected BPA in 51 % of the analyzed samples of river and coastal waters from Portugal; authors observed values greater than 2 mg/m^3 only in two sampling points located near industrial areas. Heemken et al. (2001) measured the concentration of BPA in the Elbe River and in some of its tributaries between 0.017 and 0.776 mg/m^3 ; BPA probably originated from an industrial plant manufacturing epoxy resins at the German-Czech border and from a municipal sewage plant. Basheer et al. (2004) collected surface seawater samples at different locations along the Singapore coastline, encompassing both inshore and offshore sampling locations; BPA concentrations were between <0.002 and 2.47 mg/m^3 . In general, BPA concentration in samples obtained from offshore locations were lower than inshore samples. Kawahata et al. (2004) measured appreciable concentrations of BPA in water samples (between 0.036 and 0.08 mg/m^3) in the most populated areas; values below detection limits (0.005 mg/m^3) were measured at other locations. Vethaak et al. (2005) measured low concentrations of BPA (<0.009 – 1 mg/m^3) in half of Dutch rainwater and surface water samples; the majority of samples had nanograms per liter, with few higher exceptions. Patrolecco et al. (2006) investigated the presence of BPA in water along the Tiber River (Italy); BPA concentrations were rather constant along the studied tract both in summer (0.06 – 0.09 mg/m^3) and in winter (<0.03 – 0.14 mg/m^3). Loos et al. (2007) monitored river waters in two textile industry regions, in Belgium (south Ghent) and in Italy (south Como). Similar values were obtained in both cases, with Belgian concentrations between <0.002 and 0.055 mg/m^3 and Italian data between <0.002 and 0.175 mg/m^3 . Fu et al. (2007) measured BPA concentrations between 0.0015 and 0.0925 mg/m^3 in water samples in the Jiaozhou Bay; higher values (an order of magnitude) were found in rivers surrounding the bay (0.0168 – 0.262 mg/m^3). Pojana et al. (2007) measured BPA in water samples in the Venice Lagoon, with values between <0.001 and 0.145 mg/m^3 ; the highest values were measured at the sampling stations in the proximity of an industrial effluent discharge point and of a municipal and industrial wastewater treatment plant. USEPA (2010a) reported a range of BPA average concentration in surface water of 0.012 – 0.14 mg/m^3 and a range of data of 0.0009 – 12 mg/m^3 . Wu et al. (2013) measured concentrations of BPA in waters from Huangpu River and its tributaries (China) in winter and in summer, but the BPA levels in surface waters showed no clear seasonal pattern. Also Xu et al. (2014) did not observe seasonal variation of BPA in the seawaters of a marine reserve in Hong Kong. Gorga et al. (2015) carried out an extensive monitoring campaign on several Spanish rivers observing concentrations of BPA ranging from <0.00011 to 0.649 mg/m^3 ; the higher values were found in well-known contaminated sites, near big cities or near industrial areas in the Llobregat and Ebro river basins.

BPA in biota and food

Many authors studied the migration of BPA and its derivatives from polymer packaging into food, especially under the effect of high temperature (e.g., microwave heating). Polycarbonate hydrolysis is the dominant mechanism responsible for BPA release from the polymer surface to the contacting liquids (Mercea 2009). Concentrations between 0.1 and $790 \text{ } \mu\text{g/kg}$ fresh weight (f.w.) were found in food and <0.00073 and 0.86 mg/m^3 in drinking waters/commercial drinks from different countries. The highest values in canned food were due to epoxy resins used as internal coating. For this reason, since 2001, epoxy resin films have been largely replaced with polyethylene terephthalate films (Huang et al. 2012; Wright-Walters et al. 2011). Basheer et al. (2004) measured concentrations of BPA in seafood samples (prawn, crab, blood cockle, white clam, squid, fish) purchased from a local supermarket in Singapore, resulting in significant values in all samples (between 13.3 and $213.1 \text{ } \mu\text{g/kg}$ f.w.); the maximum value was found in crab. Sun et al. (2006) investigated canned food samples purchased in Singapore; detectable amounts of BPA were found in all samples, with concentrations between 32.8 and $164.5 \text{ } \mu\text{g/kg}$ f.w. Shao et al. (2007) measured BPA in different types of fresh meat purchased in Beijing; BPA was detectable in 13 out of 27 samples, with concentrations between 0.33 and $7.08 \text{ } \mu\text{g/kg}$ f.w. High concentrations of BPA were found in aquatic animals (fish and duck meat), due to the contamination of the aquatic environment. Cao et al. (2011) analyzed foods from four different stores in Quebec City (Canada); among the 154 food samples, BPA was detected in 55 samples, with concentrations from 0.20 to $106 \text{ } \mu\text{g/kg}$ f.w. BPA levels in raw vegetable samples (not canned) were low, while BPA was detected at high concentrations in all the canned samples. BPA was not detected in any of the bottled water samples. Noonan et al. (2011) examined 78 canned and 2 frozen foods purchased from retail stores in Washington, DC, and in the surrounding Maryland (USA). BPA was detected in 71 out of 78 samples of canned foods, at concentration from <2 to $730 \text{ } \mu\text{g/kg}$ f.w.; BPA was not detected in the frozen foods. The concentrations in canned fruits were lower than in other canned foods; data were consistent with the general industry practice of using tin and not epoxy phenolic films in canned fruit containers. Gyllenhammar et al. (2012) analyzed the levels of BPA in food from a typical food market basket purchased from two store chains in four major Swedish cities. Concentrations above the limit of quantification ($2 \text{ } \mu\text{g/kg}$ f.w.) were found in fish (2.5 – $29 \text{ } \mu\text{g/kg}$ f.w.), meats (6.9 – $13 \text{ } \mu\text{g/kg}$ f.w.), potatoes ($2.2 \text{ } \mu\text{g/kg}$ f.w.), and dairy products ($2.4 \text{ } \mu\text{g/kg}$ f.w.). Lu et al. (2013) analyzed BPA in fresh fruits and vegetables commercially available in Florida (USA); concentrations between 0.2 and $4.3 \text{ } \mu\text{g/kg}$ f.w. and between 2 and $9 \text{ } \mu\text{g/kg}$ f.w., respectively, were found. Dodgen et al. (2013) investigated the accumulation of BPA in different parts

of lettuce and collards, pointing out the poor translocation of BPA from roots to the upper tissues after uptake: in the stem and leaves, concentration ranges 0.22–0.36 and 1.42–3.05 $\mu\text{g}/\text{kg}$ f.w. were observed for lettuce and collards, respectively, but the values measured in the roots were greater by 3–2 orders of magnitude (441.7 and 199.6 $\mu\text{g}/\text{kg}$ f.w.). Maggioni et al. (2013) evaluated concentrations of BPA in drinking waters from public fountains in 35 Italian cities and in bottled mineral waters; the highest concentration of BPA, 0.102 mg/m^3 , was detected only in one sample in water from public fountains, while in the other samples, the amount was only slightly above the quantification limit (0.00073 mg/m^3).

Based on BPA concentrations in food and food consumption, a daily dietary BPA intake of 0.02–0.081 $\mu\text{g}/\text{kg}/\text{day}$ for adults and 0.22–0.33 $\mu\text{g}/\text{kg}/\text{day}$ for infants was estimated (Basheer et al. 2004; Sun et al. 2006; Shao et al. 2007; Cao et al. 2011; Gyllenhammar et al. 2012; Noonan et al. 2011; Lu et al. 2013). At present, the dietary intake of BPA appears to be the primary source for human exposure.

Nonylphenol (NP)

NP is a term used to refer to a wide group of isomeric compounds ($\text{C}_{15}\text{H}_{24}\text{O}$) consisting of a nine-carbon alkyl chain bond to a phenol ring (Table 1). The various isomers can differ either in the degree of alkyl chain branching or in the position on the phenol ring. The NP isomers most produced and measured in the environment is 4-NP. NP is used as a formulant in pesticides, as a lubricating oil additive, as a catalyst in epoxy resins curing, at industrial laundries and, in the past, to produce nonylphenol ethoxylates (NPEs) for consumer products (e.g., surfactants, detergents, wetting agents, dispersants, defoamers, de-inkers, antistatic agents) (CCME 2002; European Parliament 2003; USEPA 2010b). As tris(4-NP)phosphite, it is an antioxidant for the stabilization of rubber, vinyl polymers, polyolefins, and styrenics. In 2010, the US demand for NP was 380 million pounds (ICIS 2007 in USEPA 2010b).

Toxicity

NP is an estrogen agonist (ECHA 2014). It is highly irritating and corrosive to skin and eyes, but it does not have significant skin-sensitizing potential. The acute (oral and dermal) toxicity is low. NP carcinogenicity data give some reason for concern, though more data are needed (ICIS 2007). NP is highly toxic to fish, aquatic invertebrates, and aquatic plants (OEHHA 2009).

Bakke (2003) proposed a RfD value of 0.10 $\text{mg}/\text{kg}/\text{day}$, which should be protective for human health under chronic exposure to NP and NPEs, and the Danish Institute of Safety

and Toxicology (DIST) derived a preliminary tolerable daily intake (TDI) value for NP of 5 $\mu\text{g}/\text{day}/\text{kg}$ body weight (Danish EPA 2000).

Environmental fate and transport

NP is a viscous liquid at room temperature (Kawahata et al. 2004; USEPA 2010b) and is primarily released into the environment through the discharge of municipal and industrial wastewater into surface waters (Writer et al. 2012). Pathways to the terrestrial environment include the spraying of pesticides containing NP or NPEs as formulates, landfilling of sludge, or the application of sewage sludge or pulp and paper mill sludge to agricultural soils (Soares et al. 2008; Brown et al. 2009). However, Brown et al. (2009) suggested that in practical field situations, where typical biosolids (NP <1000 mg/kg) are used at agronomic rates (<2.0 kg/m^2), NP does not accumulate and plant uptake or water quality impairment is minimal.

NP can also be the products of biodegradation of alkylphenol polyethoxylates. These compounds, in fact, biodegrade by stepwise loss of ethoxy groups, resulting in the formation at the end of various hydrophobic alkylphenols (Ying and Kookana 2003). Thus, NP and NPEs are constituents of untreated and insufficiently treated wastewater and are also metabolites of widely used alkylphenolic compounds. It has been shown that the formation of NP is favored under anaerobic conditions, but it was also reported under aerobic conditions (Micic and Hofmann 2009). The formation of NP from its precursors has been observed in anaerobic sludge treatment (Ahel et al. 1994), surface reservoir sediments (Micić et al. 2013), and estuarine sediments (Lee Ferguson et al. 2003).

Due to its physical-chemical properties (Table 2), such as low water solubility and high $\log(K_{OC})$ values, sorption plays an important role on NP fate and transport in soil-water systems and river sediments (Bennie et al. 1997; Sekela et al. 1999; Azevedo et al. 2001; Bester et al. 2001; Heemken et al. 2001; Fries and Puttmann 2003; Jonkers et al. 2003; Rice et al. 2003; Basheer et al. 2004; Kawahata et al. 2004; Vitali et al. 2004; Cespedes et al. 2005; Vethaak et al. 2005; Patrolecco et al. 2006; Fu et al. 2007; Loos et al. 2007; Pojana et al. 2007; Wu et al. 2007; Micic and Hofmann 2009; Chen et al. 2013), where it is moderately persistent (OEHHA 2009; Li et al. 2013a). Liao et al. (2014) and Roberts et al. (2014) reported that the amount of soil organic matter dominated the sorption capacity of 4-NP and NP to different soils though a clear linear relationship was not evident. Sorption process reached equilibrium in 6 h, with a first rapid sorption stage (30 min ahead) followed by a slow sorption stage (30 min afterward).

Similar results were also reported for sorption in aquatic sediments in Ding et al. (2014), even slowing down of the

process was observed in the presence of biofilm. Shchegolikhina et al. (2012), accordingly, observed also NP extractability from soil, with water and other agents, decreasing at increasing contact time with soil. In marine sediments, NP sorption is enhanced under high salinity (Yang et al. 2011).

NP is not likely to volatilize from soil and is rapidly degraded by hydroxyl radicals in the atmosphere (USEPA 2010b). NP is moderately bioaccumulative (OEHHA 2009).

NP undergoes photolysis in water. In Martínez-Zapata et al. (2013), it was degraded in ultrapure water due to direct photolysis under solar irradiation (300–800 nm) according to a first-order kinetic. Fe(III) and humic acids had a significant synergistic effect. Li et al. (2013b) investigated NP photolysis by sunlight. In pure water, the pseudo-first-order rate constant decreased from 6.73×10^{-3} to 1.57×10^{-3} l/min as the NP initial concentration increased from 40 mg/m³ to 5.0 g/m³; in seawater, the removal rate was slightly slower, the difference being ascribed to the presence of competing species. Similar results were observed by Neamtu and Frimmel (2006).

NP undergoes aerobic biodegradation in water, sediment, and soil systems, but high concentrations can be toxic to microorganisms (EC 2002). Mineralization has been observed in a variety of soil types, including agricultural soils of various textures, noncultivated temperate soils, and soils from the Canadian tundra (Topp and Starratt 2000 in CCME 2002). Gabriel et al. (2008) investigated the degradation of technical NP (a mixture of more than 100 isomers) by *Sphingobium xenophagum* Bayram; the strain degraded NP isomers differentially, being those with less bulkiness at the α -carbon and with four to six carbon atoms mainly alkyl chain being degraded more efficiently. Lu and Gan (2014) compared biodegradation kinetics of a large suite of NP isomers in river sediments under both oxic or anoxic conditions, reporting half-lives of NP isomers sediment ranging from 0.9 to 13.2 days under oxic conditions and from 15.1 to 20.1 days under slightly reduced conditions. Under reduced conditions, the persistence of NP isomers generally increased with estimated first-order half-lives of NP isomers greater than 200 days, with negligible dissipation under strongly reduced conditions.

Chang et al. (2004) observed anaerobic degradation of NP in a sediment-water system by sulfate-reducing bacteria, methanogens, and eubacteria. Fungi can degrade NP exclusively under aerobic conditions (Corvini et al. 2006a). Rozalska et al. 2010 tested filamentous fungi *Gliocephalotrichum simplex* to degrade 4-n-NP (50 g/m³), which was removed by 88 % after 24 h of incubation and almost completely after 48 h. In the same study, 4-n-NP at 100 g/m³ was also removed, but at a slower rate.

Degradation byproducts

Li et al. (2013b) observed the formation of 4-nonyl-catechol after natural irradiation of NP in water. The authors also

detected n-nonoic acid in irradiated pure water, but not in seawater.

Corvini et al. (2006b) studied the degradation pathways of NP by *Sphingomonas* sp. TTNP3, a microbial strain that exhibited high degradation capabilities toward NP used as sole carbon and energy source. The major metabolite in the degradation pathway was hydroquinone, which was further degraded to organic acids (succinate and 3,4-dihydroxy butanedioic acid); benzenediol and alkyloxy derivatives were the dead-end products. Rozalska et al. (2010) investigated the metabolic degradation pathway of 4-n-NP by the nonligninolytic filamentous fungi *G. simplex*, resulting in two possible routes. In one route, carbon detachment brought to 4-hydroxyphenylheptanoic acid and then to 3-(4-hydroxyphenyl)propanoic acid, which was transformed into 4-(1-hydroxyvinyl)phenol, on a side route, and to 2-(4-hydroxyphenyl)acetic acid and 4-hydroxybenzoic acid, on the main route. In the second route, hydroxylation at the ninth position (close to the aromatic ring) and carboxylation at the first position (distal carbon) of the nonyl-moiety brought to 9-hydroxy-9-(4-hydroxyphenyl)nonanoic acid. In 6 h of incubation, 4-hydroxybenzoic acid was the major metabolite. After 72 h of incubation, no toxic effects were observed.

Environmental evidence

Table 4 summarizes the major studies on NP concentrations in different environmental matrices and food.

NP in soil and sediments

Few studies are available in the literature on NP occurrence in soil. All the authors focused on agricultural soils to highlight the effects of sludge amendment and irrigation with untreated wastewater. Vikelsøe et al. (2002) studied the distribution of NP in dressed and fertilized agricultural soils in Denmark. NP concentrations between 0.01 and 0.98 $\mu\text{g}/\text{kg}$ d.w. were found in unamended soils, soils fertilized with manure, or with limited amounts of sewage sludge and 34 $\mu\text{g}/\text{kg}$ at runoff points. Higher concentrations of NP (1.45–2.43 mg/kg), persisting up to 8 years after amendment had ceased, were instead measured in soils exposed to a high addition of sewage sludge. Similar NP values were also reported for sludge-amended site in Canada (CCME 2002). In Mexican agriculture fields irrigated with untreated wastewater for 10 to 90 years, Gibson et al. (2010) measured NP ranging between <25 and 299 $\mu\text{g}/\text{kg}$ d.w., indicating only little evidence of NP accumulation in soils and poor evidence of migration through the different horizons in soil.

Analytical data of sediment samples point out a variability up to 4 orders of magnitude: the reported concentrations of NP range from 3.6 $\mu\text{g}/\text{kg}$ d.w. to 72 mg/kg d.w., with the highest values registered in lakes (Bennett and Metcalfe 1997; Wu

Table 4 NP concentrations in various environmental matrices and in food (percentages between brackets represent the detection frequency)

Reference	Location	Units	Value
Soils			
CCME (2002)	Soil amended with sludge, Canada	µg/kg d.w.	2720
Vikelsøe et al. (2002)	Unamended, manured or artificially fertilized soils and soils amended with limited amounts of sewage sludge, Denmark	µg/kg d.w.	0.01–0.98 Mean 0.37
	Soil amended with high amounts of sewage sludge, Denmark	µg/kg d.w.	1450–2430 Mean 1940
Gibson et al. (2010)	Agricultural fields irrigated with wastewater, Tula Valley, Mexico	µg/kg d.w.	<25–299
Sediments			
Bennett and Metcalfe (1997)	Great Lakes, USA and Canada	µg/kg d.w.	<46–37,800 Mean 3000
Bennie et al. (1997)	Great Lakes and St. Lawrence River in 1995, USA and Canada	µg/kg d.w.	170–72,000 Mean 10,600
Yamashita et al. (2000)	Tokyo Bay, Japan	µg/kg d.w.	<10–5540
Bester et al. (2001)	Bight in the North Sea, Germany	µg/kg d.w.	10–153 Mean 55
	Open sea, North Sea, Germany	µg/kg d.w.	<10–55 Mean 34 (40 %)
Heemken et al. (2001)	Elbe River and some of its tributaries, Germany	µg/kg d.w.	367–1378 Mean 640
Jonkers et al. (2003)	Western Scheldt and Rhine estuaries, Holland	µg/kg d.w.	<0.4–1080 Mean 19.5 (94 %)
Kannan et al. (2003)	Kalamazoo River, USA	µg/kg d.w.	<5.5–15.3
Rice et al. (2003)	Cuyahoga River, Ohio (USA)	µg/kg d.w.	75–340 Mean 180
Kawahata et al. (2004)	Estuarine and marine sediments from Okinawa, and Ishigaki Islands, Japan	µg/kg d.w.	<5–46 Mean 30.5 (47 %)
Vitali et al. (2004)	Rieti District, Italy	µg/kg d.w.	44–567 Mean 205
Vethaak et al. (2005)	Fresh, marine, and estuarine sediments, The Netherlands	µg/kg d.w.	<10–3800 Median 160 (91 %)
Lara-Martin et al. (2006)	Marine and estuarine sediments from Bay of Cadiz, Spain	µg/kg d.w.	13–225 Mean 108
Patrolecco et al. (2006)	Tiber River, Italy	µg/kg d.w.	50–970 Mean 414
Fu et al. (2007)	Estuarine and marine sediments from Jiaozhou Bay and surrounding rivers, China	µg/kg d.w.	3.6–39,700 Mean 3670
Pojana et al. (2007)	Sediments in Venice Lagoon, Italy	µg/kg d.w.	47–192 Mean 89
Wu et al. (2007)	Urban lakes in Wuhan City, China	µg/kg d.w.	3540–32,430 Mean 10,490
Micic and Hofmann (2009)	Danube River, Germany	µg/kg d.w.	<20–2830 Mean 130
Gong et al. (2011)	Major tributaries in Pearl River system, China	µg/kg d.w.	31–21,885 Mean 3686
Klosterhaus et al. (2013)	San Francisco Bay, USA	µg/kg d.w.	21.5–86.3 Mean 34.7
Micić et al. (2013)	Iron Gate I Reservoir on the Danube River, Romania	µg/kg d.w.	80–470
Wu et al. (2013)	Huangpu River and its tributaries, China	µg/kg d.w.	10.34–337.73
Koniecko et al. (2014)	Surface sediments of the Gulf of Gdansk, Poland—rivers	µg/kg d.w.	<0.08–4.93
	Surface sediments of the Gulf of Gdansk, Poland—coastal stations	µg/kg d.w.	<0.08–13.56
	Surface sediments of the Gulf of Gdansk, Poland—stations below 4 m depth	µg/kg d.w.	<0.08–249.08
Duan et al. (2014)	Surface sediments of the Yellow Sea and East China Sea, China	µg/kg d.w.	349.5–1642.8

Table 4 (continued)

Reference	Location	Units	Value
			Mean 890.1
Gorga et al. (2014)	Ebro River basin, Spain	μg/kg d.w.	36–538 Mean 177
Stewart et al. (2014)	Estuarine sediments in Auckland, New Zealand	μg/kg d.w.	<100–32,000 Median 153 4-n-NP <100
Gorga et al. (2015)	Sediments from Ebro, Llobregat, Júcar and Guadalquivir rivers, Spain	μg/kg d.w.	<0.24–1693
Groundwater			
Lacorte et al. (2002)	Agricultural area in Catalonia, Spain	mg/m ³	<0.01–0.35
Latorre et al. (2003)	Agricultural areas in northern Spain	mg/m ³	<0.036–0.9 (92 %)
Félix-Cañedo et al. (2013)	Groundwater in Mexico City, Mexico	mg/m ³	<0.001–0.047 (43 %)
Loos et al. (2010), Luo et al. (2014)	23 European countries, Europe	mg/m ³	<0.030–3.85 Mean 0.083 (11 %) 90th percentile 0.039
Surface Water			
Bennie et al. (1997)	Great Lakes and St. Lawrence River in 1995, USA and Canada	mg/m ³	<0.01–0.92 Mean 0.21 (24 %)
Sekela et al. (1999)	Upstream of a WWTP, Fraser River, Canada	mg/m ³	0.0066–0.0074
	Downstream of a WWTP, Fraser River, Canada	mg/m ³	0.032–0.13
Azevedo et al. (2001)	River and coastal waters, Portugal	mg/m ³	<0.01–30 Mean 1.2 (79 %)
Bester et al. (2001)	Bight of the North Sea, Germany	mg/m ³	0.0007–0.033
Heemken et al. (2001)	Elbe River and its tributaries, Germany	mg/m ³	0.0008–0.221 Mean 0.059
	North Sea	mg/m ³	0.0003–0.084 Mean
Fries and Puttmann (2003)	Rhine, Elbe, Main, Oder, Nidda, and Schwarzbach rivers, Germany	mg/m ³	<0.025–1.22 Mean 0.43
Jonkers et al. (2003)	Western Scheldt and Rhine estuaries, Holland	mg/m ³	0.031–0.934 Mean 0.17
Kannan et al. (2003)	Kalamazoo River, USA	mg/m ³	<2.6
Rice et al. (2003)	Cuyahoga River, Ohio (USA)	mg/m ³	0.1–0.5 Mean 0.24
Basheer et al. (2004)	Surface coastal water, Singapore	mg/m ³	0.02–2.76 Mean 0.95
Kawahata et al. (2004)	Estuarine and marine waters from Okinawa, and Ishigaki Islands, Japan	mg/m ³	<0.05–0.17 Mean 0.14 (29 %)
Vitali et al. (2004)	Rieti District, Italy	mg/m ³	<0.1–1.6
Cespedes et al. (2005)	Llobregat River basin, Catalonia, Spain	mg/m ³	<0.15–37.3 Mean 5.7 (90 %)
Vethaak et al. (2005)	The Netherlands	mg/m ³	<0.11–4.1 Median 0.99 (10 %)
Patrolecco et al. (2006)	Tiber River, Italy	mg/m ³	0.13–0.58 Mean 0.28
Vousta et al. (2006)	Glatt river, Switzerland	mg/m ³	0.068–0.326
Fu et al. (2007)	Marine water from the Jiaozhou Bay, China	mg/m ³	0.02–0.269
	Jiaozhou Bay inflowing rivers, China	mg/m ³	0.0906–28.6
Loos et al. (2007)	River water, Belgium	mg/m ³	0.32–2.50 Mean 1.44
Loos et al. (2007)	River water, Italy	mg/m ³	0.46–0.70 Mean 0.56
Pojana et al. (2007)	Venice Lagoon, Italy	mg/m ³	<0.0005–0.21

Table 4 (continued)

Reference	Location	Units	Value
Wu et al. (2007)	Urban lakes in Wuhan City, China	mg/m ³	1.94–32.85 Mean 11.96
Micic and Hofmann (2009)	Danube River, Germany	mg/m ³	<0.1–0.13 (18 %)
Félix-Cañedo et al. (2013)	Surface water (dams) in Mexico City, Mexico	mg/m ³	<0.001–0.655 (75 %)
Klosterhaus et al. (2013)	San Francisco Bay, USA	mg/m ³	<0.00252–0.0729 (60 %)
Wu et al. (2013)	Huangpu River and its tributaries, China—July 2010	mg/m ³	0.0202–0.1075 Mean 0.074
	Huangpu River and its tributaries, China—November 2010	mg/m ³	0.0926–0.3317 Mean 0.1606
Esteban et al. (2014)	Manzanares and Jarama rivers, Spain	mg/m ³	0.096–1.483
Luo et al. (2014)	China	mg/m ³	0.036–33.231
	Greece	mg/m ³	0.558–2.704
	Korea	mg/m ³	0.115–0.336
Xu et al. (2014)	Seawater Cape D’Aguilar Marine Reserve, Hong Kong—wet season	mg/m ³	0.14–0.50 Mean 0.39
	Seawater Cape D’Aguilar Marine Reserve, Hong Kong—dry season	mg/m ³	0.061–0.33 Mean 0.11
Zhang et al. (2014)	North Tai Lake Basin, Eastern China	mg/m ³	0.089–1.189 Mean 0.388
Gorga et al. (2015)	Iber Ebro, Llobregat, Júcar, and Guadalquivir rivers (Ebro, Llobregat, Júcar, and Guadalquivir)	mg/m ³	<0.00013–0.391
Food and biota			
Guenther et al. (2002)	Packed foodstuff from supermarkets, Germany	µg/kg f.w.	0.1–19.4 Mean 6.0
Rice et al. (2003)	Carp from Cuyahoga River, Ohio (USA)	µg/kg f.w.	6.6–110 Mean 53.4
Basheer et al. (2004)	Seafood from supermarkets, Singapore	µg/kg f.w.	46.6–197 Mean 87.7
Loyo-Rosales et al. (2004)	Spring water bottled in HDPE and PVC from supermarkets, USA	mg/m ³	0.015–0.300 Mean 0.104
Ferrara et al. (2005)	Edible marine species from Adriatic Sea, Italy	µg/kg f.w.	2.7–1286 Mean 413
Vethaak et al. (2005)	Edible freshwater specie (bream), The Netherlands	µg/kg f.w.	<10–160 Median 135 (24 %)
Vethaak et al. (2005)	Edible marine specie (flounder) from North Sea Canal, The Netherlands	µg/kg f.w.	<10–10 Median 10 (10 %)
Isobe et al. (2007)	Green mussel from India, Indonesia, Singapore, Malaysia, Thailand, Cambodia, Vietnam, and the Philippines during 1994–1999	µg/kg d.w.	18–663 (79 %)
	Tokyo Bay during 1994–1999	µg/kg d.w.	47–1347
Shao et al. (2007)	Meat/seafood from supermarkets in Beijing, China	µg/kg f.w.	<0.05–55.98 Mean 6.87
Ferrara et al. (2008)	Edible marine species from Tyrrhenian Sea, Italy	µg/kg f.w.	5–1220 Mean 147
Cacho et al. (2012)	Plastic packed vegetables from local supermarkets, Spain	µg/kg f.w.	<14.5–48 (14 %)
Diehl et al. (2012)	Marine organisms California estuary, Morro Bay, USA	µg/kg f.w.	122–2380
Gyllenhammar et al. (2012)	Fruits, cereal products, and vegetables commercially available, Sweden	µg/kg f.w.	<10–71
Dodgen et al. (2013)	Lettuce and collards, stem and leaves	µg/kg f.w.	1.18–6.95
Dodgen et al. (2013)	Lettuce and collards, roots	µg/kg f.w.	339.2–926.9
Li et al. (2013c)	Soft commercial drinks	mg/m ³	<0.03–0.22 (25 %)
Lu et al. (2013)	Vegetables and fruits, from local commercial sources, Florida (USA)	µg/kg f.w.	<0.3–11.0 4-n-NP <0.1–18.5
Maggioni et al. 2013	PET - bottled water	mg/m ³	<0.0077

Table 4 (continued)

Reference	Location	Units	Value
Maggioni et al. (2013)	Drinking water from public drinking fountains, Italy	mg/m ³	<0.0077–0.084 (23 %)
Dodder et al. (2014)	Mussels along the California coast, USA	µg/kg d.w.	96–3000 Mean 470 Median 200

et al. 2007). In sediments, high NP concentrations were generally associated to specific point sources such as industrial plants, or large amount of domestic wastewater entering the river as it flows through urban areas, especially in most populous regions (Fu et al. 2007; Wu et al. 2007; Duan et al. 2014; Stewart et al. 2014). Typically increasing from upstream to downstream was also reported, with higher values often registered in estuarine and coastal sediments (Fu et al. 2007; Gong et al. 2011; Gorga et al. 2015). The presence of NP in freshwater sediments was primarily ascribed to domestic and industrial wastes and to a lesser extent by agricultural activities (pesticide applications, sludge amendment, and irrigation with wastewater). Koniacko et al. (2014) recognized the rivers and surface runoff as the main sources of NP in coastal sediments of the Gulf of Gdansk; however, they also indicated the possibility of atmospheric transportation of black carbon originating from combustion processes on land along with adsorbed alkylphenols. In Central Europe, Micic and Hofmann (2009) detected concentrations of NP between <0.02 and 2.83 mg/kg d.w. in sediment samples collected along the Danube River; important point sources were industrial sites, especially oil refineries, and drains of untreated wastewater. Similar ranges were also measured along the Elbe River and some of its tributaries in Germany (Heemken et al. 2001) and for Rhine and the Western Scheldt estuaries in heavily industrialized areas or in areas receiving both treated and untreated domestic wastewater in the Netherlands (Jonkers et al. 2003). In Italy, concentrations of NP were measured in areas with different soil uses (urban, industrial, agricultural, open country) in the Rieti District, along the Tiber River and in the Venice Lagoon with similar values, ranging between 44 and 970 µg/kg d.w. (Vitali et al. 2004; Patrolecco et al. 2006; Pojana et al. 2007).

In marine sediments, a decrease in NP concentrations with respect to the levels measured in estuarine sediments and also with increasing distances from the coast was often observed (Bester et al. 2001; Vethaak et al. 2005). As an example, Fu et al. (2007) reported NP concentrations between 3.6 and 299 µg/kg d.w. in the sediments of Jiaozhou Bay (China) and between 23.8 and 39700 µg/kg d.w. in sediments of inflowing rivers.

In open-sea sediments, some offshore oil/gas drilling platforms were identified as a likely source of NPEs and NP, even if

it is not clear if these are due to drilling and production activities or discharges from ships (e.g., cleaning activities) in the areas (Vethaak et al. 2005). Similar to non- and slightly polar organics, a positive correlation of NP with organic carbon content of the sediments was reported, confirming a role of organic carbon for sorption (Jonkers et al. 2003; Gorga et al. 2015). For NP concentrations in sediment, a clear seasonal pattern was not reported as samples may be deposited during a long time.

NP in groundwater and surface water

NP was measured in groundwater by Lacorte et al. (2002) and Latorre et al. (2003) in two agricultural areas of Spain, with values between below 0.01 (detection limit) and 0.9 mg/m³. More recent monitoring of groundwater resulted in measured NP concentration ranging between the limit of detection, 0.001 mg/m³, and 3.85 mg/m³ (Loos et al. 2010; Félix-Cañedo et al. 2013; Luo et al. 2014). NP contamination of groundwater was mainly associated to landfill leachate, water from agricultural land, or seepage of septic tanks and sewer systems (Luo et al. 2014).

The measured concentrations of NP in surface waters span between 3×10^{-4} and 37.3 mg/m³. Considering NP is highly hydrophobic and liable to adsorb to suspended solids and eventually to accumulate in sediments, authors comparing concentrations of sediments and surface water samples from the same location often observed higher concentration in surface sediments than in surface water (Wu et al. 2007, 2013; Micić et al. 2013). Vitali et al. (2004) also pointed out that recorded dissolved NP high levels were limited to a short distance (a few kilometers) downstream from the source of contamination, as due to chemical-physical characteristics and their adsorption on particulate matter, sediments represent the final sink for NP. As a consequence, high NP in sediments and low concentrations in water was often ascribed to past emissions (Jonkers et al. 2003).

Nevertheless, the spatial distribution of NP in surface water was quite the same of the NP concentrations in sediments, with the highest levels closely related to the input of industrial or domestic wastewater discharges or wastewater treatment plant effluents (Bennie et al. 1997; Sekela et al. 1999; Kannan et al. 2003; Cespedes et al. 2005) and a general increase from

the upper rural portions of the river to the urbanized and industrialized segments (Esteban et al. 2014; Luo et al. 2014). Azevedo et al. (2001) in Portugal registered especially high (up to 30 mg/m^3) NP concentrations in industrial districts where tannery and textile industries are located. Loos et al. (2007), in Belgium (south Ghent) and in Italy (south Como), analyzed wastewater treatment plant (WWTP) effluents of textile industries and the receiving rivers and found NP concentrations in the receiving waters upstream the effluent discharge ($0.32\text{--}2.50 \text{ mg/m}^3$) comparable to NP levels in WWTP effluents ($0.37\text{--}0.73 \text{ mg/m}^3$). Despite agriculture is recognized as a minor source of NP in freshwater, Patrolecco et al. (2006) measured NP above 0.30 mg/m^3 in water samples collected from the Tiber River (Italy) at heavily exploited rural areas.

In freshwaters in Central Europe, NP up to about 1.3 mg/m^3 has been reported (Jonkers et al. 2003; Fries and Puttmann 2003; Micic and Hofmann 2009; Vethaak et al. 2005). By comparing the results of past water monitoring with more recent data, it was also highlighted a decrease in dissolved NP concentrations in European rivers in the last two decades, probably as consequence of the various voluntary restrictions or legislation on the use of NP in household cleaning products and industrial applications together with possible relocation of industrial activities (Fries and Puttmann 2003; Vousta et al. 2006; Micic and Hofmann 2009; Gorga et al. 2014).

In coastal areas, a significant decreasing trend in NP concentrations with the distance from the coast is often reported (Basheer et al. 2004; Heemken et al. 2001; Fu et al. 2007). Significant levels of NP have been occasionally reported in offshore sampling points in the vicinity of industrial areas, jetties, shipyards, marinas and recreational beaches, as well as in shipping lanes, anchorages, and near petroleum refineries (Basheer et al. 2004).

Fu et al. (2007) and Xu et al. (2008) reported of a seasonal trend of dissolved NP concentrations with higher values in summer than in winter. Such finding was ascribed firstly to high temperatures and associated microbial activity, leading to an enhanced degradation of NPEs in marine sediments and hence an increased NP concentrations in water column during summer. In the Hong Kong area, as the summer is also the wet season, it was also assumed a possible effect on NP level in water due to increased surface runoff during rain events (Xu et al. 2008). Kueh and Lam (2008) in Hong Kong measured in storm water NP concentrations between 0.08 and 12 mg/m^3 .

NP in biota, food, and bottle water

Significant NP concentrations were found in different foods, with values between 0.1 and $100 \text{ }\mu\text{g/kg f.w.}$ and $<7.7 \text{ }\mu\text{g/m}^3$ and 0.30 mg/m^3 in drinking waters/commercial drinks from different countries. In Guenther et al. (2002), NP concentrations in packed foodstuff purchased from supermarkets in Germany spanned between 0.1 and $19.4 \text{ }\mu\text{g/kg f.w.}$ The

concentration was not related to the food fat content and NP migration into food occurred at different stages of the food production.

High NP concentrations in seafood and various edible marine species were observed in Asia (Basheer et al. 2004; Isobe et al. 2007; Shao et al. 2007), Europe (Ferrara et al. 2005, 2008), and North America (Dodder et al. 2014) at comparable levels, considering the differences in species examined in each study, number of collected samples, period, analytical methods and reporting units, and also different periods.

Measured values of NP in different commercially available vegetables and fruits in Sweden (Gyllenhammar et al. 2012), Spain (Cacho et al. 2012), and Florida (Lu et al. 2013) varied roughly between 5 and $50 \text{ }\mu\text{g/kg f.w.}$

Different accumulation of NP in each species was observed. Significant values were found in carrots and pumpkins (10.4 and $11.3 \text{ }\mu\text{g/kg f.w.}$, respectively) and in apples and citruses (17.1 and $29.5 \text{ }\mu\text{g/kg f.w.}$, respectively), whereas NP was not detected in strawberries, lettuce, potato, and tomatoes. Dodgen et al. (2013) investigated the accumulation of NP in different parts of lettuce and collards, pointing out the poor translocation of NP from the roots to the upper tissues after uptake: in the stem and leaves, concentration ranges $1.18\text{--}4.31$ and $3.79\text{--}6.95 \text{ }\mu\text{g/kg f.w.}$ were observed for lettuce and collards, respectively, but the values measured in the roots were greater by 3 orders of magnitude (926.9 and $339.2 \text{ }\mu\text{g/kg f.w.}$ for lettuce and collards, respectively).

Loyo-Rosales et al. (2004) investigated the presence of NP in commercial water bottled in different materials (high-density polyethylene (HDPE), polyethylene terephthalate (PET), polyvinyl chloride (PVC)). NP was found in water contained in HDPE and PVC bottles at values of $29\text{--}180$ and $15\text{--}300 \text{ }\mu\text{g/m}^3$, respectively. Maggioni et al. (2013) evaluated concentrations of NP in drinking waters from public fountains in 35 Italian cities and in bottled mineral waters; in all samples, the amount varied from below the quantification limit ($7.7 \text{ }\mu\text{g/m}^3$) up to a maximum of $84 \text{ }\mu\text{g/m}^3$. These values are similar to the reported NP range in commercial soft drinks (Li et al. 2013c).

Based on the concentrations measured in food and the expected consumption rates, the average daily intake of NP varies between 0.067 and $0.370 \text{ }\mu\text{g/kg/day}$ for adults (60 kg body weight) (Guenther et al. 2002; Ferrara et al. 2005, 2008; Shao et al. 2007; Gyllenhammar et al. 2012; Lu et al. 2013). Loyo-Rosales et al. (2004) calculated an average NP daily intake from drinking bottled water of $0.36\text{--}0.60 \text{ }\mu\text{g/day}$. Diet seems the major exposure route for humans.

Benzophenones (BPs)

Benzophenone (diphenyl ketone, BP, $(\text{C}_6\text{H}_5)_2\text{CO}$) is composed of two aromatic rings and a carbonyl group (Table 1)

(NTP 2006). Based on this parental structure, a group of different compounds can be generated through substitutions of hydrogen atoms in the aromatic rings. The physical-chemical properties and the environmental behavior of these derived compounds are not significantly different from those of the parent compound, except for benzophenone-3 (2-hydroxy-4-methoxybenzophenone, BP-3, Table 1), which is also the most commercialized compound of the group (León et al. 2010; Gago-Ferrero et al. 2012; Liu et al. 2012a; Zhang et al. 2013b).

BPs are used as a flavor ingredient, a fragrance enhancer, a perfume fixative, and an additive for plastics, coatings, and adhesive formulations. They are also used in laundry and household cleaning products and in the manufacture of insecticides, agricultural chemicals, hypnotic drugs, antihistamines, and other pharmaceuticals. BPs are used as an ultraviolet (UV)-curing agent in sunglasses and to prevent UV light from damaging scents and colors in products such as perfumes and soaps. BP-3 is commonly used worldwide as a UV filter in cosmetic formulations, such as sunscreens and skin care products, body lotions, hair sprays, hair dyes, and shampoos (Zhang et al. 2011; Liu et al. 2012a). Moreover, they can be added to plastic packaging as a UV blocker, which allows manufacturers to package their products in clear glass or plastic rather than opaque or dark packaging. BPs are widely used as a photoinitiator for inks and varnishes that are cured with UV light (Ricking et al. 2003; Zhang et al. 2011). In 2003, BP production exceeded 453 t in the USA and 10,000 t in the European Union (NTP 2006).

Toxicity

BPs have adverse effects on reproduction and hormonal functions of fish (IARC 2010). They can alter endocrine signaling through multiple effects on receptors. The estrogenic activity of BP-3 and BP-1 (2,4-dihydroxybenzophenone) was determined by the estrogenic recombinant yeast assay, resulting in the half-maximal effective concentrations (EC_{50}) of 12.5 and 0.058 g/m^3 , respectively, and a lowest observed effect concentrations (LOEC) of 1.6 and 0.015 g/m^3 , respectively (Gago-Ferrero et al. 2012). No data are available on BP carcinogenicity to humans, though they are classified as group 2B substances, “possible carcinogenic to humans” (NTP 2006; IARC 2010).

Because of their use as an additive in fragrances, cosmetics, pharmaceuticals, insecticides, and household cleaning products, exposure to BPs through dermal contact may be significant. In León et al. (2010), BP-3 applied on the skin was absorbed and readily biotransformed into BP-1, 2,2'-dihydroxy-4-methoxybenzophenone (BP-8), and 2,3,4-trihydroxybenzophenone (THB), whose decrease over time was much slower than the parent compound. Dietary sources of exposure include food and drinking water, where BPs

might be present due to the addition as a flavoring or the migration from packaging (IARC 2010). The European Commission Scientific Committee on Food set a RfD for oral exposure of 10 $\mu g/kg/day$ (EC 2005).

Environmental fate and transport

BPs can enter the environment through solid-waste landfill leachate and wastewater treatment plants effluents (Ricking et al. 2003; Jeon et al. 2006). BP is insoluble in water (Table 2). Because of the high K_{OC} value, BP sorption on soil and sediment organic matter is significant (USEPA 1984). Volatilization can occur to some extent, but due to low vapor pressure, BP is not expected at significant levels in ambient air (USEPA 1984). BP in water can be photodegraded under sunlight exposure (Hayashi et al. 2006). Fujii and Kituchi (2005) observed BP biodegradation in activated sludge caused by a specific microbial strain. BP is persistent in the environment and susceptible to bioaccumulation (Brooks et al. 2009; IARC 2010). Based on the estimated K_{OC} value (Table 2), BP-3 has slight mobility in soil and sorbs significantly on suspended solids and sediments. Volatilization from soil surface and water is not expected to be an important process, though BP-3 in ambient air can exist in both vapor and particulate phase. The potential for BP-3 bioconcentration in aquatic organisms is moderate to high (TOXNET 2014).

BP-3 absorbs light at 288 and 326 nm and, therefore, is susceptible to direct photolysis by sunlight. However, Rodil et al. (2009) assessed the photostability of BP-3 under sunlight exposure (290 to 800 nm) of water samples spiked at 100–4000 mg/m^3 , without any significant decrease of BP-3 concentration over a 72 h irradiation period. Also, Gago-Ferrero et al. (2012) did not observe photodegradation of BP-3 over a 24 h irradiation period. Vapor-phase BP-3 is degraded in the atmosphere by reaction with photochemically produced hydroxyl radicals, with an estimated half-life of 1.9 h.

BP-3 can be biodegraded in water and soil systems. Liu et al. (2011a) and Liu et al. (2012a) investigated BP-3 biodegradation under oxic and anoxic conditions in water, resulting in BP-3 complete removal in both kinds of microcosms after 42 days of incubation. Anoxic conditions were more favorable than oxic conditions, with measured half-lives of 4.2 and 10.7 days, respectively. Gago-Ferrero et al. (2012) observed high biodegradation rates for BP-3 by white rot fungi *Trametes versicolor*, down to nondetectable levels in 8 h of incubation.

Degradation byproducts

Hayashi et al. (2006) investigated BP byproducts after exposure of an aqueous solution to UV or sunlight irradiation; two-ring hydroxylated derivatives were observed (3-hydroxybenzophenone and 4-hydroxybenzophenone, 4HB).

According to Liu et al. (2011a) and Liu et al. (2012a), after 42 days of incubation in microcosms, biodegradation of BP-3 under Fe(III)-reducing conditions produced 4-cresol and BP-1, while BP-1 was detected under oxic, nitrate-reducing and sulfate-reducing conditions.

BP-1 was identified by Gago-Ferrero et al. (2012) as a metabolite produced during degradation of BP-3 by *T. versicolor*. BP-1 was then degraded to 4,4'-dihydroxybenzophenone (4DHB) and 4HB. More investigation is necessary to identify transformation products formed in environmental matrices in order to assess the potential environmental risk for BPs compounds (Jurado et al. 2014).

Environmental evidence

The major papers in the literature reporting of BPs in environmental matrices and food are summarized in Table 5. Aquatic ecosystems in highly urbanized areas are the most important pathway for BPs to enter in the environmental matrices. Shortcomings in wastewater treatment were reported with a removal efficiency for BP-3 ranging between 55–96 % (Kim and Choi 2014); recently, Ávila et al. (2014) observed a value equal to 93 %. Oxygen availability promote CEC degradation via aerobic pathways (example for BPA and BP-3); however, the releases of BP products are high due to human and industry activity (Kasprzyk-Hordern et al. 2008). Recently, for BP-3, the most ubiquitous compound, some authors paid attention to the direct diffusion due to the release of sunscreen products in water (Sánchez-Brunete et al. 2011; Kim and Choi 2014). In fact, maximum levels of contamination were observed in swimming pool or, in summer, during bath in recreational areas. In addition to the mentioned compounds (BP, BP-3, BP-1, 4HB, 4DHB, BP-8, THB), in the matrices, benzophenone-4 (BP-4), 2,2-dihydroxy-4,4-dimethoxybenzophenone (BP-6), and benzhydrol (BH) were also found.

BPs in soil and sediments

Concentrations of BP types in soils range between 0.07 and 16.55 µg/kg d.w. Sánchez-Brunete et al. (2011) investigated the concentration of different BPs in soil samples from two agricultural fields and from one industrial site in Spain. Only BP-1 was found at 5.7 µg/kg in the industrial soil and BP-6 at 0.6 µg/kg in the agricultural soil amended with sewage sludge. In Jeon et al. (2006), the parent compound BP among seven UV filters showed both high concentration (approximately 5 µg/kg) and frequency in ground soil; BP-3 was detected in 5 soil samples out of 33, at concentrations between 0.73 and 3.88 µg/kg d.w. Due to the importance of adsorption on solid matrices, more investigations would be required; for soil, only few studies are reported in the literature, so it is difficult illustrate the real presence of BPs.

Concentrations of BPs in sediments vary between 0.1 and 110 µg/kg d.w., containing BP-3 between 0.27 and 4.7 µg/kg d.w. Samples of sediments were collected from aquatic ecosystems highly urbanized and industrialized in Songhua River (China), Saginaw River and Detroit River (USA), Magdalena River (Colombia), and Biobio region (Chile) (Zhang et al. 2011; Barón et al. 2013). Nineteen sampling stations in Chile were located close to some important discharges of chemicals, food processing, and urban discharges and 13 sampling stations in Colombia included natural and urban areas of Barranquilla City that has an important industrial district and a sea-river port. Highest levels of BP-3 were detected in Colombia; the contrast with the levels found in Chile is explained with the difference in solar radiation levels. In fact, especially in Caribbean Colombian beaches, high tourism activity and use of personal care products could justify the higher BP-3 concentrations in water compared to the Chilean case (Barón et al. 2013). Kameda et al. (2011) measured BP concentration in sediments collected from rivers (polluted by industrial and domestic wastewaters) and lakes (used as the background sites) in Saitama Prefecture, Japan. Values between 2.7 and 58 µg/kg d.w. were found at the polluted sites and between 3.8 and 105 µg/kg d.w. at the background sites. Dry and wet deposition from atmosphere and the recreational activities were thought to be the source of BP-3 at the background sites. Jeon et al. (2006) measured concentrations of different UV filters in sediment samples collected from lakes and rivers in Korea affected by municipal and recreational wastewaters. The parent compound BP was the most detected and had the highest concentrations (1.52 to 9.73 µg/kg d.w.) among the investigated BPs; BP-3 was not detected in any sediment sample. Sánchez-Brunete et al. (2011) analyzed sediments collected in rivers and along the Mediterranean coast in Spain at bathing or recreational areas. BP-6 was at values between 1.2 and 6.1 µg/kg d.w. The authors that compared concentrations of sediments and surface water samples from the same location noticed that BPs in water often appeared to be below the quantification limit: most of contaminants were accumulated in sediment (Jeon et al. 2006; Kim and Choi 2014). The contamination of sediments may pose an unacceptable risk to aquatic organisms, which tend to bioaccumulate the molecules, and to humans through the ingestion of contaminated fish (Barón et al. 2013).

BPs in groundwater and surface water

Groundwater can be an important water supply resource; however, only few information, reported in this review, on contamination of BPs are available. Stuart et al. (2011) monitored BP concentration in groundwater in England, resulting in values between 0.03 and 2.8 mg/m³. Jurado et al. (2014) evaluated the presence of BPs in the Barcelona aquifers (water is used for street cleaning and to irrigate city gardens) for the

Table 5 BP concentrations in various environmental matrices and in food (percentages between brackets represent the detection frequency)

Reference	Compound and location	Units	Value
Soils			
Jeon et al. (2006)	BPs, South Korea	µg/kg d.w.	BP 0.82–16.55, mean 4.55 (97 %) BP-3 0.73–3.88, mean 2.65 (15 %) BP-1 <0.5 BP-6 0.5–4.17, mean 1.67 (15 %) BH 0.51–6.95, mean 1.8 (39 %) 4HB 1.06–4.91, Mean 3.01 (9 %) THB <0.5
Sánchez-Brunete et al. (2011)	BPs, industrial, and agricultural areas, Spain	µg/kg d.w.	BP-1 <0.1–5.7 (ind) BP-6 <0.09–0.6 (agr) BP-3 <0.1 BP-8 <0.07 4HB <0.07
Sediments			
Jeon et al. (2006)	BPs, South Korea	µg/kg d.w.	BP 1.52–9.73, mean 4.73 (93 %) BP-3 <0.1 BP-1 <0.1 BP-6 0.5–2.14, mean 0.95 (80 %) BH 0.53, mean 0.53 (7 %) 4HB 18.38, mean 18.38 (7 %) THB <0.1
Pojana et al. (2007)	BP, Venice Lagoon, Italy	µg/kg d.w.	14–110 Mean 39.4
Sánchez-Brunete et al. (2011)	BPs, river and coastal sediments, Spain	µg/kg d.w.	BP-6 <0.15–6.1, mean 1.6 BP-1 <0.21 BP-3 <0.28 BP-8 <0.14 4HB <0.23
Kameda et al. (2011)	BP, rivers and lakes, Saitama Prefecture, Japan	µg/kg d.w.	2.7–105 Mean 34.7
Zhang et al. (2011)	BP-3, Songhua River, China	µg/kg d.w.	BP-3 0.272–0.545, mean 0.380 (100 %) BP-1 <0.14 BP-6 <0.22 4HB <0.22
Zhang et al. (2011)	BPs, Saginaw and Detroit River, USA	µg/kg d.w.	BP-3 0.728–4.66, mean 2.34 (100 %) BP-1 0.259–0.607, mean 0.454 (67 %) BP-6 0.133–0.796, mean 0.424 (67 %) 4HB 0.312–0.951, mean 0.53 (50 %)
Barón et al. (2013)	BP-3, river areas, estuary and coastal bays, Biobio region, Chile	µg/kg d.w.	<0.4–2.96
Barón et al. (2013)	BP-3, Magdalena River, Colombia	µg/kg d.w.	<0.4–5.38
Kim and Choi (2014)	BP-3, rivers, worldwide	µg/kg d.w.	<0.5–27
Groundwater			
Stuart et al. (2011)	BP, England	µg/m ³	<10–2780
Jurado et al. (2014)	BPs, Barcelona urban groundwater	µg/m ³	BP-1 mean 0.9 (16 %), max 19.4 BP-3 mean 2.3 (32 %), max 19.2 BP-4 mean 2.8 (19 %), max 36.6 4HB mean 0.2 (6 %), max 3.5 4DHB mean 0.13 (6 %), max 4.1
Jurado et al. (2014)	BPs, Mallorca street zone	µg/m ³	BP-1 mean 0.78 (43 %), max 3.2 BP-3 mean 7.9 (71 %), max 19.2 BP-4 mean 1.1 (25 %), max 6.4 4HB mean 0.38 (14 %), max 2.6 4DHB mean 0.58 (14 %), max 4.1
Jurado et al. (2014)	BPs, Poble Sec zone	µg/m ³	BP-3 mean 0.66 (25 %), max 3.4 BP-4 mean 1.8 (10 %), max 21.3
Jurado et al. (2014)	BPs, Beson River Delta zone	µg/m ³	BP-1 mean 1.9 (17 %), max 19.4 BP-3 mean 0.64 (17 %), max 4.4

Table 5 (continued)

Reference	Compound and location	Units	Value
			BP-4 mean 3.8 (25 %), max 36.6 4HB mean 0.29 (8 %), max 3.5
Surface water			
Balmer et al. (2005)	BP-3, Swiss Lakes	µg/m ³	<2–35 Mean 16.1
Jeon et al. (2006)	BPs, rivers and lakes, South Korea	µg/m ³	BP <25 BP-3 <25 BP-1 47 (4 %) BP-6 <25 BH <25 4HB 85 (17 %) THB <10
Pojana et al. (2007)	BP, Venice Lagoon, Italy	µg/m ³	<2.6–136 Mean 30
Kasprzyk-Hordern et al. (2008)	BPs, river Taff, UK	µg/m ³	BP-1 <0.3–17 BP-2 <0.5–284 BP-3 <15–44 BP-4 <3–371
Kasprzyk-Hordern et al. (2008)	BPs, river Ely, UK	µg/m ³	BP-1 <0.3–13 BP-2 <0.5–26 BP-3 <15 BP-4 <3–323
Fent et al. 2010	BP-3, river Glatt, Swiss	µg/m ³	56–68
Yoon et al. (2010)	BP, Han River, South Korea	µg/m ³	<50–59 Mean 52 (33 %)
Yoon et al. (2010)	BP, effluent-dominated creeks discharging into Han River, South Korea	µg/m ³	56–130 Mean 102
Kameda et al. (2011)	BP, rivers and lakes, Saitama Prefecture, Japan	µg/m ³	1–68 Mean 32.2
Kameda et al. (2011)	BP-3, rivers and lakes, Saitama Prefecture, Japan	µg/m ³	2–12 Mean 7
Rodil et al. (2012)	BP-4, rivers in Galicia, Spain	µg/m ³	2.5–70 Mean 25
Grabicova et al. (2013)	BP-3, recreational areas (ponds, rivers) in South Bohemia, Czech Republic	µg/m ³	12–550
Grabicova et al. (2013)	BP-4, recreational areas (ponds, rivers) in South Bohemia, Czech Republic	µg/m ³	4.0–390
Kim and Choi (2014)	BP-3, freshwater, worldwide	µg/m ³	<0.3–125
Food			
Balmer et al. (2005)	BP-3, fish from Swiss lakes	µg/kg f.w.	0.49–3.3 Mean 1.17
Gago-Ferrero et al. (2013)	BP-3, Guadalquivir River, Spain	µg/kg d.w.	<10–24.3 Median 20.4

first time; BP-3 was the most detected with frequency of 32 % of the samples (other compounds with frequency less than 16 %). The area under study had presented different levels in terms of concentrations; hot spot concentrations were 19.4 µg/m³ (BP-1), 19.2 µg/m³ (BP-3), and 36.6 µg/m³ (BP-4). Due to large amounts of WWTP effluents, Besòs River Delta was found to be the most polluted in terms of UV filter compounds. Under different redox conditions, BPs could be removed (Jurado et al. 2014); in fact, concentrations in

groundwater were lower than those expected from mixing balance of the recharge sources that contributed in Barcelona aquifers.

Concentrations of BPs in surface waters are reported between 0.3 and 550 µg/m³. Wastewater is the most important source (Zenker et al. 2008; Fent et al. 2010; Rodil et al. 2012; Jeon et al. 2006); levels decreased with higher river water flow. Balmer et al. (2005) detected concentrations of BP-3 above the detection limit (2 µg/m³) in water samples from four

Swiss lakes out of five (10–35 $\mu\text{g}/\text{m}^3$); these lakes were selected because of the near municipal wastewater plants or the recreational activities. Comparable level was detected in river Glatt and lake Greifen catchment in Switzerland (Fent et al. 2010). River Glatt, near the city of Zurich, was heavily impacted by human activities and wastewater effluents of 160,000 inhabitants. The relative amount of treated wastewater in the river water was about 10–20 %. Rodil et al. (2012) measured the concentration of many emerging pollutants (including BP-3 and BP-4) in water samples collected in the metropolitan area of La Coruna (Spain). BP-4 was detected in 75 % of the surface water samples at concentrations between 2.5 and 70 $\mu\text{g}/\text{m}^3$. BP-4 was also detected in several drinking water samples, up to a maximum level of 62 $\mu\text{g}/\text{m}^3$. BP-3 was not found in samples of surface or drinking water. Yoon et al. (2010) investigated the occurrence of wastewater-derived contaminants including BP in water samples collected along the Han River, near the northwest Seoul metropolitan area. BP was detected in six sampling sites out of ten, at concentration between 50 and 130 $\mu\text{g}/\text{m}^3$. Kameda et al. (2011) measured the concentrations of BP and BP-3 in water samples at polluted and background sites in the Saitama Prefecture (Japan). Values in the range 2–68 $\mu\text{g}/\text{m}^3$ and 1–57 $\mu\text{g}/\text{m}^3$ were found for BP and 4–12 $\mu\text{g}/\text{m}^3$ and 2–10 $\mu\text{g}/\text{m}^3$ for BP-3. Direct inputs (due to removal from the skin during recreational activities) may not be negligible (Kim and Choi 2014); Grabicova et al. (2013) investigated the occurrence of BPs in water samples from ponds and rivers near recreational area in the South Bohemia, comparing them with unpolluted background areas. The concentrations of both BP-3 and BP-4 in the recreational areas (12–550 and 4.0–390 $\mu\text{g}/\text{m}^3$, respectively) were higher than those in the background areas (14–20 and 3.4–37 $\mu\text{g}/\text{m}^3$, respectively).

BPs in biota and food

Few cases are available in the literature on BPs in food or aquatic biota. Balmer et al. (2005) detected BP-3 in fish from three Swiss lakes out of five; in particular, roach fish had values between 0.66 and 3.31 $\mu\text{g}/\text{kg}$ f.w. Results on fish samples (common carp and Andalusian barbel) collected along the Guadalquivir River basin (Gago-Ferrero et al. 2013) confirmed the bioaccumulation of BP-3. More investigations are necessary; a limit concentration of 0.6 mg/kg in food was set for the sum of BP and 4-methylbenzophenone (IARC 2010).

Benzotriazoles (BTs)

BTs are bicyclic heterocyclic compounds containing three nitrogen atoms and a fused benzene ring. 1H-benzotriazole

(BT, $\text{C}_6\text{H}_5\text{N}_3$, Table 1) is the reference compound of the group because it is the most used and measured in the environment compartments (DECOS 2000; Hart et al. 2004).

BTs are used as additives in fluids to inhibit corrosion of cooling systems and metal machines and in defrosting liquids (DECOS 2000); in particular, the use for defrosting purpose for aircrafts causes significant release on soils (Cancilla et al. 2003; Jia et al. 2006). Some BTs are present in pesticides and herbicides, like UV stabilizers also used in plastic devices to prevent yellowing and degradation of the products (Bhatarai and Gramatica 2011; Wolschke et al. 2011; Zhang et al. 2011). The global production is high, about 9000 t/year in the USA (Liu et al. 2011a).

Toxicity

Although clear antiestrogenic activity of BTs was demonstrated in vitro, no evidence of antiestrogenic activity was observed in vivo assays (Harris et al. 2007). A subchronic, 21 day reproduction toxicity tests using *Daphnia magna* resulted in a no observed effect concentration of 3 g/m^3 ; based on these results, a predicted no effect concentration of 0.06 g/m^3 was calculated (Breedveld et al. 2002). BT was classified as toxic to aquatic organisms and can cause long-term adverse effects in the aquatic environment, but it has low toxicity to humans (La Farré et al. 2008; Breedveld et al. 2002). Limited ecotoxicological data are available, mostly from acute toxicity tests on aquatic species. The EC_{50} values for fish and bacteria are 130 and 41 g/m^3 , respectively (Hem et al. 2003). Based on its molecular weight and partition coefficient, dermal absorption might be expected. Contact dermatitis was observed in metalworkers after skin exposure to BT. Based on acute toxicity data in rats (inhalation LC_{50} 2153 mg/m^3 ; oral LD_{50} 500–965 mg/kg), BT should be classified as harmful for inhalation and oral exposure (DECOS 2000).

No occupational exposure limits/standards for BT have been established or recommended (DECOS 2000). USEPA (2010c) calculated a RfD of 0.03 $\text{mg}/\text{kg}/\text{day}$ for both dermal contact and inhalation. As for carcinogenicity, based on studies in rats and mice, BT was classified as a suspected carcinogen.

Environmental fate and transport

The discharge of treated municipal wastewater is the greatest potential source for BTs in the environment; nevertheless, overruns of wastewater sewers and atmospheric deposition can be regarded as other possible input sources (Kiss and Fries 2009).

BT has high water solubility (Table 2) and is less sorbable on organic matter than the other emerging contaminants considered in this review (Giger et al. 2006). No sorption of BTs was observed in sandy soils, while peat and compost exhibited

a certain affinity with the compounds (Breedveld et al. 2002). BT has a weak hydrophobic nature (Cornell et al. 2000 in Breedveld et al. 2002) and is a weak organic acid with pKa of 8.6 (Andreozzi et al. 1998 in Breedveld et al. 2002).

BTs are slightly sensitive to light and photodegradable (Andreozzi et al. 1998). In Hem et al. (2003), approximately 65 % abatement of BTs was achieved at a dose of 320 mWs/cm². Benitez et al. (2013) suggested a first-order kinetic for the photodegradation process in water, dependent on the pH value.

BTs are generally resistant to biodegradation; therefore, they are highly persistent in the aquatic environment (Hogenboom et al. 2009; Liu et al. 2012b). Many studies reported no evidence of microbial degradation of BT and its derivatives. In Liu et al. (2011a), BT half-lives under different redox conditions were between 114 days (aerobic conditions) and 315 days (sulfate-reducing conditions). Breedveld et al. (2002) observed no degradation during a 5 month test under anaerobic conditions, while BT removal under aerobic conditions was ascribed to evaporation due to aeration rather than biodegradation.

Degradation byproducts

A UV degradation study of BT in aqueous solutions (Hem et al. 2003) evidenced the formation of aniline (1 % w/w of photodegradation byproducts), phenazine (10–20 % w/w), and other unknown byproducts (89–79 % w/w). The amount of byproducts increased with the UV dose. Irradiation of BT leads to two different routes of degradation: nitrogen elimination (with the production of aniline) followed by hydroxylation, and dimerization (phenazine was the main byproduct) (Benitez et al. 2013).

In Liu et al. (2012b), the degradation of four BTs in water samples was studied in aerobic and anaerobic microcosms, by using activated and digested sludge from a wastewater treatment plant. Five degradation byproducts were identified under aerobic conditions (phenol, phthalic acid, 1-methylbenzotriazole, 4-methoxy,1H-benzotriazole, and 5-methoxy,1H-benzotriazole) and four degradation byproducts (phenol, 1-methylbenzotriazole, dimethyl benzylamine, and carbazole) under anaerobic conditions.

Environmental evidence

Table 6 summarizes the major studies on BT concentrations in different environmental matrices. WWTP effluents are important source of BTs and important direct impact is due to deicing activities. BT was detected in water and soil samples from airports, especially during the deicing season; concentrations are greatest just after the spring snowmelt (Loos et al. 2010; Breedveld et al. 2002; Heeb et al. 2012).

BTs in soil and sediments

Few data are available on BT soil and sediment contamination; the ranges of concentration are 3.1–1700 and 0.4–13,000 µg/kg, respectively. McNeill and Cancilla (2009) investigated soils at three different US airports, but BT was detected just in one sample (4.1 µg/kg d.w.). In the old Oslo airport (Fornebu) (Breedveld et al. 2002), BT was detected in 19 out of 20 topsoil samples beside the runway, with an average concentration of 0.33 mg/kg d.w. A soil sample taken at 1.2 m depth had a concentration of 0.51 mg/kg d.w.

Breedveld et al. (2002) observed high concentrations of BT (13,000 µg/kg d.w.) in a sediment sample of a small drainage ditch draining the snow disposal site of the Fornebu Oslo airport. BT was detected at the average concentration of 420 µg/kg d.w. in three surface sediment samples and not detected (<100 µg/kg d.w.) in three other samples collected at a depth of 0.1 m in the upper organic layer of a wetland area receiving the drainage water from the snow disposal site. Regarding sediment from industrialized areas, Zhang et al. (2011) detected BT just in one sampling location (out of six) at a value of 0.39 µg/kg d.w.; on the contrary, BT was detected in all sediment samples from the Saginaw River and the Detroit River (USA) at values between 0.424 and 33.4 µg/kg d.w.

BTs in groundwater and surface water

BT concentrations in groundwater range between <0.001 and 1100 mg/m³, but the values measured by Breedveld et al. (2002) at the Oslo airports were 3 orders of magnitude higher than the maximum value measured by other authors. Excluding this paper, BT in groundwater ranges between <0.001 and 3.242 mg/m³. Breedveld et al. (2002) detected concentrations of BT between 1.2 and 1100 mg/m³ and between 0.11 and 20 mg/m³ in groundwater samples from the old (Fornebu) and new (Gardermoen) Oslo airports, respectively. Groundwater samples were taken from the snow disposal site, the drainage ditch, the deicing pad, and the regeneration plant of aircraft deicing/anti-icing fluids. The highest values were measured in groundwater samples from the sand and gravel deposits that were used to backfill the deicing pad during construction. The areas with the original peat and clay deposits had the lowest BT values.

Loos et al. (2010) performed a survey at European scale; BT was detected in 53 % groundwater samples collected at contaminated sites. The maximum concentration was of 1.032 mg/m³ and the average of 0.024 mg/m³. Kahle et al. (2009) detected BT in four out of six groundwater pumping stations located in a densely populated area of the Canton of Zurich (Switzerland); the detection occurred in the aquifers affected by significant water infiltration from rivers receiving considerable discharge from WWTPs. Reh et al. (2013) measured significant BT concentrations in 67 out of 163

Table 6 BTs concentration in various environmental matrices

Reference	Location	Units	Value
Soils			
Breedveld et al. (2002)	Oslo Airport, Fornebu, Norway	µg/kg d.w.	100–1700 Mean 500
McNeill and Cancilla (2009)	Three USA airports	µg/kg d.w.	<3.1–4.1
Sediments			
Breedveld et al. (2002)	Oslo Airport, Fornebu, Norway	µg/kg d.w.	<100–13,000 Mean 4500
Zhang et al. (2011)	Songhua River, China	µg/kg d.w.	0.385
	Saginaw and Detroit rivers, USA	µg/kg d.w.	0.424–33.4 Mean 9.43
Groundwater			
Breedveld et al. (2002)	Oslo Airport, Fornebu, Norway	mg/m ³	1.2–1100 Mean 371
	Oslo Airport, Gardermoen, Norway	mg/m ³	0.11–20 Mean 4.75
Kahle et al. (2009)	Canton of Zurich, Switzerland	mg/m ³	0.016–0.077 Mean 0.047
Loos et al. (2010)	Europe	mg/m ³	<0.001–1.032 Mean 0.024
Liu et al. (2011b)	Next to a wastewater treatment plant, Adelaide, Australia	mg/m ³	0.280±0.018
Reh et al. 2013	Karstified aquifer, Germany	mg/m ³	0.0049–3.2418 Median 0.0434
Surface Water			
Breedveld et al. (2002)	Oslo Airport, Fornebu, Norway	mg/m ³	1.5–33 Mean 9.0
Weiss and Reemtsma (2005)	Lake Tegel, Berlin region, Germany	mg/m ³	0.9
Giger et al. (2006)	Rivers in Zurich District, Switzerland	mg/m ³	0.06–6.3 Mean 0.94
	Lake in Zurich District, Switzerland	mg/m ³	0.02–1.2 Mean 0.55
Vousta et al. (2006)	Glatt river, Switzerland	mg/m ³	0.636–3.69
Kahle et al. (2009)	Lakes in the Midland region, Switzerland	mg/m ³	0.011–0.917 Mean 0.211
Kiss and Fries (2009)	Main, Hengstbach, and Hegbach rivers, Germany	mg/m ³	0.038–1.47 Mean 0.35
Nodler et al. (2011)	Leine River, upstream of a wastewater treatment plant, Germany	mg/m ³	0.034–0.176 Mean 0.095
	Leine River, downstream of a wastewater treatment plant, Germany	mg/m ³	0.248–0.845 Mean 0.510
Heeb et al. (2012)	Haihe River, China	mg/m ³	0.5–4.5 Median 1.09
Esteban et al. (2014)	Manzanares and Jarama rivers, Spain	mg/m ³	0.097–1.184

groundwater samples collected from a karstified aquifer in Germany under an urban area of about 65 km²; this area was characterized by intensive industrial activities and several waste disposal sites.

In South Australia, Liu et al. (2011b) measured four BTs in groundwater collected near a WWTP, and BT was at a concentration of 0.280±0.0175 mg/m³. The presence of the BTs

in groundwater was expected due to pumping of WWTP effluents into aquifer for water reuse scheme. Often BTs were detected both in surface water and in aquifer system; in this cases, it seemed that BTs were partially eliminated through soil and subsurface passage (Kahle et al. 2009).

The concentrations of BT in surface water are reported between 0.011 and 33 mg/m³. Breedveld et al. (2002)

Table 7 Orders of magnitude of concentrations reported in the literature for the investigated pollutants in the different environmental matrices and food

Pollutant	Soils and sediments (µg/kg d.w.)	Groundwater and surface water (mg/m ³)	Foodstuff (µg/kg f.w.)
BPA	10 ⁻¹ –10 ²	10 ⁻³ –10 ²	10 ⁻¹ –10 ³
NP	10 ⁻² –10 ⁴	10 ⁻³ –10	10 ⁻¹ –10 ³
BPs	10 ⁻¹ –10 ²	10 ⁻³ –10	10 ⁻¹ –10
BTs	10 ⁻¹ –10 ⁴	10 ⁻³ –10 ³	–

measured concentrations of BT between 1.5 and 33 mg/m³ in samples collected in the area of the Fornebu Oslo airport. Giger et al. (2006) and Vousta et al. (2006) investigated the Glatt River and its valley, a densely populated region of 260 km² with 240,000 inhabitants in the northeastern part of Switzerland. The Glatt River catchment included the northern

Table 8 Available environmental quality standards for pollutants of concern in the investigated environmental matrices

Reference	Description	Units	Value
BPA—soils and sediments			
BCLAWS (2014)	Soil—standard, agricultural, urban park, residential soil (Canada)	mg/kg d.w.	3100
BCLAWS (2014)	Soil—standard, commercial, industrial soil (Canada)	mg/kg d.w.	31,000
USEPA (2014b)	Soil—regional screening level, residential soil (USA)	mg/kg d.w.	310
USEPA (2014b)	Soil—regional screening level, industrial soil (USA)	mg/kg d.w.	4100
BPA—waters			
CMEE (1994)	Surface water—proposed water quality objective (Canada)	mg/m ³	5
BCLAWS (2014)	Drinking water—standard (Canada)	mg/m ³	1800
USEPA (2014b)	Water—regional screening level, tapwater (USA)	mg/m ³	77
NP—soils and sediments			
DanishEPA (2000)	Soil—proposed quality criteria (Denmark)	mg/kg d.w.	25
CCME (2002); Alberta (2014)	Soil—quality guidelines, residential, agricultural soil (Canada)	mg/kg d.w.	5.7
CCME (2002); Alberta (2014)	Soil—quality guidelines, commercial, industrial soil (Canada)	mg/kg d.w.	14
ISS (2011)	Soil—proposed limit for residential soil (Italy)	mg/kg d.w.	0.05
ISS (2011)	Soil—proposed limit for commercial, industrial soil (Italy)	mg/kg d.w.	12.5
CCME (2002)	Sediment—quality guidelines (Canada)	mg/kg d.w.	1.4
CIRCABC (2005)	Sediment—proposed quality standard (Europe)	mg/kg d.w.	0.18
NP—waters			
ISS (2011)	Groundwater—proposed limit (Italy)	mg/m ³	0.3
Alberta (2014)	Groundwater—tier 1 remediation guidelines (Canada)	mg/m ³	6.6
CMEE (1994)	Surface water—proposed water quality objective (Canada)	mg/m ³	0.04
EC (2013)	Surface water—maximum acceptable concentration, environmental quality standard (Europe)	mg/m ³	2
European Parliament (2013)	Surface water—maximum annual average concentration, environmental quality standard (Europe)	mg/m ³	0.3
BPs—soils and sediments			
USEPA (2007)	Soil—remediation level Superfund Swannanoa (USA)	mg/kg d.w.	9.3
BPs—waters			
USEPA (2007)	Groundwater—remediation level Superfund Swannanoa (USA)	mg/m ³	152
BTs—soils and sediments			
No value available in the literature			
BTs—waters			
Kase et al. (2011)	Surface water—proposed maximum acceptable concentration, environmental quality standard	mg/m ³	120
Kase et al. (2011)	Surface water—proposed maximum annual average concentration, environmental quality standard	mg/m ³	30

part of the city of Zurich; ten municipal wastewater treatment plants discharged their final effluents into the river. Furthermore, the international airport of Zurich was located on the east of the Glatt River. Similar values were found in the two studies (0.06–6.3 mg/m³ in Giger et al. (2006) and 0.636–3.69 mg/m³ in Vousta et al. (2006)). The mass balance clearly indicated the input from the Zurich airport (28 % of the total BT load), where BT was used as an anticorrosive component. Giger et al. (2006) also analyzed waters from three prealpine lakes in the Canton of Zurich (Greifensee, Zurich, and Geneva). Greifensee lake is located northeast of Zurich and outflows into the Glatt River; a population of about 100,000 inhabitants live in the catchment discharging to the Greifensee. Eight municipal WWTPs discharged their treated effluents into Lake Zurich, while Lake Geneva received wastewaters from Lausanne City. BT concentrations measured in the three lakes were 0.9–1.1, 0.12–0.4, and 0.18–0.2 mg/m³, respectively. Kahle et al. (2009) measured BT concentration in water samples from eight lakes of the Swiss Midland region at values between 0.11 and 0.917 mg/m³; the BT concentrations indicate that this compound, regularly discharged to surface waters, was suitable quantitative markers of domestic wastewater in surface waters. Nodler et al. (2011) measured an increase of BT of 1 order of magnitude between water samples collected upstream (0.034–0.0176 mg/m³) and downstream (0.248–0.845 mg/m³) of the WWTP discharge in the Leine River (Germany). Kiss and Fries (2009) assessed the occurrence of BT in three German rivers (Main, Hengstbach, and Hegbach). Main and Hengstbach rivers received effluents from domestic WWTPs and were influenced by the Frankfurt International Airport; the concentrations ranged between 0.025 and 1.474 mg/m³. Due to the absence of wastewater effluents in the Hegbach River, a significant concentration (0.038 mg/m³) was observed just in one sample. In surface water samples of Spain, BT hot spots were found with a maximum concentration of 1184 µg/m³ in Manzanares River, 345 µg/m³ in Jarama River, and 1120 µg/m³ in Ter River (Esteban et al. 2014). Heeb et al. (2012) investigated the 175 km stretch of the Wenyu River and the North Canal from Beijing to Tianjin (China), resulting in BT concentrations between 0.5 and 4.5 mg/m³. Wastewaters from industrial areas and from the Beijing International Airport were the most probable sources of pollution.

Conclusions

The CECs discussed in this review are widespread in the environment due to consistent industrial production and persistence. Table 7 summarizes the concentration orders of magnitude reported in the literature for the different environmental matrices and foodstuff; Table 8 resumes the environmental quality standards worldwide for the pollutants of concern available in the literature. By comparing the environmental quality

standards in Table 8 with the ranges in Table 7, BPA levels in soils and sediments are not critical, even if the lowest value proposed for BPA in solid matrix (310 mg/kg, residential soil USA) is considered, as the maximum reported values are somewhat 3 orders of magnitude lower. However, in the future, it is possible that more stringent limits are defined. The Canadian water quality objective of 5 mg/m³ is quite stringent as in several situations up to an order of magnitude, and higher concentrations have been reported in the literature. For the NP, rigorous limits have been set by the European Commission. In Italy also, provisional limits 50 µg/kg for residential soil and 0.3 mg/m³ for groundwater exist. A more stringent water quality standard of 0.04 mg/m³ was proposed in Canada. As the widespread presence of NP at high levels in the various media is reported, it seems quite important to continue and intensify environmental monitoring. For BPs and BTs, the proposed standards are few; therefore, further studies are necessary for defining useful screening values to compare with. At present, the worldwide measured concentrations of BPs reported in the literature are in compliance with the values proposed for BPs by USEPA in 2007 (9.3 mg/kg for soil and 152 mg/m³ for groundwater). Finally, significant standards for BTs were not reported in the literature; the value of 30 mg/m³ proposed for surface water is about 2 orders of magnitude lower than the maximum BT concentrations. As for foodstuff and drinking water, the range of BPA concentrations in water fulfills the proposed limit for drinking water (100 mg/m³) (Willhite et al. 2008). The concentration of BPA in food is significant, especially in canned food, but the estimated daily intake via food consumption does not exceed the proposed RfD for oral exposure of 10 µg/kg/day (Rykowska and Wasiak 2006). In general, the concentration of NP in water does not pose risk to human health, but some values exceed the proposed limits by USEPA for acute exposure (6.6 mg/m³) and for chronic exposure (0.7–1.7 mg/m³) (USEPA 2010b). NP concentrations should be monitored in the output stream of treatment plants. High NP concentrations in food have been measured, in particular in edible fish, but in general, these values do not exceed the proposed RfD for oral exposure (100 µg/kg/day) or Danish TDI (5 µg/kg/day) (Bakke 2003). No concentration limits or guidelines have been proposed for BPs and BTs in the different environmental matrices. More investigation on the occurrence of BPs in food should be useful to exclude exceeding the proposed RfD for oral exposure (10 µg/kg/day) and the concentration limit in food (0.6 mg/kg f.w.), although the daily intake might be heavily increased by dermal sorption from cosmetics (Jeon et al. 2006; IARC 2010). Particular attention should be paid to BTs because of the high values measured in soils and sediments that might lead to exceed the proposed RfD for inhalation and dermal contact (30 µg/kg/day) (Giger et al. 2006).

Further research should be carried out on these CECs. Treatment technologies for waters have to improve the removal efficiency of BPA and NP in order to reduce their discharge

in water bodies from wastewater treatment plants. More investigation should be carried out to assess the presence and distribution of all these CECs in soils and of BPs and BTs in both surface water and groundwater. Specific studies should be carried out about possible pollution of soils due to the use of biosolids and/or treated wastewater in agriculture practices, in particular to understand the transport of CECs in the edible parts of plants and the risk for humans due to their consumption. The packaging composition for foodstuff should be reformulated to avoid contamination from BPA, NP, and BPs; moreover, the use of BPs as a flavor additive should be avoided. The knowledge of the toxicology of these compounds should be improved.

Acknowledgments This work was financially supported by the Italian Ministry of Education, University and Research in the National Research Program 2010–2011 “Emerging contaminants in air, soil, and water: from source to the marine environment.”

Open Access This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

References

Ahel M, Giger W, Koch M (1994) Behavior of alkylphenol polyethoxylate surfactants in the aquatic environment. 1. Occurrence and transformation in sewage-treatment. *Water Res* 28: 1131–1142

Alberta (2014) Alberta Soil and Groundwater Remediation Guidelines. <http://esrd.alberta.ca/lands-forests/land-industrial/inspections-and-compliance/alberta-soil-and-groundwater-remediation-guidelines.aspx>. Accessed Nov 2014

Anderson PD, Denslow ND, Drewes JE, Olivieri AW, Schlenk D, Scott GI, Snyder SA (2012) Monitoring strategies for chemicals of emerging concern (CECs) in California’s aquatic ecosystems—recommendations of a science advisory panel. Technical Report 692. California Water Resources Control Board

Andreozzi R, Caprio V, Insola A, Longo G (1998) Photochemical degradation of benzotriazole in aqueous solution. *J Chem Technol Biot* 73:93–98

Arias-Estévez M, López-Periágo E, Martínez-Carballo E, Simal-Gándara J, Mejuto JC, García-Río L (2008) The mobility and degradation of pesticides in soils and the pollution of groundwater resources. *Agric Ecosyst Environ* 123:247–260

Ávila C, Nivala J, Olsson L, Kassa K, Headley T, Mueller RA, Bayona JM, Garcia J (2014) Emerging organic contaminants in vertical subsurface flow constructed wetlands: influence of media size, loading frequency and use of active aeration. *Sci Total Environ* 494–495:211–217

Azevedo DA, Lacorte S, Viana P, Barcelo D (2001) Occurrence of nonylphenol and bisphenol A in surface waters from Portugal. *J Brazil Chem Soc* 12:532–537

Bakke D (2003) Human and ecological risk assessment of nonylphenol polyethoxylate-based (NPE) surfactants in forest service herbicide

applications. Technical report. USDA Forest Service. Pacific Southwest Region (Region 5)

Balmer ME, Buser HR, Muller MD, Poiger T (2005) Occurrence of some organic UV filters in wastewater, in surface waters, and in fish from Swiss lakes. *Environ Sci Technol* 39:953–962

Barón E, Gago-Ferrero P, Gorga M, Rudolph I, Mendoza G, Zapata AM, Díaz-Cruz S, Barra R, Ocampo-Duque W, Páez M, Darbra RM, Eljarrat E, Barceló D (2013) Occurrence of hydrophobic organic pollutants (BFRs and UV-filters) in sediments from South America. *Chemosphere* 92:309–316

Basheer C, Lee HK, Tan KS (2004) Endocrine disrupting alkylphenols and bisphenol-A in coastal waters and supermarket seafood from Singapore. *Mar Pollut Bull* 48:1145–1167

BCLAWS (2014). Environmental Management Act - CONTAMINATED SITES REGULATION – British Columbia, Canada. http://www.bclaws.ca/civix/document/id/complete/statreg/375_96_06#Schedule4. Accessed Nov 2014

Benítez FJ, Acero JL, Real FJ, Roldan G, Rodriguez E (2013) Photolysis of model emerging contaminants in ultra-pure water: kinetics, by-products formation and degradation pathways. *Water Res* 47:870–880

Bennett ER, Metcalfe CD (1997) Distribution of alkylphenol compounds in great lakes sediments, United States and Canada. *Environ Toxicol Chem* 17(7):1230–1235

Bennie DT, Sullivan CA, Lee HB, Peart TE, Maguire RJ (1997) Occurrence of alkylphenols and alkylphenol mono- and diethoxylates in natural waters of the Laurentian Great Lakes basin and the upper St. Lawrence River. *Sci Total Environ* 193:263–275

Bester K, Theobald N, Schroder HF (2001) Nonylphenols, nonylphenol-ethoxylates, linear alkylbenzenesulfonates (LAS) and bis(4-chlorophenyl)-sulfone in the German Bight of the North Sea. *Chemosphere* 45:817–826

Bhatarai B, Gramatica P (2011) Modelling physico-chemical properties of (benzo)triazoles, and screening for environmental partitioning. *Water Res* 45:1463–1471

Bi E, Schmidt TC, Haderlein SB (2007) Environmental factors influencing sorption of heterocyclic aromatic compounds to soil. *Environ Sci Technol* 41:3172–3178

Breedveld GD, Roseth R, Hem L (2002) Triazoles in the terrestrial environment. 20001103–1. Final Report. Norwegian Geotechnical Institute, Oslo, Norway

Brooks AC, Gaskell PN, Maltby LL (2009) Importance of prey and predator feeding behaviors for trophic transfer and secondary poisoning. *Environ Sci Technol* 43:7916–7923

Brown S, Devin-Clarke D, Doubrava M, O’Connor G (2009) Fate of 4-nonylphenol in a biosolids amended soil. *Chemosphere* 75:549–554

Buser HR, Poiger T, Müller MD (1998) Occurrence and fate of the pharmaceutical drug diclofenac in surface waters: rapid photodegradation in a lake. *Environ Sci Technol* 33:3449–3456

Cacho JI, Campillo N, Vinas P, Hernandez-Cordoba M (2012) Determination of alkylphenols and phthalate esters in vegetables and migration studies from their packages by means of stir bar sorptive extraction coupled to gas chromatography–mass spectrometry. *J Chromatogr A* 1241:21–27

Cancilla DA, Baird JC, Rosa R (2003) Detection of aircraft deicing additives in groundwater and soil samples from Fairchild Air Force Base, a small to moderate user of deicing fluids. *B Environ Contam Tox* 70:868–875

Cao XL, Perez-Locas C, Dufresne G, Clement G, Popovic S, Beraldin F, Dabeka RW, Feeley M (2011) Concentrations of bisphenol A in the composite food samples from the 2008 Canadian total diet study in Quebec City and dietary intake estimates. *Food Addit Contam A* 28(6):791–798

CCME (2002) Canadian soil quality guidelines for the protection of environmental and human health: nonylphenol and its ethoxylates. In: Canadian environmental quality guidelines, Canadian Council of Ministers of the Environment, Winnipeg

- Cespedes R, Lacorte S, Raldua D, Ginebreda A, Barcelo D, Pina B (2005) Distribution of endocrine disruptors in the Llobregat River basin (Catalonia, NE Spain). *Chemosphere* 61:1710–1719
- Chang BV, Yu CH, Yuan SY (2004) Degradation of nonylphenol by anaerobic microorganisms from river sediment. *Chemosphere* 55: 493–500
- Chang BV, Liu JH, Liao CS (2014) Aerobic degradation of bisphenol-A and its derivatives in river sediment. *Environ Technol* 35(4):416–424
- Chefetz B, Mualem T, Ben-Ari J (2008) Sorption and mobility of pharmaceutical compounds in soil irrigated with reclaimed wastewater. *Chemosphere* 73(8):1335–1343
- Chen W, Xu J, Lu S, Jiao W, Wu L, Chang AC (2013) Fates and transport of PPCPs in soil receiving reclaimed water irrigation. *Chemosphere* 93:2621–2630
- Christian T, Schneider R, Farber HA, Skutlarek D, Meyer MT, Goldbach HE (2003) Determination of antibiotic residues in manure, soil, and surface waters. *Acta Hydrochim Hydrobiol* 31:36–44
- CIRCABC (2005). Common Implementation Strategy for the Water Framework Directive, Environmental Quality Standards (EQS) Substance Data Sheet Priority Substance No. 24 4-Nonylphenol (branched) and Nonylphenol, CAS-No. 84852-15-3 and 25154-52-3 -Final version Brussels, 31 July 2005 at the CIRCABC server. https://circabc.europa.eu/sd/a/af1b09f2-ff9a-46f6-ba2d-d4bc2adfee0/24_Nonylphenol_EQSdatasheet_310705.pdf. Accessed Dec 2014
- CME (1994) Canadian Ministry of Environment and Energy. Policies guidelines provincial water quality objectives – Ontario. PIBS 3303E
- Cornell JS, Pillard DA, Hernandez MT (2000) Comparative measures of the toxicity of component chemicals in aircraft deicing fluids. *Environ Toxicol Chem* 19:1465–1472
- Corvini PFX, Schäffer A, Schlosser D (2006a) Microbial degradation of nonylphenol and other alkylphenols—our evolving view. *Appl Microbiol Biot* 72:223–243
- Corvini PFX, Hollender J, Ji R, Schumacher S, Prell J, Hommes G, Priefer U, Vinken R, Schäffer A (2006b) The degradation of α -quaternary nonylphenol isomers by *Sphingomonas* sp. strain TTNP3 involves a type II ipso-substitution mechanism. *Appl Microbiol Biot* 70:114–122
- DanishEPA (2000) Toxicological evaluation and limit values for nonylphenol, nonylphenol ethoxylates, tricresyl, phosphates and benzoic acid - Environmental Project no. 512 2000 <http://www.statensnet.dk/pligtarkiv/fremvis.pl?vaerkid=6944&repid=0&flid=0&iarkiv=1>. Accessed Dec 2014
- Daughton CG, Ternes TA (1999) Pharmaceuticals and personal care products in the environment: agents of subtle change? *Environ Health Perspect* 107:907–937
- DECOS (2000) 1,2,3-Benzotriazole. 2000/14OSH. Technical report. The Hague: Health Council of the Netherlands, Dutch Expert Committee on Occupational Standards
- Diehl J, Johnson SE, Xia K, West A, Tomanek L (2012) The distribution of 4-nonylphenol in marine organisms of North American Pacific Coast estuaries. *Chemosphere* 87(5):290–297
- Ding H, Li Y, Hou J, Wang Q, Wu Y (2014) Sorption behavior and modeling of endocrine-disrupting chemicals on natural sediments: role of biofilm covered on surface. *Environ SciPollut Res*. doi:10.1007/s11356-014-3449-8
- Dodder NG, Maruya KA, Ferguson PL, Grace R, Klosterhaus S, La Guardia MJ, Lauenstein GG, Ramirez J (2014) Occurrence of emerging contaminants of emerging concern in mussels (*Mytilus* spp.) along the California coast, and the influence of land use, storm water discharge, and treated wastewater effluent. *MarPollut Bull* 81: 340–346
- Dodgen LK, Li J, Parker D, Gan JJ (2013) Uptake and accumulation of four PPCP/EDCs in two leafy vegetables. *Environ Pollut* 182:150–156
- Dodgen LK, Li J, Wu X, Lu Z, Gan JJ (2014) Transformation and removal pathways of four common PPCP/EDCs in soil. *Environ Pollut* 193:29–36
- Duan X, Li Y, Li X, Zhang DH, Gao Y (2014) Alkylphenols in surface sediments of the Yellow Sea and East China Sea inner shelf: Occurrence, distribution and fate. *Chemosphere* 107:265–273
- EC (2002) European Union Risk Assessment Report—4-Nonylphenol (branched) and nonylphenol, Volume 10. EUR 20387 EN. European Commission - Joint Research Centre
- EC (2003) Directive 2003/53/EC of the European Parliament and the Council of 18 June 2003 amending for the 26th time Council Directive 76/769/EEC relating to restrictions on the marketing and use of certain dangerous substances and preparations (nonylphenol, nonylphenol ethoxylate and cement). *Official Journal of the European Union* 17.7.2003. L 178/24
- EC (2005) Provisional list of monomers and additives notified to European Commission as substances which may be used in the manufacture of plastics or coatings intended to come into contact with foodstuff. SANCO D3/AS D(2005). Technical report. European Commission Health & Consumer Protection Directorate-general
- EC (2013) Directive 2013/39/EU of the European Parliament and of the Council of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy. *Official Journal of the European Union* 24.8.2013. L 226/1
- ECHA (2014) Nonylphenol and nonylphenol ethoxylates. Technical report. European Chemicals Agency, Committee for Risk Assessment (RAC), Committee for Socio-economic Analysis (SEAC)
- Esteban S, Gorga M, Petrovic M, González-Alonso S, Barceló D, Valcárcel Y (2014) Analysis and occurrence of endocrine-disrupting compounds and estrogenic activity in the surface waters of Central Spain. *Sci Total Environ* 466–467:939–951
- EUGRIS (2013) Portal for soil and water management in Europe. Emerging pollutants. <http://www.eugris.info/FurtherDescription.asp?Ca=2&Cy=0&T=Emerging%20Pollutants&e=95>. Accessed May 2014
- Félix-Cañedo TE, Durán-Álvarez JC, Jiménez-Cisneros B (2013) The occurrence and distribution of a group of organic micropollutants in Mexico City's water sources. *Sci Total Environ* 454–455:109–118
- Fent G, Hein W, Moendel M, Kubiak R (2003) Fate of 14C-bisphenol A in soils. *Chemosphere* 51:735–746
- Fent K, Zenker A, Rapp M (2010) Widespread occurrence of estrogenic UV-filters in aquatic ecosystems in Switzerland. *Environ Pollut* 158: 1817–1824
- Ferrara F, Fabietti F, Delise M, Funari E (2005) Alkylphenols and alkylphenol ethoxylates contamination of crustaceans and fishes from the Adriatic Sea (Italy). *Chemosphere* 59:1145–1150
- Ferrara F, Ademollo N, Delise M, Fabietti F, Funari E (2008) Alkylphenols and their ethoxylates in seafood from the Tyrrhenian Sea. *Chemosphere* 72:1279–1285
- Flint S, Markle T, Thompson S, Wallace E (2012) Bisphenol A exposure, effects and policy: a wildlife perspective. *J Environ Manage* 104: 19–34
- Focazio MJ, Kolpin DW, Barnes KK, Furlong ET, Meyer MT, Zaugg SD, Barber LB, Thurman EM (2008) A national reconnaissance of pharmaceuticals and other organic wastewater contaminants in the United States—II. Untreated drinking water sources. *Sci Total Environ* 402(2–3):201–216
- Fries E, Puttmann W (2003) Occurrence and behaviour of 4-nonylphenol in river water of Germany. *J Environ Monit* 5:598–603
- Fu M, Li Z, Gao H (2007) Distribution characteristics of nonylphenol in Jiaozhou Bay of Qingdao and its adjacent rivers. *Chemosphere* 69: 1009–1016

- Fujii K, Kituchi S (2005) Degradation of benzophenone, a potential xenoestrogen, by a yeast isolated from the activated sludge of a sewage treatment plant in Hokkaido. *World J Microb Biot* 21:1311–1315
- Gabriel F, Routledge E, Heidberger A, Rentsch D, Guenther K, Giger W, Sumpter J, Kohler H (2008) Isomer-specific degradation and endocrine disrupting activity of nonylphenols. *Environ Sci Technol* 42:6399–6408
- Gago-Ferrero P, Badia-Fabregat M, Olivares A, Piña B, Blázquez P, Vicent T, Caminal G, Díaz-Cruz M, Barceló D (2012) Evaluation of fungal- and photo-degradation as potential treatments for the removal of sunscreens BP3 and BP1. *Sci Total Environ* 427–428:355–363
- Gago-Ferrero P, Diaz-Cruz MS, Barcelo D (2013) Multi-residue method for trace level determination of UV filters in fish based on pressurized liquid extraction and liquid chromatography quadrupole-linear ion trap-mass spectrometry. *J Chromatogr A* 1286:93–101
- Gao JA, Pedersen JA (2005) Adsorption of sulfonamide antimicrobial agents to clay minerals. *Environ Sci Technol* 39(24):9509–9516
- Gibson R, Durán-Álvarez J, León Estrada K, Chávez A, Jiménez Cisneros B (2010) Accumulation and leaching potential of some pharmaceuticals and potential endocrine disruptors in soils irrigated with wastewater in the Tula Valley, Mexico. *Chemosphere* 81:1437–1445
- Giger W, Schaffner C, Petere H, Kohler H-P (2006) Benzotriazole and tolyltriazole as aquatic contaminants. Input and occurrence in rivers and lakes. *Environ Sci Technol* 40:7186–7192
- Godejohann M, Heintz L, Daolio C, Berset J, Muff D (2009) Comprehensive non-targeted analysis of contaminated groundwater of a former ammunition destruction site using 1H-NMR and HPLC-SPE-NMR/TOF-MS. *Environ Sci Technol* 43:7055–7061
- Gong J, Yang Y, Chen DY, Ran Y (2011) Sequential ASE extraction of alkylphenols from sediments: occurrence and environmental implications. *J Hazard Mater* 192:643–650
- Gorga M, Insa S, Petrovic M, Barceló D (2014) Analysis of endocrine disruptors and related compounds in sediment sand sewage sludge using on-line turbulent flow chromatography–liquid chromatography–tandem mass spectrometry. *J Chromatogr A* 1352:29–37
- Gorga M, Insa S, Petrovic M, Barceló D (2015) Occurrence and spatial distribution of EDCs and related compounds in waters and sediments of Iberian rivers. *Sci Total Environ* 503–504:69–86
- Grabicova K, Fedorova G, Burkina V, Steinbach C, Schmidt-Posthaus H, Zlabek V, Kroupova HK, Grabic R, Randak T (2013) Presence of UV filters in surface water and the effects of phenylbenzimidazole sulfonic acid on rainbow trout (*Oncorhynchus mykiss*) following a chronic toxicity test. *Ecotox Environ Safe* 96:41–47
- Guenther K, Heinke V, Thiele B, Kleist E, Prast H, Raecker T (2002) Endocrine disrupting nonylphenols are ubiquitous in food. *Environ Sci Technol* 36:1676–1680
- Guidechem (2013) Oxybenzone. <http://www.guidechem.com/reference/dic-2893.html> Accessed May 2014
- Gyllenhammar I, Glynn A, Damerud PO, Lignell S, van Delft R, Aune M (2012) 4-Nonylphenol and bisphenol A in Swedish food and exposure in Swedish nursing women. *Environ Int* 43:21–28
- Harris C, Routledge E, Schaffner C, Brian J, Giger W, Sumpter J (2007) Benzotriazole is antiestrogenic in vitro but not in vivo. *Environ Toxicol Chem* 26(11):2367–2372
- Hart DS, Davis LC, Erickson LE, Callender TM (2004) Sorption and partitioning parameters of benzotriazole compounds. *Microchem J* 77:9–17
- Hayashi T, Okamoto Y, Ueda K, Kojima N (2006) Formation of estrogenic products from benzophenone after exposure to sunlight. *Toxicol Lett* 167:1–7
- Heeb F, Singer H, Pernet-Coudrier B, Qi W, Liu H, Longree P, Muller B, Berg M (2012) Organic micropollutants in rivers downstream of the megacity Beijing: sources and mass fluxes in a large-scale wastewater irrigation system. *Environ Sci Technol* 46:8680–8688
- Heemken OP, Reincke H, Stachel B, Theobald N (2001) The occurrence of xenoestrogens in the Elbe river and the North Sea. *Chemosphere* 45(3):245–259
- Hem LJ, Hartnik T, Roseth R, Breedveld GD (2003) Photochemical degradation of benzotriazole. *J Environ Sci Heal A* 38(3):471–481
- Hogenboom A, Van Leerdam J, Voogt P (2009) Accurate mass screening and identification of emerging contaminants in environmental samples by liquid chromatography–hybrid linear ion trap Orbitrap mass spectrometry. *J Chromatogr A* 1216:510–519
- Hou SG, Sun HW, Gao Y (2006) Sorption of small metabolites of nonylphenol polyethoxylates in single and complex systems on aquatic suspended particulate matter. *Chemosphere* 63:31–38
- Huang YQ, Wong CKC, Zheng JS, Bouwman H, Barra R, Wallhstrom B, Neretin L, Wong MH (2012) Bisphenol A (BPA) in China: a review of sources, environmental levels, and potential human health impacts. *Environ Int* 42:91–99
- IARC (1999) Monographs on the evaluation of carcinogenic risks to humans. Re-evaluation of some organic chemicals, hydrazine and hydrogen peroxide. IARC Monographs vol. 71. International Agency for Research on Cancer, Working Group on the Evaluation of Carcinogenic Risks to Humans, Lyon, France
- IARC (2010) Benzophenone. IARC Monographs 101. International Agency for Research on Cancer
- ICIS (2007) Nonylphenol. <http://www.icis.com/Articles/2007/06/25/4503346/nonylphenol.html>. Accessed May 2014
- Ike M, Chen M, Jin C, Fujita M (2002) Acute toxicity, mutagenicity, and estrogenicity of biodegradation products of bisphenol-A. *Environ Toxicol* 17(5):457–461
- Isobe T, Takada H, Kanai M, Tsutsumi S, Isobe KO, Boonyatumanond R, Zakaria MP (2007) Distribution of polycyclic aromatic hydrocarbons (PAHs) and phenolic endocrine disrupting chemicals in South and Southeast Asian mussels. *Env Monit Ass* 135:423–440
- ISS (2011) Richiesta Parere sulle concentrazioni soglie di contaminazione per le sostanze non normate, riscontrate durante l'esecuzione del piano di caratterizzazione dell'area Chemtura Manufacturing Italy di Pedrelungo (Bg). 48353 AMPP.IA.06 (15/11/2011)
- Jeon H, Chung Y, Ryu J (2006) Simultaneous determination of benzophenone-type UV filters in water and soil by gas chromatography–mass spectrometry. *J Chromatogr A* 1131:192–202
- Jia Y, Bakken L, Breedveld G, Aagaard P, Frosteg A (2006) Organic compounds that reach subsoil may threaten groundwater quality; effect of benzotriazole on degradation kinetics and microbial community composition. *Soil Biol Biochem* 38:2543–2556
- Jonkers N, Laane R, De Voogt P (2003) Fate of nonylphenol ethoxylates and their metabolites in two Dutch estuaries: evidence of biodegradation in the field. *Environ Sci Technol* 37:321–327
- Jurado A, Gago-Ferrero P, Vázquez-Suné E, Carrera J, Pujades E, Díaz-Cruz MS, Barceló D (2014) Urban groundwater contamination by residues of UV filters. *J Hazard Mater* 271:141–149
- Kahle M, Buerge IJ, Muller MD, Poiger T (2009) Hydrophilic anthropogenic markers for quantification of wastewater contamination in ground- and surface waters. *Environ Toxicol Chem* 28(12):2528–2536
- Kalmykova Y, Björklund K, Strömvall AM, Blom L (2013) Partitioning of polycyclic aromatic hydrocarbons, alkylphenols, bisphenol A and phthalates in landfill leachates and stormwater. *Water Res* 47:1317–1328
- Kameda Y, Kimura K, Miyazaki M (2011) Occurrence and profiles of organic sun-blocking agents in surface waters and sediments in Japanese rivers and lakes. *Environ Pollut* 159:1570–1576
- Kannan K, Keith TL, Naylor CG, Staples CA, Snyder SA, Giesy JP (2003) Nonylphenol and nonylphenol ethoxylates in fish, sediment, and water from the Kalamazoo River, Michigan. *Arch Environ Con Tox* 44:77–82

- Kase R, Eggen RIL, Junghans M, Gotz C, Juliane H (2011) Assessment of micropollutants from municipal wastewater—combination of exposure and ecotoxicological effect data for Switzerland. In: Einschlag FSG (ed) Waste water—evaluation and management. InTech, Rijeka, pp 31–54
- Kasprzyk-Hordern B, Dinsdale RM, Guwy AJ (2008) The occurrence of pharmaceuticals, personal care products, endocrine disruptors and illicit drugs in surface water in South Wales, UK. *Water Res* 42: 3498–3518
- Kawahata H, Ohta H, Inoue M, Suzuki A (2004) Endocrine disruptor nonylphenol and bisphenol A contamination in Okinawa and Ishigaki Islands, Japan—within coral reefs and adjacent river mouths. *Chemosphere* 55:1519–1527
- Kibbey TCG, Paruchuri R, Sabatini DA, Chen L (2007) Adsorption of beta-blockers to environmental surfaces. *Environ Sci Technol* 41: 5349–5356
- Kim S, Choi K (2014) Occurrences, toxicities, and ecological risks of benzophenone-3, a common component of organic sunscreen products: a mini-review. *Environ Int* 70:143–157
- Kimura K, Hara H, Watanabe Y (2007) Elimination of selected acidic pharmaceuticals from municipal wastewater by an activated sludge system and membrane bioreactors. *Environ Sci Technol* 41:3708–3714
- Kinney CA, Furlong ET, Zaugg SD, Burkhard MR, Werner SL, Cahill JD, Jorgensen GR (2006) Survey of organic wastewater contaminants in biosolids destined for land application. *Environ Sci Technol* 40(23):7207–7215
- Kinney CA, Furlong ET, Kolpin DW, Burkhardt MR, Zaugg SD, Werner SL, Bossio JP, Benotti MJ (2008) Bioaccumulation of pharmaceuticals and other anthropogenic waste indicators in earthworms from agricultural soil amended with biosolid or swine manure. *Environ Sci Technol* 42:1863–1870
- Kiss A, Fries EF (2009) Occurrence of benzotriazoles in the rivers main, Hengstbach, and Hegbach (Germany). *Environ Sci Pollut Res* 16: 702–710
- Klosterhaus S, Grace R, Hamilton MC, Yee D (2013) Method validation and reconnaissance of pharmaceuticals, personal care products, and alkylphenols in surface waters, sediments, and mussels in an urban estuary. *Environ Int* 54:92–99
- Kolpin DW, Furlong ET, Meyer MT, Thurman EM, Zaugg SD, Barber LB, Buxton HT (2002) Pharmaceuticals, hormones, and other organic wastewater contaminants in U.S. streams 1999–2000: a national reconnaissance. *Environ Sci Technol* 36:1202–1211
- Koniecko I, Staniszewska M, Falkowska L, Burska D, Kielczewska J, Jasinska A (2014) Alkylphenols in surface sediments of the Gulf of Gdansk (Baltic Sea). *Water Air Soil Pollut* 225:2040
- Koschorreck J, Koch C, Rönnefahrt I (2002) Environmental risk assessment of veterinary medicinal products in the EU—a regulatory perspective. *Toxicol Lett* 131:117–124
- Kueh CSW, Lam JYC (2008) Monitoring of toxic substances in the Hong Kong marine environment. *Mar Pollut Bull* 57(6):744–757
- Kumar K, Singh AK, Chander Y, Gupta SC (2005) Antibiotic use in agriculture and its impact on the terrestrial environment. *Adv Agron* 87:1–54
- La Farré M, Perez S, Kantiani L, Barcelo D (2008) Fate and toxicity of emerging pollutants, their metabolites and transformation products in the aquatic environment. *Trac-Trend Anal Chem* 27(11)
- Lacorte S, Latorre A, Guillamon M, Barcelò D (2002) Nonylphenol, octylphenol, and bisphenol a in groundwaters as a result of agromonic practices. *Sci World J* 2:1095–1100
- Lara-Martin PA, Gomez-Parra A, Barcelo D, Gonzalez-Mazo E (2006) Presence of surfactants and their degradation intermediates in sediment cores and grabs from the Cadiz Bay area. *Environ Pollut* 144: 483–491
- Latorre A, Lacorte S, Barcelo D (2003) Presence of nonylphenol, octylphenol and bisphenol-A in two aquifers close to agricultural, industrial and urban areas. *Chemosphere* 57:111–116
- Lee Ferguson P, Bopp RF, Chillrud SN, Aller RC, Brownawell BJ (2003) Biogeochemistry of nonylphenol ethoxylates in urban estuarine sediments. *Environ Sci Technol* 37:3499–3506
- León Z, Chisvert A, Tarazona I, Salvador A (2010) Solid-phase extraction liquid chromatography–tandem mass spectrometry analytical method for the determination of 2-hydroxy-4-methoxybenzophenone and its metabolites in both human urine and semen. *Anal Bioanal Chem* 398: 831–843
- Li J, Jiang L, Liu X, Lv J (2013a) Adsorption and aerobic biodegradation of four selected endocrine disrupting chemicals in soil water system. *Int Biodeter Biodegr* 76:3–7
- Li Y, Duan X, Li X, Zhang D (2013b) Photodegradation of nonylphenol by simulated sunlight. *Mar Pollut Bull* 66:47–52
- Li Y, Zhang S, Song C, You J (2013c) Determination of Bisphenol A and Alkylphenols in Soft Drinks by High-Performance Liquid Chromatography with Fluorescence Detection. *Food Anal Methods* 6:1284–1290
- Liao X, Zhang C, Yao L, Li J, Liu M, Xu L, Evalde M (2014) Sorption behavior of nonylphenol (NP) on sewage-irrigated soil: kinetic and thermodynamic studies. *Sci Total Environ* 473–474:530–536
- Lin A, Reinhard M (2005) Photodegradation of common environmental pharmaceuticals and estrogens in river water. *Environ Toxicol Chem* 24(6):1303–1309
- Liu YS, Ying GG, Shareef A, Kookana RS (2011a) Biodegradation of three selected benzotriazoles under aerobic and anaerobic conditions. *Water Res* 45:5005–5014
- Liu YS, Ying GG, Shareef A, Kookana RS (2011b) Simultaneous determination of benzotriazoles and ultraviolet filters in ground water, effluent and biosolid samples using gas chromatography–tandem mass spectrometry. *J Chromatogr A* 1218:5328–5335
- Liu YS, Ying GG, Shareef A, Kookana RS (2012a) Biodegradation of the ultraviolet filter benzophenone-3 under different redox conditions. *Environ Toxicol Chem* 31(2):289–295
- Liu YS, Ying GG, Shareef A, Kookana RS (2012b) Occurrence and removal of benzotriazoles and ultraviolet filters in a municipal wastewater treatment plant. *Environ Pollut* 165:225–232
- Loos R, Hanke G, Umlauf G, Eisenreich SJ (2007) LC-MS-MS analysis and occurrence of octyl- and nonylphenol, their ethoxylates and their carboxylates in Belgian and Italian textile industry, waste water treatment plant effluents and surface waters. *Chemosphere* 66:690–699
- Loos R, Locoro G, Comero S, Contini S, Schwesig D, Werres F, Balsaa P, Gans O, Weiss S, Blaha L, Bolchi M, Gawlik B (2010) Pan-European survey on the occurrence of selected polar organic persistent pollutants in ground water. *Water Res* 44:4115–4126
- Loyo-Rosales JE, Rosales-Rivera GC, Lynch AM, Rice CP, Torrents A (2004) Migration of nonylphenol from plastic containers to water and a milk surrogate. *J Agr Food Chem* 52:2016–2020
- Lu Z, Gan J (2014) Isomer-specific biodegradation of nonylphenol in river sediments and structure-biodegradability relationship. *Environ Sci Technol* 48:1008–1014
- Lu J, Wu J, Stoffella PJ, Wilson PC (2013) Analysis of bisphenol A, nonylphenol, and natural estrogens in vegetables and fruits using gas chromatography–tandem mass spectrometry. *J Agr Food Chem* 61: 84–89
- Luo Y, Guo W, Ngo HH, Nghiem LD, Hai FI, Zhang J, Liang S, Wang XC (2014) A review on the occurrence of micropollutants in the aquatic environment and their fate and removal during wastewater treatment. *Sci Total Environ* 473–474:619–641
- Maggioni S, Balaguer P, Chiozzotto C, Benfenati E (2013) Screening of endocrine-disrupting phenols, herbicides, steroid estrogens, and estrogenicity in drinking water from the waterworks of 35 Italian cities and from PET-bottled mineral water. *Environ Sci Pollut Res* 20:1649–1660
- Martínez-Zapata M, Aristizábal C, Penuela G (2013) Photodegradation of the endocrine-disrupting chemicals 4n-nonylphenol and triclosan by simulated solar UV irradiation in aqueous solutions with Fe(III) and

- in the absence/presence of humic acids. *J Photoch Photobio A* 251: 41–49
- McNeill KS, Cancilla DA (2009) Detection of triazole deicing additives in soil samples from airports with low, mid, and large volume aircraft deicing activities. *B Environ Contam Tox* 82(3):265–269
- Melo SM, Brito MN (2014) Analysis and occurrence of endocrine disruptors in Brazilian water by HPLC-fluorescence detection. *Water Air Soil Pollut* 225:1783
- Mercea P (2009) Physicochemical processes involved in migration of bisphenol A from polycarbonate. *J Appl Polym Sci* 112(2):579–593
- Michałowicz J (2014) Bisphenol A—sources, toxicity and biotransformation. *Env Toxicol Pharm* 37:738–758
- Micic V, Hofmann T (2009) Occurrence and behaviour of selected hydrophobic alkylphenolic compounds in the Danube River. *Environ Pollut* 157:2759–2768
- Micić V, Krüge MA, Hofmann T (2013) Variations of common riverine contaminants in reservoir sediments. *Sci Total Environ* 458–460: 90–100
- Molnar M, Gruiz K, Hajdu CS, Nagy ZS, Fenyvesi E (2013) Tiered approach for environmental risk assessment of emerging pollutants in aquatic systems. *Proceedings, AquaConSoil 2013*, 12th International UFZ-Delta Conference on Groundwater-Soil-Systems and Water Resource Management, Barcelona, Spain, April 16–19 2013, session C14
- Neamtu M, Frimmel FH (2006) Photodegradation of endocrine disrupting chemical nonylphenol by simulated solar UV-irradiation. *Sci Total Environ* 369:295–306
- Nodler K, Licha T, Fischer S, Wagner B, Sauter M (2011) A case study on the correlation of micro-contaminants and potassium in the Leine River (Germany). *Appl Geochem* 26:2172–2180
- Noonan GO, Ackerman LK, Begley TH (2011) Concentration of bisphenol A in highly consumed canned foods on the U.S. market. *J Agr Food Chem* 59:7178–7185
- NORMAN (2013) Network of reference laboratories for monitoring of emerging environmental pollutants. http://www.norman-network.net/index_php.php?module=public/about_us/home. Accessed May 2014
- NTP (2006) Toxicology and carcinogenesis studies of benzophenone (CAS No. 119–61–9) in F344/N rats and B6C3F1 mice (feed studies). NTP TR 533. Technical report. National Toxicology Program, U.S. Department of Health and Human Services. NIH Publication no. 06–4469
- OEHHA (2009) Toxicological profile for nonylphenol. Office of Environmental Health Hazard Assessment. California Environmental Protection Agency
- Patrolocco L, Capri S, De Angelis S, Pagnotta R, Polesello S, Valsecchi S (2006) Partition of nonylphenol and related compounds among different aquatic compartments in Timber River (Central Italy). *Water Air Soil Poll* 172:151–166
- Pojana G, Jonkers N, Marcomini A (2007) Natural and synthetic endocrine disrupting compounds (EDCs) in water, sediment and biota of a coastal lagoon. *Environ Int* 33:929–936
- Rajendra S.C (2000) NTP technical report on the toxicity studies of benzophenone. Toxicity Report Series Number 61. U.S. Department of Health and Human Services Public Health Service, National Institutes of Health
- Reh R, Licha T, Geyer T, Nodler K, Sauter M (2013) Occurrence and spatial distribution of organic micro-pollutants in a complex hydrogeological karst system during low flow and high flow periods, results of a two-year study. *Sci Total Environ* 443:438–445
- Rice CP, Schmitz-Afonso I, Loyo-Rosales JE, Link E, Thoma R, Fay L (2003) Alkylphenol and alkylphenol-ethoxylates in carp, water, and sediment from the Cuyahoga River, Ohio. *Environ Sci Technol* 37: 3747–3754
- Richardson ML, Bowron JM (1985) The fate of pharmaceutical chemicals in the aquatic environment. *J Pharm Pharmacol* 37:1–12
- Ricking M, Schwarzbauer J, Franke S (2003) Molecular markers of anthropogenic activity in sediments of the Havel and Spree Rivers (Germany). *Water Res* 37:2607–2617
- RIKZ (2001) Chemical study on bisphenol A. RIKZ/2001.027. Technical report. Rijksinstituut voor Kust en Zee, Ministerie van Verkeer en Waterstaat, The Netherlands
- Roberts S, Higgins C, McCray J (2014) Sorption of emerging organic wastewater contaminants to four soils. *Water* 6:1028–1042
- Rodil R, Moeder M, Altenburger R, Schmitt-Jansen M (2009) Photostability and phytotoxicity of selected sunscreen agents and their degradation mixtures in water. *Anal Bioanal Chem* 395:1513–1524
- Rodil R, Quintana JB, Concha-Grana E, Lopez-Mahia P, Muniategui-Lorenzo S, Prada-Rodriguez D (2012) Emerging pollutants in sewage, surface and drinking water in Galicia (NW Spain). *Chemosphere* 86:1040–1049
- Rozalska S, Szewczyk R, Długoski J (2010) Biodegradation of 4-n-nonylphenol by the non-ligninolytic filamentous fungus *Glioccephalotrichum simplex*: a proposal of a metabolic pathway. *J Hazard Mater* 180:323–331
- Rykowska I, Wasiak (2006) Properties, threats, and methods of analysis of bisphenol A and its derivatives. *Acta Chromatogr* 16:7–27
- Sánchez-Brunete C, Miguel E, Albero B, Tadeo J (2011) Analysis of salicylate and benzophenone-type UV filters in soils and sediments by simultaneous extraction cleanup and gas chromatography–mass spectrometry. *J Chromatogr A* 1218:4291–4298
- Sekela M, Brewer R, Moyle G, Tuominen T (1999) Occurrence of an environmental estrogen (4-nonylphenol) in sewage treatment plant effluent and the aquatic receiving environment. *Water Sci Technol* 39(10–11):217–220
- Shao B, Han H, Li D, Ma Y, Tu X, Wu Y (2007) Analysis of alkylphenol and bisphenol A in meat by accelerated solvent extraction and liquid chromatography with tandem mass spectrometry. *Food Chem* 105: 1236–1241
- Shchegolikhina A, Schulz S, Marchner B (2012) Interacting effects of cation saturation and drying, freezing, or aging on the extractability of nonylphenol and phenanthrene from a sandy soil. *J Soil Sediment* 12:1280–1291
- Soares A, Guieysse B, Jefferson B, Cartmell E, Lester JN (2008) Nonylphenol in the environment: a critical review on occurrence, fate, toxicity and treatment in wastewaters. *Environ Int* 34:1033–1049
- Spivack J, Leib TK, Lobos JH (1994) Novel pathway for bacterial metabolism of bisphenol A, rearrangements and stilbene cleavage in bisphenol A metabolism. *J Biol Chem* 269:7323–7329
- Staples C, Friederich U, Hall T, Klecka G, Mihaich E, Ortego L, Caspers N, Hentges S (2010) Estimating potential risks to terrestrial invertebrates and plants exposed to bisphenol A in soil amended with activated sludge biosolids. *Environ Toxicol Chem* 29(2):467–475
- Stewart M, Olsena G, Hickey WC, Ferreira B, Jelić A, Petrović M, Barcelo D (2014) A survey of emerging contaminants in the estuarine receiving environment around Auckland, New Zealand. *Sci Total Environ* 468–469:202–210
- Stuart ME, Manamsa K, Talbot JC, Crane EJ (2011) Emerging contaminants in groundwater. OR/11/013. British Geological Survey Open Report. Keyworth, Nottingham (UK), 123 pp
- Sun C, Leong LP, Barlow PJ, Chan SH, Bloodworth BC (2006) Single laboratory validation of a method for the determination of bisphenol a, bisphenol A diglycidyl ether and its derivatives in canned foods by reversed-phase liquid chromatography. *J Chromatogr A* 1129: 145–148
- Ternes TA, Bonerz M, Herrmann N, Teiser B, Andersen HR (2007) Irrigation of treated wastewater in Braunschweig, Germany: an

- option to remove pharmaceuticals and musk fragrances. *Chemosphere* 66:894–904
- Thomaidis NS, Asimakopoulos AG, Bletsou AA (2012) Emerging contaminants: a tutorial mini-review. *Global NEST J* 14(1):72–79
- Tolls J (2001) Sorption of veterinary pharmaceuticals in soils: a review. *Environ Sci Technol* 35(17):3397–3406
- Topp E, Starratt A (2000) Rapid mineralization of the endocrine-disrupting chemical 4-nonylphenol in soil. *Environ Toxicol Chem* 19:313–318
- TOXNET (2013a) Toxicology Data Network—benzotriazole. <http://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+4143>. Accessed May 2014
- TOXNET (2013b) Toxicology Data Network—benzophenone. <http://toxnet.nlm.nih.gov/cgi-bin/sis/search/a?dbs+hsdb:@term+@DOCNO+6809>. Accessed May 2014
- TOXNET (2014) Toxicology Data Network—2-hydroxy-4-methoxybenzophenone. <http://toxnet.nlm.nih.gov/cgi-bin/sis/search2/r?dbs+hsdb:@term+@m+@rel+131-57-7>. Accessed Nov 2014
- USEPA (1984) Information review: benzophenone. U.S. Environmental Protection Agency, Washington
- USEPA (2007) Second superfund five-year review report Chemtronics Superfund Site Swannanoa, Buncombe County, NC. EPA ID: NCD 095 459 392
- USEPA (2010a) Bisphenol A action plan. U.S. Environmental Protection Agency, Washington
- USEPA (2010b) Nonylphenol (NP) and nonylphenol ethoxylates (NPEs) action plan. RIN 200-ZA09. Technical report. U.S. Environmental Protection Agency, Washington, DC
- USEPA (2010c) 2-(2'-Hydroxy-3',5'-di-tert-amyphenyl) benzotriazole and phenol, 2-(2H-benzotriazole-2-yl)-6-dodecyl-4-methyl; exemption from the requirement of a tolerance. U.S. Environmental Protection Agency, Washington, DC. Federal Register volume 75 no 159
- USEPA (2014a) Persistent organic pollutants: a global issue, a global response. <http://www2.epa.gov/international-cooperation/persistent-organic-pollutants-global-issue-global-response>. Accessed Nov 2014
- USEPA (2014b). Regional screening level. http://www.epa.gov/reg3hwmd/risk/human/rb-concentration_table/Generic_Tables/. Accessed Nov 2014
- USEPA-IRIS (2014) Bisphenol A. Quickview <http://cfpub.epa.gov/ncea/iris/index.cfm?fuseaction=iris.show> Quickview&substance_nmbr=0356. Accessed Nov 2014.
- Vethaak AD, Lahr J, Schrap SM, Belfroid AC, Rijs GBJ, Gerritsen A, de Boer J, Bulder AS, Grinwis GCM, Kuiper RV, Legler J, Murk TAJ, Peijnenburg W, Verhaar HJM, de Voogt P (2005) An integrated assessment of estrogenic contamination and biological effects in the aquatic environment of The Netherlands. *Chemosphere* 59: 511–524
- Vikelsøe J, Thomsen M, Carlsen L (2002) Phthalates and nonylphenols in profiles of differently dressed soils. *Sci Total Environ* 296:105–116
- Vitali M, Ensabella F, Stella D, Guidotti M (2004) Nonylphenols in freshwaters of the hydrologic system of an Italian district: association with human activities and evaluation of human exposure. *Chemosphere* 57:1637–1647
- Vousta D, Hartmann P, Schaffner C, Giger W (2006) Benzotriazoles, alkylphenols and bisphenol A in municipal wastewaters and in the Glatt River, Switzerland. *Environ Sci Pollut R* 13(5):333–341
- Weber S, Khan S, Hollender J (2005) Human risk assessment of organic contaminants in reclaimed wastewater used for irrigation. In: Khan SJ, Muston MH, Schafer AI (eds) *Integrated concepts in water recycling*. University of Wollongong, Wollongong, pp 724–735
- Wehrhan A, Kasteel R, Simunek J, Groeneweg J, Vereecken H (2007) Transport of sulfadiazine in soil columns—experiments and modeling approaches. *J Contam Hydrol* 89(1–2):107–135
- Weiss S, Reemtsma T (2005) Determination of benzotriazole corrosion inhibitors from aqueous environmental samples by liquid chromatography-electrospray ionization-tandem mass spectrometry. *Anal Chem* 77(22):7415–7420
- Willhite CC, Ball GL, McLellan CJ (2008) Derivation of a bisphenol A oral reference dose (RfD) and drinking-water equivalent concentration. *J Toxicol Env Heal B* 11(2):69–146
- Williams CF, Adamsen FJ (2006) Sorption–desorption of carbamazepine from irrigated soils. *J Environ Qual* 35:1779–1783
- Wolschke H, Xie Z, Moller A, Sturm R, Ebinghaus R (2011) Occurrence, distribution and fluxes of benzotriazoles along the German large river basins into the North Sea. *Water Res* 45:6259–6266
- Wright-Walters M, Volz C, Talbot E, Davis D (2011) An updated weight of evidence approach to the aquatic hazard assessment of bisphenol A and the derivation a new predicted no effect concentration (Pnec) using a non-parametric methodology. *Sci Total Environ* 409:676–685
- Writer JH, Ryan JN, Keefe SH, Barber LB (2012) Fate of 4-nonylphenol and 17 β -estradiol in the Redwood River of Minnesota. *Environ Sci Technol* 46:860–868
- Wu ZB, Zhang Z, Chen SP, He F, Fu GP, Liang W (2007) Nonylphenol and octylphenol in urban eutrophic lakes of the subtropical China. *Fresen Environ Bull* 16:227–234
- Wu M, Wang L, Xu G, Liu N, Tang L, Zheng J, Bu T, Lei B (2013) Seasonal and spatial distribution of 4-tert-octylphenol, 4-nonylphenol and bisphenol A in the Huangpu River and its tributaries, Shanghai, China. *Environ Monit Assess* 185:3149–3161
- Xu J, Wu L, Chen W, Chang A (2008) Simultaneous determination of pharmaceuticals, endocrine disrupting compounds and hormone in soils by gas chromatography–mass spectrometry. *J Chromatogr A* 1202(2):189–195
- Xu J, Chen W, Wu L, Green R (2009) Leachability of some emerging contaminants in reclaimed municipal wastewater-irrigated turf grass fields. *Environ Toxicol Chem* 28(9):1842–1850
- Xu EGB, Liu S, Ying GG, Zheng GJS, Lee JHW, Leung MYK (2014) The occurrence and ecological risks of endocrine disrupting chemicals in sewage effluents from three different sewage treatment plants, and in natural seawater from a marine reserve of Hong Kong. *Mar Pollut Bull* 85:352–362
- Xuan R, Blassengale AA, Wang Q (2008) Degradation of estrogenic hormones in a silt loam soil. *J Agr Food Chem* 56:9152–9158
- Yamamoto H, Liljestrand HM (2003) The fate of estrogenic compounds in the aquatic environment: sorption onto organic colloids. *Water Sci Technol* 47(9):77–84
- Yamashita N, Kannan K, Imagawa T, Villeneuve DL, Hashimoto S, Miyazaki A, Giesy JP (2000) Vertical profile of polychlorinated dibenzo-p-dioxins, dibenzofurans, naphthalenes, biphenyls, polycyclic aromatic hydrocarbons, and alkylphenols in a sediment core from Tokyo Bay, Japan. *Environ Sci Technol* 34:3560–3567
- Yang GP, Ding HY, Cao XY, Ding QY (2011) Sorption behavior of nonylphenol on marine sediments: effect of temperature, medium, sediment organic carbon and surfactant. *Mar Pollut Bull* 62:2362–2369
- Yang Y, Wang Z, Xie S (2014) Aerobic biodegradation of bisphenol A in river sediment and associated bacterial community change. *Sci Total Environ* 470–471:1184–1188
- Ye X, Zhou X, Needham LL, Calafat AM (2011) In-vitro oxidation of bisphenol A: is bisphenol A catechol a suitable biomarker for human exposure to bisphenol A? *Anal Bioanal Chem* 399:1071–1079
- Ying GG, Kookana RS (2003) Degradation of five selected endocrine disrupting chemicals in seawater and marine sediment. *Environ Sci Technol* 37:1256–1260
- Ying G, Kookana R (2005) Sorption and degradation of estrogen-like-endocrine disrupting chemicals in soil. *Environ Toxicol Chem* 24(10):2640–2645

- Yoon Y, Ryu J, Oh J, Choi BG, Snyder SA (2010) Occurrence of endocrine disrupting compounds, pharmaceuticals, and personal care products in the Han River (Seoul, South Korea). *Sci Total Environ* 408:636–643
- Yu JT, Bouwer EJ, Coelhan M (2006) Occurrence and biodegradability studies of selected pharmaceuticals and personal care products in sewage effluent. *Agr Water Manage* 86:72–80
- Yu L, Fink G, Wintgens T, Melin T, Ternes TA (2009) Sorption behavior of potential organic wastewater indicators with soils. *Water Res* 43: 951–960
- Yu Y, Liu Y, Wu L (2013) Sorption and degradation of pharmaceuticals and personal care products (PPCPs) in soils. *Environ Sci Pollut Res* 20:4261–4267
- Zenker A, Schmutz H, Fent K (2008) Simultaneous trace determination of nine organic UV-absorbing compounds (UV filters) in environmental samples. *J Chromatogr A* 1202:64–74
- Zhan M, Yang X, Xian Q, Kong L (2006) Photosensitized degradation of bisphenol A involving reactive oxygen species in the presence of humic substances. *Chemosphere* 63:378–386
- Zhang C, Zeng G, Yuan L, Yu J, Li J, Huang G, Xi B, Liu H (2007) Aerobic degradation of bisphenol A by *Achromobacter xylosoxidans* strain B-16 isolated from compost leachate of municipal solid waste. *Chemosphere* 68:181–190
- Zhang Z, Ren N, Li YF, Kunisue T, Gao D, Kannan K (2011) Determination of benzotriazole and benzophenone UV filters in sediment and sewage sludge. *Environ Sci Technol* 45:3909–3916
- Zhang W, Yin K, Chen L (2013a) Bacteria-mediated bisphenol A degradation. *Appl Microbiol Biotechnol* 97:5681–5689
- Zhang T, Sun H, Qin X, Wu Q, Zhang Y, Ma J, Kannan K (2013b) Benzophenone-type UV filters in urine and blood from children, adults, and pregnant women in China: partitioning between blood and urine as well as maternal and fetal cord blood. *Sci Total Environ* 461–462:49–55
- Zhang Y, Meng W, Zhang Y (2014) Occurrence and partitioning of phenolic endocrine-disrupting chemicals (EDCs) between surface water and suspended particulate matter in the North Tai Lake Basin, Eastern China. *Bull Environ Contam Toxicol* 92:148–153
- Zhao J, Li Y, Zhang C, Zeng Q, Zhou Q (2008) Sorption and degradation of bisphenol A by aerobic activated sludge. *J Hazard Mater* 155: 305–311
- Zhou JL, Liu R, Wilding A, Hibberd A (2007) Sorption of selected endocrine disrupting chemicals to different aquatic colloids. *Environ Sci Technol* 41:206–213
- Zwiener C, Frimmel FH (2003) Short-term test with a pilot sewage plant and biofilm reactors for the biological degradation of pharmaceutical compounds clofibrac acid, ibuprofen, and diclofenac. *Sci Total Environ* 309:201–211