

1 **The contribution of phenolic acids to the anti-inflammatory activity of**
2 **mushrooms: screening in phenolic extracts, individual parent molecules**
3 **and synthesized glucuronated and methylated derivatives**

4
5 Oludemi Taofiq^{a,b}, Ricardo C. Calhelha^{a,c}, Sandrina Heleno^{a,c}, Lillian Barros^a, Anabela
6 Martins^a, Celestino Santos-Buelga^b, Maria João R.P. Queiroz^c, Isabel C.F.R. Ferreira^{a,*}

7
8 ^a*Mountain Research Centre (CIMO), ESA, Polytechnic Institute of Bragança, Campus de*
9 *Santa Apóstola, 1152, 5301-855 Bragança, Portugal*

10 ^b*GIP-USAL, Faculty of Pharmacy, University of Salamanca, Campus Miguel de Unamuno,*
11 *37007 Salamanca, Spain*

12 ^c*Center of Chemistry, University of Minho, Campus de Gualtar 4710-057 Braga, Portugal*

13
14

ABSTRACT

15 In the present study, the ethanolic extracts of fourteen edible mushrooms were investigated
16 for their anti-inflammatory potential in LPS (lipopolysaccharide) activated RAW 264.7
17 macrophages. Furthermore the extracts were chemically characterized in terms of phenolic
18 acids and related compounds. The identified molecules (*p*-hydroxybenzoic, *p*-coumaric and
19 cinnamic acids) and their glucuronated and methylated derivatives obtained by chemical
20 synthesis were also evaluated for the same bioactivity, in order to establish structure-activity
21 relationships and to comprehend the effects of *in vivo* metabolism reactions in the activity of
22 the compounds. The extracts of *Pleurotus ostreatus*, *Macrolepiota procera*, *Boletus impolitus*
23 and *Agaricus bisporus* revealed the strongest anti-inflammatory potential (EC₅₀ values 96 ± 1
24 to 190 ± 6 µg/mL, and also the highest concentration of cinnamic acid (656 to 156 µg/g),
25 which was also the individual compound with the highest anti-inflammatory activity. The
26 derivatives of *p*-coumaric acid revealed the strongest properties, specially the derivative
27 methylated in the carboxylic group (CoA-M1) that exhibited similar activity to the one
28 showed by dexamethaxone used as anti-inflammatory standard; by contrast, the derivatives of
29 *p*-hydroxybenzoic revealed the lowest inhibition of NO production. All in all, whereas the
30 conjugation reactions change the chemical structure of phenolic acids and may increase or
31 decrease their activity, the glucuronated and methylated derivatives of the studied compounds
32 are still displaying anti-inflammatory activity.

33

34 **Keywords:** Edible Mushrooms; Phenolic acids; glucuronated and methylated derivatives;
35 Anti-inflammatory; Nitric oxide production; HPLC-PDA

36

37 1. INTRODUCTION

38 Inflammation is considered to be part of the complex biological response to remove injury or
39 harmful stimuli such as pathogens, damaged cells, or irritation and this is a central feature of
40 many pathophysiological conditions such as atherosclerosis, obesity, metabolic syndrome,
41 diabetes ([Pradhan, 2007](#)) and even several types of cancers ([Moro et al., 2012](#)).

42 When cells are exposed to immune stimulants, the pro-inflammatory cells, such as
43 macrophages, monocytes, or other host cells, start to produce cytokines and other mediators,
44 which initiate the inflammation process. Among the various inflammatory mediators, the
45 most common are interleukins (IL-1 β , IL-6, IL-8), tumour necrosis factor (TNF- α), nuclear
46 factor- κ B (NF- κ B), intercellular adhesion molecule-1 (ICAM-1), inducible type
47 cyclooxygenase-2 (COX-2), prostaglandin E2 (PGE2), 5-lipoxygenase (5-LOX), and
48 inducible nitric oxide synthase (iNOS) that leads to the production of reactive nitrogen
49 species such as nitric oxide (NO). Overproduction of these inflammatory mediators leads to
50 different kinds of cell damage ([Kanwar, Kanwar, Burrow, & Baratchi, 2009](#)).

51 Recently, the nonsteroidal anti-inflammatory drugs (NSAIDs) are usually the most
52 commonly administered drugs to reduce inflammation in the body. Many studies, however,
53 have shown that the long-term administration of NSAIDs has the potential for significant side
54 effects on the gastrointestinal tract with numerous harmful effects such as mucosal lesions,
55 bleeding, peptic ulcers, and intestinal perforation ([Dugowson & Gnanashanmugam, 2006](#)).

56 Recent studies show that NSAIDs are also associated with a relatively high incidence of renal
57 adverse drug reactions, nephrotic syndrome, high blood pressure, acute tubular necrosis and
58 cardiovascular toxicity ([Elsayed, Hesham, Mohammad, & Ramlan, 2014](#)).

59 As a result, research studies are now being channelled towards discovery of bioactive
60 compounds with ability to suppress the production of inflammatory mediators. A good model
61 to test potential anti-inflammatory drugs are macrophages, which are large specialized cells

62 that engulf and digest cellular debris, microbes or cancer cells in a process called
63 phagocytosis. They play an important role in non-specific host defence mechanisms and help
64 to initiate other defence mechanisms. Beyond stimulating the immune system, macrophages
65 play a crucial role in the inflammatory response through the release of a variety of factors,
66 such as NO, TNF- α , IL-1 β , IL-6, in response to an activating stimulus, e.g.
67 lipopolysaccharide (Moro et al., 2012).

68 Mushrooms are widely appreciated all over the world not only for their culinary and
69 nutritional properties (Kalac, 2009), but also for their pharmacological value as sources of
70 important bioactive molecules, such as antioxidant (Puttaraju, Venkateshaiah, Dharmesh,
71 Urs, & Somasundaram, 2006; Ferreira, Barros, & Abreu, 2009; Heleno, Martins, Queiroz, &
72 Ferreira, 2015), antitumor (Moradali, Mostafavi, Ghods, & Hedjaroude, 2007; Ferreira, Vaz,
73 Vasconcelos, & Martins, 2010, Carochó & Ferreira, 2013), antimicrobial (Alves, Ferreira,
74 Dias, Teixeira, Martins, & Pintado, 2012; Alves, Ferreira, Froufe, Abreu, Martins, & Pintado,
75 2013), immunomodulator (Borchers, Krishnamurthy, Keen, Meyers, & Gershwin, 2008),
76 antiatherogenic (Mori, Kobayashi, Tomita, Inatomi, & Ikeda, 2008) and hypoglycemic
77 compounds (Hu, Wang, Lien, Liaw, & Lee, 2006). Due to these properties, they have been
78 recognized as functional foods, as well as valuable sources of natural medicines and
79 nutraceuticals (Lindequist, Niedermeyer, Jülich, 2005; Guillamón et al., 2010).

80 Moreover, mushrooms have also demonstrated some anti-inflammatory potential based on
81 their ability to reduce the production of inflammatory mediators (Padilha, Avila, Sousa,
82 Cardoso, Perazzo, & Carvalho, 2009; Elsayed et al., 2014). Previous research studies have
83 been carried out on several mushroom species, mainly in methanolic and ethanolic extracts
84 (Table 1). Different compounds have been pointed out as the responsible for the anti-
85 inflammatory activity such as β -glucans (Nosálóva, Bobek, Cerna, Galbavy, & Tvrtina,
86 2001), triterpenes (Ma, Chen, Dong, & Lu, 2013), glycoproteins (Gunawardena et al., 2014)

87 and even phenolic compounds (Moro et al., 2012). However, not much is known about the
88 phenolic bioactive forms *in vivo*; these compounds are metabolized and circulate in the
89 organism as glucuronated, sulfated and methylated metabolites, displaying higher or lower
90 bioactivity (Heleno et al., 2015).

91 Therefore, the present study aimed to investigate the anti-inflammatory activity of selected
92 mushroom species from the Northeast of Portugal, using ethanolic extracts in LPS activated
93 RAW 264.7 macrophages. Furthermore, after characterization of the extracts in terms of
94 phenolic acids and related compounds, the identified individual parent molecules and their
95 synthesised glucuronated and methylated derivatives were evaluated for the same bioactivity,
96 in order to establish structure-activity relationships.

97

98 2. MATERIALS AND METHODS

99 2.1. Mushroom species and extracts preparation

100 Ten wild mushroom species (*Amanita caesaria* (Scop.) Pers., *Boletus aereus* Bull., *B. edulis*
101 Bull., *B. flagrans* Vittad., *B. impolitus* Fr., *B. reticulatus* Schaeff., *Cantharellus cibarius* Fr.,
102 *Lactarius deliciosus* (L. ex Fr.) S.F. Gray, *Macrolepiota procera* (Scop.) Singer and
103 *Morchella esculenta* Fr.), collected in the Northeast of Portugal, and four cultivated species
104 (*Agaricus bisporus* (J.E. Lange) Emil J. Imbach, *A. bisporus* Portobello (J.E. Lange) Emil J.
105 Imbach, *Pleurotus eryngii* (DC.) Quél. and *Pleurotus. ostreatus* (Jacq. ex Fr.) P. Kumm.)
106 were used in the present study. All species were deposited in the herbarium of the School of
107 Agriculture in Polytechnic Institute of Bragança, and were previously characterized by the
108 research group in terms of nutritional value and chemical composition (including primary and
109 secondary metabolites) (Barros, Dueñas, Ferreira, Baptista, & Santos-Buelga, 2009,
110 Grangeia, Heleno, Barros, Martins, & Ferreira, 2011; Heleno, Barros, Sousa, Martins, Santos-
111 Buelga, & Ferreira., 2011; Reis et al., 2011; Pereira, Barros, Martins, & Ferreira, 2012; Reis,

112 Barros, Martins, & Ferreira, 2012; Heleno et al., 2013b). Their antioxidant, antimicrobial and
113 antitumor properties were also previously evaluated by the group
114 (<http://esa.ipb.pt/biochemcore/index.php/studied-mushrooms>). In the present work, the *in*
115 *vitro* anti-inflammatory activity was evaluated in ethanolic extracts prepared as follows.
116 Lyophilized (Ly-8-FM-ULE, Snijders, Holland) mushroom powder (20 mesh) of each species
117 (0.5 g) was extracted with ethanol (15 mL), by maceration with stirring for 1 h. Then, the
118 extract was filtered through Whatman no 4 filter paper and the extraction procedure was
119 repeated one more time. The filtrate was rotary evaporated to remove ethanol and the
120 extraction yield was calculated by measuring the extract weight.

121

122 **2.2. Reagents**

123 Acetonitrile 99.9% was of high-performance liquid chromatography (HPLC) grade from Lab-
124 Scan (Lisbon, Portugal). *p*-Hydroxybenzoic acid, *p*-coumaric acid, cinnamic acid, Dulbecco's
125 modified Eagle's minimum essential medium (DMEM), fetal bovine serum (FBS), penicillin,
126 streptomycin, Griess reagent system (Promega), DMSO, sulphorodamine B (SRB) and
127 lipopolysaccharide (LPS) were obtained from Sigma-Aldrich Co. (Saint Louis, MO, USA).
128 All other chemicals and solvents were of analytical grade and purchased from common
129 suppliers.

130

131 **2.3. Chemical characterization of the extracts**

132 The dry mass of each mushroom extract was re-dissolved in water/ethanol (50:50, v/v) and
133 filtered through a 0.22 µm nylon disposable filter for HPLC analysis. The analysis was
134 performed using a Shimadzu 20A series ultra-fast liquid chromatograph (UFLC, Shimadzu
135 Cooperation, Kyoto, Japan). Separation was achieved on a Waters Spherisorb S3 ODS2 C₁₈
136 column (3 µm, 150 mm x 4.6 mm) column thermostatted at 35 °C. The solvents used were:

137 (A) 0.1% formic acid in water, (B) acetonitrile. The elution gradient established was: 10% B
138 to 15% B over 5 min, 15–25% B over 5 min, 25–35% B over 10 min, isocratic 50% B for 10
139 min, and re-equilibration of the column, using a flow rate of 0.5 mL/min. Detection was
140 carried out in a photodiode array detector (PDA), using 280 nm as the preferred wavelength.
141 The phenolic acids (group of phenolic compounds identified in the samples) were quantified
142 by comparison of the area of their peaks recorded at 280 nm with calibration curves obtained
143 from commercial standards of each compound: protocatechuic acid ($y = 164741x$, $R^2=0.999$),
144 *p*-hydroxybenzoic acid ($y = 113523x$, $R^2=0.999$), *p*-coumaric acid ($y = 433521x$, $R^2=0.998$)
145 and cinnamic acid ($y = 583527x$, $R^2=0.998$), 5 to 80 µg/mL. The results were expressed as µg
146 per g of extract.

147

148 **2.4. Phenolic acids and synthesised derivatives**

149 *p*-Hydroxybenzoic acid, *p*-coumaric acid and cinnamic acid, identified in the mushroom
150 extracts, were used as starting reagents for the synthesis of methylated and glucuronated
151 derivatives of the identified compounds (**Figure 1**); these compounds were previously
152 synthesized and completely characterized by the authors ([Heleno et al., 2013a](#); [Heleno et al.,](#)
153 [2014b](#)). Briefly, the glucuronated derivatives (HA-GP, CoA-GP and CA-GP) were obtained
154 by reacting the parent molecules with acetobromo- α -*D*-glucuronic acid methyl ester under
155 argon and using DMSO (dimethylsulfoxide) as solvent at room temperature. The methylated
156 derivatives (HA-M1, CoA-M1 and CA-M) were prepared using methanol and sulphuric acid
157 at room temperature; HA-M2 and CoA-M2 were synthesised using dimethyl sulphate in
158 acetone at room temperature. Finally, HA-M3 and CoA-M3 were obtained by the hydrolysis
159 of compounds HA-M2 and CoA-M2 using ethanol at 65°C and adjusting the pH to 3. All the
160 synthesised compounds were fully characterized by ^1H NMR, ^{13}C NMR, HRMS (high
161 resolution mass spectrometry) and melting point.

162

163 **2.5. Evaluation of the anti-inflammatory activity**

164 *2.5.1. Cells treatment*

165 The anti-inflammatory activity was carried out according to [Moro et al. \(2012\)](#) and [García-](#)
166 [Lafuente et al. \(2014\)](#) with some modifications. The mouse macrophage-like cell line RAW
167 264.7 was cultured in DMEM medium supplemented with 10% heat-inactivated foetal bovine
168 serum, glutamine and antibiotics at 37 °C under 5% CO₂, in humidified air. For each
169 experiment, cells were detached with a cell scraper. A cell density of 5 x 10⁵ cells/mL was
170 used, and the proportion of dead cells was less than 5%, according to Trypan blue dye
171 exclusion test.

172 Cells were seeded in 96-well plates at 150,000 cells/well and allowed do attach to the plate
173 overnight. Then, cells were treated with the different concentrations of each one of the
174 extracts for 1h. Dexamethasone (50 µM) was used as a positive control for the experiment.
175 The following step was the stimulation with LPS (1 µg/mL) for 18h. The effect of all the
176 tested samples in the absence of LPS was also evaluated, in order to observe if they induced
177 changes in nitric oxide (NO) basal levels. In negative controls, no LPS was added. Both
178 extracts and LPS were dissolved in supplemented DMEM.

179

180 *2.5.2. Nitric oxide determination*

181 Both the extracts, and the pure identified compounds and their synthesized derivatives were
182 submitted to the anti-inflammatory activity assay. The ethanolic extracts were dissolved in
183 water (non cytotoxic solvent) at 8 mg/mL, while the identified individual compounds and
184 their synthesised methylated and glucuronated derivatives were dissolved in DMSO at 50%
185 concentration in stock solutions. These solutions were then submitted to further dilutions
186 (400 µg/mL to 50 µg/mL and 2500 µM to 39 µM, for the extracts and compounds,

187 respectively) in order to determine effective concentrations (Moro et al., 2012; García-
188 Lafuente et al., 2014).

189 For the determination of nitric oxide, Griess Reagent System kit was used, which contains
190 sulphanilamide, N-(1-naphthyl)ethylenediamine hydrochloride (NED) and nitrite solutions. A
191 reference curve of the nitrite (sodium nitrite 100 μ M to 1.6 μ M; $y=0.0066x+0.1349$;
192 $R^2=0.9986$) was prepared in a 96-well plate. The cell culture supernatant (100 μ L) was
193 transferred to the plate and mixed with Sulphanilamide and NED solutions, 5-10 minutes
194 each, at room temperature. The nitric oxide produced was determined by measuring the
195 absorbance at 540 nm (microplate reader ELX800 Biotek), and by comparison with the
196 standard calibration curve.

197

198 **2.6. Statistical analysis**

199 For all the experiments three samples were analyzed and all the assays were carried out in
200 triplicate. The results are expressed as mean values \pm standard deviation (SD). The
201 differences between the different samples were analyzed using one-way analysis of variance
202 (ANOVA) followed by Tukey's honestly significant difference post hoc test with $\alpha = 0.05$,
203 coupled with Welch's statistic. This treatment was carried out using SPSS v. 22.0 program.

204

205 **3. RESULTS AND DISCUSSION**

206 ***3.1. Chemical characterization of the extracts***

207 Ethanollic extracts were prepared from fourteen different edible mushroom species: *Agaricus*
208 *bisporus*, *A. bisporus* Portobello, *Amanita caesaria*, *Boletus aereus*, *B. edulis*, *B. flagrans*, *B.*
209 *impolitus*, *B. reticulatus*, *Cantharellus cibarius*, *Lactarius deliciosus*, *Macrolepiota procera*,
210 *Pleurotus eryngii*, *Pleurotus ostreatus* and *Morchella esculenta*. One phenolic acid (*p*-
211 hydroxybenzoic acid) and two cinnamic acids (*p*-coumaric and cinnamic acids) were detected
212 by HPLC-PDA in the different extracts; their concentrations are presented in **Table 2**. The

213 phenolic composition of the mushrooms seems to be characterised by the presence of
214 phenolic acids, being cinnamic acid the major compound in most cases. No peaks were found
215 in the extracts whose UV spectra could be associated to hydroxycinnamic acids or their
216 tartaric or quinic esters, nor were flavonoids found. The fact that no flavonoids were
217 identified is not unexpected since, it is assumed that only plants possess the biosynthetic
218 ability to produce flavonoids and not animals and fungi (Iwashina, 2000). Furthermore, these
219 compounds have been extensively reported in mushroom species (Barros et al., 2009; Vaz et
220 al., 2011).

221 The three compounds were only identified and quantified in three of the analysed species, *B.*
222 *aereus*, *B. impolitus* and *P. ostreatus*, while *A. caesarea*, *C. cibarius* and *L. deliciosus*
223 presented *p*-hydroxybenzoic and cinnamic acids. The other mushroom samples only revealed
224 the presence of cinnamic acid. *P. ostreatus* revealed the highest concentration of phenolic
225 acids mainly due to the high contribution of cinnamic ($619 \pm 3 \mu\text{g/g}$) and *p*-hydroxybenzoic
226 ($295 \pm 5 \mu\text{g/g}$) acids. *M. procera* and *B. impolitus* also presented high amounts of cinnamic
227 acid ($522 \pm 1 \mu\text{g/g}$ and $505 \pm 12 \mu\text{g/g}$, respectively). The samples *A. bisporus* Portobello, *B.*
228 *edulis* and *B. flagrans* revealed only the presence of cinnamic acid, but in very low
229 concentration. Reis et al. (2011) and Heleno et al. (2013b) also reported the presence of
230 protocatechuic acid in *A. caesaria* and *M. esculenta*, respectively, although they performed an
231 extraction with methanol:water, while in the present study ethanolic extracts were prepared.
232 Therefore, the different extraction conditions could be responsible for the dissimilarity
233 observed in the phenolic profiles.

234

235 **3.2. In vitro anti-inflammatory activity of the extracts**

236 The effects of the ethanolic extracts on the production of inflammatory mediators (NO) in
237 RAW 264.7 macrophages, upon stimulation with LPS, are shown in **Figure 2**. The results

238 indicate that most of the assayed extracts inhibited LPS-induced NO production in a dose-
239 dependent manner. For an easier comparison of the results, EC₅₀ values were calculated
240 based on 50% of inhibition of NO production (**Table 3**). The most efficient species was *P.*
241 *ostreatus* (96 ± 1 µg/mL), followed by *M. procera* (162 ± 2 µg/mL), *B. impolitus* (166 ± 10
242 µg/mL) and *A. bisporus* (190 ± 6 µg/mL). These results are in agreement with the reports by
243 [Moro et al. \(2012\)](#) and [Gunawardena et al. \(2014\)](#) that described anti-inflammatory activity,
244 by decreasing NO levels in RAW 264.7 cells, of ethanolic and methanolic extracts of *A.*
245 *bisporus*, *C. cibarius*, *L. deliciosus* and *P. ostreatus* (**Table 1**). The ethanolic extracts from *A.*
246 *bisporus* Portobello, *B. edulis* and *B. flagrans* appeared as the less active, showing EC₅₀
247 values above 400 µg/mL. [Moro et al. \(2012\)](#) reported some activity in the case of methanolic
248 extracts of *B. edulis* with 10% inhibition of NO production at concentrations of 500 µg/mL.
249 Although many substances may participate in the anti-inflammatory activity, phenolic
250 compounds have been largely recognised as natural molecules with anti-inflammatory
251 effects. Positive correlations have been found between phenolic compounds and anti-
252 inflammatory effects ([Cheung, Cheung, & Ooi, 2003](#); [Kim et al., 2008](#)). In the present study,
253 it was also observed that the extract with the highest anti-inflammatory activity showed the
254 highest levels of cinnamic and phenolic acids.

255

256 **3.3. *In vitro* anti-inflammatory activity of the compounds identified in the extracts**

257 To demonstrate this supposition, the activity of the individual compounds present in the
258 extracts were further evaluated (**Table 4**). Phenolic acids, and cinnamic acid, that are the
259 main responsible molecules for the bioactivities exhibited by mushrooms are well known for
260 their biological properties, mainly due to the OH groups present in their chemical structure,
261 either in the carboxylic group or in the phenolic ring ([Heleno et al., 2015](#)). Cinnamic acid
262 (CA) showed the highest anti-inflammatory activity presenting the lowest EC₅₀ values (182 ±

263 16 μM), followed by *p*-hydroxybenzoic ($239 \pm 29 \mu\text{M}$) and *p*-coumaric ($442 \pm 33 \mu\text{M}$) acids.
264 Cinnamic acid presents a carboxylic group and no OH groups in the benzene ring. *p*-
265 Hydroxybenzoic and *p*-coumaric acids present an OH group in the *para* position that is
266 usually a position described as having biological properties. However, in the present study
267 and comparing the activity exhibited by these three molecules, the OH group in the *para*
268 position seems to decrease the anti-inflammatory ability, maybe due to the mechanism of
269 action in this specific cell line. Several authors suggest that the anti-inflammatory activity is
270 related with the ability of the compounds to inhibit the activity of the cyclooxygenase (COX)
271 enzyme, which is responsible for the synthesis of prostaglandins, mediators with a great
272 ability to induce inflammation (Lee et al., 2006; Tanaka et al., 2009). Studies on the
273 mechanism of action of these molecules are important to better understand their behaviour.
274 Nevertheless, these results are in agreement with the ones reported by our research group for
275 the antimicrobial activity of these compounds, where cinnamic acid also revealed the
276 strongest activity followed by *p*-hydroxybenzoic and *p*-coumaric acids (Heleno et al., 2014a).

277

278 ***3.4. In vitro anti-inflammatory activity of glucuronated and methylated derivatives of the*** 279 ***identified compounds***

280 As phenolic acids are metabolized in the organism and suffer conjugation reactions
281 originating different metabolites such as glucuronated and methylated derivatives, a change
282 in their effects or activity may also occur. Thus, the bioactivity of the parent molecule can be
283 increased, decreased or maintained (Heleno et al., 2015). Hereby the glucuronated and
284 methylated derivatives of the considered acids were analysed and compared to the one of the
285 parent molecule. Among the glucuronated derivatives, CoA-GP (glucuronated derivative of
286 *p*-coumaric acid) presented strong anti-inflammatory activity ($58 \pm 5 \mu\text{M}$), being comparable
287 to the activity of the standard dexamethaxone ($40 \pm 4 \mu\text{M}$), followed by the glucuronated

288 derivatives of cinnamic (CA-GP) ($179 \pm 71 \mu\text{M}$) and *p*-hydroxybenzoic (HA-GP) ($1901 \pm$
289 $104 \mu\text{M}$) acids. Contrarily to the results verified for the anti-inflammatory activity of the
290 three parental molecules, the glucuronated derivative of *p*-coumaric acid exhibited the
291 strongest activity, presenting an acetylated glucuronide in the carboxylic group and an OH
292 group in the *para* position equal to the glucuronated derivative of *p*-hydroxybenzoic acid,
293 but, with a double bond in the middle increasing the lateral chain. Thus, the strongest activity
294 can be related with the double bound that stabilizes the molecule and the reactivity of the
295 benzenic ring to be able to inhibit the activity of COX. Regarding the methylated derivatives,
296 those of *p*-coumaric acid presented higher activity than the ones of cinnamic or *p*-
297 hydroxybenzoic acids; in particular, the methylated derivative CoA-M1, with an ester instead
298 of the carboxylic group, revealed very strong activity ($35 \pm 2 \mu\text{M}$), very close to the
299 dexamethaxone value ($40 \pm 4 \mu\text{M}$). Also in the case of methylated derivatives it seems that
300 the double bound of *p*-coumaric acid allows a higher ability to interact with the tested cell
301 line, while in the case of *p*-hydroxybenzoic acid that has no double bound, the ester and OH
302 groups are more close to each other and can chemically interact changing the molecule
303 stability and decreasing the bioactivity.

304 Comparing the activity of each parent molecule and the corresponding glucuronated and
305 methylated derivatives, the order was as follows: *p*-hydroxybenzoic acid: HA > HA-M3 >
306 HA-M2 > HA-M1 > HA-GP; *p*-coumaric acid: CoA-M1 > CoA-GP > CoA-M2 > CoA-M3 >
307 CoA; and cinnamic acid: CA-GP > CA > CA-M. *p*-Hydroxybenzoic acid showed higher
308 activity than the corresponding derivatives, with HA-M3 as the most active compound; the
309 decreased anti-inflammatory activity observed for HA-M1, HA-M2 and HA-GP could be
310 explained by the esterification of the carboxylic group. On the contrary, all *p*-coumaric acid
311 derivatives showed higher activity than the parent molecule (CoA), particularly the
312 methylated compound CoA-M1, suggesting that in this case the esterification of the

313 carboxylic group together with the free OH group in *para* position increased their anti-
314 inflammatory activity. The glucuronated derivative of cinnamic acid (CA-GP) maintained its
315 activity in comparison with the parent molecule (CA), while the methylated one (CA-M)
316 decreased it.

317 These results are in relatively good agreement with the ones previously reported by our
318 research group comparing the antimicrobial activity of *p*-hydroxybenzoic, *p*-coumaric and
319 cinnamic acids and their respective methylated and glucuronated derivatives. The
320 methylation of *p*-hydroxybenzoic and cinnamic acids decreased the antimicrobial activity,
321 while the methylation of *p*-coumaric acid increased it (Heleno et al., 2014a); moreover, the
322 glucuronidation of the parent molecules also decreased the antimicrobial activity, except for
323 HA-GP that showed higher antifungal activity against some pathogenic strains (Heleno et al.,
324 2013a).

325

326 **4. Conclusion**

327 Overall, the mushroom species: *P. ostreatus*, *M. procera*, *B. impolitus* and *A. bisporus*
328 revealed the strongest anti-inflammatory potential presenting the highest inhibition of NO
329 production. These mushroom species also revealed the highest concentration in cinnamic
330 acid, which was also the individual compound that presented the strongest anti-inflammatory
331 activity and, therefore, could play an important role in the observed activity. However, the
332 conjugation reactions occurring in the organism can change the chemical structure of
333 cinnamic and phenolic acids increasing or decreasing their *in vivo* anti-inflammatory activity.
334 The possible metabolites previously synthesised by the authors and tested in the present work
335 are still displaying activity, in some cases like CoA-GP and CoA-M1 higher than the parent
336 compound and very close to the activity exhibited by dexamethaxone used as anti-
337 inflammatory standard.

338

339 **Acknowledgements**

340 The authors are grateful to Fundação para a Ciência e a Tecnologia (FCT, Portugal) and
341 FEDER-COMPETE/QREN/EU for the financial support through the research projects PEst-
342 OE/AGR/UI0690/2011 and PEst-C/QUI/UI0686/2011. S.A. Heleno (BD/70304/2010) and
343 R.C. Calhella (BPD/68344/2010) also thank FCT, POPH-QREN and FSE. C. Santos-Buelga
344 is also thankful to the Spanish MINECO for financial support through the project BFU2012-
345 35228.

346

347 **References**

- 348 Alves, M.J., Ferreira, I.C.F.R., Dias, J., Teixeira, V., Martins, A., & Pintado, M.A. (2012).
349 Review on antimicrobial activity of mushroom (Basidiomycetes) extracts and isolated
350 compounds. *Planta Medica*, 78, 1707–1711.
- 351 Alves, M.J., Ferreira, I.C.F.R., Froufe, H.J.C., Abreu, R.M.V., Martins, A., & Pintado, M.
352 (2013). Antimicrobial activity of phenolic compounds identified in wild mushrooms, SAR
353 analysis and docking studies. *Journal of Applied Microbiology*, 115, 346–357.
- 354 Barros, L., Dueñas, M., Ferreira, I.C.F.R., Baptista, P., & Santos-Buelga, C. (2009). Phenolic
355 acids determination by HPLC-DAD-ESI/MS in sixteen different Portuguese wild mushrooms
356 species. *Food and Chemical Toxicology*, 47, 1076–1079.
- 357 Borchers, A.T., Krishnamurthy, A., Keen, C.L., Meyers, F.J., & Gershwin, M.E. (2008). The
358 immunobiology of mushrooms. *Experimental Biology and Medicine*, 233, 259–276.
- 359 Carocho, M., & Ferreira, I.C.F.R. (2013). The role of phenolic compounds in the fight against
360 cancer – A review. *Anti-Cancer Agents in Medicinal Chemistry*, 13, 1236–1258.
- 361 Cheung, L.M., Cheung, P.C.K., & Ooi, V.E.C. (2003). Antioxidant activity and total
362 phenolics of edible mushroom extracts. *Food Chemistry*, 81, 249–255.
- 363 Dugowson, C.E., & Gnanashanmugam, P. (2006). Nonsteroidal anti-inflammatory drugs.
364 *Physical Medicine and Rehabilitation Clinics of North America*, 17, 347–354.

365 Elsayed, A. E., Hesham, E.L.E., Mohammad, A.M.W., & Ramlan Aziz. (2014). Mushrooms:
366 A Potential Natural Source of Anti-Inflammatory Compounds for Medical Applications.
367 *Mediators of Inflammation*, 2014, 1-15.

368 Ferreira, I.C.F.R., Barros, L., & Abreu, R.M.V. (2009). Antioxidants in wild mushrooms.
369 *Current Medicinal Chemistry*, 16, 1543–1560.

370 Ferreira, I.C.F.R., Vaz, J.A., Vasconcelos, M.A., & Martins, A. (2010). Compounds from
371 wild mushrooms with antitumor potential. *Anti Cancer Agents in Medicinal Chemistry*, 10,
372 424–436.

373 García-Lafuente, A., Moro, C., Manchón, N., Gonzalo-Ruiz, A., Villares, A., Guillamón, E.,
374 Rostagno, M., & Mateo-Vivaracho, L. (2014). In vitro anti-inflammatory activity of phenolic
375 rich extracts from white and red common beans. *Food Chemistry*, 161, 216–223.

376 Grangeia, C., Heleno, S.A., Barros, L., Martins, A., & Ferreira, I.C.F.R. (2011). Effects of
377 trophism on nutritional and nutraceutical potential of wild edible mushrooms. *Food Research*
378 *International*, 44, 1029–1035.

379 Guillamón, E., García-Lafuente, A., Lozano, M., D'Arrigo, M., Rostagno, M.A., Villares, A.,
380 & Martinez, J.A. (2010). Edible mushrooms: role in the prevention of cardiovascular
381 diseases, *Fitoterapia*, 81, 715–723.

382 Gunawardena, D., Bennett, L., Shanmugam, K., King, K., Williams, R., Zabarás, D, Head R,
383 Ooi, L., Gyengesi, E., & Münch, G. (2014). Anti-inflammatory effects of five commercially
384 available mushroom species determined in lipopolysaccharide and interferon γ activated
385 murine macrophages. *Food Chemistry*, 148, 92–96.

386 Heleno, S.A., Barros, L., Sousa, M.J., Martins, A., Santos-Buelga, C., & Ferreira, I.C.F.R.
387 (2011). Targeted metabolites analysis in wild Boletus species. *Food Science and Technology*,
388 44, 1343-1348.

389 Heleno, S.A., Ferreira, I.C.F.R., Esteves, A.P., Cirić, A., Glamočlija, J., Martins, A., Soković,
390 & M., Queiroz, M.-J.R.P. (2013a). Antimicrobial and demelanizing activity of *Ganoderma*
391 *lucidum* extract, *p*-hydroxybenzoic and cinnamic acids and their synthetic acetylated
392 glucuronide methyl esters. *Food and Chemical Toxicology*, 58, 95–100.

393 Heleno, S.A, Stojković, D., Barros, L., Glamočlija, J., Soković, M., Martins, A., Queiroz, M.-
394 J.R.P., & Ferreira, I.C.F.R. (2013b). A comparative study of chemical composition,

395 antioxidant and antimicrobial properties of *Morchella esculenta* (L.) Pers. from Portugal and
396 Serbia. *Food Research International*, 51, 236-243.

397 Heleno, S.A., Ferreira, I.C.F.R., Cirić, A., Glamočlija, J., Martins, A., Queiroz, M.-J. R.P., &
398 Sokovic, M. (2014a). *Coprinopsis atramentaria* extract, organic acids, synthesized
399 glucuronated and methylated derivatives as antibacterial and antifungal agents. *Food &*
400 *Function*, 5, 2521–2528.

401 Heleno, S.A., Ferreira, I.C.F.R., Calhelha, R.C., Esteves, A.P., & Queiroz, M.-J.R.P. (2014b).
402 Cytotoxicity of *Coprinopsis atramentaria* extract, organic acids and their synthesized
403 methylated and glucuronate derivatives. *Food Research International*, 55, 170–175.

404 Heleno, S.A., Martins, A., Queiroz, M.-J.R.P., & Ferreira, I.C.F.R. (2015). Bioactivity of
405 phenolic acids: Metabolites versus parent compounds: A review. *Food Chemistry*, 173, 501–
406 513.

407 Hu, S.H., Wang, J.C., Lien, J.L., Liaw, E.T., & Lee, M.Y. (2006). Antihyperglycemic effect
408 of polysaccharide from fermented broth of *Pleurotus citrinopileatus*. *Applied Microbiology*
409 *and Biotechnology*, 70, 107–113.

410 Iwashina, T. (2000). The structure and distribution of the flavonoids in plants. *Journal of*
411 *Plant Research*, 13, 287-299.

412 Kalac, P. (2009). Chemical composition and nutritional value of European species of wild
413 growing mushrooms: a review. *Food Chemistry*, 113, 9–16.

414 Kanwar, J.R., Kanwar, R.K., Burrow, H., & Baratchi, S. (2009). Recent advances on the roles
415 of NO in cancer and chronic inflammatory disorders. *Current Medicinal Chemistry*, 16,
416 2373- 2394.

417 Kim, M.Y., Seguin, P., Ahn, J.K., Kim, J.J., Chun, S.C., Kim, E.H., Seo, S.H., Kang, E.Y.,
418 Kim, S.L., Park, Y.J., Ro, H.M., & Chung, I.M. (2008). Phenolic compound concentration
419 and antioxidant activities of edible and medicinal mushrooms from Korea. *Journal of*
420 *Agricultural and Food Chemistry*, 56, 7265–7270.

421 Lee, J.Y., Jang, Y.W., Kang, H.S., Moon, H., Sim, S.S., & Kim, C.J. (2006). Anti-
422 inflammatory Action of Phenolic Compounds from *Gastrodia elata* Root. *Archives of*
423 *Pharmacal Research*, 29, 849-858.

424 Lim, H.-W., Yoon, J.-H., Kim, Y.-S., Lee, M.-W., Park, S.-Y., & Choi, H.-K. (2007). Free
425 radical-scavenging and inhibition of nitric oxide production by four grades of pine mushroom
426 (*Tricholoma matsutake* Sing.). *Food Chemistry*, *103*, 1337–1342.

427 Lindequist, U., Niedermeyer, T.M.J., & Jülich, W.D. (2005). The pharmacological potential
428 of mushrooms. *Evidence-based Complementary and Alternative Medicine*, *2*, 285–299.

429 Liu, Y.W., Su, Y.W., Ong, W.K., Chen, Mei, C.H., C.C., Fan, T.H & Tsai, Y.C. (2014).
430 Inhibitory effects of *Pleurotus tuber-regium* mycelia and bioactive constituents on LPS-
431 treated RAW264.7 cells. *Journal of Functional Foods*, *7*, 662 – 670.

432 Ma, L., Chen, H., Dong, P., & Lu, X. (2013). Anti-inflammatory and anticancer activities of
433 extracts and compounds from the mushroom *Inonotus obliquus*. *Food Chemistry*, *139*, 503–
434 508.

435 Moradali, M.F, Mostafavi, H., Ghods, S., & Hedjaroude, G.-A. (2007). Immunomodulating
436 and anticancer agents in the realm of macromycetes fungi (macrofungi), *International*
437 *Immunopharmacology*, *7*, 701–724.

438 Mori, K., Kobayashi, C., Tomita, T., Inatomi, S., & Ikeda, M. (2008). Antiatherosclerotic
439 effect of the edible mushrooms *Pleurotus eryngii* (Eringi), *Grifola frondosa* (Maitake), and
440 *Hypsizygus marmoreus* (Bunashimeji) in apolipoprotein Edeficient mice. *Nutrition Research*,
441 *28*, 335–342.

442 Moro, C., Palacios, I., Lozano, M., D'Arrigo, M., Guillamón, E., Villares, A., et al. (2012).
443 Anti-inflammatory activity of methanolic extracts from edible mushrooms in LPS activated
444 RAW 264.7 macrophages. *Food Chemistry*, *130*, 350–355.

445 Nosálóva, V., Bobek, P., Cerna, S., Galbavy, S., & Tvrtina, G.S. (2001). Effects of pleuran
446 (β -glucan isolated from *Pleurotus ostreatus*) on experimental colitis in rats. *Physiological*
447 *Research*, *50*, 575–81.

448 Park, Y.M.J, Won, J.H, Kim, Y.H, Choi, J.W, Park, H.J, & Lee K.T. (2005). *In vivo* and *in*
449 *vitro* anti-inflammatory and anti-nociceptive effects of the methanol extract of *Inonotus*
450 *obliquus*. *Journal of Ethnopharmacology*, *101*, 120–128.

451 Pereira, E., Barros, L., Martins, A., & Ferreira, I.C.F.R., (2012). Towards chemical and
452 nutritional inventory of Portuguese wild edible mushrooms in different habitats. *Food*
453 *Chemistry*, *130*, 394–403.

- 454 Pradhan, A. (2007). Obesity, metabolic syndrome, and type 2 diabetes: Inflammatory basis of
455 glucose metabolic disorders. *Nutrition Reviews*, *65*, 152–156.
- 456 Padilha, M.M., Avila, A.A., Sousa, P.J., Cardoso, L.G., Perazzo, F.F., & Carvalho, J.C.
457 (2009). Anti-inflammatory activity of aqueous and alkaline extracts from mushrooms
458 (*Agaricus blazei* Murill). *Journal of Medicinal Food*, *12*, 359–364.
- 459 Puttaraju, N.G., Venkateshaiah, S.U., Dharmesh, S.M., Urs, S.M. & Somasundaram, R.
460 (2006). Antioxidant activity of indigenous edible mushrooms. *Journal of Agricultural and*
461 *Food Chemistry*, *54*, 9764–9772.
- 462 Queiroz, L.S, Nascimento, M.S, Cruz, A.K.M., Castro, A.J.G., Moura, M.F.V., Baseia, I.G.,
463 Araújo, R.M., Benevides, N.M.B., Lima, L.F.A., & Leite, E.L. (2010). Glucans from the
464 *Caripia montagnei* mushroom present anti-inflammatory activity. *International*
465 *Immunopharmacology*, *10*, 34–42.
- 466 Reis, F.S., Barros, L., Martins, A., & Ferreira, I.C.F.R. (2012). Chemical composition and
467 nutritional value of the most widely appreciated cultivated mushrooms: an inter-species
468 comparative study. *Food Chemical and Toxicology*, *50*, 191–197.
- 469 Reis, F.S., Heleno, S.A., Barros, L., Sousa, M.J., Martins, A., Santos-Buelga, C., & Ferreira
470 I.C.F.R. (2011). Toward the antioxidant and chemical characterization of mycorrhizal
471 mushrooms from northeast Portugal. *Journal of Food Science*, *76*, C824–C830.
- 472 Vaz, J.A., Barros, L., Martins, A., Moraise, J.S., Vasconcelos, M.H., & Ferreira, I.C.F.R.
473 (2011). Phenolic profile of seventeen Portuguese wild mushrooms. *LWT - Food Science and*
474 *Technology*, *44*, 343-346.
- 475 Tanaka, K.I., Suemasu, S., Ishihara, T., Tasaka, Y., Arai, Y., & Misushima, T. (2009).
476 Inhibition of both COX-1 and COX-2 and resulting decrease in the level of prostaglandins E2
477 is responsible for non-steroidal anti-inflammatory drug (NSAID)-dependent exacerbation of
478 colitis. *European Journal of Pharmacology*, *603*, 120–132.

Table 1. Some previous studies on anti-inflammatory activity of different mushroom species evaluated through NO assay.

Species	Country	Extract	Inhibition of NO production	References
<i>Agaricus bisporus</i>	Spain	Methanol	30% at 0.5 mg/mL	Moro et al., 2012
	Australia	Ethanol	50% at 0.032 mg/mL	Gunawardena et al., 2014
<i>Boletus edulis</i>	Spain	Methanol	10% at 0.5 mg/mL	Moro et al., 2012
<i>Cantharellus cibarius</i>	Spain	Methanol	70% at 0.5 mg/mL	Moro et al., 2012
<i>Caripia montagnei</i>	Brazil	Acetone 80%, methanol	43% at 10 mg/kg 54% at 30 mg/kg 49% at 50 mg/kg	Queiroz et al., 2010
<i>Cratarellus cornucopoides</i>	Spain	Methanol	55% at 0.5 mg/mL	Moro et al., 2012
<i>Flammulina velutipes</i>	Australia	Ethanol	50% at 0.024 mg/mL	Gunawardena et al., 2014
<i>Inonotus obliquus</i>	China	Ethanol	65% at 40 µg/mL	Ma et al., 2013
	South Korea	Ethanol	50% at 89 µg/mL	Park et al., 2005
<i>Lactarius deliciosus</i>	Spain	Methanol	40% at 0.5 mg/mL	Moro et al., 2012
<i>Lentinus edodes</i>	Australia	Ethanol	50% at 0.027 mg/mL	Gunawardena et al., 2014
<i>Pleurotus ostreatus</i>	Australia	Ethanol	50% at 0.077 mg/mL	Gunawardena et al., 2014
	Spain	Methanol	15% at 0.5 mg/mL	Moro et al., 2012
<i>Pleurotus tuber-regium</i>	Belgium	Ethanol	70% at 0.5 mg/mL	Liu et al., 2014
<i>Tricholoma matsutake</i>	South Korea	Dichloromethane	47% at 2 mg/mL	Lim et al., 2007

Table 2. Cinnamic and phenolic acids identified and quantified by HPLC-PDA in the ethanolic extracts of the analysed mushrooms.

Mushroom species	Extraction yield (%)	<i>p</i> -Hydroxybenzoic acid ($\mu\text{g/g}$)	<i>p</i> -Coumaric acid ($\mu\text{g/g}$)	Cinnamic acid ($\mu\text{g/g}$)
<i>Agaricus bisporus</i>	6.9 \pm 0.7 ^{abcd}	nd	nd	149 \pm 2 ^d
<i>Agaricus bisporus</i> Portobello	6.0 \pm 0.1 ^{bcd}	nd	nd	11 \pm 1 ^{hi}
<i>Amanita caesaria</i>	5.4 \pm 0.5 ^{cd}	57 \pm 3 ^e	nd	156 \pm 3 ^d
<i>Boletus aereus</i>	7.7 \pm 0.6 ^{abc}	43 \pm 1 ^f	74 \pm 1 ^b	50 \pm 3 ^f
<i>Boletus edulis</i>	10.4 \pm 0.5 ^a	nd	nd	14.2 \pm 0.4 ^{gh}
<i>Boletus flagrans</i>	4.9 \pm 0.3 ^{cd}	nd	nd	6.1 \pm 0.3 ⁱ
<i>Boletus impolitus</i>	4.1 \pm 0.5 ^d	125 \pm 9 ^c	45 \pm 2 ^c	505 \pm 12 ^c
<i>Boletus reticulatus</i>	9.5 \pm 0.3 ^{ab}	nd	nd	20.3 \pm 0.1 ^g
<i>Cantharellus cibarius</i>	4.0 \pm 0.3 ^d	151 \pm 2 ^b	nd	71 \pm 1 ^e
<i>Lactarius deliciosus</i>	3.7 \pm 0.4 ^d	108 \pm 5 ^d	nd	67 \pm 1 ^e
<i>Morchella esculenta</i>	4.0 \pm 0.8 ^d	nd	nd	71 \pm 3 ^e
<i>Macrolepiota procera</i>	3.7 \pm 0.4 ^d	nd	nd	522 \pm 1 ^b
<i>Pleurotus eryngii</i>	10.1 \pm 0.6 ^a	nd	nd	16 \pm 1 ^{gh}
<i>Pleurotus ostreatus</i>	4.0 \pm 0.6 ^d	297 \pm 5 ^a	171 \pm 1 ^a	619 \pm 3 ^a

nd- not detected. ANOVA In each column, different letters mean statistical significant differences ($p < 0.05$) between samples.

Table 3. Extract concentrations responsible for 50% of reduction of NO production (EC₅₀ values) in RAW 264.7 cell line.


Mushroom species	EC ₅₀ values (µg/mL)	Mushroom species	EC ₅₀ values (µg/mL)
<i>Agaricus bisporus</i>	190 ± 6 ^{ef}	<i>Boletus reticulatus</i>	378 ± 28 ^{ab}
<i>Agaricus bisporus</i> portobelo	>400 ^a	<i>Cantharellus cibarius</i>	202 ± 17 ^e
<i>Amanita caesaria</i>	186 ± 7 ^{ef}	<i>Lactarius deliciosus</i>	253 ± 14 ^d
<i>Boletus aereus</i>	357 ± 3 ^b	<i>Macrolepiota procera</i>	162 ± 2 ^g
<i>Boletus edulis</i>	>400 ^a	<i>Morchella esculenta</i>	287 ± 9 ^c
<i>Boletus flagrans</i>	>400 ^a	<i>Pleurotus eryngii</i>	388 ± 17 ^a
<i>Boletus impolitus</i>	166 ± 10 ^{fg}	<i>Pleurotus ostreatus</i>	96 ± 1 ^h

EC₅₀ values correspond to 50% of inhibition of the NO production in comparison with the negative control (100% of NO production). In the columns, different letters mean statistical significant differences (p<0.05) between samples. Dexamethaxone EC₅₀ value = 16 ± 2 µg/mL.

Table 4. Concentrations of the studied acids and their glucuronated and methylated derivatives responsible for 50% of reduction of NO production (EC₅₀ values, μM) in RAW 264.7 cell line.


<i>p</i> -Hydroxybenzoic acid and derivatives		<i>p</i> -Coumaric acid and derivatives		Cinnamic acid and derivatives	
HA	239 ± 29 ^c	CoA	442 ± 33 ^a	CA	182 ± 16 ^b
HA-GP	1901 ± 104 ^a	CoA-GP	58 ± 5 ^c	CA-GP	179 ± 17 ^b
HA-M1	1825 ± 120 ^a	CoA-M1	35 ± 2 ^c	CA-M	224 ± 16 ^a
HA-M2	526 ± 26 ^b	CoA-M2	128 ± 10 ^b		
HA-M3	509 ± 47 ^b	CoA-M3	129 ± 6 ^b		

EC₅₀ values correspond to 50% of inhibition of the NO production in comparison with the negative control (100% of NO production). In each column, different letters mean statistical significant differences (p<0.05) between compounds. Dexamethaxone EC₅₀ value = 40 ± 4 μM.




i) K_2CO_3 1.5 equivs, DMSO, Ar, RT, 24h;
ii) MeOH; H_2SO_4 , 5 days, R.T.

A



i) K_2CO_3 1.5 equivs, DMSO, Ar, RT, 24h;
ii) MeOH; H_2SO_4 , 5 days, R.T.
iii) K_2CO_3 , 2 equivs, acetone, 50°C, 24h;
iv) NaOH, 3 equivs, EtOH, 65°C, 3h

B



C

Figure 1. Synthesis of methylated and glucuronated derivatives of *p*-hydroxybenzoic, *p*-coumaric and cinnamic acids. **A**) i) Glucuronidation of cinnamic acid (CA). CA-GP- cinnamic acid glucuronide protected form, 2,3,4-tri-*O*-acetyl-1-cinnamoyl-*D*-glucuronic acid methyl ester (Heleno et al., 2013a); ii) Methylation of CA. CAM- methyl 3-phenylacrylate (Heleno et al., 2014b). **B**) i) Glucuronidation of *p*-coumaric acid (CoA). CoA-GP- *p*-Coumaric acid glucuronide protected form, 2,3,4-tri-*O*-acetyl-1-*p*-coumaroyl-*D*-glucuronic acid methyl ester (Heleno et al., 2014b); ii-iv) Methylations of CoA. CoA-M1- 3-(4-hydroxyphenyl) acrylate, CoA-M2- methyl-(4-methoxyphenyl) acrylate, CoA-M3- 3-(4-methoxyphenyl) acrylic acid (Heleno et al. 2014b). **C**) i) Glucuronidation of *p*-hydroxybenzoic acid (HA). HA-GP- *p*-hydroxybenzoic acid protected form, 2,3,4-tri-*O*-acetyl-1-*p*-hydroxybenzoyl-*D*-glucuronic acid methyl ester (Heleno et al., 2013a); ii-iv) Methylations of *p*-hydroxybenzoic acid, HA-M1- methyl 4-hydroxybenzoate, HA-M2- methyl-*p*-anisate, HA-M3- 4-methoxybenzoic acid (Heleno et al., 2014b).




Figure 2. Effects of the ethanolic mushroom extracts in the NO produced by RAW 264.7 cell line. Mean values and standard errors, expressed in relation to the negative control (without extract)- 100% of production. ■ 0 μg/mL; ■ 50 μg/mL; ■ 100 μg/mL; ■ 200 μg/mL; □ 400 μg/mL.