

University of Groningen

Genetics of Atrial Fibrillation in 2020 GWAS, Genome Sequencing, Polygenic Risk, and Beyond

Roselli, Carolina; Rienstra, Michiel; Ellinor, Patrick T.

Published in:
Circulation research

DOI:
[10.1161/CIRCRESAHA.120.316575](https://doi.org/10.1161/CIRCRESAHA.120.316575)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2020

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Roselli, C., Rienstra, M., & Ellinor, P. T. (2020). Genetics of Atrial Fibrillation in 2020 GWAS, Genome Sequencing, Polygenic Risk, and Beyond. *Circulation research*, 127(1), 21-33.
<https://doi.org/10.1161/CIRCRESAHA.120.316575>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

COMPENDIUM ON ATRIAL FIBRILLATION

Genetics of Atrial Fibrillation in 2020

GWAS, Genome Sequencing, Polygenic Risk, and Beyond

Carolina Roselli¹, Michiel Rienstra², Patrick T. Ellinor³

ABSTRACT: Atrial fibrillation is a common heart rhythm disorder that leads to an increased risk for stroke and heart failure. Atrial fibrillation is a complex disease with both environmental and genetic risk factors that contribute to the arrhythmia. Over the last decade, rapid progress has been made in identifying the genetic basis for this common condition. In this review, we provide an overview of the primary types of genetic analyses performed for atrial fibrillation, including linkage studies, genome-wide association studies, and studies of rare coding variation. With these results in mind, we aim to highlighting the existing knowledge gaps and future directions for atrial fibrillation genetics research.

Key Words: atrial fibrillation ■ exome ■ genetics ■ genome-wide association study ■ mutation

Atrial fibrillation (AF) is a common heart rhythm disorder with an estimated 33 million people affected worldwide.¹ Reported risk factors for AF include advancing age, obesity, hypertension, diabetes mellitus, and cardiovascular diseases.² Studies have also shown an increased risk for men to develop AF, compared with women.³ As discussed in the accompanying article in this Compendium,⁴ AF can lead to many serious medical consequences, including stroke, heart failure, cognitive impairment, and increased mortality.

The treatment of AF remains challenging. Although there are effective medications for anticoagulation to reduce the risk of stroke, antiarrhythmic medications are limited by a lack of efficacy to reduce symptoms and have potential side effects. Alternatively, catheter ablation procedures can be effective in reducing the burden of AF, but these procedures are invasive, can be associated with complications, and may require a repeat procedure for the long-term management of AF. Thus, there is a pressing need to develop new therapies for AF.

Similar to other common cardiovascular diseases, such as hypertension and myocardial infarction, AF is a complex disease with shared environmental and genetic factors that contribute to disease pathogenesis. Over the last decade, multiple studies have observed familial aggregation of individuals with lone AF.⁵ Similarly, the

heritability of AF has been elegantly demonstrated in the Icelandic population.⁶ Based on a study on monozygotic twins, the heritability of AF has been estimated as high as 62%, indicating a strong genetic component.⁷ In aggregate, these studies have consistently observed an increased risk of AF particularly when a first-degree family member is affected and among individuals with early-onset forms of arrhythmia.⁸ In more recent work, Weng et al⁹ used genetic data to estimate that the heritability of AF based on common genetic variants in individuals of European ancestry is $\approx 22\%$.

In the current review, we will provide an overview of multiple approaches used to examine the genetic basis of AF. We will present the most relevant results from these analyses and discuss emerging technological advances that could be leveraged to expand our understanding of the field. We discuss 3 broad genetic approaches applied to AF including (1) linkage analysis using families with Mendelian forms of AF, (2) genome-wide association studies (GWAS) using genotyping array data, and (3) coding variation from genome sequence data (Figure 1). These approaches are not mutually exclusive but are helpful as a framework to consider when reviewing the genetic studies of AF published to date. We will subsequently describe the application of GWAS data to clinical risk prediction. Finally, we will discuss the knowledge

Correspondence to: Patrick T. Ellinor, MD, PhD, Cardiovascular Disease Initiative, The Broad Institute of MIT and Harvard, Cambridge, MA 02142. Email ellinor@mgh.harvard.edu

For Sources of Funding and Disclosures, see page 31.

© 2020 American Heart Association, Inc.

Circulation Research is available at www.ahajournals.org/journal/res

Nonstandard Abbreviations and Acronyms

AF	atrial fibrillation
eQTL	expression quantitative trait locus
GWAS	genome-wide association studies
LOF	loss-of-function
PRS	polygenic risk score
SNP	single-nucleotide polymorphism

gaps in the field of AF genetics and describe emerging technologies that may shape the future of the field.

THREE BROAD GENETIC APPROACHES APPLIED TO AF

Familial AF and Linkage Analysis

Linkage analysis is typically performed in families with many affected individuals and a clear hereditary pattern. The analytic approach leverages genetic linkage or the tendency for a genetic marker near a disease-causing gene to be inherited together. The earliest application of this approach used a few hundred highly informative markers scattered throughout the genome. By matching transmission of the genetic marker with disease status in the family, a disease-causing region or locus can be linked to a given marker. In turn, the genes in this region can be sequenced to identify mutations that associate with disease. This approach has been elegantly used to identify many of the early genes implicated in hypertrophic cardiomyopathy and long-QT syndrome. As summarized in Figure 2, a number of causative mutations for AF have been identified in large families or populations. Specific examples include mutations in the ion channel *KCNQ1*, the cardiac peptide *NPPA*, the transcription factor *TBX5*, and a motor protein *MYL4*.

The first mutation linked to familial AF was found in the ion channel *KCNQ1*, a gene that encodes the α subunit of the I_{Ks} current. The AF-related mutations result in a gain of channel function and likely shortening of the atrial refractory period, a finding that would make it easier for reentry to continue and increase the susceptibility to AF.¹⁰ In a distinct family, a mutation was identified in *NPPA*, the gene encoding the atrial natriuretic peptide, a protein that is highly expressed in the heart.¹¹ A frameshift mutation was found to remove a stop codon and lead to an extended mutant protein that had appeared to be protected from degradation resulting in greater circulating levels and increased activity.¹² In vivo experiments from the same study have shown that mutations in *NPPA* lead to a shortening of the monophasic action potential duration as well as the effective refractory period. These changes on atrial electrophysiology could increase susceptibility to AF.

A gain-of-function mutation in the transcription factor, *TBX5*, was associated with familial AF in the setting of Holt-Oram syndrome,¹³ a developmental disorder that leads to heart and limb malformations.¹⁴ In vitro studies demonstrated that the mutated *TBX5* had enhanced binding to DNA and could lead to upregulation of downstream targets, such as *NPPA* and *CX40*.^{13,15} In a large-scale study of Icelanders, autosomal recessive mutations in *MYL4* were identified among individuals with early-onset AF.¹⁶ This atrial-specific myosin light chain was found to have a mutation in an F-actin binding region, and modeling the mutation in zebrafish resulted in disruption of the sarcomere and atrial enlargement.

Thus, individual families with hereditary forms of AF can be incredibly helpful in informing disease biology. However, the mutations identified to date are by nature rare and therefore have a small impact on the overall scope of this common arrhythmia.

Genome-Wide Association Studies for AF

In contrast to studies in families, GWAS permit the analysis of entire populations by comparing individuals with and without AF at a large scale. Each individual is genetically fingerprinted using a low cost, high throughput genotyping array. These arrays are used to determine the status of hundreds of thousands of genetic variants or single-nucleotide polymorphisms (SNPs) throughout the genome. While individual SNPs contain relatively little information, in aggregate, the use of hundreds of thousands of markers can capture the majority of the genetic diversity between individuals. To compare the data from one genotyping platform to another, the data are imputed or harmonized to a common reference panel consisting of millions of genetic markers.¹⁷ Comparisons of all the genetic markers are then made between cases and controls to identify regions associated with disease (Figure 1, middle). Importantly, unlike the analysis in families in which a single causative mutation is identified, in GWAS a region or locus is linked to disease. This region may or may not contain any genes. Many causative variants identified by GWAS are in noncoding regions of the genome and have an effect on the regulation of a nearby gene. In recent years, the cost of genotyping arrays has fallen to <\$50 per individual so it is now feasible undertake large-scale studies of common diseases and studies with >100 000 cases are increasingly common. Large sample sizes help to both ensure the validity of the results and to define the full extent of the genetic basis of the disease.

The first GWAS for AF was reported in 2007 and remarkably started with only a few hundred AF cases in the initial discovery.¹⁸ The small number of individuals needed in this first report was a reflection of the strength of the genetic association which is considerably greater than that observed for most other genome-wide




Figure 1. Three primary types of genetic analyses for atrial fibrillation (AF).

Linkage analyses primarily focus on large families with hereditary forms of AF. The disease-linked regions can include multiple candidate genes one of which will contain a disease-causing mutation. Genome-wide association studies (GWAS) analyses are based on genotype array data that consists largely of noncoding variants that are presumed to regulate genes in the region or locus. Analyses of coding variation are derived from whole-exome or whole-genome sequencing data. Rare coding or loss-of-function variants are grouped and jointly tested in AF cases vs controls to identify specific disease-causing genes. Please note that these approaches are not mutually exclusive and are often combined depending upon the study design. LOF indicates loss-of-function; SNP, single-nucleotide polymorphism; and STR, short tandem repeat.

studies. Carrying a single variant at this 4q25 locus near the gene *PITX2* conferred over a 60% increased risk of disease in the general population and an even greater risk in younger individuals. As with many disease-associated regions identified by GWAS, the association with AF at the *PITX2* locus is in a noncoding region of the

genome. As reviewed in other articles in this issue,¹⁹ the AF risk variants modulate the expression of *PITX2* and loss of this region or the *PITX2* gene itself can result in AF.^{20,21} In the ensuing years, many other GWAS for AF have been completed in predominantly European ancestry participants.²²⁻²⁶ Initially novel loci were identified at

	<i>KCNQ1</i>	<i>NPPA</i>	<i>TBX5</i>	<i>MYL4</i>	<i>TTN</i>
EVIDENCE					
Familial AF gene	●	●	●	●	●
Coding variant			●		●
GWAS locus			●		●
Gene-based test			●		●
Functional study	●	●	●	●	

Figure 2. Major atrial fibrillation (AF)-associated genes and lines of evidence.

The figure illustrates AF-associated genes that were discovered through family based or gene-based studies. For each gene the lines of evidence are listed. The table includes the evidence from familial AF genetic analysis, whether coding variants in the gene are associated with AF, if the gene lies within an AF genome-wide association studies (GWAS) locus, whether loss-of-function variation is associated with AF, and functional evidence that has been reported for the gene in the context of AF. TF indicates transcription factor.

16q22 close to the gene *ZFHX3* and 1q21 close to the *KCNN3*.^{27–29} In 2 recent studies that both included over 60 000 cases,^{30,31} nearly 140 AF loci have been identified to date (Table 1). Although rapid progress has been made in AF genetics, it is important to note that the vast majority of participants are of European descent (Figure 3). The 2 largest non-European GWAS were published in 2017 and report results from Korean³² and Japanese³³ cohorts. At least 3 variants for AF show significant heterogeneity across different ancestries (rs2129977 at *PITX2*, rs11598047 at *NEURL*, and rs2359171 at *ZFHX3*). Additionally, while a signal at the *PITX2* locus can be found across ancestries, the top variant is not always the same nor strongly correlated.³⁰ Two AF loci from a GWAS in Japanese ancestry, at the genes *NEBL* and *SH3PXD2A*, do not replicate in European ancestry and may be ancestry specific.³³

Identifying Genes at AF Loci

Genetic variants identified in GWAS studies are largely located in noncoding regions of the genome. These noncoding variants are presumed to alter the activity of a transcriptional regulatory element such as an enhancer or repressor that, in turn, results in modifying the transcription of a nearby gene. Importantly, for most GWAS variants there is usually no straightforward path from an association by GWAS to a gene and a disease mechanism. A recent study used STARR-seq (self-transcribing active regulatory region sequencing) to identify regulatory elements and their target genes at multiple GWAS loci for AF. Furthermore, they found that the loss of a regulatory element at the *HCN4* locus led to reduced gene expression.³⁴ A myriad of analyses can follow a large-scale GWAS ranging from computational analyses to the derivation of polygenic risk scores (PRSs) for AF risk prediction and the incorporation of GWAS data into other analyses (Table 2). In the subsequent sections, we will touch on a few helpful approaches with respect to AF, but for other potential directions please also see these recent reviews.^{47–49} One straightforward application of GWAS data is to perform a pathway analysis to evaluate the collective effect of the genetic association on different biological functions. Globally the AF-associated genes represent distinct functional groups, including those underlying cardiac development, cellular electrophysiology, cardiomyocyte contractility, and structure.^{26,30,31} A similar approach was taken in a GWAS of the Japanese population and implicated suppression of the mTOR (mammalian target of rapamycin) signaling pathway in AF.⁵⁰

As noted above, a major challenge with GWAS is that the analyses usually identify a region of interest rather than a specific causative gene. Bridging the gap from variants to genes remains a major challenge in disease genetics, particularly for complex polygenic traits such as AF. One common and helpful approach is to use expression

quantitative trait loci (eQTL) mapping. An eQTL analysis links the genotype of a SNP at an AF locus to the expression of genes in the region. If an AF-associated SNP is strongly linked to the expression of a single gene that gene is likely to be the causative gene at the locus. While this approach is simple in theory, practically there are 2 primary limitations. First, eQTL analyses are often tissue specific. Although there are terrific publicly available resources such as GTEx or the Genotype-Tissue Expression project (<https://gtexportal.org>) that includes gene expression profiles from many tissues, unfortunately the cardiac analyses were limited to the left ventricle and right atrial appendage.¹¹ Recent work in the Cleveland Clinic Atrial Tissue Bank^{24,51} and the Myocardial Applied Genomics Network³⁰ have addressed this limitation by investigating the gene expression profiles of left atrial tissue. A second limitation of using eQTLs is that it requires large sample sizes and this is not always possible when tissues are hard to obtain such as the left atrium or the pulmonary veins in the case of AF. Finally, while eQTLs are powerful, they only explain a fraction of the disease loci. For example, in the latest 2 GWAS of AF only 13% of the variants at AF loci could be linked to the expression of a single gene, and at 22% of AF loci the variants linked to one or more genes. These results are summarized in Table 1.

A complimentary approach to eQTL analyses is to use the 3-dimensional architecture of the genome to identify causative AF genes. Since many AF SNPs are in noncoding regions of the genome, they are presumed to alter regulatory elements such as enhancers or repressors that, in turn, bind to the promoter of a nearby gene to regulate its expression. The contact points between AF-associated regulatory elements and gene promoters can be assessed using chromosome conformation capture technologies. For example, Hi-C is a genome-wide chromosome conformation capture technique that allows the unbiased detection of chromatin interactions across the entire genome. Similar to eQTL analyses, enhancer-promoter contacts can be tissue specific and a recent Hi-C study from the human heart roughly doubled the number of AF-associated genes derived from GWAS data.⁵² Other analyses of AF GWAS data such as the incorporation of multi-omic data sets⁵³ or epigenetic analyses including STARR-seq⁵⁴ are described in detail in the accompanying review by van Ouwkerk and colleagues.⁵⁵ Ultimately, although any gene implicated by these methods will require further validation *in vitro* and *in vivo*.

Assessing Polygenic Risk From GWAS

Since we now have very dense GWAS data sets for AF, it is natural to wonder whether this data could be used in a clinical setting to identify high risk individuals, stratify screening efforts or look for differential treatment outcomes. The polygenic nature of AF as captured in GWAS can be transformed into a genetic risk score for each individual. An AF PRS summarizes the cumulative

Table 1. To Date, There Are at Least 138 AF Loci Identified in Single Variant Testing With $P < 5 \times 10^{-8}$

Rsid	Nearest Gene(s) or eGene*	Rsid	Nearest Gene(s) or eGene*	Rsid	Nearest Gene(s) or eGene*
rs187585530	UBE4B	rs716845	KCNN2	rs1822273	NAV2
rs880315	CASZ1	rs2012809	FBN2, SLC27A6	rs949078	SORL1, MIR100HG
rs7529220	HSPG2, CELA3B	rs34750263	WNT8A, NME5	rs76097649	KCNJ5
rs2885697	SCMH1*	rs174048	ARHGAP26, NR3C1	rs6490029	CUX2
rs11590635	AGBL4	rs12188351	SLIT3	rs10842383	LINC00477, BCAT1
rs56202902	FAF1	rs6882776	NKX2-5	rs113819537	SSPN*
rs146518726	C1orf185	rs73366713	ATXN1	rs12809354	PKP2
rs12044963	KCND3	rs34969716	KDM1B	rs7978685	NACA
rs4484922	CASQ2*	rs1307274	C6orf1, NUDT3	rs35349325	BEST3
rs79187193	GJA5	rs3176326	CDKN1A	rs11180703	KRR1, PHLDA1
rs11264280	KCNN3, PMVK	rs6907805	CGA, ZNF292	rs883079	TBX5
rs72700114	METTL11B, LINC01142	rs210632	GOPC	rs12810346	TBX5-AS1, TBX3
rs608930	GORAB, PRRX1	rs17079881	SLC35F1	rs10773657	HIP1R
rs10753933	PPFIA4*	rs13191450	GJA1, HSF2	rs12298484	DNAH10
rs4951261	NUCKS1	rs12208899	LINC00326, EYA4	rs6560886	FBRSL1
rs6546620	KIF3C	rs117984853	UST	rs9580438	LINC00540, BASP1P1
rs6742276	XPO1	rs11768850	SUN1	rs35569628	CUL4A
rs2540949	CEP68*	rs55734480	DGKB	rs28631169	MYH7
rs10165883	SNRNP27	rs6462078	CREB5	rs2145587	AKAP6
rs72926475	REEP1, KDM3A	rs74910854	PMS2P2*	rs73241997	SNX6, CFL2
rs28387148	GYPC*	rs11773884	CDK6	rs2738413	SYNE2
rs67969609	TEX41	rs62483627	COG5	rs74884082	DPF3
rs12992412	MBD5	rs11773845	CAV1	rs10873299	LRRC74, IRF2BPL
rs56181519	WIPF1*	rs55985730	OPN1SW	rs147301839	MYZAP
rs2288327	FKBP7*	rs7789146	KCNH2	rs62011291	USP3
rs3820888	SPATS2L*	rs35620480	LINC00208, GATA4	rs12591736	TLE3, UACA
rs35544454	ERBB4	rs7508	ASAH1*	rs74022964	HCN4, REC114
rs6810325	MKRN2*	rs7846485	XPO7	rs12908004	LINC00927, ARNT2
rs73032363	THRB	rs62521286	FBXO32	rs12908437	IGF1R*
rs6790396	SCN10A	rs35006907	MTSS1, LINC00964	rs2286466	RPL3L*
rs34080181	SLC25A26*	rs7460121	MIR30B	rs2359171	ZFHX3
rs17005647	FRMD4B	rs6993266	PTK2	rs7225165	YWHAE, CRK
rs7632427	EPHA3	rs4977397	SLC24A2, MLLT3	rs8073937	POLR2A, TNFSF12
rs17490701	PHLDB2	rs4385527	C9orf3	rs72811294	MYOCD
rs1278493	PPP2R3A	rs4743034	ZNF462	rs11658278	ZPBP2
rs4855075	GNB4	rs10760361	PSMB7	rs242557	MAPT
rs60902112	XXYL1	rs2274115	LHX3	rs76774446	GOSR2
rs9872035	PAK2	rs2296610	NEBL	rs7219869	KCNJ2, CASC17
rs3822259	WDR1	rs7919685	NRBF2*	rs12604076	CYTH1
rs1458038	PRDM8, FGF5	rs7096385	SIRT1	rs9953366	SMAD7
rs3960788	UBE2D3*	rs60212594	SYNPO2L	rs8088085	MEX3C
rs2129977	PITX2, C4orf32	rs11001667	C10orf11	rs2145274	CASC20, BMP2
rs55754224	CAMK2D	rs1044258	C10orf76	rs7269123	C20orf166
rs10213171	ARHGAP10	rs11598047	NEURL	rs2834618	LOC100506385
rs10520260	HAND2-AS1*	rs2047036	SH3PXD2A	rs465276	TUBA8
rs6596717	LOC102467213, EFNA5	rs10749053	RBM20	rs133902	MYO18B

The table above includes the sentinel variant at each locus from AF GWAS publications. The listed genes are either the nearest gene(s) or the eGene. The eGene is listed when it is the only eGene at that locus reported by Roselli et al.³⁰ or Nielsen et al.³¹ Notably, only 18 out of 138 variants are associated with only one eGene. AF indicates atrial fibrillation; eQTL, expression quantitative trait locus; and GWAS, genome-wide association studies.

*The eGene is defined as a gene with an eQTL to the variant at an AF GWAS locus.




Figure 3. Ancestry of the cases in genome-wide association studies (GWAS) for atrial fibrillation (AF).

European ancestry sample is plotted towards the left in white and non-European ancestry is plotted towards the right highlighted in different colors. Plotted is the number of cases included in each published AF GWAS study or meta-analysis. Eleven out of 13 studies include predominantly European ancestry samples, shown in white. Two studies are Japanese only and Korean only. Within the multi-ancestry meta-analyses Roselli et al³⁰ included the largest proportion of non-European cases, including Japanese, Brazilian, African American, and Hispanic samples. AFR indicates African-American; BRA, Brazilian; EUR, European; HISP, Hispanic; JAP, Japanese; and KOR, Korean.

genetic risk and can be computed using anywhere from just a few variants at the top loci or the data from millions of SNPs across the entire genome (Figure 4).

An initial AF PRS used the top 12 genome-wide significant genetic variants,⁵⁶ while more contemporary iterations incorporate over 6.6 million variants.⁵⁷ In the latter approach, the top 1.5% individuals with a high PRS had a more than a 4-fold increased risk for AF. The variants and weights included in the latest AF PRS are publicly available (<http://www.broadcvdi.org>). In one interesting application of an AF PRS, Weng et al⁵⁸ found that individuals in the highest tertile of polygenic risk had a higher lifetime risk for AF (47%) compared with the individuals in the lowest tertile (26%). The combination of clinical and genetic data permitted additional refinement of risk. For example, individuals with a low clinical risk but a high genetic risk had an overall lifetime risk for AF of 44%. A complementary review in this Compendium focuses on the potential clinical applications of AF genetics and PRSs.⁵⁹

Exome and Genome Sequencing to Identify AF Genes

While GWAS has been deployed at great scale, a major limitation of the method is that it is only able to capture

known variants in the genome. Yet many diseases arise from unique mutations in the coding region of genes that emerge spontaneously in an individual of family. Although these mutations can be identified using a targeted approach of a single or small number of genes, the availability of genome-wide approaches has largely replaced these earlier studies. Currently, it is possible to sequence the entire protein-coding region of the genome for about \$200 or the entire genome for <\$800, and prices are continuing to fall. These technological advancements and price reductions have started to enable the application of large-scale sequencing studies for AF.

Once a population of cases and controls has been sequenced, common genetic variants are analyzed using the approach previously described for GWAS. In contrast, rare coding genetic markers, often defined as present in <1% of the population, are analyzed in gene-based tests. For a gene-based test, coding variants are analyzed jointly across a gene unit for an association with disease, as depicted in Figure 1. The most commonly used approach restricts the analysis to variants predicted to lead to a loss-of-function (LOF) of the encoded protein. A considerable advantage of testing rare LOF variants over GWAS is establishing a direct link from gene function to disease. In addition, this analysis provides a clear direction

Table 2. Overview of Studies That Integrated AF GWAS Data With Other Clinical, Epigenetic or Genetic Data Sets

Study Type	Description and Reference
Mendelian randomization	Obesity ³⁵
	Thyroid function ³⁶
	Serum albumin ³⁷
	Body mass index/body composition and cardiovascular conditions including AF ³⁸
	Body composition ³⁹
Methylation	Genome-wide DNA methylation analysis ⁴⁰
	Methylome-Wide Association Study ⁴¹
Heritability	Heritability of AF ⁹
Ancestry-specific analyses	European ancestry as a risk factor for AF ⁴²
AF recurrence after ablation	Common variants predict AF recurrence ⁴³
Genetic interaction	Gene-gene interactions ⁴⁴
	Genetic interactions with risk factors ⁴⁵
Associations with mitochondrial DNA	Mitochondrial DNA and incident AF ⁴⁶

AF indicates atrial fibrillation; and GWAS, genome-wide association studies.

of effect between LOF of the encoded protein and disease. Although many sequencing analyses focus on LOF variants, it is important to note that gain-of-function mutations such as those identified in families with *TBX5* or *KCNQ1* would not be identified with this approach.

In 2017, Thorolfsdottir et al⁶⁰ analyzed whole-genome sequencing data from the Icelandic population including 14 255 AF cases and 374 939 controls. The Icelandic population provides a unique resource for genetic discovery, because it is a relatively homogeneous and genetically isolated population that is enriched with rare LOF coding variants. The study identified a low-frequency missense mutation in the gene *PLEC* that encodes the cytoskeletal protein plectin. The missense mutation in *PLEC* is associated with increased risk for AF. Additionally, a missense mutation in the myosin gene *MYH6* was significantly associated with AF, a gene that has previously been associated with sick sinus syndrome.⁶¹

Within the past 2 years, at least 4 studies have identified LOF mutations in the gene *TTN* among individuals with AF. Ahlberg et al⁶² found an enrichment of LOF mutations in *TTN* among families and individuals in Denmark with early-onset AF. Shortly thereafter, Choi et al⁶³ observed a similar finding among unrelated individuals with early-onset AF. Using nearly 2800 AF cases and 5000 controls, they found that *TTN* LOF mutations were present in $\approx 2\%$ of individuals with an onset of AF before 65 years of age. With younger ages of onset, the frequency of *TTN* mutations rose to a high of over 7% of individuals younger than 30 years of age. As with dilated cardiomyopathy, the association between *TTN* and AF had a stronger effect when the analysis was restricted to *TTN* exons that were highly expressed in cardiac tissue.

Although these studies pointed to an increased frequency of *TTN* mutations in individuals with early-onset

AF, the question then arose whether *TTN* mutations were detectable in the general population with AF. To address this, Choi et al⁶⁴ used exome sequencing data in 1400 AF cases and >40 000 controls from the UK Biobank. They observed a similar strong association between LOF variation in *TTN* and AF.⁶⁴ Furthermore, the penetrance of AF among *TTN* mutation carriers was markedly higher among individuals with an increased AF polygenic risk.

FUTURE DIRECTIONS IN AF GENETICS

While rapid progress has been made in our understanding of the genetic basis of AF over the last decade, it is important to realize that we are currently in the midst of an explosive growth in the scale of genetic data available worldwide. In the following sections, we have sought to put these emerging resources in the context of future potential studies in AF genetic research (Figure 5). While there are many other potential avenues of exploration, hopefully this will serve as a broad framework for the reader.

Expanding Genetic Studies in Non-Europeans

As poignantly illustrated in Figure 3, the vast majority of genetic analyses for AF have been performed in individuals of European descent. The historical tendency to focus on European populations has led to a nonrepresentative distribution of ancestries in genetic studies compared with the real-world diversity. As we move forward, it will be critical to expand our genetic resources throughout the world, not only for AF but for all common diseases. Expanding beyond Europeans has 2 primary advantages. First, it is clear that there are unique lessons that can be learned about common diseases among different races and ethnicities. For example, in a large Japanese GWAS for AF, only 85% of the top hits overlapped with the results from individuals of European ancestry. Second, as we think about applying genetic risk scores to clinical care, it will be important that we do not blindly apply a score developed in Europeans to other races and ethnicities. Such an approach may further exacerbate health care disparities.⁶⁵ Multiple programs in the United States are trying to address some of the disparity in genetic research including the NHLBI TOPMed Program (<https://www.nhlbiwgs.org>), Million Veteran Program (<https://www.mvp.va.gov>), and All of Us (<https://www.allofus.nih.gov>). The All of Us project is an ongoing longitudinal collection of over 1 million individuals and is being intentionally structured to ensure ethnic and racial diversity. The expansion of biobanks throughout the world including in China (<https://www.ckbiobank.org>), India⁶⁶ and Africa (<https://h3africa.org>) will also be essential to this effort.

Sequencing, Sequencing, and More Sequencing

An exciting development in AF genetics has been the continued expansion of the availability of exome and




Figure 4. Overview of polygenic risk scores (PRS) for atrial fibrillation (AF).

A PRS is calculated for each individual as a sum of the product of genetic dosage and a weight. The weights are derived from the effect estimates of a genome-wide association study. The PRS of individuals in a population follows a gaussian distribution. Individuals in the highest percentile of the distributions show an increased risk for AF vs the remaining population. Potential applications of an AF PRS can include improving risk prediction, prioritizing high-risk individuals for screening, and examining differential outcomes of AF.

genome sequencing. As discussed above, the major advantage of sequencing over GWAS is that sequencing directly implicates the gene as causally related to AF. There are multiple international efforts that will coalesce to enhance our understanding of common disease genetics. These include ongoing work in the UK Biobank, deCODE, Estonia, Finland, Japanese Biobank, Million Veterans Program, the NHGRI CCDG, and NHLBI TOPMed Programs among many others. As a result, within the next 2 years, we can anticipate having data sets for AF consisting of >50 000 cases with either exome or genome sequencing. The power of such large data sets was nicely illustrated for autism where many additional disease-causing genes were identified with the enhanced power of these large-scale studies.⁶⁷

The utility in harnessing sequence data for AF can perhaps be exemplified best by the investigators at deCODE genetics.⁶⁸ The Icelandic population is relatively homogenous and can be traced back to a small set of common ancestors, making it one of the few bottleneck populations present in the current day world. The resulting enrichment for rare mutations can

manifest in naturally occurring genetic knockouts in humans.⁶⁹ The uniqueness of the population structure combined with extensive whole-genome sequencing has already led to the identification of multiple new AF genes. Similarly, in the coming years, sequencing of other bottleneck populations such as Finland⁷⁰ and Sardinia⁷¹ will be a valuable resource for studying common diseases including AF.

Structural Variation From Whole-Genome Sequencing

Beyond gene-based tests, it is also clear that current whole-genome sequencing data sets have not been used to their fullest potential. Structural variants, such as inversions, duplications, translocations, as well as large deletions and insertions may be associated with AF. It has been estimated that structural variants can have a larger than expected impact on the genomic differences between individuals.⁷² Identification of these structural modifications from the raw sequencing data is more challenging than identifying single




Figure 5. Future directions in atrial fibrillation (AF) genetics.

Overview of emerging technologies and analyses that could shape the field of AF genetics for the next decade. Large-scale rare coding sequence data: With dropping sequencing costs large-scale exome sequencing data sets will become available and accelerate the detection of rare and ultra rare coding variation that associate with AF. Structural genetic variation: Whole-genome sequencing data allows the detection of structural variation such as inversions, translocations, and large insertions and deletions. Methods to detect structural variation are improving and could lead to uncovering novel structural variant contributions to AF. Polygenic risk for diverse ethnicities: Increasing the contribution of non-European samples in AF genome-wide association studies (GWAS) will improve the polygenic risk prediction for diverse ethnicities. Functional cellular knockout assays: Gene knockout studies in relevant cell types, such as atrial cardiomyocytes, will enable the evaluation of AF candidate genes from GWAS loci in the context of functionally relevant readouts. Single-cell RNA sequencing: Next generation sequencing technologies such as the transcriptional profiling of individual cells from cardiac tissue will transform AF genetics and increase the resolution of gene expression profiles to a cell-type-specific level. Cell-type-specific expression quantitative trait loci could resolve the causal gene at AF GWAS loci. Machine learning on big data: Machine learning can facilitate the integration of big data sources such as gene expression profiles, proteomics data, protein-protein interaction networks, methylation data, regulatory regions, and spatial organization of the DNA. Machine learning algorithms will support the goal to identify causal genes for AF, resolve regulatory mechanisms at AF GWAS loci and uncover patterns that imply disease mechanisms of AF. LOF indicates loss-of-function.

variants, short insertions, or short deletions. It requires data reprocessing and the application of specialized algorithms to the sequencing data. To date, no single method can identify every type of structural variant with high confidence.⁷³ This suggests that a combination of methods may generate the highest yield. Structural variants can have a large impact on the function of genes and have been associated with diseases such as cancer⁷⁴ and complex diseases, including Crohn disease, rheumatoid arthritis, and diabetes mellitus.⁷⁵ We can anticipate that the systematic assessment of structural variation will identify novel genetic mechanisms for AF.

Expanding Gene Expression and Epigenetic Analyses to Tissues Relevant to AF

To move from associations to mechanisms for the AF GWAS loci, it will be essential to expand our repertoire of disease relevant eQTL and epigenetic data sets. Currently, there are modestly sized transcriptional data sets

from the human left atrium and very limited data from pulmonary venous tissues. Expanding these data from the 100 to 200 available samples by an order of magnitude will dramatically increase the availability of eQTLs that link AF disease variants to causative genes.

It will also be essential to move beyond the analyses of bulk tissues and to focus on the analyses of single-cells for transcriptional and epigenetic profiling,⁷⁶ techniques which have only rarely been applied to AF to date. Recent work by Tucker et al⁷⁷ in which they performed single nucleus RNA sequencing of the healthy human heart provides an example of the benefit from this approach. In a study of nearly 280 000 single nuclei from the 4 chambers of the human heart, a number of findings emerged that are relevant to AF. They were able to identify at least 10 major cell types in the heart, chamber specific transcription in nonmyocyte populations, and an cardiomyocyte enrichment of the genes at AF GWAS loci.⁷⁷

A logical extension of this work will be to compare the single-cell transcriptional and epigenetic profiles of left

atrial and pulmonary vein samples from AF cases versus controls. These results would also enable the discovery of changes of the cellular compositions, transcription, and cell-type-specific eQTLs in AF versus healthy individuals.

Developing Large-Scale Functional Screens for AF Genes

Even with additional expression and epigenetic data sets, it will be critical to expand our throughput for the functional assessment of the genes at the AF GWAS loci. The current state of our AF GWAS results can illustrate the scope of the problem. At present, there are close to 140 genetic loci for AF and within these loci there are >1000 genes or transcripts. Of these many loci, only a minority have a single gene that can be convincingly linked by eQTL or Hi-C analyses. This disconnect between our expansive knowledge of disease-associated variants and limited understanding of the mechanisms is not unique to AF but is present for essentially all common diseases.

To help address this challenge, the International Common Disease Alliance was founded in 2019. This partnership between academia, governments, pharmaceutical, and technology companies is a collaborative initiative with the goal to improve diagnosis, prevention, and treatment of disease through accelerating research that focuses on translating genetic findings into disease biology. While a lofty goal, the implementation of high throughput functional studies to elucidate the missing link between noncoding genetic variants, causal gene, and gene function was identified as one of the key priorities by this effort (<https://www.icda.bio>).

How could such functional screens be implemented for AF? It will clearly be impossible to characterize more than a small number of mouse or even zebrafish models and as a result, we will have to turn to cell-based models as an intermediate step. While cell-based models have many potential limitations, the scale of screening is the primary strength of this approach. Combining stem cell-derived cardiomyocytes and CRISPR-Cas9 technology will facilitate high throughput gene knockout studies for cellular assays. Potential cell readouts could include electrophysiological measurements of calcium signaling or action potential duration,^{78,79} structural assessments of sarcomere assembly,⁸⁰ contractility,⁸¹ and transcription.^{82,83} Given that current AF GWAS loci represent a cross section of transcription factors, ion channels and sarcomeric proteins, cell-based screening will likely require multiple readouts.

It will also be important that more consideration is given to the development of cell models that more fully recapitulate the diverse etiologies of AF. For example, there are at least 10 major cell types in the human atria and the study of myocytes will not be helpful if an AF gene is predominantly expressed in another cell type such as fibroblasts⁸⁴ or in a cell subtype arising from the conduction system or pulmonary veins.

Implementing Machine Learning to Identify Endophenotypes of AF

The rapid evolution in machine learning methods has already started to transform the medical field. The availability of multi-dimensional data sets ranging from electronic health records, imaging data, clinical measurements, and genetics is providing the basis for algorithm-based clinical research that has the power to improve risk prediction, response to treatment and clinical diagnostics for AF. Particularly exciting recent work has applied machine learning to the ECG to identify subclinical markers or endophenotypes that predict the future development of AF.^{85,86} In the future, it will be interesting to study the genetics of these AF endophenotypes and to use machine learning to enhance the prediction of LOF impact of coding variants.⁸⁷ Furthermore, it is clear that novel computational approaches will be required to integrate the existing and emerging data sources in AF genetics. Predicting the most likely causal genes at an AF GWAS locus or identify gene networks relevant for AF will require the integration of large data sources. The combination of GWAS results with gene expression profiles, proteomics data, protein-protein interaction networks, methylation data, regulatory regions, and spatial organization of the DNA requires efficient computational solutions that can deal with multi-dimensional data. For a broader discussion on this topic please see the accompanying review in this Compendium.⁸⁸

CONCLUSIONS

AF is a complex disease with a combination of both environmental and genetic factors that contribute to the pathogenesis of the arrhythmia. Rapid progress has been made in identifying many common variant loci in GWAS for AF, yet major challenges remain in moving from disease associations to specific mechanisms. Recent genome and exome-based sequencing studies have identified *TTN* as the gene most commonly associated with mutations in individuals with AF. Future studies will seek to explore the application of PRSs to clinical care, build out genetic studies in non-European populations, and further expand single-cell sequencing and omic technologies in cells and tissues relevant to AF. Further refinement of the genetic basis of AF will ultimately facilitate the identification of new therapeutic targets and enable more precise risk stratification for this common arrhythmia.

ARTICLE INFORMATION

Affiliations

From the Cardiovascular Disease Initiative, Broad Institute of MIT and Harvard, Cambridge, MA (C.R., P.T.E.); Department of Cardiology, University of Groningen, University Medical Center Groningen, the Netherlands (C.R., M.R.); and Cardiovascular Research Center and Cardiac Arrhythmia Service, Massachusetts General Hospital, Boston (P.T.E.).

Sources of Funding

Dr Ellinor is supported by the National Institutes of Health (1R01HL092577, R01HL128914, and K24HL105780), the American Heart Association (18SFRN34110082), and by the Foundation Leducq (14CVD01).

Disclosures

Dr Ellinor is supported by a grant from Bayer AG to the Broad Institute focused on the genetics and therapeutics of cardiovascular disease. Dr Ellinor has consulted for Bayer AG, Novartis, MyoKardia, and Quest Diagnostics. C. Roselli is supported by a grant from Bayer AG to the Broad Institute focused on the development of therapeutics for cardiovascular disease. The other author reports no conflicts.

REFERENCES

- Chugh SS, Havmoeller R, Narayanan K, Singh D, Rienstra M, Benjamin EJ, Gillum RF, Kim YH, McAnulty JH Jr, Zheng ZJ, et al. Worldwide epidemiology of atrial fibrillation: a global burden of disease 2010 Study. *Circulation*. 2014;129:837–847. doi: 10.1161/CIRCULATIONAHA.113.005119
- Huxley RR, Lopez FL, Folsom AR, Agarwal SK, Loefer LR, Soliman EZ, Maclehorse R, Konety S, Alonso A. Absolute and attributable risks of atrial fibrillation in relation to optimal and borderline risk factors: the Atherosclerosis Risk in Communities (ARIC) study. *Circulation*. 2011;123:1501–1508. doi: 10.1161/CIRCULATIONAHA.110.009035
- Magnussen C, Niiranen TJ, Ojeda FM, Gianfagna F, Blankenberg S, Njølstad I, Vartiainen E, Sans S, Pasterkamp G, Hughes M, et al. Sex differences and similarities in atrial fibrillation epidemiology, risk factors, and mortality in community cohorts. *Circulation*. 2017;136:1588–1597. doi: 10.1161/CIRCULATIONAHA.117.028981
- Kornej J, Börschel CS, Benjamin EJ, Schnabel RB. Epidemiology of atrial fibrillation in the 21st century: novel methods and new insights. *Circ Res*. 2020;127:4–20. doi: 10.1161/CIRCRESAHA.120.316340
- Ellinor PT, Yoerger DM, Ruskin JN, MacRae CA. Familial aggregation in lone atrial fibrillation. *Hum Genet*. 2005;118:179–184. doi: 10.1007/s00439-005-0034-8
- Arnar DO, Thorvaldsson S, Manolio TA, Thorgeirsson G, Kristjansson K, Hakonarson H, Stefansson K. Familial aggregation of atrial fibrillation in Iceland. *Eur Heart J*. 2006;27:708–712. doi: 10.1093/eurheartj/ehi727
- Christophersen IE, Ravn LS, Budtz-Joergensen E, Skytthe A, Haunsoe S, Svendsen JH, Christensen K. Familial aggregation of atrial fibrillation: a study in danish twins. *Circ Arrhythm Electrophysiol*. 2009;2:378–383. doi: 10.1161/CIRCEP.108.786665
- Alzahrani Z, Ornelas-Loredo A, Darbar SD, Farooqui A, Mol D, Chalazan B, Villagrana NE, McCauley M, Lazar S, Wissner E, et al. Association between family history and early-onset atrial fibrillation across racial and Ethnic Groups. *JAMA Netw Open*. 2018;1:e182497. doi: 10.1001/jamanetworkopen.2018.2497
- Weng LC, Choi SH, Klarin D, Smith JG, Loh PR, Chaffin M, Roselli C, Hulme OL, Lunetta KL, Dupuis J, et al. Heritability of atrial fibrillation. *Circ Cardiovasc Genet*. 2017;10:e001838. doi: 10.1161/CIRCGENETICS.117.001838
- Chen YH, Xu SJ, Bendahhou S, Wang XL, Wang Y, Xu WY, Jin HW, Sun H, Su XY, Zhuang QN, et al. KCNQ1 gain-of-function mutation in familial atrial fibrillation. *Science*. 2003;299:251–254. doi: 10.1126/science.1077771
- Lonsdale J, Thomas J, Salvatore M, Phillips R, Lo E, Shad S, Hasz R, Walters G, Garcia F, Young N, et al; GTEx Consortium. The Genotype-Tissue Expression (GTEx) project. *Nat Genet*. 2013;45:580–585. doi: 10.1038/ng.2653
- Hodgson-Zingman DM, Karst ML, Zingman LV, Heublein DM, Darbar D, Herron KJ, Ballew JD, de Andrade M, Burnett JC Jr, Olson TM. Atrial natriuretic peptide frameshift mutation in familial atrial fibrillation. *N Engl J Med*. 2008;359:158–165. doi: 10.1056/NEJMoa0706300
- Postma AV, van de Meerakker JB, Mathijssen IB, Barnett P, Christoffels VM, Ilgun A, Lam J, Wilde AA, Lekanke Deprez RH, Moorman AF. A gain-of-function TBX5 mutation is associated with atypical holt-oram syndrome and paroxysmal atrial fibrillation. *Circ Res*. 2008;102:1433–1442. doi: 10.1161/CIRCRESAHA.107.168294
- Holt M, Oram S. Familial heart disease with skeletal malformations. *Br Heart J*. 1960;22:236–242. doi: 10.1136/hrt.22.2.236
- Ma JF, Yang F, Mahida SN, Zhao L, Chen X, Zhang ML, Sun Z, Yao Y, Zhang YX, Zheng GY, et al. TBX5 mutations contribute to early-onset atrial fibrillation in Chinese and Caucasians. *Cardiovasc Res*. 2016;109:442–450. doi: 10.1093/cvr/cvw003
- Orr N, Arnaout R, Gula LJ, Spears DA, Leong-Sit P, Li O, Tarhuni W, Reischauer S, Chauhan VS, Borkovich M, et al. A mutation in the atrial-specific myosin light chain gene (MYL4) causes familial atrial fibrillation. *Nat Commun*. 2016;7:11303. doi: 10.1038/ncomms11303
- Kowalski MH, Qian H, Hou Z, Rosen JD, Tapia AL, Shan Y, Jain D, Argos M, Arnett DK, Avery C, et al. Use of >100,000 NHLBI Trans-Omics for Precision Medicine (TOPMed) consortium whole genome sequences improves imputation quality and detection of rare variant associations in admixed African and Hispanic/Latino populations. *PLOS Genet*. 2019;15:e1008500. doi: 10.1371/journal.pgen.1008500
- Gudbjartsson DF, Arnar DO, Helgadóttir A, Gretarsdóttir S, Holm H, Sigurdsson A, Jonasdóttir A, Baker A, Thorleifsson G, Kristjansson K, et al. Variants conferring risk of atrial fibrillation on chromosome 4q25. *Nature*. 2007;448:353–357. doi: 10.1038/nature06007
- Nattel S, Heijman J, Zhou L, Dobrev D. Molecular basis of atrial fibrillation pathophysiology and therapy: a translational perspective. *Circ Res*. 2020;127:51–72. doi: 10.1161/CIRCRESAHA.120.316363
- Kirchhof P, Kahr PC, Kaese S, Piccini I, Vokshi I, Scheld HH, Rotering H, Fortmueller L, Laakmann S, Verheule S, et al. PITX2c is expressed in the adult left atrium, and reducing Pitx2c expression promotes atrial fibrillation inducibility and complex changes in gene expression. *Circ Cardiovasc Genet*. 2011;4:123–133. doi: 10.1161/CIRCGENETICS.110.958058
- Syeda F, Kirchhof P, Fabritz L. PITX2-dependent gene regulation in atrial fibrillation and rhythm control. *J Physiol*. 2017;595:4019–4026. doi: 10.1113/JP273123
- Sinner MF, Tucker NR, Lunetta KL, Ozaki K, Smith JG, Trompet S, Bis JC, Lin H, Chung MK, Nielsen JB, et al; METASTROKE Consortium; AFGen Consortium. Integrating genetic, transcriptional, and functional analyses to identify 5 novel genes for atrial fibrillation. *Circulation*. 2014;130:1225–1235. doi: 10.1161/CIRCULATIONAHA.114.009892
- Ellinor PT, Lunetta KL, Albert CM, Glazer NL, Ritchie MD, Smith AV, Arking DE, Müller-Nurasyid M, Krijthe BP, Lubitz SA, et al. Meta-analysis identifies six new susceptibility loci for atrial fibrillation. *Nat Genet*. 2012;44:670–675. doi: 10.1038/ng.2261
- Christophersen IE, Rienstra M, Roselli C, Yin X, Geelhoed B, Barnard J, Lin H, Arking DE, Smith AV, Albert CM, et al. Large-scale analyses of common and rare variants identify 12 new loci associated with atrial fibrillation. *Nat Genet*. 2017;49:946–952. doi: 10.1038/ng.3843
- Thorólfsson RB, Sveinbjörnsson G, Sulem P, Nielsen JB, Jonsson S, Halldorsson GH, Melsted P, Ivarsdóttir EV, Davidsson OB, Kristjansson RP, et al. Coding variants in RPL3L and MYZAP increase risk of atrial fibrillation. *Commun Biol*. 2018;1:68. doi: 10.1038/s42003-018-0068-9
- Nielsen JB, Fritsche LG, Zhou W, Teslovich TM, Holmen OL, Gustafsson S, Gabrielsen ME, Schmidt EM, Beaumont R, Wolford BN, et al. Genome-wide Study of atrial fibrillation identifies seven risk loci and highlights biological pathways and regulatory elements involved in cardiac development. *Am J Hum Genet*. 2018;102:103–115. doi: 10.1016/j.ajhg.2017.12.003
- Gudbjartsson DF, Holm H, Gretarsdóttir S, Thorleifsson G, Walters GB, Thorgeirsson G, Gulcher J, Mathiesen EB, Njølstad I, Nyrnes A, et al. A sequence variant in ZFXH3 on 16q22 associates with atrial fibrillation and ischemic stroke. *Nat Genet*. 2009;41:876–878. doi: 10.1038/ng.417
- Benjamin EJ, Rice KM, Arking DE, Pfeufer A, van Noord C, Smith AV, Schnabel RB, Bis JC, Boerwinkle E, Sinner MF, et al. Variants in ZFXH3 are associated with atrial fibrillation in individuals of European ancestry. *Nat Genet*. 2009;41:879–881. doi: 10.1038/ng.416
- Ellinor PT, Lunetta KL, Glazer NL, Pfeufer A, Alonso A, Chung MK, Sinner MF, de Bakker PI, Mueller M, Lubitz SA, et al. Common variants in KCNN3 are associated with lone atrial fibrillation. *Nat Genet*. 2010;42:240–244. doi: 10.1038/ng.537
- Roselli C, Chaffin MD, Weng LC, Aeschbacher S, Ahlberg G, Albert CM, Almgren P, Alonso A, Anderson CD, Aragam KG, et al. Multi-ethnic genome-wide association study for atrial fibrillation. *Nat Genet*. 2018;50:1225–1233. doi: 10.1038/s41588-018-0133-9
- Nielsen JB, Thorólfsson RB, Fritsche LG, Zhou W, Skov MW, Graham SE, Herron TJ, McCarthy S, Schmidt EM, Sveinbjörnsson G, et al. Biobank-driven genomic discovery yields new insight into atrial fibrillation biology. *Nat Genet*. 2018;50:1234–1239. doi: 10.1038/s41588-018-0171-3
- Lee JY, Kim TH, Yang PS, Lim HE, Choi EK, Shim J, Shin E, Uhm JS, Kim JS, Joung B, et al. Korean atrial fibrillation network genome-wide association study for early-onset atrial fibrillation identifies novel susceptibility loci. *Eur Heart J*. 2017;38:2586–2594. doi: 10.1093/eurheartj/ehx213
- Low SK, Takahashi A, Ebana Y, Ozaki K, Christophersen IE, Ellinor PT, Ogishima S, Yamamoto M, Satoh M, Sasaki M, et al; AFGen Consortium. Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. *Nat Genet*. 2017;49:953–958. doi: 10.1038/ng.3842

34. van Ouwkerk AF, Bosada F, Liu J, Zhang J, van Duijvenboden K, Chaffin M, Tucker N, Pijnappels DA, Ellinor PT, Barnett P, et al. Identification of functional variant enhancers associated with atrial fibrillation [published online April 6, 2020]. *Circ Res*. doi: 10.1161/CIRCRESAHA.119.316006
35. Chatterjee NA, Giulianini F, Geelhoed B, Lunetta KL, Misialek JR, Niemeijer MN, Rienstra M, Rose LM, Smith AV, Arking DE, et al. Genetic obesity and the risk of atrial fibrillation: causal estimates from mendelian randomization. *Circulation*. 2017;135:741–754. doi: 10.1161/CIRCULATIONAHA.116.024921
36. Ellervik C, Roselli C, Christophersen IE, Alonso A, Pietzner M, Sitlani CM, Trompet S, Arking DE, Geelhoed B, Guo X, et al. Assessment of the relationship between genetic determinants of thyroid function and atrial fibrillation: a Mendelian Randomization Study. *JAMA Cardiol*. 2019;4:144–152. doi: 10.1001/jamacardio.2018.4635
37. Liao L, Zhang S, Li W, Liu Y, Li J, Zhuang X, Liao XX. Serum albumin and atrial fibrillation: insights from epidemiological and mendelian randomization studies. *Eur J Epidemiol*. 2019;35:113–122. doi: 10.1007/s10654-019-00583-6
38. Larsson SC, Bäck M, Rees JMB, Mason AM, Burgess S. Body mass index and body composition in relation to 14 cardiovascular conditions in UK Biobank: a Mendelian randomization study. *Eur Heart J*. 2020;41:221–226. doi: 10.1093/eurheartj/ehz388
39. Tikkanen E, Gustafsson S, Knowles JW, Perez M, Burgess S, Ingelsson E. Body composition and atrial fibrillation: a Mendelian randomization study. *Eur Heart J*. 2019;40:1277–1282. doi: 10.1093/eurheartj/ehz003
40. Zhao G, Zhou J, Gao J, Liu Y, Gu S, Zhang X, Su P. Genome-wide DNA methylation analysis in permanent atrial fibrillation. *Mol Med Rep*. 2017;16:5505–5514. doi: 10.3892/mmr.2017.7221
41. Lin H, Yin X, Xie Z, Lunetta KL, Lubitz SA, Larson MG, Ko D, Magnani JW, Mendelson MM, Liu C, et al. Methylome-wide Association Study of atrial fibrillation in Framingham Heart Study. *Sci Rep*. 2017;7:40377. doi: 10.1038/srep40377
42. Marcus GM, Alonso A, Peralta CA, Lettrec G, Vittinghoff E, Lubitz SA, Fox ER, Levitzky YS, Mehra R, Kerr KF, et al; Candidate-Gene Association Resource (CARE) Study. European ancestry as a risk factor for atrial fibrillation in African Americans. *Circulation*. 2010;122:2009–2015. doi: 10.1161/CIRCULATIONAHA.110.958306
43. Zhao LQ, Zhang GB, Wen ZJ, Huang CK, Wu HQ, Xu J, Qi BZ, Wang ZM, Shi YY, Liu SW. Common variants predict recurrence after nonfamilial atrial fibrillation ablation in Chinese Han population. *Int J Cardiol*. 2017;227:360–366. doi: 10.1016/j.ijcard.2016.11.057
44. Lin H, Mueller-Nurasyid M, Smith AV, Arking DE, Barnard J, Bartz TM, Lunetta KL, Lohman K, Kleber ME, Lubitz SA, et al. Gene-gene interaction analyses for atrial fibrillation. *Sci Rep*. 2016;6:35371. doi: 10.1038/srep35371
45. Weng LC, Lunetta KL, Müller-Nurasyid M, Smith AV, Thériault S, Weeke PE, Barnard J, Bis JC, Lyytikäinen LP, Kleber ME, et al. Genetic interactions with age, sex, body mass index, and hypertension in relation to atrial fibrillation: the AFGen consortium. *Sci Rep*. 2017;7:11303. doi: 10.1038/s41598-017-09396-7
46. Zhao D, Bartz TM, Sotoodehnia N, Post WS, Heckbert SR, Alonso A, Longchamps RJ, Castellani CA, Hong YS, Rotter JI, et al. Mitochondrial DNA copy number and incident atrial fibrillation [published online November 20, 2019]. *BioRxiv*. doi: 10.1101/848085
47. Gallagher MD, Chen-Plotkin AS. The post-GWAS era: from association to function. *Am J Hum Genet*. 2018;102:717–730. doi: 10.1016/j.ajhg.2018.04.002
48. Broekema RV, Bakker OB, Jonkers IH. A practical view of fine-mapping and gene prioritization in the post-genome-wide association era. *Open Biol*. 2020;10:190221. doi: 10.1098/rsob.190221
49. Tucker NR, Clauss S, Ellinor PT. Common variation in atrial fibrillation: navigating the path from genetic association to mechanism. *Cardiovasc Res*. 2016;109:493–501. doi: 10.1093/cvr/cvv283
50. Ebana Y, Sun Y, Yang X, Watanabe T, Makita S, Ozaki K, Tanaka T, Arai H, Furukawa T. Pathway analysis with genome-wide association study (GWAS) data detected the association of atrial fibrillation with the mTOR signaling pathway. *Int J Cardiol Heart Vasc*. 2019;24:100383. doi: 10.1016/j.ijcha.2019.100383
51. Deshmukh A, Barnard J, Sun H, Newton D, Castel L, Pettersson G, Johnston D, Roselli E, Gillinov AM, McCurry K, et al. Left atrial transcriptional changes associated with atrial fibrillation susceptibility and persistence. *Circ Arrhythm Electrophysiol*. 2015;8:32–41. doi: 10.1161/CIRCEP.114.001632
52. Bianchi V, Geeven G, Tucker N, Hilvering CRE, Hall AW, Roselli C, Hill MC, Martin JF, Margulies KB, Ellinor PT, et al. Detailed regulatory interaction map of the human heart facilitates gene discovery for cardiovascular disease [published online July 17, 2019]. *BioRxiv*. doi: 10.1101/705715. <https://www.biorxiv.org/content/10.1101/705715v1>.
53. Wang B, Lunetta KL, Dupuis J, Lubitz SA, Trinquart L, Yao L, Ellinor PT, Benjamin EJ, Lin H. Integrative omics approach to identifying genes associated with atrial fibrillation. *Circ Res*. 2020;126:350–360. doi: 10.1161/CIRCRESAHA.119.315179
54. van Ouwkerk AF, Bosada FM, van Duijvenboden K, Hill MC, Montefiori LE, Scholman KT, Liu J, de Vries AAF, Boukens BJ, Ellinor PT, et al. Identification of atrial fibrillation associated genes and functional non-coding variants. *Nat Commun*. 2019;10:4755. doi: 10.1038/s41467-019-12721-5
55. van Ouwkerk AF, Hall AW, Kadow ZA, Lazarevic S, Reyat JS, Tucker NR, Nadador RD, Bosada FM, Bianchi V, Ellinor PT, et al. Epigenetic and transcriptional networks underlying atrial fibrillation. *Circ Res*. 2020;127:34–50. doi: 10.1161/CIRCRESAHA.120.316574
56. Lubitz SA, Lunetta KL, Lin H, Arking DE, Trompet S, Li G, Krijthe BP, Chasman DJ, Barnard J, Kleber ME, et al. Novel genetic markers associate with atrial fibrillation risk in Europeans and Japanese. *J Am Coll Cardiol*. 2014;63:1200–1210. doi: 10.1016/j.jacc.2013.12.015
57. Khera AV, Chaffin M, Aragam KG, Haas ME, Roselli C, Choi SH, Natarajan P, Lander SA, Lubitz SA, Ellinor PT, et al. Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations. *Nat Genet*. 2018;50:1219–1224. doi: 10.1038/s41588-018-0183-z
58. Weng LC, Preis SR, Hulme OL, Larson MG, Choi SH, Wang B, Trinquart L, McManus DD, Staerk L, Lin H, et al. Genetic predisposition, clinical risk factor burden, and lifetime risk of atrial fibrillation. *Circulation*. 2018;137:1027–1038. doi: 10.1161/CIRCULATIONAHA.117.031431
59. Shoemaker MB, Shah RL, Roden DM, Perez MV. How will genetics inform the clinical care of atrial fibrillation? *Circ Res*. 2020;127:111–127. doi: 10.1161/CIRCRESAHA.120.316365
60. Thorolfsdottir RB, Sveinbjornsson G, Sulem P, Helgadóttir A, Gretarsdóttir S, Benonisdóttir S, Magnúsdóttir A, Davidsson OB, Rajamani S, Roden DM, et al. A missense variant in PLEC increases risk of atrial fibrillation. *J Am Coll Cardiol*. 2017;70:2157–2168. doi: 10.1016/j.jacc.2017.09.005
61. Holm H, Gudbjartsson DF, Sulem P, Masson G, Helgadóttir HT, Zanon C, Magnússon OT, Helgason A, Saemundsdóttir J, Gylfason A, et al. A rare variant in MYH6 is associated with high risk of sick sinus syndrome. *Nat Genet*. 2011;43:316–320. doi: 10.1038/ng.781
62. Ahlberg G, Refsgaard L, Lundegaard PR, Andreassen L, Ranthe MF, Linscheid N, Nielsen JB, Melbye M, Haunsoe S, Sajadiet A, et al. Rare truncating variants in the sarcomeric protein titin associate with familial and early-onset atrial fibrillation. *Nat Commun*. 2018;9:4316. doi: 10.1038/s41467-018-06618-y
63. Choi SH, Weng LC, Roselli C, Lin H, Haggerty CM, Shoemaker MB, Barnard J, Arking DE, Chasman DJ, Albert CM, et al; DiscovEHR study and the NHLBI Trans-Omics for Precision Medicine (TOPMed) Consortium. Association between titin loss-of-function variants and early-onset atrial fibrillation. *JAMA*. 2018;320:2354–2364. doi: 10.1001/jama.2018.18179
64. Choi SH, Jurgens SJ, Weng LC, Pirruccello JP, Roselli C, Chaffin M, Lee CJ, Hall AW, Khera AV, Lunetta KL, et al. Monogenic and polygenic contributions to atrial fibrillation risk: results from a national biobank. *Circ Res*. 2020;126:200–209. doi: 10.1161/CIRCRESAHA.119.315686
65. Duncan L, Shen H, Gelaye B, Meijssen J, Ressler K, Feldman M, Peterson R, Domingue B. Analysis of polygenic risk score usage and performance in diverse human populations. *Nat Commun*. 2019;10:3328. doi: 10.1038/s41467-019-11112-0
66. Chatterjee S, Majumder PP. Kalyani cohort - the first platform in Eastern India for longitudinal studies on health and disease parameters in peri-urban setting. *Glob Health Epidemiol Genom*. 2017;2:e2. doi: 10.1017/ghg.2016.19
67. Satterstrom FK, Kosmicki JA, Wang J, Breen MS, De Rubeis S, An JY, Peng M, Collins R, Grove J, Klei L, et al; Autism Sequencing Consortium; iPSYCH-Broad Consortium. Large-Scale Exome Sequencing Study implicates both developmental and functional changes in the neurobiology of autism. *Cell*. 2020;180:568–584.e23. doi: 10.1016/j.cell.2019.12.036
68. Gudbjartsson DF, Helgason H, Gudjonsson SA, Zink F, Oddson A, Gylfason A, Besenbacher S, Magnusson G, Halldórsson BV, Hjartarson E, et al. Large-scale whole-genome sequencing of the Icelandic population. *Nat Genet*. 2015;47:435–444. doi: 10.1038/ng.3247
69. Chheda H, Palta P, Pirinen M, McCarthy S, Walter K, Koskinen S, Salomaa V, Daly M, Durbin R, Palotie A, et al. Whole-genome view of the consequences of a population bottleneck using 2926 genome sequences from Finland and United Kingdom. *Eur J Hum Genet*. 2017;25:477–484. doi: 10.1038/ejhg.2016.205

70. Lim ET, Würtz P, Havulinna AS, Palta P, Tukiainen T, Rehnström K, Esko T, Mägi R, Inouye M, Lappalainen T, et al; Sequencing Initiative Suomi (SISu) Project. Distribution and medical impact of loss-of-function variants in the Finnish founder population. *PLoS Genet*. 2014;10:e1004494. doi: 10.1371/journal.pgen.1004494
71. Sidore C, Busonero F, Maschio A, Porcu E, Naitza S, Zoledziwska M, Mulas A, Pistis G, Steri M, Danjou F, et al. Genome sequencing elucidates sardinian genetic architecture and augments association analyses for lipid and blood inflammatory markers. *Nat Genet*. 2015;47:1272–1281. doi: 10.1038/ng.3368
72. Lafrate AJ, Feuk L, Rivera MN, Listewnik ML, Donahoe PK, Qi Y, Scherer SW, Lee C. Detection of large-scale variation in the human genome. *Nat Genet*. 2004;36:949–951. doi: 10.1038/ng1416
73. Kosugi S, Momozawa Y, Liu X, Terao C, Kubo M, Kamatani Y. Comprehensive evaluation of structural variation detection algorithms for whole genome sequencing. *Genome Biol*. 2019;20:117. doi: 10.1186/s13059-019-1720-5
74. Nik-Zainal S, Davies H, Staaf J, Ramakrishna M, Glodzik D, Zou X, Martincorena I, Alexandrov LB, Martin S, Wedge DC, et al. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature*. 2016;534:47–54. doi: 10.1038/nature17676
75. Craddock N, Hurles ME, Cardin N, Pearson RD, Plagnol V, Robson S, Vukcevic D, Barnes C, Conrad DF, Giannoulata E, et al; Wellcome Trust Case Control Consortium. Genome-wide association study of CNVs in 16,000 cases of eight common diseases and 3,000 shared controls. *Nature*. 2010;464:713–720. doi: 10.1038/nature08979
76. Method of the year 2019: single-cell multimodal omics. *Nat Methods*. 2020;17:1. doi: 10.1038/s41592-019-0703-5
77. Tucker NR, Chaffin M, Fleming SJ, Hall AW, Parsons VA, Bedi K, Akkad AD, Herndon CN, Arduini A, Papangeli I, et al. Transcriptional and cellular diversity of the human heart [published online May 14, 2020]. *Circulation*. doi: 10.1161/CIRCULATIONAHA.119.045401
78. Leyton-Mange JS, Mills RW, Macri VS, Jang MY, Butte FN, Ellinor PT, Milan DJ. Rapid cellular phenotyping of human pluripotent stem cell-derived cardiomyocytes using a genetically encoded fluorescent voltage sensor. *Stem Cell Reports*. 2014;2:163–170. doi: 10.1016/j.stemcr.2014.01.003
79. Bedut S, Seminatore-Nole C, Lamamy V, Caignard S, Boutin JA, Nosjean O, Stephan JP, Coge F. High-throughput drug profiling with voltage- and calcium-sensitive fluorescent probes in human iPSC-derived cardiomyocytes. *Am J Physiol Heart Circ Physiol*. 2016;311:H44–H53. doi: 10.1152/ajpheart.00793.2015
80. Bray MA, Singh S, Han H, Davis CT, Borgeson B, Hartland C, Kost-Alimova M, Gustafsdottir SM, Gibson CC, Carpenter AE. Cell Painting, a high-content image-based assay for morphological profiling using multiplexed fluorescent dyes. *Nat Protoc*. 2016;11:1757–1774. doi: 10.1038/nprot.2016.105
81. Toepfer CN, Sharma A, Cicconet M, Garfinkel AC, Mücke M, Neyazi M, Willcox JAL, Agarwal R, Schmid M, Rao J, et al. SarcTrack. *Circ Res*. 2019;124:1172–1183. doi: 10.1161/CIRCRESAHA.118.314505
82. Subramanian A, Narayan R, Corsello SM, Peck DD, Natoli TE, Lu X, Gould J, Davis JF, Tubelli AA, Asiedu JK, et al. A next generation connectivity map: L1000 platform and the first 1,000,000 profiles. *Cell*. 2017;171:1437–1452.e17. doi: 10.1016/j.cell.2017.10.049
83. Dixit A, Parnas O, Li B, Chen J, Fulco CP, Jerby-Arnon L, Marjanovic ND, Dionne D, Burks T, Raychowdhury R, et al. Perturb-seq: dissecting molecular circuits with scalable single-cell RNA profiling of pooled genetic screens. *Cell*. 2016;167:1853–1866.e17. doi: 10.1016/j.cell.2016.11.038
84. Palano G, Jansson M, Backmark A, Martinsson S, Sabirsh A, Hultenby K, Åkerblad P, Granberg KL, Jennbacken K, Müllers E, et al. A high-content, in vitro cardiac fibrosis assay for high-throughput, phenotypic identification of compounds with anti-fibrotic activity. *J Mol Cell Cardiol*. 2020;142:105–117. doi: 10.1016/j.yjmcc.2020.04.002
85. Attia ZI, Kapa S, Lopez-Jimenez F, McKie PM, Ladewig DJ, Satam G, Pellikka PA, Enriquez-Sarano M, Noseworthy PA, Munger TM, et al. Screening for cardiac contractile dysfunction using an artificial intelligence-enabled electrocardiogram. *Nat Med*. 2019;25:70–74. doi: 10.1038/s41591-018-0240-2
86. Raghunath S, Pfeifer JM, Cerna AEU, Nemani A, Carbonati T, Jing L, vanMaanen D, McCarty BE, Hartzel DN, Ruhl JA, et al. Deep neural networks can predict incident atrial fibrillation from the 12-lead electrocardiogram and may help prevent associated strokes [published online April 27, 2020]. *MedRxiv*. doi: 10.1101/2020.04.23.20067967. <https://www.medrxiv.org/content/10.1101/2020.04.23.20067967v1>.
87. Pagel KA, Pejaver V, Lin GN, Nam HJ, Mort M, Cooper DN, Sebat J, Iakoucheva LM, Mooney SD, Radivojac P. When loss-of-function is loss of function: assessing mutational signatures and impact of loss-of-function genetic variants. *Bioinformatics*. 2017;33:i389–i398. doi: 10.1093/bioinformatics/btx272
88. Siontis KC, Yao X, Pirruccello JP, Philippakis AA, Noseworthy PA. How will machine learning inform the clinical care of atrial fibrillation? *Circ Res*. 2020;127:155–169. doi: 10.1161/CIRCRESAHA.120.316401