



UNIVERSITY OF LEEDS

This is a repository copy of *Toxoplasma gondii* major surface antigen (SAG1): *in vitro* analysis of host cell binding.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/1245/>

---

**Article:**

Robinson, S.A., Smith, J.E. and Millner, P.A. (2004) *Toxoplasma gondii* major surface antigen (SAG1): *in vitro* analysis of host cell binding. *Parasitology*, 128 (4). pp. 391-396. ISSN 0031-1820

<https://doi.org/10.1017/S0031182003004736>

---

**Reuse**

See Attached

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

# *Toxoplasma gondii* major surface antigen (SAG1): *in vitro* analysis of host cell binding

S. A. ROBINSON<sup>1</sup>, J. E. SMITH<sup>2</sup> and P. A. MILLNER<sup>1\*</sup>

<sup>1</sup>*School of Biochemistry and Molecular Biology, University of Leeds, Leeds LS2 9JT, UK*

<sup>2</sup>*School of Biology, University of Leeds, Leeds LS2 9JT, UK*

(Received 1 August 2003; revised 9 October 2003; accepted 23 October 2003)

## SUMMARY

Previous studies have indicated that SAG1, the major surface molecule of the protozoan parasite *Toxoplasma gondii*, is an important attachment ligand for the host cell. However, the research data that supports this claim comes largely from studies investigating tachyzoite binding, and not SAG1 binding *per se*. In this study we successfully developed an *in vitro* attachment assay to directly evaluate the mechanism of SAG1-host cell binding. Competition experiments were then performed using SAG1 that had been pre-treated with the neoglycoprotein BSA-glucosamide or with antibody. Soluble BSA-glucosamide blocked SAG1 attachment to MDBK cells in a dose-dependent manner, implying that SAG1 binding is mediated, in part, via attachment to host cell surface glucosamine. Interestingly, pre-incubation of SAG1 in polyclonal sera from chronically infected mice failed to block binding. This challenges the assumption that anti-SAG1 antibodies block parasite attachment through the masking of SAG1 host cell binding domains. Taken together, this evidence presents new strategies for understanding SAG1-mediated attachment.

Key words: *Toxoplasma gondii*, attachment, SAG1, glucosamide, antibody.

## INTRODUCTION

*Toxoplasma gondii* is a globally prevalent protozoan parasite affecting all warm-blooded animals (Miller, Frenkel & Dubey, 1972; Jackson & Hutchinson, 1989). It causes the disease toxoplasmosis which, although generally asymptomatic, can be life threatening in immunocompromised patients, such as those suffering from AIDS (Luft & Remington, 1992). The stage of the parasite responsible for dissemination of acute infection, termed the tachyzoite, is promiscuous *in vitro*, invading every mammalian cell line. While evidence that molecules secreted from the microneme organelles direct the process of host cell attachment is accumulating (Garcia-Réguet *et al.* 2000; Reiss *et al.* 2001; Brecht *et al.* 2001; Lourenco *et al.* 2001), the role of parasite surface molecules in this process remains equivocal.

The tachyzoite surface, which presumably represents the primary interface with the host cell surface, bears a number of glycoposphatidylinositol (GPI)-anchored surface proteins (Nagel & Boothroyd, 1989; Striepen *et al.* 1997), the most abundant of which is a 30 kDa protein named SAG1 (Kasper, Crabb & Pfefferkorn, 1983; Burg *et al.* 1988). Four lines of approach indirectly indicate that SAG1 binds

host cell surfaces. *In vitro* antibody neutralization invasion assays demonstrate that a subset of anti-SAG1 MAbs block tachyzoite invasion, implying that SAG1 might function as a specific parasite attachment ligand (Grimwood & Smith, 1992, 1996; Mineo *et al.* 1993; Mineo & Kasper, 1994). Competitive invasion assays have shown that the neoglycoprotein (NG) BSA-glucosamide blocks the ability of wild-type parasites to bind host cells, whereas those expressing truncated SAG1 remain unaffected (Mineo *et al.* 1993). In more studies, solubilized SAG1 has been shown to attach directly to host cell surfaces (Fourmaux *et al.* 1996; Channon *et al.* 1999; Velge-Roussel *et al.* 2001). These studies have not, however, investigated the specificity of this binding. Finally, the solution of the structure of SAG1 reveals the presence of a basic groove at the interface, formed during SAG1 dimerization (He *et al.* 2002). This domain faces away from the parasite surface and is therefore perfectly positioned for interaction with host cell surface ligands.

Although anti-SAG1 MAbs and BSA-glucosamide inhibit tachyzoite attachment *in vitro*, it is not known whether their blocking effects are explicitly mediated through masking host cell binding domains on SAG1. For instance, antibody neutralization experiments were used to investigate the ability of anti-SAG1 MAbs to prevent tachyzoite binding, rather than SAG1 binding. To address this question we here investigate the influence of antibodies and sugar-conjugates directly on SAG1.

\* Corresponding author: School of Biochemistry and Molecular Biology, University of Leeds, Leeds LS2 9JT, UK. Tel: +0113 343 3149. Fax: +0113 343 3167. E-mail: p.a.millner@bmb.leeds.ac.uk

## MATERIALS AND METHODS

*Cell lines*

All reagents for mammalian tissue culture were purchased from GibcoBRL. HS27 Human foreskin fibroblast cells (ATCC, USA) were grown in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% (v/v) fetal bovine serum (FBS) and incubated at 37 °C and 10% CO<sub>2</sub> in air. MDBK cells (adult bovine kidney; Madin & Darby, 1958; Porton Down, UK) were grown in Ham's F-12 nutrient mixture containing L-glutamine supplemented with 10% FBS and incubated at 37 °C and 5% CO<sub>2</sub> in air.

*Antibodies*

The production of MAb C1E3 has been previously described by Grimwood & Smith (1992). Polyclonal immune mouse serum (IMS) was taken by cardiac puncture from mice infected with the cystic strain ME49. Anti-mouse IgG-HRP conjugate was from Sigma-Aldrich, UK. Polyclonal IMS and MAb C1E3 were affinity-purified using a 1 ml Hi Trap Protein G column (Amersham) following manufacturer's instructions.

*Preparation of tachyzoite proteins*

Tachyzoites of the *T. gondii* RH strain were maintained by twice-weekly intraperitoneal passage in Tucks number 1 mice. Exudates were washed 3 times by centrifugation at 3000 g for 10 min followed with resuspension in phosphate-buffered saline (PBS), pH 7.2. Parasite pellets were finally resuspended in 4 ml of solubilization buffer (50 mM NaHCO<sub>3</sub>, pH 9.0 containing 0.1% (v/v) NP40), incubated at 4 °C for 1 h then freeze-thawed 3 times. Material was sonicated on ice 5 times for 20 sec, centrifuged at 10000 g for 5 min at 4 °C and the supernatant containing solubilized tachyzoite lysate was used for SAG1 purification and binding assays.

*Purification of SAG1*

SAG1 was purified in a one-step anion-exchange high performance liquid chromatography (HPLC) procedure. Four mg of 0.22 µm filtered tachyzoite lysate prepared in solubilization buffer was loaded onto an AP1 column (Waters; diameter=0.8 cm, height=7.5 cm) packed with 3.5 ml of DEAE fast-flow Sepharose (Amersham) and pre-incubated with the same buffer. Solubilization buffer (10 column volumes) was flushed through the column at a flow rate of 1 ml/min. Non-adherent protein, comprising pure SAG1, was collected manually in 1 ml fractions. Bound protein was eluted from the column using 2 M NaCl in solubilization buffer and dialysed against solubilization buffer at 4 °C overnight. Sodium

dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE) and immuno-blotting with anti-SAG1 mAb C1E3 were used to assess the protein profiles of fractions. Fractions containing purified SAG1 were pooled, flash-frozen in liquid N<sub>2</sub> and stored in 100 µl aliquots at -20 °C.

*SDS-PAGE and immuno-blotting*

SDS-PAGE was performed according to the method of Laemmli (1970) using 10% separating gels. Protein samples were not reduced. Following electrophoresis, samples were either silver-stained (Heukeshoven & Dernick, 1988), or transferred to PVDF membrane (Schleicher and Schuell, Dassel, Germany) as described by Towbin, Staehelin & Gordon (1978). Following blocking for 1 h with 2% (w/v) non-fat dry milk powder in PBST buffer (PBS, 0.1% (v/v) Tween 20, pH 7.4), membranes were incubated for 1 h at 37 °C in MAb C1E3 or polyclonal IMS diluted 1:500 in PBST containing 5% (w/v) non-fat dry milk. After copious washes in PBST buffer, proteins were revealed using horseradish peroxidase (HRP)-conjugated anti-mouse IgG (diluted to 1:1000) followed by 3,3-diaminobenzidine tetrahydrochloride substrate (DABS) staining (Sigma-Aldrich).

*Cell binding assays*

The *in vitro* protein attachment assay was based on a procedure by Fourmaux *et al.* (1996). Cells were grown on 6-well dishes (9.5 cm<sup>2</sup> wells) until confluent. Medium was discarded and cells were blocked by 3 × 10 min washes with 10 ml of medium containing 2% (w/v) bovine serum albumin (BSA). After blocking, 0.5 mg of soluble tachyzoite lysate prepared in 5 ml of medium was incubated on the confluent monolayer for 30 min at 37 °C. Non-adherent lysate was discarded and following 6 washes in 10 ml of medium, cells were solubilized by the addition of 1 ml of 1 × SDS-PAGE sample loading buffer (75 mM Tris-HCl, pH 6.8, 5% (w/v) SDS, 0.0025% (w/v) bromophenol blue) then detached using a cell scraper. Samples were immediately subjected to analysis by SDS-PAGE and immuno-blotting.

## RESULTS

*Purification of SAG1 by anion-exchange chromatography*

Anion-exchange chromatography separates molecules according to charge differences. Using the ExPASy Molecular Biology Server (<http://www.ncbi.nlm.nih.gov>), the theoretical pI of mature SAG1 was calculated to be 7.5. This moderately high pI was exploited to purify SAG1 by anion-exchange HPLC. Tachyzoite lysates were fractionated by




Fig. 1. Anion-exchange HPLC purification of SAG1. Tachyzoite lysates were fractionated by anion exchange at pH 9.0, 8.5 and 8.0. Flow-through (F and Fs) and bound (B and Bs) fractions were separated on a 12% (w/v) polyacrylamide gel. Lanes T and Ts are non-fractionated tachyzoite lysates. Lanes Ts, Fs and Bs were silver-stained, lanes T, F and B were analysed by immuno-blotting using anti-SAG1 MAb C1E3 as primary antibody. Cross-reacting bands were detected using anti-mouse IgG HRP conjugate followed by DABS staining. Values on the right side indicate molecular sizes of marker proteins in kDa.




Fig. 2. Binding of tachyzoite molecules to host cell surfaces. Immuno-blot analysis of (A) tachyzoite proteins binding to MDBK and HS27 cells in the *in vitro* protein attachment assay, and (B) crude tachyzoite cell lysate. Host cells were incubated with (lanes 2 and 4) or without (lanes 1 and 3) soluble tachyzoite lysate (0.5 mg), then washed and host cell proteins separated by SDS-PAGE on a 12% (w/v) polyacrylamide gel. For immuno-blotting, proteins were analysed with IMS (A, lanes 1 and 2; and B) or MAb C1E3 (A, lanes 3 and 4) used as primary antibody. Cross-reacting bands were detected using anti-mouse IgG HRP conjugate followed by DABS staining. Values on the right side indicate molecular sizes of marker proteins in kDa.

anion-exchange HPLC in column buffer at pH 9.0. Silver staining and Western analysis (with MAb C1E3) did not detect SAG1 in the bound tachyzoite fraction (Fig. 1). At this pH SAG1 was the only parasite protein that eluted from the ion-exchange resin. Thus anion-exchange HPLC proved to be very effective in separating virtually all SAG1 from tachyzoite lysates. Interestingly, purification at pH 8.5 and 8.0 fractionated SAG1 from most, but not all, parasite proteins. From this, the actual pI of solubilized SAG1 was deduced as being above 9.0.

*SAG1 from total cell lysate binds to MDBK cells and HS27 cell surfaces*

The first step was to investigate the binding of proteins from tachyzoite lysates to the host cell surface. Immune mouse sera (IMS) identified major bands of approximately 28 kDa and 75 kDa in MDBK cell extracts, and 28 kDa and 48 kDa in HS27 cell extracts, following incubation of monolayers with tachyzoite lysate (Fig. 2A). Comparing these fractions to the protein profile of tachyzoite total cell




Fig. 3. Inhibition of SAG1 attachment to MDBK cells using BSA-glucosamide. SAG1 was pre-incubated with the NG BSA-glucosamide prior to the *in vitro* protein attachment assay. Incubated cells were fractionated by SDS-PAGE on a 12% (w/v) polyacrylamide gel and proteins detected by silver staining. Lane 1 contained non-incubated MDBK cells, lane 2 is SAG1 and lane 3 contained MDBK cells incubated with 0.1 mg of SAG1. Lanes 4–7 contained MDBK cells incubated in SAG1 that had been pre-treated with serial dilutions of BSA-glucosamide (0.1  $\mu$ M, 1  $\mu$ M, 10  $\mu$ M and 100  $\mu$ M, respectively). Lane 8 is SAG1 pre-treated with BSA alone. Values on the right side indicate molecular sizes of marker proteins in kDa.

lysate (Fig. 2B) revealed that many abundant tachyzoite proteins did not adhere to the host cell surfaces, indicating that the binding of the 28 kDa, 48 kDa and 75 kDa proteins was specific. Probing with MAb C1E3 confirmed the 28 kDa molecule as SAG1. No bands were present in control, unincubated cell extracts.

To minimize the likelihood that tachyzoite proteins were present in the host cell fractions due to internalization or receptor recycling, the *in vitro* protein attachment assay was repeated at 4 °C to retard host cell endocytotic machinery. Comparing the results of attachment at 4 °C with attachment at 37 °C revealed no differences (data not shown), suggesting that the tachyzoite proteins in host cell fractions arose as a consequence of binding to the host cell surfaces.

#### *Effect of neoglycoprotein BSA-glucosamide on attachment of SAG1 to MDBK cells*

We wanted to investigate the putative binding of SAG1 to host cell-surface glucosamine. Competition experiments were performed using SAG1 that had been pre-treated with serial dilutions of the neoglycoprotein BSA-glucosamide. Soluble BSA-glucosamide blocked SAG1 attachment to MDBK cells in a dose-dependent manner; virtually all SAG1 binding was prevented with 100  $\mu$ M BSA-glucosamide (Fig. 3). The attachment blockade was shown to be specific for glucosamide since control incubations




Fig. 4. The role of SAG1 epitopes in SAG1 binding. The binding of SAG1 epitopes to host cells was assessed by pre-incubating SAG1 with either MAb C1E3 or IMS. Incubated cells were fractionated by SDS-PAGE on a 12% (w/v) polyacrylamide gel and proteins were analysed by Western blotting using anti-SAG1 MAb C1E3 as primary antibody. Cross-reacting bands were detected using anti-mouse IgG HRP conjugate followed by DABS staining. Lane 1 contained MDBK cells incubated with 0.5 mg purified antibody. Lane 2 contained MDBK cells incubated with 0.1 mg SAG1 that had been pre-treated with 0.5 mg purified antibody. Values on the right side indicate molecular sizes of marker proteins in kDa. Arrows indicate the presence of higher molecular weight complexes of SAG1 detected by MAb C1E3.

with BSA had no effect on SAG1 binding (Fig. 3, lane 8). Pre-incubation of host cells with BSA-glucosamide or BSA had no effect on their integrity or morphology (data not shown). Taken together these results illustrate that BSA-glucosamide competes with host cell surfaces for SAG1 binding.

#### *Effect of neoglycoprotein antibody on attachment of SAG1 to MDBK cells*

SAG1 was pre-incubated with MAb C1E3 or polyclonal IMS to determine whether these agents perturb the binding of SAG1 to host cell surface. Preliminary incubation of SAG1 with a 4-fold molar excess of MAb C1E3 did not inhibit attachment of SAG1 to MDBK cell surfaces, and so it appears that the MAb C1E3 epitope does not overlap with the SAG1 domain for host cell binding (Fig. 4). SAG1 could also bind host cell surfaces following pre-incubation with a 4-fold molar excess of IMS. These results demonstrate that immunodominant B cell epitopes on SAG1 do not directly bind to the surfaces of host cells. Interestingly, Western analysis with MAb C1E3 detected minor bands of approximately 55 kDa, 120 kDa and 180 kDa in MDBK cell extracts that had been incubated with the SAG1-antibody mixtures (Fig. 4, lanes 2). These molecules were resistant to reduction in 25 mM

DTT, and were therefore not oligomers of IgG heavy chain (data not shown).

#### DISCUSSION

We have shown that SAG1 specifically binds the surface of bovine kidney cells and human foreskin fibroblasts, thus adding to the collection of host cells to which SAG1 attaches, i.e. Vero cells (Fourmaux *et al.* 1996), monocytes (Channon *et al.* 1999) and epithelial cells (Velge-Roussel *et al.* 2001).

The finding that attachment of SAG1 to bovine kidney cells could be blocked by pre-incubation of SAG1 in BSA-glucosamide is in agreement with a previous study which found that the ability of BSA-glucosamide to block invasion was markedly reduced when SAG1-deficient mutants were used in place of wild-type parasites (Mineo *et al.* 1993). From this result it would be tempting to suggest that SAG1 binding to host cells is mediated through interaction with host cell-surface glucosamine. Even so, direct binding of SAG1 to glucosamide has not been confirmed, and further work is required to assess the specificity of interaction with other sugar-conjugates.

Interestingly, it has been established that attachment of *T. gondii* to host cells is mediated, in part, by the binding of parasite lectins to complex sugars such as glycosaminoglycans, e.g. heparin (Ortega-Barria & Boothroyd, 1999; Carruthers *et al.* 2000; Jacquet *et al.* 2001). Recently, recombinant SAG3 (rSAG3) was shown to recognize sulphated proteoglycans on the surface of CHO-K1 (Jacquet *et al.* 2001). Modelling the SAG3 structure reveals that the outside surface of the SAG3 groove is negatively charged, which may apparently function to direct specific binding of negatively charged heparin, a sulphated glycosaminoglycan, to the positively charged groove (He *et al.* 2002). It is tempting to speculate that the absence of this feature in rSAG1 allows the protein to ubiquitously attach to sulphated and non-sulphated cell surface glycans. Even so, whilst Jacquet *et al.* (2001) demonstrated that rSAG3 binds with high affinity to heparin, these authors found no evidence for the binding of rSAG1 to heparin. However, host cell binding regions may not be correctly folded on rSAG1, and so the binding of native SAG1 to sulphated proteoglycans cannot be ruled out.

It is unlikely that all host cells will exhibit the same surface receptors or that a single parasite surface protein could mediate invasion of every cell type. Thus, it is likely that there is redundancy in these ligands. For example, in this paper we have shown that SAG1 can bind human foreskin fibroblasts, although Dzierszinski *et al.* (2000) have shown that SAG3 also binds this cell type. Disruption of the SAG3 gene reduces, but does not abolish invasion, and thus SAG1 could be responsible for mediating parasite attachment to this cell type. Even so, the

presence of the diverse family of SAG1-related sequence (SRS) proteins may circumvent the need for both SAG1 and SAG3, making the parasite even more versatile. Interestingly, efforts to create a SAG1/SAG3 knockout have so far been unsuccessful.

One intriguing finding was the discovery that SAG1 forms high molecular weight complexes by self-association. These data corroborate the finding that rSAG1 forms dimers in solution (He *et al.* 2002), though the observations presented here also imply that SAG1 can also form heteromers and hexamers. The behaviour of SAG1 could be characterized in two ways. First, higher complexes were not present subsequent to purification, only appearing once SAG1 had been stored at  $-20^{\circ}\text{C}$ . Second, the absence of a band of approximately 80–90 kDa indicates that SAG1 does not form trimers, thus each complex is a multimeric association of SAG1 dimers.

*In vitro* invasion assays using anti-SAG1 MAb C1E3 have previously indicated that SAG1 has a role in host cell attachment (Grimwood & Smith, 1992, 1996). In these experiments, MAb C1E3 was used to perturb parasite invasion, and since only a subset of anti-SAG1 MAbs conveyed this effect it has been postulated that the blocking ability of MAb C1E3 was due to the masking of a specific epitope of SAG1. With the sole purpose of understanding the mechanism of anti-SAG1 MAb neutralization, the stoichiometry of these antibody–antigen interactions was addressed. Unexpectedly, the results presented here indicate that MAb C1E3 does not in fact block SAG1 binding to host cell surfaces. To comprehend this peculiar finding, the interaction between MAb C1E3 and SAG1 requires further investigation. Since MAb C1E3 binding appears to be implicitly linked to the process of host cell attachment, the identification of the epitope for this MAb on SAG1 could provide a valuable insight into the role of SAG1 epitopes in host cell attachment.

We thank Debra Evans and Sinead Moore for their expert technical assistance.

#### REFERENCES

- BRECHT, S., CARRUTHERS, V. B., FERGUSON, D. J., GIDDINGS, O. K., WANG, G., JAKLE, U., HARPER, J. M., SIBLEY, L. D. & SOLDATI, D. (2001). The *Toxoplasma* micronemal protein MIC4 is an adhesin composed of six conserved apple domains. *Journal of Biological Chemistry* **276**, 4119–4127.
- BURG, J. L., PERELMAN, D., KASPER, L. H., WARE, P. L. & BOOTHROYD, J. C. (1988). Molecular analysis of the gene encoding the major surface antigen of *Toxoplasma gondii*. *Journal of Immunology* **141**, 3584–3593.
- CARRUTHERS, V. B., HAKANSSON, S., GIDDINGS, O. K. & SIBLEY, L. D. (2000). *Toxoplasma gondii* uses sulfated proteoglycans for substrate and host cell attachment. *Infection and Immunity* **68**, 4005–4011.

- CHANNON, J. Y., SUH, E. I., SEGUIN, R. M. & KASPER, L. H. (1999). Attachment ligands of viable *Toxoplasma gondii* induce soluble immunosuppressive factors in human monocytes. *Infection and Immunity* **67**, 2547–2551.
- DZIERZYSKI, F., MORTUAIRE, M., CESBRON-DELAUW, M. F. & TOMAVO, S. (2000). Targeted disruption of the glycosylphosphatidylinositol-anchored surface antigen SAG3 gene in *Toxoplasma gondii* decreases host cell adhesion and drastically reduces virulence in mice. *Molecular Microbiology* **37**, 574–582.
- FOURMAUX, M. N., ACHBAROU, A. & MERCEREAU-PUIJALON, O., BIDERRE, C., BRICHE, I., LOYENS, A., ODBERG-FERRAGUT, C., CAMUS, D. & DUBREMETZ, J. F. (1996). The MIC1 microneme protein of *Toxoplasma gondii* contains a duplicated receptor-like domain and binds to host cell surface. *Molecular and Biochemical Parasitology* **83**, 201–210.
- GARCIA-RÉGUET, N., LEBRUN, M., FOURMAUX, M. N., MERCEREAU-PUIJALON, O., MANN, T., BECKERS, C. J., SAMYN, B., VAN BEEUMEN, J., BOUT, D. & DUBREMETZ, J. F. (2000). The microneme protein MIC3 of *Toxoplasma gondii* is a secretory adhesin that binds to both the surface of the host cells and the surface of the parasite. *Cellular Microbiology* **2**, 353–364.
- GRIMWOOD, J. & SMITH, J. E. (1992). *Toxoplasma gondii*: The role of a 30 kDa surface protein in host cell invasion. *Experimental Parasitology* **74**, 106–111.
- GRIMWOOD, J. & SMITH, J. E. (1996). *Toxoplasma gondii*: The role of parasite and secreted proteins in host cell invasion. *International Journal for Parasitology* **2**, 169–173.
- HE, X. L., GRIGG, M. E., BOOTHROYD, J. C. & GARCIA, K. C. (2002). Structure of the immunodominant surface antigen from *Toxoplasma gondii* SRS superfamily. *Nature, Structural Biology* **9**, 606–611.
- HEUKESHOVEN, J. & DERNICK, R. (1988). Improved silver staining procedure for fast staining in PhastSystem Development Unit. I. Staining of sodium dodecyl sulfate gels. *Electrophoresis* **9**, 28–32.
- JACKSON, M. H. & HUTCHINSON, W. M. (1989). The prevalence and source of *Toxoplasma* infection in the community. *Advances in Parasitology* **91**, 237–249.
- JACQUET, A., COULON, L., DE NEVE, J., DAMINET, V., HAUMONT, M., GARCIA, L., BOLLEN, A., JURADO, M. & BIEMANS, R. (2001). The surface antigen SAG3 mediates the attachment of *Toxoplasma gondii* to cell-surface proteoglycans. *Molecular and Biochemical Parasitology* **116**, 35–44.
- KASPER, L. H., CRABB, J. H. & PFEFFERKORN, E. R. (1983). Purification of a major membrane protein of *Toxoplasma gondii* by immuno-adsorption with a monoclonal antibody. *Journal of Immunology* **130**, 2407–2412.
- LAEMMLI, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature, London* **227**, 680–685.
- LOURENCO, E. V., PERIERA, S. R., FAÇA, V. M., COELHO-CASTELO, A. A. M., MINEO, J. R., ROQUE-BARREIRA, M.-C., GREENE, L. J. & CASTEOL-PANUNTO, A. (2001). *Toxoplasma gondii* micronemal protein MIC1 is a lactose-binding lectin. *Glycobiology* **11**, 541–547.
- LUFT, B. F. & REMINGTON, J. S. (1992). Toxoplasmic encephalitis in AIDS. *Clinical Infection and Disease* **15**, 211–222.
- MADIN, S. H. & DARBY, N. B. (1958). Established kidney cell lines of normal adult bovine and ovine origin. *Proceedings of the Society of Experimental Biology and Medicine* **98**, 574–576.
- MILLER, N. L., FRENKEL, J. K. & DUBEY, J. P. (1972). Oral infections with *Toxoplasma* cysts and oocysts in felines, other mammals, and in birds. *Journal of Parasitology* **58**, 928–937.
- MINEO, J. R. & KASPER, L. H. (1994). Attachment of *Toxoplasma gondii* to host cells involves major surface protein, SAG1-1 (P30). *Experimental Parasitology* **79**, 351–361.
- MINEO, J. R., McLEOD, R., MACK, D., SMITH, J. E., KHAN, I. A., ELY, K. H. & KASPER, L. H. (1993). Antibodies to *Toxoplasma gondii* major surface protein (SAG-1, P30) inhibit infection of host cells and are produced in murine intestine after peroral infection. *Journal of Immunology* **150**, 3951–3964.
- NAGEL, S. D. & BOOTHROYD, J. C. (1989). The major surface antigen of *Toxoplasma gondii* is anchored by a glycolipid. *Journal of Biological Chemistry* **264**, 5569–5574.
- ORTEGA-BARRIA, E. & BOOTHROYD, J. C. (1999). A *Toxoplasma* lectin-like activity specific for sulfated polysaccharides is involved in host cell infection. *Journal of Biological Chemistry* **274**, 1267–1276.
- REISS, M., VIEBIG, N., BRECHT, S., FOURMAUX, M.-F., SOËTE, M., DI CRISTINA, M., DUBREMETZ, J. F. & SOLDATI, D. (2001). Identification and characterisation of an escorter for two secretory adhesions in *Toxoplasma gondii*. *Journal of Cell Biology* **152**, 563–578.
- STRIEPEN, B., ZINECKER, C. F., DAMM, J. B. L., MELGERS, P. A., GERWIG, G. J., KOOLEN, M., VLIEGENTHART, J. F., DUBREMETZ, J. F. & SCHWARTZ, T. T. (1997). Structure of low molecular weight antigen of *Toxoplasma gondii*: a glucose alpha-1-4 N-acetylgalactosamine makes free glycosyl-phosphatidylinositols highly immunogenic. *Journal of Molecular Biology* **266**, 797–813.
- TOWBIN, H., STAHELIN, T. & GORDON, J. (1978). Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proceedings of the National Academy of Sciences, USA* **76**, 4350–4354.
- VELGE-ROUSSEL, F., DIMIER-POISSON, I., BUZONI-GATEL, D. & BOUT, D. (2001). Anti-SAG1 peptide antibodies inhibit the penetration of *Toxoplasma gondii* tachyzoites into enterocyte cell lines. *Parasitology* **123**, 225–233.