



Title	Ceramide Aminoethylphosphonate in the Egg of the JANBOTANISHI, Ampullarius urceus
Author(s)	Tamari, Masato
Citation	長崎大学教育学部自然科学研究報告. vol.40, p.91-97; 1989
Issue Date	1989-02-28
URL	<a href="http://hdl.handle.net/10069/32450">http://hdl.handle.net/10069/32450</a>
Right	

This document is downloaded at: 2020-09-18T12:47:36Z

## Ceramide Aminoethylphosphonate in the Egg of the JANBOTANISHI, *Ampullarius urceus*

Masato TAMARI

Laboratory of Food and Nutrition, Faculty of Education,  
Nagasaki University, Nagasaki 852, JAPAN

(Received Oct. 31, 1988)

### ABSTRACT

Ceramide aminoethylphosphonate was isolated from the egg of the JANBOTANISHI, *Ampullarius urceus*. The compound was purified with a combination of mild alkaline hydrolysis of the total lipids, QAE-Sephadex-A 25 column chromatography and two dimensional thin layer chromatography.

The infrared spectrum showed an absorption band at  $1180\text{ cm}^{-1}$  due to c-p bond, and was essentially identical with that of the standard ceramide aminoethylphosphonate. Upon hydrolysis of the substance by strong acid, neither change in the chromatographic behaviors of this substance nor liberation of inorganic phosphate was observed. The stability of the compound to acid hydrolysis suggested the presence of a c-p bond.

On comparison with synthetic compound, the aqueous hydrolysis product behaved like 2-aminoethylphosphonic acid on thin layer chromatograms.

The predominant fatty acids were palmitic acid, eicosapentaenoic acid, erucic acid and docosatrienoic acid.

### INTRODUCTION

Natural compounds containing a carbon-phosphorus bond were first detected when Horiguchi and Kandatsu<sup>(1)</sup> isolated 2-aminoethylphosphonic acid from ciliate protozoa in the rumen of sheep. Since then this compound has been shown to be present either in the free form or in the bound to proteins or lipids in the protozoan *Tetrahymena pyriformis*, fresh water molluscus, terrestrial mammals, and marine animals in the phyla coelenterate, mollusca, echinoderamata. So far, phosphonate-containing lipids (termed phosphonolipid<sup>(2)</sup>) have been found as analogs of phosphatidylethanolamine, phosphatidylcholine and sphingoethanolamine from tissue of lower<sup>(3)</sup> and higher animals<sup>(4)</sup>. Rouser et al<sup>(5)</sup> first isolated a characteristic lipid from

sea anemone by means of column chromatography, and it was assumed tentatively to consist of a ceramide attached to 2-aminoethylphosphonic acid by an ester linkage. Since then this ceramide aminoethylphosphonate has been shown to be widely present in shellfish both in fresh water and sea water<sup>(6)</sup>, and Fungi<sup>(7)</sup>.

In our previous paper<sup>(8)</sup>, we reported that ceramide aminoethylphosphonate was isolated from the AGEMAKI, *Sinonovacula constricta*. Until the present, there is no report on the phosphonolipid and fatty acid composition of the JANBOTANISHI, *Ampullarius urceus*. Therefore, in this paper is reported the isolation and fatty acid composition of the ceramide aminoethylphosphonate from egg of the JANBOTANISHI, *Ampullarius urceus*.

## MATERIALS AND METHODS

### *Materials:*

QAE-Sephadex-A25 was the product of Pharmacia. 2-aminoethylphosphonic acid was prepared by the method described by Kosolapoff<sup>(9)</sup>.

### *Fractionation of Phosphorus Fraction of the Egg:*

Fractionation of phosphorus fraction (A: acetone soluble, B: chloroform/methanol soluble, C: TCA soluble, D: TCA insoluble) from the freeze dried egg of the JANBOTANISHI, *Ampullarius urceus* was carried out by the method as previously described<sup>(8)</sup>: A portion of each fraction was hydrolyzed with 6N HCl at 120°C for 24 hrs. The hydrolysate was treated several times with ether. The aqueous layer was filtered, evaporated to dryness, and redissolved with water for analysis of phosphonate- and total-phosphorus.

### *Isolation of Ceramide Aminoethylphosphonate:*

Lipid of the freeze dried egg (10g) of the *Ampullarius urceus* was extracted by the method as previously described<sup>(8)</sup>. The combined extracts was washed according to the method described by Folch et al.<sup>(10)</sup> The chloroform layer was evaporated to dryness in a N<sub>2</sub> stream, and the isolation and purification of the sphingophosphonolipid from the lipid was achieved by three methods as previously described<sup>(8)</sup>: by the mild alkaline hydrolysis according to Dowson<sup>(11)</sup> and Hori<sup>(12)</sup>, by the QAE-Sephadex column chromatography and by the preparative thin layer chromatography.

### *Infrared Analysis:*

The infrared absorption spectrum was determined on a pellet of potassium bromide using a Nihonbunko IR-S Spectrophotometer.

### *Qualitative Analysis:*

The ceramide aminoethylphosphonate was hydrolyzed with 6N HCl at 120°C for 24 hrs. The liberated fatty acids were extracted with light petroleum and the water soluble compounds were extracted by Folch's partition.

The aqueous layer was evaporated to dryness under reduced pressure.

The presence of aminoethylphosphonate in the hydrolysate was demonstrated by thin layer chromatography (Kiesel gel 60 F<sub>254</sub>, 20 × 20 cm, 0.25 mm) using 99% ethanol: 7% aqueous ammonia (1 : 2, v/v)<sup>(7, 13)</sup>.

Authentic 2-amioethylphosphonic acid was co-chromatographed and the spots were detected by both ninhydrine and Rosenberg's method<sup>(14)</sup>.

#### Quantitative Analysis:

Fatty acid ester were obtained from the ceramide aminoethylphosphonate by methanolysis (5% HCl in methanol at 100°C for 4 hrs) and subjected to gas chromatography. Phosphonate phosphorus was estimated by the method of Tamari et al<sup>(15)</sup>. Total phosphorus was estimated by the method of Chen et al<sup>(16)</sup>.

#### Gas Chromatography of Fatty Acid:

Fatty acid methyl esters were separated on a Shimadzu GC-8A apparatus equipped with a flame ionization detector. The column was a 3 mm × 2 m stainless column packed with 5% Shinchron E 71 on 80-100 mesh Chimalite, and the instrument was operated at 200°C. The peaks were identified by comparison with those of standard methyl esters on the basis of individual retention time.

## RESULTS

In Table 1 is reported the phosphonate content of the phosphorus fractions from the egg of the *Ampullarius urceus*. The phosphonate content in the A, B, C and D fractions were 4.48, 46.23, 8.00 and 615.61  $\mu\text{g}$  per g of the freeze dried eggs of the *Ampullarius urceus*, respectively.

The B and D fractions contained 6.9% and 91.3% of the phosphonate in the freeze dried egg. About 28, 4, 1 and 28% of the total phosphorus was found as phosphonate in the A, B, C and D fractions. A typical thin layer chromatographic compositions of phospholipid classes was identical to that of other invertebrates<sup>(17)</sup>.

Table 1 Phosphorus Contents of the Phosphorus Fraction Obtained from Egg of the JANBOTANISHI, *Ampullarius urceus*.

Fraction	C-P ( $\mu\text{g/g}$ )	T-P ( $\mu\text{g/g}$ )	C-P/T-P (%)
A	4.48 ( 0.66)	15.76	28.4
B	46.23 ( 6.86)	976.01	4.7
C	7.99 ( 1.19)	977.06	0.8
D	615.61 (91.29)	2181.10	28.2

C-P: Phosphonate-phosphorus, T-P: Total-phosphorus,  
( ) : as % of total C-P




Fig. 1 Two Dimensional Thin Layer Chromatogram of the Alkaline Stable Lipids Obtained by the Mild Alkaline Hydrolysis as Previously Described<sup>(8)</sup>.

The chromatogram was developed as previously described<sup>(8)</sup>. Spot are: 1, ceramide aminoethylphosphonate (34.8%); 2, probably hydroxy fatty acid containing ceramide aminoethylphosphonate (12.0%); 3, sphingomyelin (53.2%)




Fig. 2 Infrared Spectrum of the Ceramide Aminoethylphosphonate Purified by a Combination of the Mild Alkaline Hydrolysis, the Column Chromatography and the Preparative Thin Layer Chromatography as Previously Described<sup>(8)</sup>.

Figure 1 shows the thin layer chromatogram of the sphingolipid obtained by the mild alkaline hydrolysis method. Ceramide aminoethylphosphonate (spot 1), and probably what is hydroxy fatty acid containing ceramide aminoethylphosphonate (spot 2) and sphingomyelin (spot 3), consisted of 34.8, 12.0 and 53.2% of the sphingolipid phosphorus, respectively. The result obtained here indicates that ceramide aminoethylphosphonate is the major sphingolipid component of the lipid moiety of the egg of the JANBOTANISHI, *Ampullarius urceus*.

Further, ceramide aminoethylphosphonate was purified by QAE-sephadex column chromatography from the sphingolipids obtained by mild alkaline hydrolysis of the phospholipid fraction, and pure ceramide aminoethylphosphonate was then obtained by preparative thin layer chromatography.

Figure 2 shows the infrared spectrum of the ceramide aminoethylphosphonate obtained by the alkaline

Table 2 Fatty Acid Composition of Total Lipid, Sphingolipid and Ceramide Aminoethylphosphonate from Egg of the JANBOTANISHI, *Ampullarius urceus*.

Fatty acid	Total lipid (%)	Sphingolipid (%)	Ceramide aminoethylphosphonate (%)
14 : 0	4.66	2.91	3.07
16 : 0	35.56	60.36	16.90
18 : 0	11.72	21.82	6.47
18 : 1	8.52	1.94	6.82
18 : 2	12.92	2.55	2.22
18 : 3	3.33	0.61	1.50
20 : 3	0.40	0.61	6.83
20 : 4	14.52	2.79	1.53
20 : 5	4.93	4.00	30.51
22 : 1	1.60	0.36	13.58
22 : 3	3.86	2.06	10.58

hydrolysis method. The infrared spectrum lacked a band at 1730-1750  $\text{cm}^{-1}$ , indicating the absence of ester bonds. A band was present at 1180  $\text{cm}^{-1}$ , indicating the presence of carbon-phosphorus bond, and was essentially identical with that of ceramide aminoethylphosphonate, obtained by Rouser et al<sup>(5)</sup>, Hayashi et al<sup>(18)</sup>, and Hori et al<sup>(11, 19)</sup>.

The carbon-phosphorus compound in the acid hydrolysate of the ceramide aminoethylphosphonate was found to be 2-aminoethylphosphonic acid by thin layer chromatography. The  $R_f$  (0.72) of the spot agreed with that of authentic 2-aminoethylphosphonic acid. Inorganic phosphorus and N-methyl-derivatives of 2-aminoethylphosphonic acid were not detected in the hydrolysate. The results of gas chromatographic analysis of fatty acid composition are shown in Table 2. The fatty acid composition was very simple. The fatty acids of the ceramide aminoethylphosphonate were characterized by relatively high percentages for palmitic acid, eicosapentaenoic acid, erucic acid and docosatrienoic acid. The sum of these four acids comprised about 72% of the total fatty acids. Oleic acid (6.8%), stearic acid (6.5%) and arachidic acid (1.5%) were also found as minor components. From the above results, the isolated phosphonolipid was identified as the ceramide 2-aminoethylphosphonate.

## DISCUSSION

The distribution of phosphonolipids in nature is primarily limited to lower animals such as mollusca, coelenterates and protozoa, although the lipids have also been found in small quantities in mammalian tissues<sup>(20-22)</sup>.

In the phosphonolipids, it has been established that glycerol-aminoethylphosphonate (GPnL) occurs as the major phospholipid of the ciliary membranes of protozoan species<sup>(23, 24)</sup>. To date, there is still no report on the detection of the GPnL in

shellfish and sea anemones.

On the other hand, ceramide aminoethylphosphonate are found in high concentration in mollusca, coelenterates and shellfish, while the concentration is low in *Tetrahymena*. Matsubara<sup>(25)</sup> found that the oyster adductor muscle contained the highest concentration (45% of the total sphingolipids) of ceramide aminoethylphosphonate. Komai et al<sup>(26)</sup> found that ceramide aminoethylphosphonate occurs in approximately 11% of the phospholipids in the nervous system of *A. Kurodai* a marine gastropod. Moreover, Komai et al speculate that ceramide aminoethylphosphonate may be indispensable in shellfish for neuronal function. Kittredge et al<sup>(27)</sup> indicate that the presence of a covalent carbon-phosphorus linkage in the anemone has been postulated as functionally analogous to a fixation process.

Mason<sup>(28)</sup> indicates the possibility that the presence of a highly ionic lipid such as ceramide aminoethylphosphonate could play a direct role in facilitating the transport of small ions from the aqueous environment into the intercellular space of the anemone.

On the other hand, the occurrence of the ceramide aminoethylphosphonate in the sea-urchin gonad<sup>(29)</sup> and fresh water mussel spermatozoa<sup>(17)</sup> and egg<sup>(30, 31)</sup> has been reported. While this sphingophosphonolipid is unique to the invertebrates, its exact function is unknown, but their resistance to endogenous hydrolytic enzymes is highly suggestive of a protective function.

It will be of further interest to determine of the presence of ceramide aminoethylphosphonate is localized in the egg of shellfishes.

## REFERENCES

- (1) M. Horiguchi and M. Kandatsu, *Nature, Lond.*, 184, 901 (1959)
- (2) E. Baer and N. Z. Stanacov, *J. Biol. Chem.*, 239, 3209 (1965)
- (3) J. S. Kittredge and E. Robert, *Science*, 164, 37 (1969)
- (4) S. Hasegawa, M. Tamari and M. Kametaka, *J. Biochem.*, 80, 531 (1976)
- (5) G. Rouser, D. H. Kitchevsky, D. Heller and E. Lieber, *J. Am. Oil Chemists Soc.*, 40, 425 (1963)
- (6) T. Hori, O. Itasaka and H. Inoue, *J. Biochem.*, 56, 477 (1964)
- (7) M. K. Wassef and J. W. Hendrix, *Biochim. Biophys. Acta*, 486, 172 (1977)
- (8) M. Tamari and M. Kandatsu, *Agric. Biol. Chem.*, 50, 1495 (1986)
- (9) G. M. Kosolapoff, *J. Am. Chem. Soc.*, 69, 2112 (1947)
- (10) J. Folch, M. Lee and G. H. Sloan-Stanley, *J. Biol. Chem.*, 226, 497 (1957)
- (11) R. M. C. Dowson, *Biochem. J.*, 75, 45 (1960)
- (12) T. Hori, M. Sugita and O. Itasaka, *J. Biochem.*, 65, 451 (1969)
- (13) J. A. Brushaber, J. J. Child and R. H. Haskins, *Can. J. Microbiol.*, 18, 1059 (1972)
- (14) H. Rosenberg, *J. Chromatog.*, 2, 487 (1959)
- (15) M. Tamari, M. Horiguchi and M. Kandatsu, *Nippon Nogeikagaku Kaishi*, 45, 433 (1971)
- (16) P. S. Chen, T. Y. Toribara and H. Warner, *Anal. Chem.*, 28, 1756 (1956)

- (17) S. Higashi and T. Hori, *Biochim. Biophys. Acta*, 152, 568 (1968)
- (18) A. Hayashi, T. Matsubara and Y. Mishima, *J. Fac. Sci. Tech. Kinki Univ.*, 2, 39 (1967)
- (19) T. Hori, O. Itasaka and H. Inoue, *J. Biochem.*, 59, 570 (1966)
- (20) M. Tamari, M. Ogawa, S. Hasegawa and M. Kametaka, *Agric. Biol. Chem.*, 40, 2057 (1976)
- (21) S. Hasegawa, M. Tamari and M. Kametaka, *J. Biochem.*, 80, 531 (1976)
- (22) M. C. Moschidis, *J. Chromatogr.*, 268, 485 (1983)
- (23) Y. Nozawa and G. A. Thompson, *J. Cell Biol.*, 49, 712 (1971)
- (24) K. E. Kennedy and G. A. Thompson, *Science*, 168, 989 (1970)
- (25) T. Matsubara, *Biochim. Biophys. Acta*, 388, 353 (1975)
- (26) Y. Komai, S. Matsukawa and M. Satake, *Biochim. Biophys. Acta*, 316, 271 (1973)
- (27) J. Kittredge, E. Roberts and D. Simonsen, *Biochemistry*, 1, 624 (1962)
- (28) W. T. Mason, *Biochim. Biophys. Acta*, 280, 538 (1972)
- (29) H. Ikezoe, *Bull. Obihiro Ohtani Junior College*, No. 10, 5 (1973)
- (30) T. Hori, M. Sugita, J. Kambayashi and O. Itasaka, *Yukagaku*, 24, 181 (1975)
- (31) T. Hori, M. Sugita, T. Nishimori and O. Itasaka, *Yukagaku*, 24, 611 (1975)