



This is a repository copy of *Heat shock proteins in health and disease: therapeutic targets or therapeutic agents?* .

White Rose Research Online URL for this paper:

<http://eprints.whiterose.ac.uk/585/>

Article:

Pockley, A.G. (2001) Heat shock proteins in health and disease: therapeutic targets or therapeutic agents? *Expert Reviews in Molecular Medicine*. ISSN 1462-3994

Reuse

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.

Heat shock proteins in health and disease: therapeutic targets or therapeutic agents?

A. Graham Pockley

For many years, heat shock or stress proteins have been regarded as intracellular molecules that have a range of housekeeping and cytoprotective functions, only being released into the extracellular environment in pathological situations such as necrotic cell death. However, evidence is now accumulating to indicate that, under certain circumstances, these proteins can be released from cells in the absence of cellular necrosis, and that extracellular heat shock proteins have a range of immunoregulatory activities. The capacity of heat shock proteins to induce pro-inflammatory responses, together with the phylogenetic similarity between prokaryotic and eukaryotic heat shock proteins, has led to the proposition that these proteins provide a link between infection and autoimmune disease. Indeed, both elevated levels of antibodies to heat shock proteins and an enhanced immune reactivity to heat shock proteins have been noted in a variety of pathogenic disease states. However, further evaluation of heat shock protein reactivity in autoimmune disease and after transplantation has shown that, rather than promoting disease, reactivity to self-heat shock proteins can downregulate the disease process. It might be that self-reactivity to heat shock proteins is a physiological response that regulates the development and progression of pro-inflammatory immunity to these ubiquitously expressed molecules. The evolving evidence that heat shock proteins are present in the extracellular environment, that reactivity to heat shock proteins does not necessarily reflect adverse, pro-inflammatory responses and that the promotion of reactivity to self-heat shock proteins can downregulate pathogenic processes all suggest a potential role for heat shock proteins as therapeutic agents, rather than as therapeutic targets.

A. Graham Pockley

Reader in Immunobiology, Division of Clinical Sciences (North), Clinical Sciences Centre (University of Sheffield), Northern General Hospital, Herries Road, Sheffield, S5 7AU, UK. Tel: +44 114 271 4450; Fax: +44 114 261 9246; E-mail: g.pockley@sheffield.ac.uk.

It was in 1962 that Ritossa and co-workers first discovered that subjecting *Drosophila melanogaster* larvae to temperature shock induced specific gene activation (Ref. 1); however, it was not until 1974 that the first products of these genes were identified and the term 'heat shock protein' was adopted (Ref. 2). Subsequent work has demonstrated that heat shock proteins are present, and can be induced, in all species and that they are among the most phylogenetically

conserved proteins. Heat shock proteins are categorised into several families that are named on the basis of their approximate molecular mass (e.g. the 70 kDa Hsp70; Table 1). Under physiological conditions, some of these proteins function as molecular chaperones or proteases that have a number of intracellular functions. Chaperones are involved in the assembly and folding of oligomeric proteins, whereas proteases such as the ubiquitin-dependent proteasome

Table 1. Major mammalian heat shock proteins and their function (tab001gps)

Major family and members	Cellular localisation	Cellular function
Small		
α B-crystallin	Cytoplasm	Cytoskeletal stabilisation
Hsp27	Cytoplasm/nucleus	Actin dynamics
Heme oxygenase, Hsp32	Cytoplasm	Haeme catabolism, antioxidant properties
Hsp60 or chaperonins		
Hsp60	Mitochondria	Both: bind to partially folded polypeptides and assist correct folding; assemble multimeric complexes
TCP-1	Cytoplasm	
Hsp70		
Hsp70 (inducible)	Cytoplasm/nucleus	All: bind to extended polypeptides; prevent aggregation of unfolded peptides; dissociate some oligomers; bind ATP and show ATPase activity
Hsc70 (cognate)	Cytoplasm/peroxisome	
Grp78/BiP	ER	
mtHsp70/Grp75	Mitochondria	Hsp70 is involved in regulation of HSF1 activity and the repression of heat shock protein gene transcription
Hsp90		
Hsp90 (α and β)	Cytoplasm	All: bind to other proteins; regulate protein activity; prevent aggregation of re-folded peptide; correct assembly and folding of newly synthesised protein
Grp94/gp96/Hsp100	ER	
Hsp90 appears to be involved in maintaining the HSF1 monomeric state in non-stressful conditions; represents 1–2% of total protein		
Hsp110		
Hsp110 (human)	Nucleolus/cytoplasm	Thermal tolerance
Apg-1 (mouse)	Cytoplasm	Protein refolding
Hsp105	Cytoplasm	

Abbreviations: ER, endoplasmic reticulum; TCP-1, tailless complex polypeptide; Grp, glucose-regulated protein; Hsp, heat shock protein; BiP, immunoglobulin heavy chain binding protein; mtHsp70, mitochondrial Hsp70; HSF1, heat shock factor 1; Apg-1, protein kinase essential for autophagy.

mediate the degradation of damaged proteins (Refs 3, 4).

The term heat shock proteins is somewhat of a misnomer, as they are not induced solely by heat shock. Indeed, in addition to being constitutively expressed (making up 5–10% of the total protein content under normal growth conditions), these proteins can be markedly induced (up to 15% of the total cellular protein content) by a range of cellular insults including increased temperature, oxidative stress, nutritional deficiencies, ultraviolet irradiation, exposure to chemicals (e.g. ethanol), viral infection, and ischaemia–reperfusion injury (Refs 5, 6). Stressors that cause protein unfolding, misfolding or aggregation trigger a stress response that leads to the induction of gene transcription for proteins with the capacity to stabilise and re-fold proteins, thereby re-establishing the balance between protein synthesis, assembly and degradation.

Regulation of heat shock protein gene transcription is mediated by the interaction of the heat shock factor (HSF) transcription factors (of which the principal one in vertebrates is HSF1) with heat shock elements (HSEs) in the heat shock protein gene promoter regions (Refs 7, 8). In the unstressed state, HSF1 is present in the cytoplasm as a latent monomeric molecule that is unable to bind to DNA. Under stressful conditions, HSF1 is hyperphosphorylated in a *ras*-dependent manner by members of the mitogen-activated protein kinase (MAPK) subfamilies (e.g. ERK1, JNK/SAPK, p38 protein kinase; Refs 9, 10). HSF1 is converted to phosphorylated trimers with the capacity to bind DNA, and translocates from the cytoplasm to the nucleus (reviewed in Ref. 11; see Fig. 1). The signal that activates HSF1 is thought to be a flux of newly synthesised non-native proteins (Ref. 8). The consequences of HSF binding to its target, and the events that result in the ensuing transcription of heat shock genes, have been reviewed previously (Ref. 12).

The generation of heat shock proteins must be only transient, even if exposure to stress is over a prolonged period, as a continued presence of heat shock proteins would adversely influence protein homeostasis and a variety of intracellular functions. One mechanism by which the activity of HSF1 is regulated is via the binding of Hsp70 to its transactivation domain, thereby leading to repression of heat shock gene transcription (Ref. 13). The interaction between Hsp70 and

HSF1 has no effect on DNA binding or the stress-induced phosphorylation state of HSF1 (Ref. 13). A second mechanism that regulates heat shock protein synthesis is the interaction between heat shock protein binding factor 1 (HSBP1) and the active trimeric form of HSF1 and Hsp70, thereby inhibiting the capacity of HSF1 to bind to DNA (Ref. 14). HSBP1 is predominantly localised in the nucleus and levels of HSBP1 mRNA have been shown to be present at high levels in a variety of cell lines and animal tissues and to be unaffected by heat shock (Ref. 14).

This article reviews the current literature on heat shock proteins and their influence on pathogenic processes such as autoimmunity, organ allograft rejection and vascular disease. It highlights the evolving evidence that rather than being exclusively intracellular, heat shock proteins are present in, and can be released into, the extracellular compartment under physiological conditions and elicit a range of functions. Given the functional versatility of heat shock proteins and their capacity to mediate both induction and regulation of immunity, further studies aimed at understanding the mechanisms underlying these functions might reveal strategies by which heat shock proteins can be used as therapeutic agents either for the generation of protective immunity or for the downregulation of deleterious inflammatory conditions.

Heat shock proteins as molecular chaperones

Molecular chaperones are defined as 'proteins that assist the correct non-covalent assembly of other protein-containing structures *in vivo* but are not permanent components of these structures when they are performing their normal biological functions' (Ref. 15). An alternative definition is that 'a molecular chaperone is a protein that binds to and stabilises an otherwise unstable conformer of another protein, and by controlled binding and release of the substrate protein facilitates its correct fate *in vivo*, be it folding, oligomeric assembly, transport to another subcellular compartment, or controlled switching between active/inactive conformations' (Ref. 16). The precise nature of peptide binding and the factors involved appear to be chaperone-dependent. Whereas ATP binding is important for the release of peptides from Hsp70, BiP (immunoglobulin heavy chain binding protein; reviewed in Ref. 17) and Hsp90 (Ref. 18), its role




Figure 1. Regulation of transcription of heat shock protein genes by heat shock factor. Heat shock factor (HSF) is present in the cytoplasm as a latent monomeric molecule that is unable to bind to DNA. Under stressful conditions, the flux of non-native proteins (which are non-functional, prone to aggregation, protease-sensitive, and bind to chaperones) leads to phosphorylation (P) and trimerisation of HSF. The trimers translocate to the nucleus, bind the promoter regions of heat shock protein (hsp) genes and mediate hsp gene transcription. The activity of HSF trimers is downregulated by hsp (e.g. Hsp70) and the heat shock binding protein 1 (HSBP1) that is found in the nucleus. Diagrams are based on those included in Refs 11 and 14 (fig001gps).

in peptide binding to and unloading from gp96, the endoplasmic reticulum parologue of Hsp90, is unclear (Ref. 19). Not all stress proteins function as molecular chaperones; however, those that do fulfil an essential intracellular role and, in the extracellular compartment, have the capacity to mediate the induction of peptide-specific immunity, as is described later.

Potential pathogenic role for heat shock proteins

Heat shock proteins, particularly those of the Hsp60 and Hsp70 families, are immunodominant molecules, and a significant element of the immune response to pathogenic microorganisms is directed towards peptides derived from heat shock proteins (Refs 20, 21). This phenomenon, together with the ubiquitous nature of human heat shock proteins and the high degree of sequence homology between mammalian and

bacterial heat shock protein cognates (~50–60% identical residues in the case of the Hsp60 family) has led to debate as to whether the immune system recognises heat shock proteins as dominant microbial antigens or potentially harmful self-antigens (Ref. 20). It has also been suggested that heat shock proteins might provide a link between infection and autoimmunity, either through recognition of conserved epitopes or via cross-reactivity / molecular mimicry (Ref. 22). Evidence for a link between heat shock protein reactivity and disease pathogenesis, particularly autoimmune disease, vascular disease and organ allograft rejection, has arisen from several studies.

Heat shock proteins and heat shock protein reactivity in autoimmune disease *Hsp60*

Several investigations have implicated Hsp60 and immune reactivity to members of the

Hsp60 family in autoimmune diseases, the best studied of which are arthritis and diabetes. Hsp60 is expressed in the synovial tissue of patients with rheumatoid arthritis (RA) and juvenile chronic arthritis (Ref. 23), and T cells derived from the synovial fluid are activated by mycobacterial Hsp65 (Refs 24, 25). T-cell reactivity to self-Hsp60 has been reported in patients with RA (Ref. 26); immortalised B cells from the synovial tissue of RA patients show specificity for bacterial Hsp60 (Ref. 27); and elevated levels of circulating antibodies to Hsp60 are present in children with juvenile chronic arthritis (Ref. 28). T-cell-mediated responses to mycobacterial Hsp65 have also been implicated in experimental models of arthritis, and disease can be initiated in rats by the transfer of T-cell clones specific for mycobacterial Hsp65 (Ref. 29). In addition, antibodies to Hsp65 are elevated in mice with pristane-induced arthritis (Ref. 30).

Evidence of a role for Hsp60 in type 1 diabetes [insulin-dependent diabetes mellitus (IDDM)] is somewhat equivocal (Ref. 31). Supporting such a role is evidence that naïve T cells from non-obese diabetic (NOD) mice can be activated by both self-Hsp60 and mycobacterial Hsp60 (Ref. 32), that anti-Hsp60 T cells can mediate insulitis and hyperglycaemia in the NOD mouse (Ref. 33), and that peripheral blood T cells from patients with IDDM demonstrate a heightened proliferative response to human Hsp60 and Hsp60 peptides (Ref. 34). However, in NOD mice, immunity to autoantigens other than heat shock proteins, such as glutamic acid decarboxylase 65 (GAD), appears much earlier than responsiveness to mycobacterial Hsp65 (Ref. 35), thereby arguing against an essential role for heat shock proteins in disease induction in this model. In addition, no evidence for serological immunity to islet cell heat shock proteins has been reported in IDDM (Ref. 36).

Evidence of a role for Hsp60 in the pathogenesis of multiple sclerosis (MS) is less apparent. Hsp60 expression has been identified in chronic MS plaques (Refs 37, 38), and a humoral response to Hsp60 has been detected in the cerebrospinal fluid of patients with MS; however, the latter is not specific for MS and is also present in a number of chronic degenerative conditions (Ref. 39). Nor is peripheral blood lymphocyte reactivity to Hsp60 altered in MS patients (Ref. 40).

Hsp70

In contrast to the findings for Hsp60, Hsp70 has been implicated as a potential autoantigen in MS (Refs 41, 42). In IDDM, the preferential expression of Hsp70 by β cells, but not α cells, in the islets of Langerhans might be important for the understanding of autoimmune destruction of β cells in this disease (Ref. 43). Autoantibodies to the constitutive form of Hsp70 (Hsc70) have been identified in a proportion of patients with primary biliary cirrhosis (45.7%) and patients with autoimmune hepatitis patients (52.9%), but not in patients with chronic hepatitis B or C infection (Ref. 44). Reactivity to Hsp70 has also been implicated in the induction of disease in toxin-induced interstitial nephritis (Ref. 45).

Heat shock proteins and heat shock protein reactivity in transplantation

In addition to autoimmune disease, heat shock proteins and reactivity to heat shock proteins have been associated with allograft rejection. Heat shock proteins are induced during graft preservation, ischaemia-reperfusion and surgery (Refs 46, 47, 48), and by the inflammatory process of the rejection response, including the localised production of cytokines by infiltrating leukocytes (Ref. 49). In rats, Hsp70 gene and protein expression are increased in rejecting cardiac allografts, and graft-infiltrating lymphocytes proliferate in response to recombinant mycobacterial Hsp65 and Hsp71 (Refs 50, 51, 52, 53). Heat shock protein expression is also induced in the intestinal epithelium and lamina propria after rat small-bowel transplantation and appears, in part at least, to be resistant to immunosuppression with tacrolimus (Ref. 54). In humans, heat shock protein expression is increased in rejecting lungs (Ref. 55); T cells from rejected renal grafts respond to Hsp72 (Ref. 56); and mycobacterial Hsp65-induced growth of graft-infiltrating lymphocytes from endomyocardial biopsies correlates with cardiac graft rejection (Ref. 52).

These findings have led to the proposition that heat shock protein expression in allograft tissue induces heat shock protein reactivity, thereby promoting the development of acute and chronic graft rejection (Refs 57, 58, 59). However, heat shock proteins are cytoprotective molecules and their induction in the peri- and immediate post-transplantation periods is likely to be a protective response targeted towards the

maintenance of cell and tissue integrity. This is supported by reports that heat shock proteins attenuate preservation and ischaemia-reperfusion injury (Refs 57, 60, 61, 62, 63), and that they protect endothelial cells from neutrophil-mediated necrosis (Ref. 64) and a variety of cell types from oxidative injury (Refs 65, 66, 67). In addition, lower levels of Hsp70 in pre-liver-transplant biopsies and organ perfusates are associated with early graft loss (Ref. 68).

The precise influence of heat shock proteins on allograft survival is currently unclear. A direct involvement of Hsp60 in the rejection process has been suggested by the observations that skin from transgenic mice overexpressing Hsp60 transplanted into allogeneic recipients is rejected more rapidly than skin transplanted from wild-type donors (Ref. 69). By contrast, skin transplanted into Hsp60-transgenic mice, in which spontaneous autoimmunity to Hsp60 is reduced, is rejected more slowly than skin grafted into wild-type recipients (Ref. 69). However, further studies are required to define more clearly a direct role for heat shock proteins in the induction and progression of allograft rejection.

Heat shock proteins and heat shock protein reactivity in vascular disease

It is now apparent that there is an inflammatory component to vascular disease that involves the accumulation of monocytes and activated T cells in atherosclerotic lesions and the localised presence of pro-inflammatory cytokines (Refs 70, 71). Evidence also suggests that the immunological component of the development of atherosclerosis might, at least in part, involve the expression of, and reactivity to, heat shock proteins (Ref. 70). The evidence for this proposition has arisen from three findings. First, the intensity of heat shock protein expression positively correlates with the severity of atherosclerosis; second, there is a localised enrichment of $\gamma\delta$ T cells in the lesion (Ref. 72), and this is of particular interest given the capacity of $\gamma\delta$ T cells to directly recognise and respond to autologous heat shock proteins (Ref. 73); and third, immunisation with recombinant mycobacterial Hsp65 can induce atherosclerotic lesions in normocholesterolaemic rabbits (Ref. 74).

A role for Hsp60 in the induction of the inflammatory response that characterises

atherosclerosis has also been suggested. Lipid-laden cells formed from the uptake of oxidised low-density lipoprotein (LDL), via a 'scavenger' receptor that does not recognise native LDL, are a principal component of the atherosclerotic plaque. Exposure of the monocytic cell lines U937 and HL60 to oxidised LDL induces marked expression of Hsp60 (Ref. 75). These findings suggest that the inflammatory response associated with atherosclerosis might in part be promoted by the activation of T cells reactive with the Hsp60 that is expressed on monocytes within the lesion or released locally. Localised expression of heat shock proteins might also be influenced by haemodynamic factors, as raised blood pressure has direct effects on the vasculature (Refs 76, 77) and vessels subjected to greater mechanical and shear stress express heat shock proteins and are more prone to the development of atherosclerosis (Refs 72, 77, 78, 79).

Humoral responses to heat shock proteins have also been implicated in vascular disease. Elevated levels of circulating antibody to the mycobacterial 65 kDa heat shock protein have been reported in carotid atherosclerosis (Ref. 80), coronary heart disease (Ref. 81) and borderline hypertension (Ref. 82). Levels of antibodies to human Hsp60 are also raised in peripheral vascular disease (Ref. 83). The *in vivo* physiological significance of antibodies to heat shock proteins in the pathogenesis of vascular disease has yet to be clearly established. However, they have been shown to mediate endothelial cell cytotoxicity (Ref. 79), and the observation that anti-Hsp65/60 antibodies in individuals with atherosclerosis recognise three distinct, self-Hsp65/60 sequences might implicate them in the initiation of atherosclerosis via an autoimmune-type mechanism (Ref. 84). Levels of anti-Hsp65 antibodies might have diagnostic value as titres have recently been shown to predict the 5-year mortality of patients with carotid atherosclerosis (Ref. 85).

Heat shock proteins in normal aging

Increasing age is associated with a reduced capacity to maintain homeostasis in all physiological systems and it might be that this results, in part at least, from a parallel and progressive decline in the ability to produce heat shock proteins. If this is so, an attenuated heat shock protein response could contribute to the increased susceptibility to environmental

challenges and the more prevalent morbidity and mortality seen in aged individuals (Refs 86, 87).

In vitro studies have shown that Hsp70 expression in heat-stressed lung cells (Ref. 88), hepatocytes and liver (Refs 89, 90), splenocytes (Ref. 91), myocardium (Ref. 92) and mononuclear cells is reduced with increasing age (Ref. 86), as is the induction of Hsp70 expression in response to ischaemia (Ref. 93) and mitogenic stimulation (Ref. 94). Hsp70 gene expression declines during normal aging in human retina (Ref. 95), and heat shock-induced Hsp70 expression is decreased in senescent and late-passage cells, both of which suggest that the process of aging itself might be associated with reduced Hsp70 production (Refs 96, 97, 98).

Although currently uncertain, possible mechanisms underlying an attenuated stress response during aging might include a reduced availability of HSF (Ref. 86) or age-associated increases in abnormal or denatured proteins that could interfere with HSF binding to HSEs (Ref. 99). Alternatively, age-related decreases in the capacity of HSF to undergo the oligomerisation that is essential for binding to HSEs might be involved.

Heat shock protein release

Heat shock proteins are typically regarded as intracellular molecules; however, it is now apparent that heat shock proteins can be released into the extracellular compartment. It was reported in the late 1980s that heat shock proteins could be released from cultured rat embryo cells (Ref. 100). Heat treatment of the cells broadened the spectrum of proteins released, from a small set of proteins including Hsc70 to a larger set including Hsp70 and Hsp110. It was suggested that the release of heat shock proteins might have resulted from changes in pH and gas tension, disruption of the diffusion layer at the cell surface or mechanical stresses associated with in vitro manipulations (Ref. 100). The release of heat shock proteins did not appear to be mediated via the common secretory pathway, as it was not blocked by the inhibitors colchicine and monensin (Ref. 100). Nor did it result from cell lysis, as exposure of cells to low concentrations of non-ionic detergents indicated that Hsp70 is not readily released from damaged cells. Instead, a selective mechanism has been suggested. Evidence cited in favour of this is the fact that Hsp70 synthesised in the presence of the lysine amino acid analogue

aminoethyl cysteine was not released from cells, probably due to an altered structure or function preventing its correct interaction with the specific release mechanism (Ref. 100). The precise mechanism(s) by which heat shock proteins are actively released by viable cells has yet to be elucidated.

Subsequent studies have shown heat shock proteins to be released from a variety of cells including cultured human islet cells (Ref. 101), rat glial cells and a human neuroblastoma cell line (Ref. 102), as well as cultured vascular smooth muscle cells exposed to reactive oxygen species (Ref. 103). In these studies, release did not appear to be a result of cellular necrosis. Myocardial injury induces Hsp60 release from rat hearts in organ culture; however, this most probably resulted from myocardial necrosis (Ref. 104). A selective release of heat shock proteins from necrotic, but not apoptotic, cells has also been described (Ref. 105). The physiological basis for heat shock protein release from intact (non-necrotic) cells has yet to be fully understood. Glia-axon transfer proteins, which include Hsp70, Hsc70 and Hsp100, are transferred from adjacent glial cells to the squid giant axon (Ref. 106), and heat shock protein release might be an altruistic response on the part of one cell for the protection of adjacent cells (Ref. 100).

Hsp60 and Hsp70 can be detected in the serum of normal individuals (Refs 107, 108), and in keeping with a reduced capacity to generate stress responses with aging, serum levels of heat shock proteins also decline with age (Ref. 109). Elevated levels of heat shock proteins have been observed in subjects with borderline hypertension (Ref. 110), as well as in patients with peripheral and renal vascular disease (Ref. 83). It is interesting to note that, in the borderline hypertension study, levels of Hsp60 correlated with the presence of atherosclerosis (Ref. 110), and a similar finding in a population-based study of clinically normal subjects has been reported (Ref. 111). Hsp70 release into the serum following myocardial infarction has also been reported in patients who have experienced preceding angina (Ref. 112).

Although it is clear that heat shock proteins are present and can be released into the peripheral circulation in response to several conditions, the physiological role of these proteins has yet to be defined. The identification of heat shock proteins and antibodies directed against heat shock proteins in normal individuals (Refs 107,


108, 111) indicates that their presence is not limited to disease. The emerging evidence that stress proteins can interact with cell-surface receptors and elicit a range of biological activities including the downregulation of autoimmune disease (see below) suggests that they might be involved in regulating immunity to the ubiquitously expressed and highly conserved heat shock proteins, and in the maintenance of the 'normal' state.

Heat shock proteins as immunomodulators and intercellular signalling molecules

Heat shock proteins have been shown to have a number of immunological effects (Fig. 2). Bacterial and mycobacterial heat shock proteins induce pro-inflammatory cytokine expression (Refs 113, 114, 115), and bacterial heat shock proteins induce intercellular cell adhesion molecule 1 (ICAM-1) and vascular cell adhesion molecule 1 (VCAM-1) expression on human vascular endothelial cells (Ref. 113). Chlamydial

and human Hsp60 activate human vascular endothelial cells to express E-selectin, ICAM-1 and VCAM-1, and activate vascular endothelial cells, smooth muscle cells and macrophages to secrete interleukin 6 (IL-6) (Ref. 116). With kinetics similar to those induced by lipopolysaccharide (LPS), mammalian Hsp60 has also been demonstrated to induce a rapid release of tumour necrosis factor α (TNF- α) and nitric oxide from macrophages, as well as the expression of IL-12 and IL-15 (Ref. 117).

Evidence that heat shock proteins can elicit a range of biological and pro-inflammatory effects has stimulated interest in finding cell-surface receptors for these molecules. The existence of specific receptors for heat shock proteins was initially confirmed in studies demonstrating that the presentation of heat shock protein-associated peptides by major histocompatibility complex (MHC) class I molecules required receptor-mediated endocytosis (Refs 118, 119, 120), and the identities of receptors for heat shock proteins are now becoming apparent.



The heat shock protein Hsp60 is an intercellular signalling molecule

Expert Reviews in Molecular Medicine © 2001 Cambridge University Press

Figure 2. The heat shock protein Hsp60 as an intercellular signalling molecule. Hsp60 has been shown to have several immunological effects, including the induction of pro-inflammatory cytokine secretion from, and adhesion molecule expression on, a number of myeloid and vascular cell types, including smooth muscle cells. Abbreviations: ICAM-1, intercellular adhesion molecule 1; IL, interleukin; TNF- α , tumour necrosis factor α ; VCAM-1, vascular cell adhesion molecule 1 (fig002gps).

Human Hsp60 activates human peripheral blood mononuclear cells and monocytes through the CD14 antigen, using the signalling pathway also utilised by LPS (Ref. 121; see Fig. 3). Signalling is also mediated by the Toll-like receptor 4 (Ref. 122), which is an important mediator of innate immunity and LPS signalling in murine cells (Ref. 123). Hsp70 has also been shown to bind with high affinity to human monocytes and the CD14 molecule is involved in Hsp70-induced activation (Ref. 124). There appears to be a CD14-dependent interaction leading to intracellular calcium fluxes and the induction of pro-inflammatory cytokines (IL-1 β , IL-6, TNF- α), and a CD14-independent, but calcium-dependent, response that leads to TNF- α production (Ref. 124; see Fig. 3).

The cell-surface receptor for gp96 on dendritic cells (DCs) is downregulated as they undergo maturation (Ref. 125). This receptor has now been identified as the CD91 molecule (which is also known as the α 2-macroglobulin receptor or the LDL-related protein), to which it binds directly (Ref. 126). In addition to gp96,

the CD91 molecule has since been shown to be a common receptor for Hsp70, Hsp90 and calreticulin, all of which can mediate the induction of peptide-specific immunity as summarised below (Ref. 127).

It is now clear that heat shock proteins have an intercellular signalling role as well as a chaperone function, and Asea and colleagues have coined the term 'chaperokine' for these ubiquitously expressed and versatile families of molecules (Ref. 124).

The potential therapeutic value of heat shock proteins

Some of the earliest evidence that heat shock proteins might have a therapeutic potential arose from the observations that exogenous members of the Hsp70 family protect spinal sensory neurons from axotomy-induced death and cultured aortic cells from heat stress (Refs 128, 129). Subsequent work demonstrated that exogenous Hsp70 could also protect rabbit arterial smooth muscle cells subjected to serum




Figure 3. The heat shock proteins Hsp60 and Hsp70 induce pro-inflammatory cytokine secretion from monocytes. Hsp60 induces several pro-inflammatory cytokines, and the signalling pathways responsible are yet to be fully clarified. It induces the secretion of interleukin 6 (IL-6) from human monocytes via signalling through CD14 and p38 mitogen-activated protein kinase, and binds to the Toll-like receptor 4 complex (Tlr4), for which CD14 is a co-receptor, to induce the expression of TNF- α and nitric oxide. It also induces the expression of a range of cytokines, including IL-12 and IL-15, through as yet uncharacterized pathways. Hsp70 acts through a CD14-dependent pathway to stimulate IL-1 β , IL-6 and TNF- α production, and also a CD14-independent pathway that leads to TNF- α production, suggesting that CD14 is also a co-receptor for an as yet uncharacterised Hsp70 receptor. Both Hsp70 pathways are calcium dependent (fig003gps).

deprivation, by a mechanism that involved cell association but not internalisation (Ref. 130).

There are a number of indications in which heat shock proteins might be of therapeutic value. To date, the majority of studies have focused on either their capacity to regulate inflammatory responses in autoimmune disease or their ability to induce peptide-specific immune responses against tumours and pathogenic organisms.

Modulation of inflammatory disease

Despite the association of heat shock protein expression and heat shock protein reactivity with autoimmunity, several observations question the proposition that self-heat shock protein reactivity has a direct pro-inflammatory role in autoimmune disease. Although the literature on the subject is less comprehensive, the situation might also be the same in the case of transplantation.

The normal T-cell repertoire includes cells reactive against autologous heat shock proteins (Refs 20, 131, 132). Although heat shock proteins have been considered to be intracellular proteins and therefore normally shielded from self-reactive T cells, it is now known that they are released from a variety of normal cells in culture (Refs 100, 101, 102, 103), expressed on the cell surface (Ref. 133) and present in the peripheral circulation of normal individuals (Refs 107, 108, 111). As discussed below, it appears that T-cell reactivity to self-heat shock proteins is a protective phenotype, and it is interesting to note that peripheral blood T-cell responsiveness to self- and non-self-heat shock proteins segregates with their expression of CD45 isotypes. Human Hsp60 activates CD45RA⁺RO⁻ (naïve) T cells, bacterial-specific peptides activate CD45RA⁻RO⁺ (memory) T cells and bacterial Hsp60 activates both CD45RA⁺RO⁻ and CD45RA⁻RO⁺ T cells (Ref. 132). The observation that both types of T-cell subset are activated by bacterial Hsp60 indicates that T cells can recognise and respond to conserved (self) epitopes on the whole bacterial molecule. What is currently not known is the cytokine-secreting profiles of cells responding to the different heat shock proteins or specific peptides derived from them. This is of particular importance given the evidence from autoimmune disease that self-heat shock protein reactivity appears to induce a regulatory phenotype, whereas reactivity to non-self induces a pro-inflammatory phenotype, as is discussed below.

Autoimmune disease

In contrast to their proposed capacity to promote pathogenic processes such as autoimmune disease, T-cell reactivity to heat shock proteins can also protect against disease, as demonstrated by the capacity of Hsp60 and Hsp70 to downregulate autoimmune disease (Refs 30, 134, 135, 136, 137, 138, 139, 140, 141).

An insight into the possible mechanisms by which self-heat shock proteins might modulate autoimmune disease has come from the work of de Graeff-Meeder and co-workers (Ref. 134). In patients with juvenile chronic arthritis, in whom the disease follows a relapsing-remitting rather than progressive course, the presence of circulating T cells responsive to human (self) Hsp60 was beneficial. These T cells were of the regulatory T helper 2 (Th2) phenotype, whereas T cells reactive with the 65 kDa mycobacterial antigen Hsp65 displayed the inflammatory Th1 phenotype and their presence correlated with disease severity (Ref. 142). It has also been shown that stimulation of T cells from the synovial fluid of RA patients with human, but not bacterial, Hsp60 can stimulate regulatory responses (Ref. 136). The apparent capacity of self-heat shock protein to modulate autoimmune disease in the clinical situation confirms data indicating that protection by Hsp60 in experimental autoimmune disease appears to be elicited by autoreactive T cells recognising specific sequences of self-stress proteins (Refs 140, 141). The ability of Hsp60 peptides to modulate adjuvant arthritis appears to reside in the capacity of induced regulatory T cells to produce IL-10, as well as IL-4 and interferon γ (IFN- γ) (Ref. 143).

Members of the Hsp70 family can also elicit protection from autoimmune disease, and Hsp71 from *Mycobacterium tuberculosis* can modulate experimental rat arthritis (Ref. 139). In a similar way to Hsp60, the capacity of peptides from mycobacterial Hsp70 to protect against the subsequent induction of adjuvant arthritis in Lewis rats appears to be mediated via the production of suppressive cytokines, including IL-10 (Refs 144, 145).

Allograft immunity

The capacity of heat shock proteins to modify allograft rejection responses is less well defined; however, immunising recipient animals with self-Hsp60, or Hsp60 peptides that have the capacity to shift Hsp60 reactivity from a Th1 to

a Th2 phenotype, can delay murine skin allograft rejection (Ref. 69). These studies indicate that rather than being pro-inflammatory, self-Hsp T-cell reactivity could be part of a normal immunoregulatory T-cell response that has the potential to control inflammatory disease (Ref. 146).

Induction of peptide-specific immunity

In addition to their capacity to downregulate pro-inflammatory conditions the potential value of heat shock proteins for inducing protective immunity has been explored by several groups, primarily in the areas of tumour immunity and infectious disease.

Tumour immunity


It has been known for some time that heat shock proteins bind peptide (Refs 4, 147) and that heat shock proteins purified from cells chaperone a large number of peptides derived from the cells from which they are isolated – the so-called ‘antigenic repertoire’ of that cell (Ref. 148). Early studies showed that fractionated tumour cell lysates have the capacity to reduce tumour cell growth in mice (Ref. 149). Since then, it has been well established that immunisation of mice with Hsp70, Hsp90 and gp96 isolated from murine tumour cells induces anti-tumour immunity and tumour-specific cytolytic T cells, and that the immunity results from tumour-derived peptides associated with the heat shock protein rather than from the heat shock proteins themselves (Ref. 150). More recently, it has been reported that calreticulin, Hsp110 and grp170 can also be used in heat shock protein-based cancer immunotherapy (Refs 151, 152). The finding that the immunological properties of heat shock proteins and the capacity of Hsp70 and gp96 to induce tumour protection as shown in rodent models are also observed in amphibians (*Xenopus*) (Ref. 153) indicates the evolutionary conserved nature of these functions, and strongly supports the successful translation of these strategies into the clinical environment. In that regard, preliminary clinical trials have demonstrated the induction of cancer-specific CD8⁺ T-cell responses in 6/12 patients immunised with gp96-peptide complexes prepared from their own tumour (Ref. 154). Clearly, the capacity of tumour-derived heat shock proteins to induce specific and protective immunity might have profound

effects on the treatment and management of patients with malignant disease.

However, the immunological effects of heat shock proteins purified from tumour cells have a dual nature. The induction of immunity to methylcholanthrene-induced fibrosarcoma by the administration of gp96 purified from the tumour displays a consistent dose restriction: two intradermal administrations of <1 µg gp96 is ineffective; two doses of 1 µg induce immunity and provide optimal protection against tumour growth; and two doses of 10 µg do not protect (Ref. 155). The lack of protection at high doses of tumour-derived gp96 is an active, antigen-specific downregulation of tumour-specific immunity that can be adoptively transferred by CD4⁺ T cells purified from animals treated with high doses of tumour-derived gp96 (Ref. 155). These findings are exciting as they suggest that immunisation with heat shock proteins that are chaperoning clinically relevant peptides might be an effective strategy for downregulating several diseases including autoimmunity. In that regard, gp96 purified from liver and pancreas of C57/B6 mice has been shown to elicit protection from autoimmune damage in NOD mice that is long term and can be adoptively transferred (Ref. 156).

The mechanisms by which heat shock proteins can mediate peptide-specific immunity are yet to be clearly defined. Antigen-presenting cells (APCs) such as DCs and monocytes play a key role, as they have been shown to internalise heat shock proteins spontaneously by receptor-mediated endocytosis via the CD91 receptor, and direct chaperoned proteins/peptides into the intracellular pathway for MHC class I-restricted presentation to CD8⁺ T cells, concomitant with the induction of DC maturation and cytokine secretion (Refs 118, 126, 127; see Fig. 4). It is interesting to note that α2-macroglobulin, the originally described ligand for CD91, is also able to channel exogenous antigens into the endogenous pathway of antigen presentation via the same receptor (Ref. 157).

The mechanism by which high doses of heat shock protein can induce immunoregulation is also unclear. In addition to inducing DC maturation and the expression of antigen-presenting and co-stimulatory molecules (Refs 105, 125), gp96 promotes the accumulation of DCs into the draining lymph node (Ref. 158). It might be that larger quantities of gp96 lead to a



The heat shock protein gp96 delivers antigenic peptides and maturation signals to antigen-presenting cells, and induces release of cytokines

Expert Reviews in Molecular Medicine ©2001 Cambridge University Press

Figure 4. The heat shock protein gp96 delivers antigenic peptides and maturation signals to antigen-presenting cells, and induces release of cytokines. (a) gp96-peptide complexes bind to CD91 and are taken up by dendritic cells via receptor-mediated endocytosis. (b) Peptides carried on gp96 are thus delivered to the major histocompatibility complex (MHC) class I presentation pathway and are (c) re-presented on the cell surface in association with MHC class I antigens for recognition by antigen-specific CD8⁺ T cells via the T-cell receptor (TCR) and associated molecules. (d) gp96 also delivers maturation signals to the dendritic cells, and induces the expression of MHC antigens, co-stimulatory molecules such as B7 (which binds to CD28), and intercellular adhesion molecule 1 (ICAM-1). This, combined with (e) the induction of pro-inflammatory cytokines, promotes the generation of immune responses to gp96-chaperoned peptides. Abbreviations: GM-CSF, granulocyte-macrophage colony-stimulating factor; IL-1, interleukin 1, TNF- α , tumour necrosis factor α (**fig004gps**).

greater APC-mediated cytokine signal and the preferential induction of regulatory CD4⁺ T cells (Ref. 155).

Induction of immunity to infectious agents

The capacity of heat shock proteins to chaperone antigenic repertoires and induce specific immunity to them has led to studies evaluating whether the administration of heat shock proteins from virally transformed cells, or cells infected by pathogenic organisms, would induce specific immunity (Ref. 159). This has been shown to be the case, and specific immunity has been induced by the administration of heat

shock proteins isolated from SV40-transformed and influenza-infected cells (Refs 160, 161). Peptide-specific cytolytic T cells and protective anti-viral immunity can also be induced by immunising mice with a mixture of gp96 or Hsp70 reconstituted with specific cytotoxic T lymphocyte epitopes from SV40, influenza virus and lymphocytic choriomeningitis virus (Refs 162, 163, 164). An alternative approach is to covalently link appropriate antigens to heat shock proteins; indeed, the immunisation of mice with the human immunodeficiency virus 1 (HIV-1) p24 protein covalently linked to mycobacterial Hsp70 elicits antibody, cytokine and lymphocyte proliferative responses (Ref. 165). Covalent linking is not




Figure 5. Potential therapeutic applications for heat shock proteins. The potential therapeutic value of heat shock proteins (Hsps) purified from appropriate tissues lies in their capacity to induce pro-inflammatory responses at low concentrations and induce regulatory immunity at high doses. The observations that recognition of non-self-heat shock proteins leads to inflammatory responses, whereas the recognition of conserved epitopes induces regulatory responses, indicate that the administration of appropriate heat shock protein peptides might have clinical efficacy in several conditions (**fig005gps**).

necessary for the induction of immunity, as non-covalently bound MHC class II influenza virus peptide can also induce immune reactivity to the Hsp70-binding peptide (Ref. 166).

Summary

There are clearly many aspects of heat shock protein biology that remain puzzling. On the one hand, reactivity to heat shock proteins appears to be associated with several pathological disease states, yet, on the other hand, heat shock proteins are ubiquitously expressed and reactivity to self-derived molecules can confer protection against a number of pro-inflammatory conditions. Future work will need to translate the experimental data on the capacity of heat shock proteins to induce tumour protection and immunity to infectious agents into the clinical environment and more fully evaluate the mechanisms by which these effects are induced and regulated. The observations that heat shock proteins can be released and that they can directly or indirectly elicit potent immunoregulatory activities give a new perspective on the roles of heat shock proteins and anti-heat shock protein reactivity in autoimmunity, transplantation, vascular disease and other conditions. It is the qualitative nature of the response to heat shock proteins rather than its presence per se that is important, and

future experimental and clinical studies attempting to associate heat shock proteins with disease pathogenesis need to be designed to address these issues. It is also important to define definitively the specificity of any responses, so that the outcome can be attributed to self- or non-self-reactivity. By doing this, the contribution of infective agents to pathogenic processes such as autoimmunity and vascular disease can be truly evaluated.

Heat shock proteins are extremely versatile and potent molecules, the importance of which to biological processes is highlighted by the high degree to which their structure and function are phylogenetically conserved. Our knowledge of the physiological role of heat shock proteins is currently limited; however, a better understanding of their function and thereby the acquisition of the capacity to harness their power might lead to their use as therapeutic agents and revolutionise clinical practice in a number of areas (Fig. 5).

Acknowledgements and funding

I thank Dr Pramod Srivastava (University of Connecticut School of Medicine, Farmington, USA) and the anonymous reviewer for their critical evaluation of this article. I am grateful to the Wellcome Trust, UK, and Vasogen Incorporated, Canada, for financial support.

Work in the author's laboratory is supported in part by the National Heart, Lung and Blood Institute, USA (R21HL 69726-01).

References

- 1 Ritossa, F.A. (1962) A new puffing pattern induced by temperature shock and DNP in *Drosophila*. *Experientia* 18, 571-573
- 2 Tissières, A., Mitchell, H.K. and Tracy, U.M. (1974) Protein synthesis in salivary glands of *Drosophila melanogaster*: relation to chromosome puffs. *J Mol Biol* 84, 389-398., PubMed ID: 75214244
- 3 Hightower, L.E. (1991) Heat shock, stress proteins, chaperones, and proteotoxicity. *Cell* 66, 191-197, PubMed ID: 91309134
- 4 Gething, M.J. and Sambrook, J. (1992) Protein folding in the cell. *Nature* 355, 33-45, PubMed ID: 92114932
- 5 Lindquist, S. and Craig, E.A. (1988) The heat-shock proteins. *Annu Rev Genet* 22, 631-677, PubMed ID: 89192244
- 6 Welch, W.J. (1993) How cells respond to stress. *Sci Am* 268, 56-64, PubMed ID: 93242380
- 7 Voellmy, R. (1994) Transduction of the stress signal and mechanisms of transcriptional regulation of heat shock/stress protein gene expression in higher eukaryotes. *Crit Rev Eukaryot Gene Expr* 4, 357-401, PubMed ID: 95252683
- 8 Morimoto, R.I. et al. (1994) Regulation of heat shock gene transcription by a family of heat shock factors. In *The Biology of Heat Shock Proteins and Molecular Chaperones* (Morimoto, R., Tissières, A. and Georgopoulos, C., eds), pp. 417-455, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, USA
- 9 Knauf, U. et al. (1996) Repression of human heat shock factor 1 activity at control temperature by phosphorylation. *Genes Dev* 10, 2782-2793, PubMed ID: 97102678
- 10 Kim, J. et al. (1997) Analysis of the phosphorylation of human heat shock transcription factor-1 by MAP kinase family members. *J Cell Biochem* 67, 43-54, PubMed ID: 97469376
- 11 Morimoto, R.I. (1998) Regulation of the heat shock transcriptional response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes Dev* 12, 3788-3796, PubMed ID: 99088027
- 12 Wu, C. (1995) Heat shock transcription factors: structure and regulation. *Annu Rev Cell Dev Biol* 11, 441-469, PubMed ID: 96244627
- 13 Shi, Y., Mosser, D.D. and Morimoto, R.I. (1998) Molecular chaperones as HSF1-specific transcriptional repressors. *Genes Dev* 12, 654-666, PubMed ID: 98167909
- 14 Satyal, S.H. et al. (1998) Negative regulation of the heat shock transcriptional response by HSPB1. *Genes Dev* 12, 1962-1974, PubMed ID: 98315075
- 15 Ellis, J.R. (1996) Stress proteins as molecular chaperones. In *Stress Proteins in Medicine* (van Eden, W. and Young, D., eds), pp. 1-26, Marcel Dekker Inc, New York, USA
- 16 Hendrick, J.P. and Hartl, F.U. (1993) Molecular chaperone functions of heat-shock proteins. *Annu Rev Biochem* 62, 349-384, PubMed ID: 93356537
- 17 Bukau, B. and Horwich, A.L. (1998) The Hsp70 and Hsp60 chaperone machines. *Cell* 92, 351-366, PubMed ID: 98135762
- 18 Grenert, J.P., Johnson, B.D. and Toft, D.O. (1999) The importance of ATP binding and hydrolysis by hsp90 in formation and function of protein heterocomplexes. *J Biol Chem* 274, 17525-17533, PubMed ID: 99292709
- 19 Wearsch, P.A. and Nicchitta, C.V. (1997) Interaction of endoplasmic reticulum chaperone GRP94 with peptide substrates is adenine nucleotide-independent. *J Biol Chem* 272, 5152-5156, PubMed ID: 97184170
- 20 Kaufmann, S.H. (1990) Heat shock proteins and the immune response. *Immunol Today* 11, 129-136, PubMed ID: 90253569
- 21 Young, R.A. (1990) Stress proteins and immunology. *Annu Rev Immunol* 8, 401-420, PubMed ID: 90262673
- 22 Lamb, J.R. et al. (1989) Stress proteins may provide a link between the immune response to infection and autoimmunity. *Int Immunol* 1, 191-196, PubMed ID: 91137505
- 23 Boog, C.J. et al. (1992) Two monoclonal antibodies generated against human hsp60 show reactivity with synovial membranes of patients with juvenile chronic arthritis. *J Exp Med* 175, 1805-1810, PubMed ID: 92268831
- 24 Res, P.C. et al. (1988) Synovial fluid T cell reactivity against 65 kD heat shock protein of mycobacteria in early chronic arthritis. *Lancet* 2, 478-480, PubMed ID: 88301688
- 25 Pope, R.M., Lovis, R.M. and Gupta, R.S. (1992) Activation of synovial fluid T lymphocytes by 60-kd heat-shock proteins in patients with inflammatory synovitis. *Arthritis Rheum* 35,

- 43-48, PubMed ID: 92118137
- 26 De Graeff-Meeder, E.R. et al. (1991) Recognition of human 60 kD heat shock protein by mononuclear cells from patients with juvenile chronic arthritis. *Lancet* 337, 1368-1372, PubMed ID: 91245862
- 27 Krenn, V. et al. (1996) Immortalized B-lymphocytes from rheumatoid synovial tissue show specificity for bacterial HSP 60. *Virchows Arch* 427, 511-518, PubMed ID: 96211701
- 28 de Graeff-Meeder, E.R. et al. (1993) Antibodies to human HSP60 in patients with juvenile chronic arthritis, diabetes mellitus, and cystic fibrosis. *Pediatr Res* 34, 424-428, PubMed ID: 94077613
- 29 van Eden, W. et al. (1988) Cloning of the mycobacterial epitope recognized by T lymphocytes in adjuvant arthritis. *Nature* 331, 171-173, PubMed ID: 88122568
- 30 Thompson, S.J. et al. (1990) Autoimmune reactions to heat-shock proteins in pristane-induced arthritis. *Eur J Immunol* 20, 2479-2484, PubMed ID: 91071287
- 31 Roep, B.O. (1996) T-cell responses to autoantigens in IDDM. The search for the Holy Grail. *Diabetes* 45, 1147-1156, PubMed ID: 96368706
- 32 Birk, O.S. et al. (1996) NOD mouse diabetes: the ubiquitous mouse hsp60 is a beta-cell target antigen of autoimmune T cells. *J Autoimmun* 9, 159-166, PubMed ID: 96328754
- 33 Elias, D. et al. (1991) Vaccination against autoimmune mouse diabetes with a T-cell epitope of the human 65-kDa heat shock protein. *Proc Natl Acad Sci U S A* 88, 3088-3091, PubMed ID: 91195297
- 34 Abulafia-Lapid, R. et al. (1999) T cell proliferative responses of type 1 diabetes patients and healthy individuals to human hsp60 and its peptides. *J Autoimmun* 12, 121-129, PubMed ID: 99158649
- 35 Tisch, R. et al. (1993) Immune response to glutamic acid decarboxylase correlates with insulitis in non-obese diabetic mice. *Nature* 366, 72-75, PubMed ID: 94050122
- 36 Atkinson, M.A. et al. (1991) No evidence for serological autoimmunity to islet cell heat shock proteins in insulin dependent diabetes. *J Clin Invest* 87, 721-724, PubMed ID: 91123473
- 37 Selmaj, K., Brosnan, C.F. and Raine, C.S. (1992) Expression of heat shock protein-65 by oligodendrocytes in vivo and in vitro: implications for multiple sclerosis. *Neurology* 42, 795-800, PubMed ID: 92228169
- 38 Raine, C.S. et al. (1996) Multiple sclerosis: a protective or a pathogenic role for heat shock protein 60 in the central nervous system? *Lab Invest* 75, 109-123, PubMed ID: 96294707
- 39 Gao, Y.L., Raine, C.S. and Brosnan, C.F. (1994) Humoral response to hsp 65 in multiple sclerosis and other neurologic conditions. *Neurology* 44, 941-946, PubMed ID: 94247671
- 40 Salvetti, M. et al. (1992) T-lymphocyte reactivity to the recombinant mycobacterial 65- and 70-kDa heat shock proteins in multiple sclerosis. *J Autoimmun* 5, 691-702, PubMed ID: 93143814
- 41 Salvetti, M. et al. (1996) The immune response to mycobacterial 70-kDa heat shock proteins frequently involves autoreactive T cells and is quantitatively deregulated in multiple sclerosis. *J Neuroimmunol* 65, 143-153, PubMed ID: 96268225
- 42 Battistini, L. et al. (1995) Gamma delta T cell receptor analysis supports a role for HSP 70 selection of lymphocytes in multiple sclerosis lesions. *Mol Med* 1, 554-562, PubMed ID: 96091379
- 43 Strandell, E. et al. (1995) Interleukin-1 beta induces the expression of hsp70, heme oxygenase and Mn-SOD in FACS-purified rat islet beta-cells, but not in alpha-cells. *Immunol Lett* 48, 145-148, PubMed ID: 96360146
- 44 Shingai, R. et al. (1995) Autoantibody against 70 kD heat shock protein in patients with autoimmune liver diseases. *J Hepatol* 23, 382-390, PubMed ID: 96116124
- 45 Weiss, R.A. et al. (1994) T cells reactive to an inducible heat shock protein induce disease in toxin-induced interstitial nephritis. *J Exp Med* 180, 2239-2250, PubMed ID: 95053758
- 46 Knowlton, A.A., Brecher, P. and Apstein, C.S. (1991) Rapid expression of heat shock protein in the rabbit after brief cardiac ischemia. *J Clin Invest* 87, 139-147, PubMed ID: 91086440
- 47 Vass, K., Welch, W.J. and Nowak, T.S., Jr. (1988) Localization of 70-kDa stress protein induction in gerbil brain after ischemia. *Acta Neuropathol* 77, 128-135, PubMed ID: 89147508
- 48 Udelsman, R., Blake, M.J. and Holbrook, N.J. (1991) Molecular response to surgical stress: specific and simultaneous heat shock protein induction in the adrenal cortex, aorta, and vena cava. *Surgery* 110, 1125-1131, PubMed ID: 92081001

Heat shock proteins in health and disease: therapeutic targets or therapeutic agents?

- 49 Qian, J. et al. (1995) Expression of stress proteins and lymphocyte reactivity in heterotopic cardiac allografts undergoing cellular rejection. *Transpl Immunol* 3, 114-123, PubMed ID: 96103420
- 50 Baba, H.A. et al. (1997) Inducible heat shock protein 70 in rat cardiac allograft and its immunohistochemical localization in cardiac myocytes. *Transplantation* 64, 1035-1040, PubMed ID: 98018673
- 51 Mehta, N.K. et al. (1997) Heat shock protein 70 expression in native and heterotopically transplanted rat hearts. *J Surg Res* 70, 151-155, PubMed ID: 97388874
- 52 Moliterno, R. et al. (1995) Heat shock protein-induced T-lymphocyte propagation from endomyocardial biopsies in heart transplantation. *J Heart Lung Transplant* 14, 329-337, PubMed ID: 95298818
- 53 Moliterno, R. et al. (1995) Heat shock protein reactivity of lymphocytes isolated from heterotopic rat cardiac allografts. *Transplantation* 59, 598-604, PubMed ID: 95184337
- 54 Ogita, K. et al. (2000) Stress responses in graft and native intestine after rat heterotopic small bowel transplantation. *Transplantation* 69, 2273-2277, PubMed ID: 20324477
- 55 Rizzo, M. et al. (1998) Increased expression of HDJ-2 (heat shock protein 40) and heat shock protein 70 in biopsy specimens of transplanted human lungs. *J Heart Lung Transplant* 17, 241-249, PubMed ID: 98223064
- 56 Trieb, K. et al. (1996) T cells from rejected human kidney allografts respond to heat shock protein 72. *Transpl Immunol* 4, 43-45, PubMed ID: 96358149
- 57 Perdrizet, G.A. (1996) The heat shock response and organ transplantation. *Transplant Rev* 10, 78-98
- 58 Duquesnoy, R.J. and Moliterno, R. (1995) Role of heat shock protein immunity in transplantation. In *Stress Proteins in Medicine* (van Eden, W. and Young, D., eds), pp. 327-343, Marcel Dekker, New York, USA
- 59 Duquesnoy, R.J. et al. (1999) Evidence for heat shock protein immunity in a rat cardiac allograft model of chronic rejection. *Transplantation* 67, 156-164, PubMed ID: 99118786
- 60 Chen, G. et al. (1997) Induction of heat shock protein 72kDa expression is associated with attenuation of ischaemia-reperfusion induced microvascular injury. *J Surg Res* 69, 435-439, PubMed ID: 97367717
- 61 Hiratsuka, M. et al. (1998) Heat shock pretreatment protects pulmonary isografts from subsequent ischemia-reperfusion injury. *J Heart Lung Transplant* 17, 1238-1246, PubMed ID: 99098235
- 62 Hiratsuka, M. et al. (1999) Gene transfer of heat shock protein 70 protects lung grafts from ischemia-reperfusion injury. *Ann Thorac Surg* 67, 1421-1427, PubMed ID: 99281555
- 63 Okubo, S. et al. (2001) Gene transfer of heat-shock protein 70 reduces infarct size in vivo after ischemia/reperfusion in the rabbit heart. *Circulation* 103, 877-881, PubMed ID: 21112935
- 64 Wang, J.H. et al. (1995) Induction of heat shock protein 72 prevents neutrophil-mediated human endothelial cell necrosis. *Arch Surg* 130, 1260-1265, PubMed ID: 96094956
- 65 Gill, R.R. et al. (1998) Heat shock provides delayed protection against oxidative injury in cultured human umbilical vein endothelial cells. *J Mol Cell Cardiol* 30, 2739-2749, PubMed ID: 99144355
- 66 Amrani, M. et al. (1998) Relative induction of heat shock protein in coronary endothelial cells and cardiomyocytes: implications for myocardial protection. *J Thorac Cardiovasc Surg* 115, 200-209, PubMed ID: 98112894
- 67 Martin, J.L. et al. (1997) Small heat shock proteins and protection against ischemic injury in cardiac myocytes. *Circulation* 96, 4343-4348, PubMed ID: 98077384
- 68 Flohé, S. et al. (1998) Expression of HSP 70 as a potential prognostic marker for acute rejection in human liver transplantation. *Transpl Int* 11, 89-94, PubMed ID: 98222459
- 69 Birk, O.S. et al. (1999) The 60-kDa heat shock protein modulates allograft rejection. *Proc Natl Acad Sci U S A* 96, 5159-5163, PubMed ID: 99238499
- 70 Wick, G. et al. (1995) Role of heat shock protein 65/60 in the pathogenesis of atherosclerosis. *Int Arch Allergy Immunol* 107, 130-131, PubMed ID: 95337703
- 71 Frostegård, J. et al. (1999) Cytokine expression in advanced human atherosclerotic plaques: dominance of pro-inflammatory (Th1) and macrophage-stimulating cytokines. *Atherosclerosis* 145, 33-43, PubMed ID: 99355247
- 72 Kleindienst, R. et al. (1993) Immunology of atherosclerosis. Demonstration of heat shock protein 60 expression and T lymphocytes bearing alpha/beta or gamma/delta receptor in human atherosclerotic lesions. *Am J Pathol* 142, 1927-1937, PubMed ID: 93282523

- 73 Born, W. et al. (1990) Recognition of heat shock proteins and gamma delta cell function. *Immunol Today* 11, 40-43, PubMed ID: 90241398
- 74 Xu, Q. et al. (1992) Induction of arteriosclerosis in normocholesterolemic rabbits by immunization with heat shock protein 65. *Arterioscler Thromb* 12, 789-799, PubMed ID: 92313999
- 75 Frostegard, J. et al. (1996) Induction of heat shock protein in monocytic cells by oxidized low density lipoprotein. *Atherosclerosis* 121, 93-103, PubMed ID: 96262517
- 76 Hollander, W. et al. (1993) The effects of hypertension on cerebral atherosclerosis in the cynomolgus monkey. *Stroke* 24, 1218-1226; discussion 1226-1217, PubMed ID: 93342710
- 77 Sawchuk, A.P. et al. (1994) A prospective, in vivo study of the relationship between blood flow hemodynamics and atherosclerosis in a hyperlipidemic swine model. *J Vasc Surg* 19, 58-63; discussion 63-54, PubMed ID: 94133333
- 78 Solberg, L.A. and Strong, J.P. (1983) Risk factors and atherosclerotic lesions. A review of autopsy studies. *Arteriosclerosis* 3, 187-198, PubMed ID: 83203653
- 79 Schett, G. et al. (1995) Autoantibodies against heat shock protein 60 mediate endothelial cytotoxicity. *J Clin Invest* 96, 2569-2577, PubMed ID: 96292278
- 80 Xu, Q. et al. (1993) Association of serum antibodies to heat-shock protein 65 with carotid atherosclerosis. *Lancet* 341, 255-259, PubMed ID: 93148716
- 81 Hoppichler, F. et al. (1996) Changes of serum antibodies to heat-shock protein 65 in coronary heart disease and acute myocardial infarction. *Atherosclerosis* 126, 333-338, PubMed ID: 97057829
- 82 Frostegard, J. et al. (1997) Association of serum antibodies to heat-shock protein 65 with borderline hypertension. *Hypertension* 29, 40-44, PubMed ID: 97191119
- 83 Wright, B.H. et al. (2000) Elevated levels of circulating heat shock protein 70 (Hsp70) in peripheral and renal vascular disease. *Heart Vessels* 15, 18-22, PubMed ID: 20455119
- 84 Metzler, B. et al. (1997) Epitope specificity of anti-heat shock protein 65/60 serum antibodies in atherosclerosis. *Arterioscler Thromb Vasc Biol* 17, 536-541, PubMed ID: 97226205
- 85 Xu, Q. et al. (1999) Association of serum antibodies to heat-shock protein 65 with carotid atherosclerosis : clinical significance determined in a follow-up study. *Circulation* 100, 1169-1174, PubMed ID: 99414126
- 86 Richardson, A. and Holbrook, N.J. (1996) Aging and the cellular response to stress: reduction in the heat shock response. In *Cellular Aging and Cell Death* (Holbrook, N., Martin, G. and Lockshin, R., eds), pp. 67-79, Wiley-Liss, New York, USA
- 87 Shelton, D.N. et al. (1999) Microarray analysis of replicative senescence. *Curr Biol* 9, 939-945, PubMed ID: 99439863
- 88 Farnoli, J. et al. (1990) Decreased expression of heat shock protein 70 mRNA and protein after heat treatment in cells of aged rats. *Proc Natl Acad Sci U S A* 87, 846-850, PubMed ID: 90138923
- 89 Heydari, A.R., Conrad, C.C. and Richardson, A. (1995) Expression of heat shock genes in hepatocytes is affected by age and food restriction in rats. *J Nutr* 125, 410-418, PubMed ID: 95182192
- 90 Hall, D.M. et al. (2000) Aging reduces adaptive capacity and stress protein expression in the liver after heat stress. *J Appl Physiol* 89, 749-759, PubMed ID: 20384744
- 91 Pahlavani, M.A. et al. (1995) The expression of heat shock protein 70 decreases with age in lymphocytes from rats and rhesus monkeys. *Exp Cell Res* 218, 310-318, PubMed ID: 95255507
- 92 Gray, C.C. et al. (2000) Age dependence of heat stress mediated cardioprotection. *Ann Thorac Surg* 70, 621-626, PubMed ID: 20424091
- 93 Nitta, Y. et al. (1994) Diminished heat shock protein 70 mRNA induction in aged rat hearts after ischemia. *Am J Physiol* 267, H1795-1803, PubMed ID: 95068445
- 94 Faassen, A.E. et al. (1989) Diminished heat-shock protein synthesis following mitogen stimulation of lymphocytes from aged donors. *Exp Cell Res* 183, 326-334, PubMed ID: 89356886
- 95 Bernstein, S.L. et al. (2000) Heat shock cognate-70 gene expression declines during normal aging of the primate retina. *Invest Ophthalmol Vis Sci* 41, 2857-2862, PubMed ID: 20424936
- 96 Liu, A.Y. et al. (1989) Attenuated induction of heat shock gene expression in aging diploid fibroblasts. *J Biol Chem* 264, 12037-12045, PubMed ID: 89308612
- 97 Luce, M.C. and Cristofalo, V.J. (1992) Reduction in heat shock gene expression correlates with increased thermosensitivity in senescent human fibroblasts. *Exp Cell Res* 202, 9-16, PubMed ID: 92380258
- 98 Effros, R.B., Zhu, X. and Walford, R.L. (1994)

- Stress response of senescent T lymphocytes: reduced hsp70 is independent of the proliferative block. *J Gerontol* 49, B65-70, PubMed ID: 94172207
- 99 Munro, S. and Pelham, H. (1985) What turns on heat shock genes? *Nature* 317, 477-478, PubMed ID: 86014394
- 100 Hightower, L.E. and Guidon, P.T., Jr. (1989) Selective release from cultured mammalian cells of heat-shock (stress) proteins that resemble glial-axon transfer proteins. *J Cell Physiol* 138, 257-266, PubMed ID: 89139596
- 101 Child, D.F. et al. (1995) Heat shock protein studies in type 1 and type 2 diabetes and human islet cell culture. *Diabet Med* 12, 595-599, PubMed ID: 96024112
- 102 Bassan, M. et al. (1998) The identification of secreted heat shock 60-like protein from rat glial cells and a human neuroblastoma cell line. *Neurosci Lett* 250, 37-40, PubMed ID: 98359249
- 103 Liao, D.F. et al. (2000) Purification and identification of secreted oxidative stress-induced factors from vascular smooth muscle cells. *J Biol Chem* 275, 189-196, PubMed ID: 20085031
- 104 Schett, G. et al. (1999) Myocardial injury leads to a release of heat shock protein (hsp) 60 and a suppression of the anti-hsp65 immune response. *Cardiovasc Res* 42, 685-695, PubMed ID: 20003802
- 105 Basu, S. et al. (2000) Necrotic but not apoptotic cell death releases heat shock proteins, which deliver a partial maturation signal to dendritic cells and activate the NF- κ B pathway. *Int Immunol* 12, 1539-1546, PubMed ID: 20514086
- 106 Tytell, M., Greenberg, S.G. and Lasek, R.J. (1986) Heat shock-like protein is transferred from glia to axon. *Brain Res* 363, 161-164, PubMed ID: 86132619
- 107 Pockley, A.G., Shepherd, J. and Corton, J.M. (1998) Detection of heat shock protein 70 (Hsp70) and anti-Hsp70 antibodies in the serum of normal individuals. *Immunol Invest* 27, 367-377, PubMed ID: 99059555
- 108 Pockley, A.G. et al. (1999) Identification of human heat shock protein 60 (Hsp60) and anti-Hsp60 antibodies in the peripheral circulation of normal individuals. *Cell Stress Chaperones* 4, 29-35, PubMed ID: 99399023
- 109 Rea, I.M., McNerlan, S. and Pockley, A.G. (2001) Serum heat shock protein and anti-heat shock protein antibody levels in aging. *Exp Gerontol* 36, 341-352, PubMed ID: 21125081
- 110 Pockley, A.G. et al. (2000) Circulating heat shock protein 60 is associated with early cardiovascular disease. *Hypertension* 36, 303-307, PubMed ID: 20407283
- 111 Xu, Q. et al. (2000) Serum soluble heat shock protein 60 is elevated in subjects with atherosclerosis in a general population. *Circulation* 102, 14-20, PubMed ID: 20342853
- 112 Teckhoff, B.J. et al. (1996) Release of inducible heat shock protein 72 in acute myocardial infarction depends on preceding angina. *Circulation* 94 (Suppl.), I184
- 113 Galdiero, M., de l'Ero, G.C. and Marcatili, A. (1997) Cytokine and adhesion molecule expression in human monocytes and endothelial cells stimulated with bacterial heat shock proteins. *Infect Immun* 65, 699-707, PubMed ID: 97162346
- 114 Retzlaff, C. et al. (1994) Bacterial heat shock proteins directly induce cytokine mRNA and interleukin-1 secretion in macrophage cultures. *Infect Immun* 62, 5689-5693, PubMed ID: 95048814
- 115 Peetermans, W.E. et al. (1994) Mycobacterial heat-shock protein 65 induces proinflammatory cytokines but does not activate human mononuclear phagocytes. *Scand J Immunol* 39, 613-617, PubMed ID: 94278442
- 116 Kol, A. et al. (1999) Chlamydial and human heat shock protein 60s activate human vascular endothelium, smooth muscle cells, and macrophages. *J Clin Invest* 103, 571-577, PubMed ID: 99146972
- 117 Chen, W. et al. (1999) Human 60-kDa heat-shock protein: a danger signal to the innate immune system. *J Immunol* 162, 3212-3219, PubMed ID: 99192732
- 118 Arnold-Schild, D. et al. (1999) Cutting edge: receptor-mediated endocytosis of heat shock proteins by professional antigen-presenting cells. *J Immunol* 162, 3757-3760, PubMed ID: 99218387
- 119 Singh-Jasuja, H. et al. (2000) Cross-presentation of glycoprotein 96-associated antigens on major histocompatibility complex class I molecules requires receptor-mediated endocytosis. *J Exp Med* 191, 1965-1974, PubMed ID: 20298902
- 120 Castellino, F. et al. (2000) Receptor-mediated uptake of antigen/heat shock protein complexes results in major histocompatibility complex class I antigen presentation via two distinct processing pathways. *J Exp Med* 191, 1957-1964, PubMed ID: 20298901

Heat shock proteins in health and disease: therapeutic targets or therapeutic agents?

- 121 Kol, A. et al. (2000) Cutting edge: heat shock protein (HSP) 60 activates the innate immune response: CD14 is an essential receptor for HSP60 activation of mononuclear cells. *J Immunol* 164, 13-17, PubMed ID: 20072722
- 122 Ohashi, K. et al. (2000) Cutting edge: heat shock protein 60 is a putative endogenous ligand of the toll-like receptor-4 complex. *J Immunol* 164, 558-561, PubMed ID: 20090838
- 123 Hoshino, K. et al. (1999) Cutting edge: Toll-like receptor 4 (TLR4)-deficient mice are hyporesponsive to lipopolysaccharide: evidence for TLR4 as the Lps gene product. *J Immunol* 162, 3749-3752, PubMed ID: 99218385
- 124 Asea, A. et al. (2000) HSP70 stimulates cytokine production through a CD14-dependant pathway, demonstrating its dual role as a chaperone and cytokine. *Nat Med* 6, 435-442, PubMed ID: 20207089
- 125 Singh-Jasuja, H. et al. (2000) The heat shock protein gp96 induces maturation of dendritic cells and down-regulation of its receptor. *Eur J Immunol* 30, 2211-2215
- 126 Binder, R.J., Han, D.K. and Srivastava, P.K. (2000) CD91: a receptor for heat shock protein gp96. *Nat Immunol* 1, 151-155, PubMed ID: 21205395
- 127 Basu, S. et al. (2001) CD91 is a common receptor for heat shock proteins gp96, hsp90, hsp70, and calreticulin. *Immunity* 14, 303-313, PubMed ID: 21186398
- 128 Houenou, L.J. et al. (1996) Exogenous heat shock cognate protein Hsc 70 prevents axotomy-induced death of spinal sensory neurons. *Cell Stress Chaperones* 1, 161-166, PubMed ID: 97365774
- 129 Johnson, A.D., Berberian, P.A. and Bond, M.G. (1990) Effect of heat shock proteins on survival of isolated aortic cells from normal and atherosclerotic cynomolgus macaques. *Atherosclerosis* 84, 111-119, PubMed ID: 91128479
- 130 Johnson, A.D. and Tytell, M. (1993) Exogenous HSP70 becomes cell associated, but not internalized, by stressed arterial smooth muscle cells. *In Vitro Cell Dev Biol Anim* 29A, 807-812, PubMed ID: 94163398
- 131 Cohen, I.R. (1996) Heat shock protein 60 and the regulation of autoimmunity. In *Stress Proteins in Medicine* (van Eden, W. and Young, D., eds), pp. 93-102, Marcel Dekker, New York, USA
- 132 Ramage, J.M. et al. (1999) T cell responses to heat-shock protein 60: differential responses by CD4+ T cell subsets according to their expression of CD45 isotypes. *J Immunol* 162, 704-710, PubMed ID: 99113727
- 133 Multhoff, G. and Hightower, L.E. (1996) Cell surface expression of heat shock proteins and the immune response. *Cell Stress Chaperones* 1, 167-176, PubMed ID: 97365775
- 134 de Graeff-Meeder, E.R. et al. (1995) Juvenile chronic arthritis: T cell reactivity to human HSP60 in patients with a favorable course of arthritis. *J Clin Invest* 95, 934-940, PubMed ID: 95190049
- 135 van Roon, J.A. et al. (1996) Prevention and reversal of cartilage degradation in rheumatoid arthritis by interleukin-10 and interleukin-4. *Arthritis Rheum* 39, 829-835, PubMed ID: 96212444
- 136 van Roon, J.A. et al. (1997) Stimulation of suppressive T cell responses by human but not bacterial 60-kD heat-shock protein in synovial fluid of patients with rheumatoid arthritis. *J Clin Invest* 100, 459-463, PubMed ID: 97364814
- 137 van den Broek, M.F. et al. (1989) Protection against streptococcal cell wall-induced arthritis by pretreatment with the 65-kD mycobacterial heat shock protein. *J Exp Med* 170, 449-466, PubMed ID: 89328323
- 138 Billingham, M.E. et al. (1990) A mycobacterial 65-kD heat shock protein induces antigen-specific suppression of adjuvant arthritis, but is not itself arthritogenic. *J Exp Med* 171, 339-344, PubMed ID: 90111626
- 139 Kingston, A.E. et al. (1996) A 71-kD heat shock protein (hsp) from *Mycobacterium tuberculosis* has modulatory effects on experimental rat arthritis. *Clin Exp Immunol* 103, 77-82, PubMed ID: 96150286
- 140 Anderton, S.M. and van Eden, W. (1996) T lymphocyte recognition of hsp60 in experimental arthritis. In *Stress Proteins in Medicine* (van Eden, W. and Young, D., eds), pp. 73-91, Marcel Dekker, New York, USA
- 141 Anderton, S.M. et al. (1995) Activation of T cells recognizing self 60-kD heat shock protein can protect against experimental arthritis. *J Exp Med* 181, 943-952, PubMed ID: 95173608
- 142 van Roon, J. et al. (1996) Reactivity of T cells from patients with rheumatoid arthritis towards human and mycobacterial hsp60. *FASEB J* 10, A1312
- 143 Paul, A.G. et al. (2000) Highly autoproliative T cells specific for 60-kDa heat shock protein produce IL-4/IL-10 and IFN-gamma and are protective in adjuvant

- arthritis. *J Immunol* 165, 7270-7277, PubMed ID: 20570974
- 144 Tanaka, S. et al. (1999) Activation of T cells recognizing an epitope of heat-shock protein 70 can protect against rat adjuvant arthritis. *J Immunol* 163, 5560-5565, PubMed ID: 20021855
- 145 Wendling, U. et al. (2000) A conserved mycobacterial heat shock protein (hsp) 70 sequence prevents adjuvant arthritis upon nasal administration and induces IL-10-producing T cells that cross-react with the mammalian self-hsp70 homologue. *J Immunol* 164, 2711-2717, PubMed ID: 20143867
- 146 van Eden, W. et al. (1998) Do heat shock proteins control the balance of T-cell regulation in inflammatory diseases? *Immunol Today* 19, 303-307, PubMed ID: 98331052
- 147 Young, D. et al. (1993) Molecular chaperones and the immune response. *Philos Trans R Soc Lond B Biol Sci* 339, 363-367; discussion 367-368, PubMed ID: 93262075
- 148 Srivastava, P.K. and Udon, H. (1994) Heat shock protein-peptide complexes in cancer immunotherapy. *Curr Opin Immunol* 6, 728-732, PubMed ID: 95127110
- 149 Srivastava, P.K., DeLeo, A.B. and Old, L.J. (1986) Tumor rejection antigens of chemically induced sarcomas of inbred mice. *Proc Natl Acad Sci U S A* 83, 3407-3411, PubMed ID: 86205907
- 150 Udon, H. and Srivastava, P.K. (1994) Comparison of tumor-specific immunogenicities of stress-induced proteins gp96, hsp90, and hsp70. *J Immunol* 152, 5398-5403, PubMed ID: 94246182
- 151 Basu, S. and Srivastava, P.K. (1999) Calreticulin, a peptide-binding chaperone of the endoplasmic reticulum, elicits tumor- and peptide-specific immunity. *J Exp Med* 189, 797-802, PubMed ID: 99160626
- 152 Wang, X.Y. et al. (2001) Characterization of heat shock protein 110 and glucose-regulated protein 170 as cancer vaccines and the effect of fever-range hyperthermia on vaccine activity. *J Immunol* 166, 490-497, PubMed ID: 20571932
- 153 Robert, J. et al. (2001) Phylogenetic conservation of the molecular and immunological properties of the chaperones gp96 and hsp70. *Eur J Immunol* 31, 186-195, PubMed ID: 21150881
- 154 Janetzki, S. et al. (2000) Immunization of cancer patients with autologous cancer-derived heat shock protein gp96 preparations: a pilot study. *Int J Cancer* 88, 232-238, PubMed ID: 20461194
- 155 Chandawarkar, R.Y., Wagh, M.S. and Srivastava, P.K. (1999) The dual nature of specific immunological activity of tumor-derived gp96 preparations. *J Exp Med* 189, 1437-1442, PubMed ID: 99242642
- 156 Srivastava, P.K. and Chandawarkar, R.Y. (1999) Method and compositions for the treatment of autoimmune disease using heat shock proteins. Patent 6 007 821, Fordham University, Bronx, NY, USA
- 157 Binder, R.J., Karimeddini, D. and Srivastava, P.K. (2001) Adjuvanticity of alpha(2)-macroglobulin, an independent ligand for the heat shock protein receptor CD91. *J Immunol* 166, 4968-4972, PubMed ID: 21186219
- 158 Binder, R.J. et al. (2000) Cutting edge: heat shock protein gp96 induces maturation and migration of CD11c+ cells in vivo. *J Immunol* 165, 6029-6035, PubMed ID: 20540054
- 159 Srivastava, P.K. and Maki, R.G. (1991) Stress-induced proteins in immune response to cancer. *Curr Top Microbiol Immunol* 167, 109-123, PubMed ID: 91275569
- 160 Blachere, N.E. et al. (1993) Heat shock protein vaccines against cancer. *J Immunother* 14, 352-356, PubMed ID: 94107875
- 161 Heikema, A. et al. (1997) Generation of heat shock protein-based vaccines by intracellular loading of gp96 with antigenic peptides. *Immunol Lett* 57, 69-74, PubMed ID: 97376318
- 162 Suto, R. and Srivastava, P.K. (1995) A mechanism for the specific immunogenicity of heat shock protein-chaperoned peptides. *Science* 269, 1585-1588, PubMed ID: 95397154
- 163 Blachere, N.E. et al. (1997) Heat shock protein-peptide complexes, reconstituted in vitro, elicit peptide-specific cytotoxic T lymphocyte response and tumor immunity. *J Exp Med* 186, 1315-1322, PubMed ID: 97477415
- 164 Ciupitu, A.M. et al. (1998) Immunization with a lymphocytic choriomeningitis virus peptide mixed with heat shock protein 70 results in protective antiviral immunity and specific cytotoxic T lymphocytes. *J Exp Med* 187, 685-691, PubMed ID: 98149861
- 165 Suzue, K. and Young, R.A. (1996) Adjuvant-free hsp70 fusion protein system elicits humoral and cellular immune responses to HIV-1 p24. *J Immunol* 156, 873-879, PubMed ID: 96133020
- 166 Roman, E. and Moreno, C. (1996) Synthetic peptides non-covalently bound to bacterial hsp 70 elicit peptide-specific T-cell responses in vivo. *Immunology* 88, 487-492, PubMed ID: 97036099

Further reading, resources and contacts

The Cell Stress Society International (CSSI) is a US-based organisation that promotes 'the organization of conferences related to stress response research and the publication and dissemination of related journals and monographs'.

<http://www.sp.uconn.edu/%7Ehightowe/cssi/>

Cell Stress & Chaperones is a 'multi-disciplinary peer-reviewed journal that captures the eclectic spirit of the cellular stress response field in a single, concentrated source of current information. The Journal publishes basic and applied research on cellular stress responses of animals, microorganisms and plants'.

<http://cest.allenpress.com/cestonline/?request=index-html>

Features associated with this article

Figures

Figure 1. Regulation of transcription of heat shock protein genes by heat shock factor (fig001gps).

Figure 2. The heat shock protein Hsp60 is an intercellular signalling molecule (fig002gps).

Figure 3. The heat shock proteins Hsp60 and Hsp70 induce pro-inflammatory cytokine secretion from monocytes (fig003gps).

Figure 4. The heat shock protein gp96 delivers antigenic peptides and maturation signals to antigen-presenting cells, and induces release of cytokines (fig004gps).

Figure 5. Potential therapeutic applications for heat shock proteins (fig005gps).

Table

Table 1. Major mammalian heat shock proteins and their function (tab001gps).

Citation details for this article

A. Graham Pockley (2001) Heat shock proteins in health and disease: therapeutic targets or therapeutic agents? *Exp. Rev. Mol. Med.* 21 September, <http://www-erm.m.cbcu.cam.ac.uk/01003556h.htm>