



This is a repository copy of *Characterization of spotted hyena, Crocuta crocuta microsatellite loci*.

White Rose Research Online URL for this paper:  
<http://eprints.whiterose.ac.uk/369/>

---

**Article:**

Wilhelm, K., Dawson, D.A., Gentle, L.K. et al. (7 more authors) (2003) Characterization of spotted hyena, *Crocuta crocuta* microsatellite loci. *Molecular Ecology Notes*, 3 (2). pp. 360-362. ISSN 1471-8278

<https://doi.org/10.1046/j.1471-8286.2003.00450.x>

---

**Reuse**

Unless indicated otherwise, fulltext items are protected by copyright with all rights reserved. The copyright exception in section 29 of the Copyright, Designs and Patents Act 1988 allows the making of a single copy solely for the purpose of non-commercial research or private study within the limits of fair dealing. The publisher or other rights-holder may allow further reproduction and re-use of this version - refer to the White Rose Research Online record for this item. Where records identify the publisher as the copyright holder, users can verify any specific terms of use on the publisher's website.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.



[eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk)  
<https://eprints.whiterose.ac.uk/>

## PRIMER NOTE

# Characterization of spotted hyena, *Crocuta crocuta* microsatellite loci

KERSTIN WILHELM,\*§ DEBORAH A. DAWSON,†¶ LOUISE K. GENTLE,† GILES F. HORSFIELD,† CHRISTIAN SCHLÖTTERER,‡\*\* CAROLYN GREIG,† MARION EAST,\*§ HERIBERT HOFER,\*§ DIETHARD TAUTZ,‡†† and TERRY BURKE,†¶

\*Max-Planck-Institut für Verhaltensphysiologie, Postfach 1564, D-82305 Seewiesen, Germany, †Department of Biology, University of Leicester, University Road, Leicester, LE1 7RH, UK, ‡Zoologisches Institut der Universität München, Luisenstr. 14, D-80333 München, Germany

## Abstract

**We have isolated 10 polymorphic microsatellite loci in the spotted hyena, *Crocuta crocuta*. The loci displayed between eight and 14 alleles in a minimum of 12 individuals tested. These loci will be used to investigate relatedness within social groups, the genetic structure of populations, sexual selection, and mate choice in spotted hyenas.**

*Keywords:* Carnivora, *Crocuta crocuta*, hyena, Hyenidae, microsatellite

*Received 7 February 2003; revision accepted 17 March 2003*

The spotted hyena, *Crocuta crocuta*, is a predator-scavenger that occupies a wide range of habitats in sub-Saharan Africa and lives in social clans that vary in size from a few to up to 80 members in which females are philopatric and males typically disperse (Mills & Hofer 1998). This species displays a suite of unusual traits, including female dominance, facultative siblicide, masculinization of the female clitoris and males that queue for social dominance (East & Hofer 2002). To investigate paternity, reproductive success and mate choice in this unique species, we developed 10 polymorphic microsatellite markers.

We collected hair and/or blood/tissue samples from 46 spotted hyenas, comprising two clans, inhabiting the Serengeti National Park, Tanzania. We extracted DNA from hair samples using Chelex-100, following the Promega DNA extraction protocol (Promega 1996), and from blood, following the protocols of Sambrook *et al.* (1989) or using the QIAamp blood and tissue kit (QIAGEN).

Correspondence: Deborah A. Dawson. Present address: ¶Department of Animal and Plant Sciences, University of Sheffield, Sheffield, S10 2TN, UK. Fax: + 44 (0)114 2220002; E-mail: D.A.Dawson@sheffield.ac.uk

§Present address: Institute for Zoo and Wildlife Research, Alfred-Kowalke-Str. 17, D-10315 Berlin, Germany.

\*\*Institut für Tierzucht und Genetik, Veterinärmedizinische Universität Wien, Josef Baumann Gasse 1, A-1210 Wien, Austria.

††Institute for Genetics, Cologne University, Weyertal 121, D-50931 Cologne, Germany.

Three genomic libraries were prepared. For the preparation of the first two libraries (A and B), pooled genomic DNA was digested with the restriction endonucleases *AluI*, *HaeIII* and *RsaI*. The digest fragments were size-selected (100–600 bp) and ligated into *SmaI*-digested M13mp18/mp19 phage (library A) or *SmaI*-digested pUC18 (library B). Transformed colonies were screened for CA.GT and GA.CT as described in Rassmann *et al.* (1991).

To prepare the third microsatellite library (C), pooled DNA from one male and one female was used. The library was prepared enriched for (TTTC.AAAG)<sub>n</sub> following Armour *et al.* (1994) with one modification: to prevent duplicate clones, the DNA fragments were not polymerase chain reaction (PCR)-amplified before the enrichment hybridization. In total, 1200 transformant clones were screened by hybridization to the sequences (TTTC.AAAG)<sub>n</sub> (Armour *et al.* 1994) and (CA.GT)<sub>n</sub> or (GA.CT)<sub>n</sub> (Amersham Pharmacia Biotech) radiolabelled with [ $\alpha^{32}$ P]-dCTP. 210 positive clones were obtained. Forty-nine positive clones were sequenced using BIG DYE Terminators on an ABI 377 Sequencer (Applied Biosystems). The majority of clones contained tetranucleotide motifs but several clones contained dinucleotide (CA.GT)<sub>n</sub> repeats, despite the library not being enriched for these.

All sequences were checked to determine whether they were unique using GENEJOCKEY software (Biosoft). In total, 38 unique microsatellite sequences were isolated from the three libraries (11, eight and 19, respectively, from libraries

**Table 1** Characteristics of 10 spotted hyena (*Crocuta crocuta*) microsatellite loci

Locus	EMBL accession no., library & clone name	Repeat motif (5'–3')†	Primer sequence (5'–3') (& 5' fluorescent primer label, underlined)	Primer $T_a$ (°C)	PCR profile used	MgCl <sub>2</sub> (mM)	Exp. allele size (bp)†	$n$	Analysis method used ( $n$ )	Obs. allele size range (bp)	$H_O$	$H_E$
<i>Ccroc01</i>	AJ512845 A, CSH64	(CA) <sub>18</sub>	(F) CCTCAATTAGGAACATAAAAAAGTG (R) GAAGGAAGGAAGCAATATGC	56 56	67–55 (B)	1.5	208	24	SS	8 150–200	0.92	0.85
<i>Ccroc02</i>	AJ512846 A, CSH65	(TG) <sub>20</sub>	(F) GCATGCGATAAATTTGAAGATG (R) (6-FAM)-CAAAAAGGAGAAATTTAGCAGA	57 56	67–55 (B & H)	1.5	118	44*	ABI	10 85–111	0.59	0.69
<i>Ccroc03</i>	AJ512850 B, HY7GH	(CA) <sub>4</sub> (GA) <sub>1</sub> (CA) <sub>1</sub> (CG) <sub>1</sub> (CA) <sub>18</sub>	(F) TTGAAACAAGGATAAAATTAATGAAAAGA (R) ACAGTAGAAAATAATCCAAGTATAAACC	60 56	60–50 (B)	2.0	133	12	SS	8 120–175	0.67	0.75
<i>Ccroc04</i>	AJ512851 B, HY15GH	(TG) <sub>3</sub> (TACG) <sub>1</sub> (TG) <sub>12</sub>	(F) TGGGCTAAGTCTTTCCCTGA (R) (TET)-ATGGGGGCTCCTTCACTC	60 60	67–55 (B) 62 (H)	1.5	307	42*	ABI	9 305–323	0.86	0.82
<i>Ccroc05</i>	AJ512855 B, HY27GH	(CA) <sub>27</sub>	(F) ACCAGTGATCTGGACTGGGA (R) (TET)-AAAAGTAATATGACTGCCAAAAGC	61 58	67–55 (B & H)	1.5	171	45*	ABI	8 155–175	0.93	0.83
<i>Ccroc06</i>	AJ512859 C, T01A03	(TTTC) <sub>22</sub> (TT) <sub>1</sub> (TTTC) <sub>18</sub>	(F) GATCGTPTTTTGACAAGTGCTG (R) (6-FAM)-AAACCCACTTCGAGCCTGAT	58 61	58 (B & H)	1.5	308	46*	ABI (38) SS (8)	10 228–328	0.98	0.85
<i>Ccroc07</i>	AJ512860 C, T01C09	(GGAA) <sub>6</sub> (GAA) <sub>1</sub> (GAAA) <sub>19</sub>	(F) TCCCTCAAGTCACTCGGTGT (R) (HEX)-TGCTAATGTTTCATTGCAGGG	61 60	62 (B & H)	1.5	281	45*	ABI	12 273–328	0.96	0.90
<i>Ccroc08</i>	AJ512866 C, T01H04	(TTTC) <sub>2</sub> (GTTC) <sub>1</sub> (TTTC) <sub>1</sub> (TT) <sub>1</sub> (TTTC) <sub>22</sub> (TTCTC) <sub>1</sub> (TTTC) <sub>15</sub>	(F) TTTTTTAAAGTTCTTAGGCAGGAC (R) (6-FAM)-GGAGCCTGTTTCAGTCTGTG	58 58	67–55 (B) 62 (H)	1.5	274	45*	ABI	14 162–276	0.89	0.89
<i>Ccroc09</i>	AJ512870 C, T02B12	(GT) <sub>19</sub>	(F) CAGAATTAATCCATATCACAACACTGC (R) (TET)-TAGGAACCTCTTGCCGTCAG	60 60	67–55 (B) 62 (H)	1.5	218	43*	ABI 310	8 208–224	0.77	0.77
<i>Ccroc10</i>	AJ512872 C, T02D10	(TTTC) <sub>5</sub> (TC) <sub>2</sub> (TTTC) <sub>18</sub> (TC) <sub>1</sub> (TTTC) <sub>1</sub> (N) <sub>9</sub> (TTTA) <sub>2</sub> (T) <sub>4</sub> (GA) <sub>7</sub>	(F) CTCAGTTTATTAACACCCGC (R) (HEX)-AAGCTCTTGGAACCTAAGCTACTTG	59 58	67–55 (B) 62 (H)	1.5	223	46*	ABI (34) SS (12)	11 184–218	0.91	0.85

†Of sequenced clone.

 $T_a$  = annealing temperature.  $n$  = number of individuals genotyped;  $A$  = number of alleles;  $H_O$  = observed heterozygosity;  $H_E$  = expected heterozygosity; B = blood/tissue, H = hair;

\*between 22 and 25 of the samples were hair samples; SS = genotypes scored on 6% polyacrylamide gels stained with silver; ABI = genotypes scored on an ABI 377 Sequencer.

PCR profiles used:

Profile 67–55 = 94 °C for 30 s, X °C for 45 s, 72 °C for 45 s 2 cycles per step (X = 67–58 °C decreasing by 3 °C each step) then 94 °C for 30 s, 55 °C for 45 s, 72 °C for 45 s for 35 cycles.

Profile 60–50 = 94 °C for 30 s, X °C for 30 s, 72 °C for 30 s 1 cycles per step (X = 60–55 °C decreasing by 1 °C each step) then 94 °C for 30 s, 50 °C for 30 s, 72 °C for 30 s for 30 cycles followed by 72 °C for 5 min.

Profile 62 = 95 °C for 30 s, 62 °C for 30 s, 72 °C for 1 min for 43 cycles.

Profile 58 = 94 °C for 30 s, 58 °C for 45 s, 72 °C for 45 s for 40 cycles.

A to C: EMBL accession numbers AJ512839–AJ512876). After submission to the EMBL database, all 38 sequences were confirmed as unique using BLASTN 2.2.4 (Altschul *et al.* 1997). Primers were designed from 17 sequences (five, three and nine, respectively, from libraries A to C) using PRIMER 3 (Rozen & Skaletsky 1996, 1997). The remaining 20 sequences were not used, but all except one (AJ512848) are suitable for primer design.

The loci were tested for polymorphism using a panel of 12–46 individuals. PCR reactions were performed in a 2400 Perkin Elmer thermocycler using 0.2 ml microreaction tubes (Perkin Elmer). DNA was amplified in a 15 µL PCR reaction volume, which contained between 0.5 and 10 ng of template hyena DNA, 200 µM dNTPs, 50 µM of each primer, 0.345 U (0.023 U/µL) of *Taq* DNA polymerase (Amersham Pharmacia Biotech) and 1 × reaction buffer (50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 10 mM Tris-HCl, pH 9.0).

Four different cycling profiles were necessary to allow for differences in primer annealing temperatures and the different tissues from which DNA was extracted (Table 1). Each PCR run, for each locus, was ensured contamination-free by including a DNA-free control (sterile ultrapure water). PCR amplification was confirmed on 1.5% agarose gels stained with ethidium bromide.

For eight loci, the genotypes were analysed on an ABI 377 or ABI 310 Sequencer (Applied Biosystems) using fluorescent primers (Table 1) and GENESCAN software (v2.0.2, Applied Biosystems). Loci *Ccroc01*, *Ccroc03*, *Ccroc06* and *Ccroc10* were analysed on 6% polyacrylamide gels stained with silver (Promega 1996). Allele sizes were assigned using a ØX174 *Hae*III DNA marker (Applied Biosystems) and two hyena PCR-amplified reference samples. Most genotypes at *Ccroc06* and *Ccroc10* (34 and 38, respectively, of the 46) were also checked, and all found to match, on an ABI 310 Sequencer.

Of the 17 primer pairs tested, seven were abandoned. Five loci were abandoned due to nonspecific amplification (CSH18-AJ512839, CSH44-AJ512842, T01D05-AJ512861, T01H06-AJ512867, T02H12-AJ512876); the products at two of these (CSH18 and T02H12) may have resulted from one primer being positioned in a SINE element identified in other carnivore species (Slattery *et al.* 2000). No product was obtained for locus T01D06 (AJ512862), and locus CSH77-AJ512849 (which shows a strong similarity to a human sequence; GenBank AC097478) was monomorphic.

The 10 polymorphic loci displayed between eight and 14 alleles in a minimum of 12 individuals (Table 1). Primer sequences, expected PCR product sizes, allele size ranges and numbers of alleles for the loci are given in Table 1. The

observed and expected heterozygosities for each locus were calculated using CERVUS v2.0 (Marshall *et al.* 1998; Table 1). There was no significant deviation from Hardy–Weinberg expected allele frequencies for the samples presented in Table 1 for any locus ( $P > 0.05$ ) and, in a pedigree analysis of 123 known mothers and 463 offspring (111 twin litters), no null alleles were detected at any of the six loci used (*Ccroc05*, *Ccroc06*, *Ccroc07*, *Ccroc08*, *Ccroc09* and *Ccroc10*; unpublished data).

## Acknowledgements

We are grateful to the Fritz-Thyssen-Stiftung, the Stifterverband der deutschen Wissenschaft, the Max-Planck-Gesellschaft and the Institute for Zoo and Wildlife Research for financial assistance. DD is supported by the Sheffield Molecular Genetics Facility, funded by the Natural Environmental Research Council, UK. We thank Olivier Hanotte for testing the primers developed from the first microsatellite library.

## References

- Altschul SF, Madden TL, Schäffer AA *et al.* (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Research*, **25**, 3389–3402.
- Armour JAL, Neumann R, Gobert S, Jeffreys AJ (1994) Isolation of human simple repeat loci by hybridization selection. *Human Molecular Genetics*, **3**, 599–605.
- East ML, Hofer H (2002) Conflict and cooperation in a female dominated society: a reassessment of the 'hyperaggressive' image of spotted hyenas. *Advances in the Study of Behavior*, **31**, 1–30.
- Marshall TC, Slate J, Kruuk LEB, Pemberton JM (1998) Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology*, **7**, 639–655 (<http://helios.bto.ed.ac.uk/evolgen/cervus/cervus.html>).
- Mills G, Hofer H (1998) *Hyaenas Status Survey and Conservation Action Plan*. International Union for the Conservation of Nature and Natural Resources, Gland, Switzerland.
- Promega (1996) *GenePrint STR Systems Technical Manual*, Part# TMD004. Revised 2/96. Promega Corp., Madison, USA.
- Rassmann K, Schlötterer C, Tautz D (1991) Isolation of simple-sequence loci for use in polymerase chain reaction-based DNA fingerprinting. *Electrophoresis*, **12**, 113–118.
- Rozen S, Skaletsky HJ (1996, 1997) *PRIMER Version 3*. MIT Center for Genome Research and Whitehead Institute for Biomedical Research, Cambridge, MA, USA ([http://www-genome.wi.mit.edu/genome\\_software/other/primer3.html](http://www-genome.wi.mit.edu/genome_software/other/primer3.html)).
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: a Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, USA.
- Slattery JP, Murphy WJ, O'Brien SJ (2000) Patterns of diversity among SINE elements isolated from three Y-chromosome genes in carnivores. *Journal of Molecular Biology and Evolution*, **17**, 825–829.