



UNIVERSITY OF LEEDS

This is a repository copy of *Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/154/>

Article:

Gong, YY orcid.org/0000-0003-4927-5526, Cardwell, K, Hounsa, A et al. (4 more authors) (2002) Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study. *British Medical Journal*, 325 (7354). pp. 20-21. ISSN 1759-2151

<https://doi.org/10.1136/bmj.325.7354.20>

Reuse

See Attached

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>



Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study

Y Y Gong, K Cardwell, A Hounsa, S Egal, P C Turner, A J Hall and C P Wild

BMJ 2002;325:20-21
doi:10.1136/bmj.325.7354.20

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/325/7354/20>

These include:

References

This article cites 4 articles, 2 of which can be accessed free at:
<http://bmj.com/cgi/content/full/325/7354/20#BIBL>

3 online articles that cite this article can be accessed at:
<http://bmj.com/cgi/content/full/325/7354/20#otherarticles>

Rapid responses

One rapid response has been posted to this article, which you can access for free at:
<http://bmj.com/cgi/content/full/325/7354/20#responses>

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/325/7354/20>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Topic collections

Articles on similar topics can be found in the following collections

- [Pharmacology and toxicology](#) (284 articles)
 - [Medicine in Developing Countries](#) (725 articles)
 - [Other Pediatrics](#) (1357 articles)
-

Correction

A correction has been published for this article. The contents of the [correction](#) have been appended to the original article in this reprint. The correction is also available online at:
<http://bmj.com/cgi/content/full/325/7372/1089/a>

Notes

To order reprints of this article go to:
<http://bmj.bmjournals.com/cgi/reprintform>

To subscribe to *BMJ* go to:
<http://www.bmjournals.com/subscriptions>

What is already known on this topic

Delay in presentation and assessment of patients with suspected stroke prevents the possible benefits from thrombolysis being achieved

Little is known about the presentation and early management of patients with acute stroke in the United Kingdom

What this study adds

Most patients with suspected stroke in the United Kingdom arrive at hospital within six hours of the onset of symptoms

Not all patients are evaluated by a senior doctor within three hours of arrival at hospital and most do not undergo computed tomography

The potential for thrombolysis in patients with acute stroke can be improved significantly by greater use of emergency services and expediting evaluation and investigations by doctors

(Sunderland Royal Hospital), M Power (Ulster Hospital, Dundonald), A K Sharma (University Hospital Aintree, Liverpool), K R Lees (Western Infirmary, Glasgow), and D G Smithard (William Harvey Hospital, Ashford, Kent).

Contributors: See bmj.com

Funding: This project was supported by an unrestricted grant from Boehringer Ingelheim.

Competing interests: LK and MMB have been reimbursed by Boehringer Ingelheim to attend conferences. AKS and KRL have been reimbursed by Boehringer Ingelheim to attend conferences and to give lectures. RIV is an employee of Boehringer Ingelheim. None of the authors stand to gain financially from publication.

- 1 American Heart Association. *Heart and stroke statistical update*. Dallas, TX: American Heart Association, 1999.
- 2 Hankey GJ, Warlow CP. Treatment and secondary prevention of stroke: evidence, costs and effects on individuals and populations. *Lancet* 1999;354:1457-63.

- 3 Grotta JC. Acute stroke therapy in the millennium: consummating the marriage between the laboratory and the bedside: the Feinberg lecture. *Stroke* 1999;30:1722-8.
- 4 Bath PM, Lees KR. ABC of arterial and venous disease: acute stroke. *BMJ* 2000;320:920-3.
- 5 Wardlaw JM, del Zoppo G, Yamaguchi T. Thrombolysis for acute ischaemic stroke. *Cochrane Library*. Issue 1. Oxford: Update software, 2000.
- 6 Lees KR. Thrombolysis. *Brit Med Bull* 2000;389-400.
- 7 Alberts MJ, Hademenos G, Latchaw RE, Jagoda A, Marler JR, Mayberg MR, et al. Recommendations for the establishment of primary stroke centers. *JAMA* 2000;283:3102-9.
- 8 Smith MA, Doliszny KM, Shahar E, McGovern PG, Arnett DK, Luepker RV. Delayed hospital arrival for acute stroke: the Minnesota stroke survey. *Ann Intern Med* 1998;129:190-6.
- 9 Katzan IL, Furlan AJ, Way LE, Farnk JJ, Harper DL, Hinchey JA, et al. A systematic audit of iv tPA in Cleveland area hospitals. *Stroke* 1999;30:266.
- 10 Evenson K, Rosamond W, Morris D. Prehospital and in-hospital delays in acute stroke care. *Neuroepidemiology* 2001;20:65-76.
- 11 Lacy C, Suh D, Bueno M, Kostis J, for the STROKE Collaborative Study Group. Delay in presentation and evaluation for acute stroke: Stroke Time Registry for Outcomes Knowledge and Epidemiology (STROKE). *Stroke* 2001;32:63-9.
- 12 Schroeder E, Rosamond W, Morris D, Evenson K, Hinn A. Determinants of emergency medical services use in a population with stroke symptoms: the Second Delay in Accessing Stroke Healthcare (DASH II) Study. *Stroke* 2000;31:2591-6.
- 13 Morris DL, Rosamond W, Madden K, Schultz C, Hamilton S. Prehospital and emergency department delays after acute stroke. The Genentech Stroke Presentation Survey. *Stroke* 2000;31:2585.
- 14 Kothari R, Jauch E, Broderick J, Brott T, Sauerbeck L, Khoury J, et al. Acute stroke: delays to presentation and emergency department evaluation. *Ann Emerg Med* 1999;33:3-8.
- 15 Rosamond W, Gorton R, Hinn A, Hohenhaus S, Morris D. Rapid response to stroke symptoms: the Delay in Accessing Stroke Healthcare (DASH) Study. *Acad Emerg Med* 1998;5:45-51.
- 16 Menon SC, Pandey DK, Morganstern LB. Critical factors in determining access to acute stroke care. *Neurology* 1998;51:427-32.
- 17 Wester P, Radberg J, Lundgren B, Peltonen M, for the Seek-Medical-Attention-in-Time Study Group. Factors associated with delay admission to hospital and in-hospital delays in acute stroke and TIA. *Stroke* 1999;30:40-8.
- 18 Morris D, Rosamond W, Hinn A, Gorton R. Time delays in accessing stroke care in the emergency department. *Acad Emerg Med* 1999;6:218-23.
- 19 Bratina P, Greenberg L, Pasteur W, Grotta J. Current emergency department management of stroke in Houston, Texas. *Stroke* 1995;26:409-14.
- 20 Intercollegiate Working Party on Stroke. *National clinical guidelines on stroke*. London: Royal College of Physicians, 2000.
- 21 *Coronary Heart Disease/Stroke Task Force report*. Edinburgh: Scottish Executive, 2001.
- 22 Barer D, Main A, Lodwick R. Practicability of early treatment of acute stroke. *Lancet* 1992;339:1540-1.
- 23 Harper GD, Haigh RA, Potter JF, Castleden CM. Factors delaying hospital admission after stroke in Leicestershire. *Stroke* 1992;23:835-8.
- 24 Hacke W. A late step in the right direction for stroke care. *Lancet* 2000;356:869-70.

(Accepted 22 January 2002)

Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study

Y Y Gong, K Cardwell, A Hounsa, S Egal, P C Turner, A J Hall, C P Wild

Molecular Epidemiology Unit, Epidemiology and Health Services Research, School of Medicine, University of Leeds, Leeds LS2 9JT
Y Y Gong
research fellow
P C Turner
research fellow
C P Wild
professor

continued over


BMJ 2002;325:20-1

Fetal and early childhood environment, including the nutritional status of the pregnant mother and the infant, are considered critical for growth and risk of disease in later life.¹ Many people in developing countries are not only malnourished but also chronically exposed to high levels of toxic fungal metabolites (mycotoxins). One family of mycotoxins, the aflatoxins, are carcinogenic and immunotoxic and cause growth retardation in animals.² Aflatoxins contaminate staple foods in West Africa, particularly maize and groundnuts, as a result of hot, humid storage conditions that promote fungal growth. High exposure to aflatoxins occurs throughout childhood in the region,^{3,4} suggest-

ing that growth and development could be critically affected. We assessed exposure to aflatoxins in relation to anthropometric measures in children in Benin and Togo.

Methods and results

We studied 480 children (aged 9 months to 5 years) from 16 villages in four geographic zones (four in each zone): Sudan savannah, north Guinea savannah, south Guinea savannah, and coastal savannah. The Ministries for Health in Benin and Togo gave ethical approval, and parents gave informed consent. We determined



Concentrations of aflatoxin-albumin adduct categorised into four groups for height for age and weight for age z scores on the basis of the WHO classification of malnutrition (z score ≤ 2) and severe malnutrition (≤ 3). Geometric mean adduct concentrations are shown, with 95% confidence intervals, adjusted for weaning status, agro-ecological zone, and socioeconomic status. Height for age and weight for age z scores were significantly associated with aflatoxin-albumin concentration (trend test: $F=15.19$, $P=0.0001$, $r^2=0.3766$; and $F=8.48$, $P=0.0038$, $r^2=0.3680$).

weight for age, height for age, and weight for height z scores, according to the median value of a World Health Organization reference population. A z score ≤ 2 is classified as malnutrition, and ≤ 3 represents severe malnutrition. We also determined weaning status and the socioeconomic status of the mother and family. We assessed aflatoxin exposure over the previous two to three months by measuring aflatoxin bound to albumin in blood.³

We detected aflatoxin-albumin adducts in 475/479 (99%) samples (one sample missing), with a geometric mean concentration of 32.8 (range 5-1064) pg/mg albumin. Aflatoxin-albumin concentration increased with age up to 3 years, after which it reached a plateau. In the 302 children aged 3 years or under, the mean concentration was 2.5-fold higher in fully weaned children (45.6 pg/mg; 95% confidence interval 38.8 to 53.7) than in those still partially breast fed (18.0 pg/mg; 15.2 to 21.3). In a multivariable model adjusting for age, sex, socioeconomic status, and agro-ecological zone, weaning status was significantly associated with aflatoxin-albumin concentration ($P=0.0001$).

Prevalence of malnutrition was 33% for stunting (height for age z score ≤ 2), 29% for being underweight (weight for age z score ≤ 2), and 6% for wasting (weight for height z score ≤ 2). Children with stunting or who were underweight had 30-40% higher mean aflatoxin-albumin concentrations. After adjustment as above, the negative correlation between individual aflatoxin-albumin concentration and each of the three growth parameters was highly significant ($P=0.001$ for height for age, $P=0.005$ for weight for age, and $P=0.047$ for weight for height). In a categorical analysis, the association with aflatoxin-albumin concentration was again significant, with clear dose-response relations with height for age and weight for age z scores (figure).

Comment

This study reveals a striking association between exposure to aflatoxin in children and both stunting

(a reflection of chronic malnutrition) and being underweight (an indicator of acute malnutrition). In West Africa, people are chronically exposed to high levels of aflatoxins starting in utero and continuing throughout life.⁴ In this study, children still partially breast fed had lower exposure, almost certainly reflecting lower toxin levels in milk than in weaning and family foods. Thus growth faltering occurs at a time of change to solid foods, when there is co-exposure to aflatoxin and a plethora of infectious hazards (for example, malaria, diarrhoea, respiratory infections). Whether the association between aflatoxin exposure and impaired growth is a direct result of aflatoxin toxicity or reflects consumption of fungus affected food of poor nutritional quality cannot be confirmed from the cross sectional design. However, these observations emphasise the need to investigate this question and to develop strategies to reduce exposure to aflatoxin, possibly involving interventions targeted at the post-weaning period in African children.⁵

We thank C. Aquereburu for participating in the planning process; M. Koube, Anik Gandjeto, Zenato Assani, Marius Adjaba, and G. Ayeni from IITA Benin, who participated in the field work; and the people of Benin and Togo who agreed to be part of this research.

Contributors: YYG, KC, AH, PCT, AJH, and CPW were all responsible for the design of the study. KC, AH, SE, and AJH took part in the fieldwork. YYG, PCT, and CPW were responsible for the laboratory analysis. YYG and SE computed the data and conducted the statistical analysis. All authors contributed to writing the manuscript. CPW is guarantor for the paper.

Funding: This study was funded by a grant from GTZ (project no 98.7860.4-001.00) and support to CPW and PCT from a grant from the NIEHS, USA (no ES06052).

Competing interests: None declared.

- 1 Terry MB, Susser E. The impact of fetal and infant exposures along the life course. *Int J Epidemiol* 2001;30:95-6.
- 2 Hall AJ, Wild CP. Epidemiology of aflatoxin-related disease. In: Eaton DA, Groopman JD, eds. *Human health, veterinary and agricultural significance*. San Diego, CA: Academic Press, 1994:233-58.
- 3 Turner PC, Mendy M, Whittle H, Fortuin M, Hall AJ, Wild CP. Hepatitis B infection and aflatoxin biomarker levels in Gambian children. *Trop Med Int Health* 2000;5:837-41.
- 4 Montesano R, Hainaut P, Wild CP. Hepatocellular carcinoma: from gene to public health. *J Natl Cancer Inst* 1997;89:1844-51.
- 5 Wild CP, Hall AJ. Primary prevention of hepatocellular carcinoma in developing countries. *Mutat Res* 2000;462:381-93.

(Accepted 20 February 2002)

What is a good doctor and how can we make one?

We want your views on this for a future theme issue of the *BMJ*.

Governmental and professional regulation now operates in most aspects of doctors' lives yet doctors themselves have had little opportunity to air their views on what they think makes a good doctor or how good doctors should be made.

Contribute to this debate now by visiting our home page (bmj.com) or going direct to the debate at www.bmj.com/gooddoctor

All submissions will be published on the website and a selection of the best letters will be published in the theme issue that will be devoted to this topic in September 2002.

International
Institute of Tropical
Agriculture,
Cotonou, Benin,
West Africa
K Cardwell
research plant
pathologist
A Hounsa
public health doctor
S Egal
junior professional
officer

London School of
Hygiene and
Tropical Medicine,
London
WC1E 7HT
A J Hall
professor

Correspondence to:
C P Wild
cp.wild@leeds.ac.uk

private practice. Top merit awards are awarded for national and international standing within the specialty and can more than double a consultant's basic salary. Any time that a hospital doctor devotes to computing does not lead to increased income potential.

Conclusions

Over many years, general practice computing has prospered, whereas hospital clinical computing has not. Differences in leadership and economic incentives partially explain this. In general practice the government and the profession worked together to remove barriers and provide incentives to computerisation. In hospitals the opposite happened. Changes are needed to provide professional leadership and economic incentives in both primary and secondary sectors. An early step would be to establish united stakeholder organisations for clinical users and information technology professionals in health care, covering all aspects of healthcare computing.

The NHS is now planning to deploy integrated patient record systems across both primary and secondary care.²⁰ The examples of Kaiser Permanente and the Veterans Administration suggest that such systems may play a critical part in improving effectiveness and efficiency.²¹ However, such a project faces several technical obstacles, mainly associated with scalability. It is much easier to computerise small general practices than large complex hospitals, let alone provide integrated services across an organisation as large as the NHS. These technical issues—which include patient record architecture, terminology, interoperability standards, security, and developments in computer technology—are the subject of my second article.

I thank Jeremy Wyatt for comments on an earlier draft of this article.

Competing interests: I have participated in many of the events described and have provided consultancy services to various NHS organisations

- 1 Wanless D. *Securing our future health: taking a long-term view. Final report.* London: HM Treasury, 2002. (www.hm-treasury.gov.uk/Consultations_and_Legislation/wanless/consult_wanless_final.cfm, accessed 16 Aug 2002.)
- 2 Department of Health. *Delivering 21st century IT support for the NHS: national strategic programme.* London: DoH, 2002. (www.doh.gov.uk/ipu/whatnew/deliveringit/nhsitimpplan.pdf, accessed 4 Aug 2002.)
- 3 Institute of Medicine. *Crossing the quality chasm: a new health system for the 21st century.* Washington DC: National Academy Press, 2001. (www.nap.edu/books/0309072808/html/, accessed 16 Aug 2002.)
- 4 Institute of Medicine. *The computer-based patient record: an essential technology for health care.* Rev ed. Washington DC: National Academy Press, 1997. (www.nap.edu/books/0309055326/html/index.html, accessed 16 Aug 2002.)
- 5 NHS Information Authority. *NHS connect programme.* Birmingham: NHSIA, 2002. (www.nhsia.nhs.uk/nhsnet/pages/connecting/nhsconnect, accessed 6 Aug 2002.)
- 6 Royal College of General Practitioners. *RCGP information sheet October 2000 No 7. General practice computerisation.* London: RCGP, 2000. (www.rcgp.org.uk/rcgp/information/publications/information/rcf0007/Rcf0007.asp, accessed 4 Aug 2002.)
- 7 Van Bemmel JH, van Ginneken AM, van der Lei J. A progress report on computer-based patient records in Europe. In: Institute of Medicine. *The computer-based patient record: an essential technology for health care.* Rev ed. Washington DC: National Academy Press, 1997:21-43.
- 8 Deloitte and Touche. *The emerging European health telematics industry: market analysis.* Version 1.1. Brussels: European Commission—Directorate General Information Society, 2000. (Reference C13.25533. ([ftp://ftp.cordis.lu/pub/ist/docs/ka1/he_hist_rep.pdf](http://ftp.cordis.lu/pub/ist/docs/ka1/he_hist_rep.pdf), accessed 18 Oct 2002.)
- 9 Kazimirska M, Renaud C, Sawaya L, Zitner D, Korman R. Computer literacy and electronic medical records. College of Family Physicians of Canada. www.cfpc.ca/programs/online/chipsurvey.asp (accessed 29 Jul 2002).
- 10 13th Annual HIMSS leadership survey "Trends in healthcare information technology." 2002. www.himss.org/2002survey (accessed 3 Aug 2002).

- 11 Preece JF, Gillings DB, Lippmann EO, Pearson NG. An on-line record maintenance and retrieval system in general practice. *Int J Biomed Comput* 1970;1:329-37.
- 12 Preece JF. *The use of computers in general practice.* 4th ed. Edinburgh: Churchill Livingstone, 2000.
- 13 Bradshaw-Smith JH. A computer record-keeping system for general practice. *BMJ* 1976;i:1395-7.
- 14 Lindberg DAB. In praise of computing. In: Blum BI, Duncan K, eds. *A history of medical informatics.* New York: ACM Press, 1990:4-13.
- 15 Herbert SI. A review of computer policy and practice. In: *A time for decision: computing policy and practice in the NHS.* London: Nuffield Provincial Hospitals Trust, 1982:5-54.
- 16 Benson T. *Medical informatics: a report for managers and clinicians.* Harlow: Longman Health Services Management, 1991.
- 17 Malcolm A, Poyser J, eds. *Computers and the general practitioner: proceedings of the GP-Info symposium, London, 1980.* Oxford: Pergamon Press for the Royal College of General Practitioners, 1982.
- 18 Collins T. *Crash: ten easy ways to avoid a computer disaster.* London: Simon and Schuster, 1997.
- 19 Fogarty L. Primary care informatics development—one view through the miasma. *J Inform Primary Care* 1997;Jan:2-11. (www.phcsg.org.uk/informatics/jan97/jan2.htm, accessed 4 Aug 2002.)
- 20 Department of Health. *Delivering 21st century IT support for the NHS. National specification for integrated care records service: consultation draft.* Leeds: DoH, 2002. (www.doh.gov.uk/ipu/whatnew/specs_12d.pdf, accessed 16 Aug 2002.)
- 21 Feachem RGA, Sekhri NK, White KL. Getting more for their dollar: a comparison of the NHS with California's Kaiser Permanente. *BMJ* 2002;324:135-43. (<http://bmj.com/cgi/content/full/324/7330/135>, accessed 22 Aug 2002.)
- 22 Department of Health. *Electronic transmission of prescriptions (ETP).* London: DoH, 2001. (www.doh.gov.uk/pharmacy/etp.htm, accessed 5 Aug 2002.)
- 23 Hodge MH. History of the TDS medical information system. In: Blum BI, Duncan K, eds. *A history of medical informatics.* New York: ACM Press, 1990:328-44.
- 24 Bakker AR, Closters L, Mol JL. *Concluding report on the NOBIN-ZIS-project 1972-1976.* Leiden: Academisch Ziekenhuis Leiden, 1978.

Corrections and clarifications

UK senior doctors' career destinations, job satisfaction, and future intentions: questionnaire survey

A lapse in concentration as we processed this paper by Jean M Davidson and colleagues (28 September, pp 685-6) led to the omission of an authors' amendment at proof stage. The paragraph that starts, "We asked respondents to score five statements about job satisfaction" was misleading. It should have read: "We asked respondents to score each of five statements about job satisfaction on a five point ordered scale from 'strongly agree' to 'strongly disagree.' The statements were 'I find enjoyment in my current post'; 'I am doing interesting and challenging work'; 'I feel dissatisfied in my current post'; 'Most days I am enthusiastic about my work'; and 'I am often bored with my work.' We calculated a job satisfaction score for each respondent over all five statements, by assigning a value of 1 to 5 for the responses, from the least to most positive answer, and totalling them: 20 or more represented a positive response, on average, to all statements, and we suggest that this shows a high level of satisfaction."

Dietary aflatoxin exposure and impaired growth in young children from Benin and Togo: cross sectional study

An error crept into this paper by Y Y Gong and colleagues (6 July, pp 20-1). Unfortunately, < -2 and < -3 (referring to z scores) were inadvertently replaced with ≤ 2 and ≤ 3 in both the text (methods and results section) and the figure caption. The correct symbols appeared in the figure.

Mental health campaigners cancel march because of fears of backlash

In this news article by Zosia Kmiotowicz (14 September, p 562), we wrongly referred to Rampton as a prison. It is of course a high security hospital.