



# Flexible Rasch mixture models with package psychomix

Hannah Frick, Carolin Strobl, Friedrich Leisch,  
Achim Zeileis

Working Papers in Economics and Statistics

2011-21

**University of Innsbruck**  
**Working Papers in Economics and Statistics**

The series is jointly edited and published by

Department of Economics

Department of Public Finance

Department of Statistics

Contact Address:

University of Innsbruck

Department of Public Finance

Universitaetsstrasse 15

A-6020 Innsbruck

Austria

Tel: + 43 512 507 7171

Fax: + 43 512 507 2970

e-mail: [eeecon@uibk.ac.at](mailto:eeecon@uibk.ac.at)

The most recent version of all working papers can be downloaded at  
<http://eeecon.uibk.ac.at/wopec/>

For a list of recent papers see the backpages of this paper.

# Flexible Rasch Mixture Models with Package **psychomix**

**Hannah Frick**  
Universität Innsbruck

**Carolin Strobl**  
Universität Zürich

**Friedrich Leisch**  
Universität für  
Bodenkultur Wien

**Achim Zeileis**  
Universität Innsbruck

---

## Abstract

Measurement invariance is an important assumption in the Rasch model and mixture models constitute a flexible way of checking for a violation of this assumption by detecting unobserved heterogeneity in item response data. Here, a general class of Rasch mixture models is established and implemented in R, using conditional maximum likelihood estimation of the item parameters (given the raw scores) along with flexible specification of two model building blocks: (1) Mixture weights for the unobserved classes can be treated as model parameters or based on covariates in a concomitant variable model. (2) The distribution of raw score probabilities can be parametrized in two possible ways, either using a saturated model or a specification through mean and variance. The function `raschmix()` in the R package **psychomix** provides these models, leveraging the general infrastructure for fitting mixture models in the **flexmix** package. Usage of the function and its associated methods is illustrated on artificial data as well as empirical data from a study of verbally aggressive behavior.

*Keywords:* mixed Rasch model, Rost model, mixture model, **flexmix**, R.

---

## 1. Introduction

In item response theory (IRT), latent traits are usually measured by employing probabilistic models for responses to sets of items. One of the most prominent examples for such an approach is the Rasch model (Rasch 1960) which captures the difficulty (or equivalently easiness) of binary items and the respondent's trait level on a single common scale. Generally, a central assumption of most IRT models (including the Rasch model) is measurement invariance, i.e., that all items measure the latent trait in the same way for all subjects. If violated, measurements obtained from such a model provide no fair comparisons of the subjects. A typical violation of measurement invariance in the Rasch model is differential item functioning (DIF), see Ackerman (1992).

Therefore, assessing the assumption of measurement invariance and checking for DIF is crucial when establishing a Rasch model for measurements of latent traits. Hence, various approaches have been suggested in the literature that try to assess heterogeneity in (groups of) subjects either based on observed covariates or unobserved latent classes. If covariates are available, classical tests like the Wald or likelihood ratio test can be employed to compare model fits between some reference group and one or more focal groups (Fischer and Molenaar 1995). Typically, these groups are defined by the researcher based on categorical covariates or arbi-

trary splits in either numerical covariates or the raw scores (Andersen 1972). More recently, extensions of these classical tests have also been embedded into a mixed model representation (Van den Noortgate and De Boeck 2005). Another recently suggested technique is to recursively define and assess groupings in a data-driven way based on all available covariates (both numerical and categorical) in so-called Rasch trees (Strobl, Kopf, and Zeileis 2011).

Heterogeneity occurring in latent classes only (i.e., not observed or captured by covariates), however, is typically addressed by mixtures of IRT models. Specifically, Rost (1990) combined a mixture model approach with the Rasch model. If any covariates are present, they can be used to predict the latent classes (as opposed to the item parameters themselves) in a second step (Cohen and Bolt 2005). More recently, extensions to this mixture model approach have been suggested that encompass this prediction, see Tay, Newman, and Vermunt (2011) for a unifying framework.

In this paper, we introduce the **psychomix** package for the R system for statistical computing (R Development Core Team 2011) that provides software for fitting a general and flexible class of Rasch mixture models along with comprehensive methods for model selection, assessment, and visualization. The package leverages the general and object-oriented infrastructure for fitting mixture models from the **flexmix** package (Leisch 2004; Grün and Leisch 2008), combining it with the function `RaschModel.fit()` from the **psychotools** package (Zeileis, Strobl, and Wickelmaier 2011) for the estimation of Rasch models. All packages are freely available from the Comprehensive R Archive Network at <http://CRAN.R-project.org/>.

The reason for using `RaschModel.fit()` as opposed to other previously existing (and much more powerful and flexible) R packages for Rasch modeling – such as **ltm** (Rizopoulos 2006) or **eRm** (Mair and Hatzinger 2007) – is reduced computational complexity: `RaschModel.fit()` is intended to provide a “no frills” implementation of simple Rasch models, useful when refitting a model multiple times in mixtures or recursive partitions (see also Strobl *et al.* 2011).

While **psychomix** was under development, another R implementation of the Rost (1990) model became available in package **mRm** (Preinerstorfer 2011). As this builds on specialized C++ code, it runs considerably faster than the generic **flexmix** approach – however, it only covers this one particular type of model and offers fewer methods for specifying, inspecting, and assessing (fitted) models. In **psychomix**, both approaches are reconciled by optionally employing the **mRm** solution as an input to the **flexmix** routines.

In the following, we first briefly review both Rasch and mixture models and combine them in a general Rasch mixture framework (Section 2). Subsequently, the R implementation in **psychomix** is introduced (Section 3), illustrated by means of simulated data, and applied in practice to a study of verbally aggressive behavior (Section 4). Concluding remarks are provided in Section 5.

## 2. Rasch mixture models

In the following, we first provide a short introduction to the Rasch model, subsequently outline the basics of mixture models in general, and finally introduce a general class of Rasch mixture models along with the corresponding estimation techniques.

### 2.1. The Rasch model

Latent traits can be measured through a set of items to which binary responses are given.

Success of solving an item or agreeing with it is coded as “1”, while “0” codes the opposite response. The model suggested by Rasch (1960) uses the person’s ability  $\theta_i$  ( $i = 1, \dots, n$ ) and the item’s difficulty  $\beta_j$  ( $j = 1, \dots, m$ ) to model the response  $y_{ij}$  of person  $i$  to item  $j$ :

$$P(Y_{ij} = y_{ij} | \theta_i, \beta_j) = \frac{\exp\{y_{ij}(\theta_i - \beta_j)\}}{1 + \exp\{\theta_i - \beta_j\}}. \quad (1)$$

Under the assumption of independence – both across persons and items within persons (see Fischer and Molenaar 1995, Chapter 1) – the likelihood for the whole sample  $y = (y_{ij})_{n \times m}$  can be written as the product of the likelihood contributions from Equation 1 for all combinations of subjects and items. It is parameterized by the vector of all person parameters  $\theta = (\theta_1, \dots, \theta_n)^\top$  and the vector of all item parameters  $\beta = (\beta_1, \dots, \beta_m)^\top$  (see Equation 2). On the basis of the number of correctly solved items, the so-called “raw” scores  $r_i = \sum_{j=1}^m y_{ij}$ , it can be factorized into a conditional likelihood of the item parameters  $h(\cdot)$  and the score probabilities  $g(\cdot)$  (Equation 3). Because the scores  $r$  are sufficient statistics for the person parameters  $\theta$ , the likelihood of the item parameters  $\beta$  conditional on the scores  $r$  does not depend on the person parameters  $\theta$  (Equation 4).

$$L(\theta, \beta) = f(y | \theta, \beta) \quad (2)$$

$$= h(y | r, \theta, \beta) g(r | \theta, \beta) \quad (3)$$

$$= h(y | r, \beta) g(r | \theta, \beta). \quad (4)$$

The conditional likelihood of the item parameters takes the form

$$h(y | r, \beta) = \prod_{i=1}^n \frac{\exp\{-\sum_{j=1}^m y_{ij} \beta_j\}}{\gamma_{r_i}(\beta)} \quad (5)$$

where  $\gamma_{r_i}(\cdot)$  is the elementary symmetric function of order  $r_i$ , capturing all possible response patterns leading to a certain score (see Fischer and Molenaar 1995, Chapter 3, for details).

There are several approaches to estimating the Rasch model: *Joint maximum likelihood (ML)* estimation of  $\beta$  and  $\theta$  is inconsistent, thus two other approaches have been established. Both are two-step approaches but differ in the way the person parameters  $\theta$  are handled. For *marginal ML* estimation a distribution for  $\theta$  is assumed and integrated out in  $L(\theta, \beta)$ , or equivalently in  $g(r | \theta, \beta)$ . In the *conditional ML* approach only the conditional likelihood of the item parameters  $h(y | r, \beta)$  from Equation 5 is maximized for estimating the item parameters. Technically, this is equivalent to maximizing  $L(\theta, \beta)$  with respect to  $\beta$  if one assumes that  $g(r | \delta) = g(r | \theta, \beta)$  does not depend on  $\theta$  or  $\beta$ , but potentially other parameters  $\delta$ .

In R, the **ltm** package (Rizopoulos 2006) uses the marginal ML approach while the **eRm** package (Mair and Hatzinger 2007) employs the conditional ML approach, i.e., uses and reports only the conditional part of the likelihood in the estimation of  $\beta$ . The latter approach is also taken by the `RaschModel.fit()` function in the **psychotools** package (Zeileis et al. 2011).

## 2.2. Mixture models

Mixture models are a generic approach for modeling data that is assumed to stem from different groups (or clusters) but group membership is unknown.

The likelihood  $f(\cdot)$  of such a mixture model is a weighted sum (with prior weights  $\pi_k$ ) of the likelihood from several components  $f_k(\cdot)$  representing the different groups:

$$f(y_i) = \sum_{k=1}^K \pi_k f_k(y_i).$$

Generally, the components  $f_k(\cdot)$  can be densities or (regression) models. Typically, all  $k$  components  $f_k(\cdot)$  are assumed to be of the same type  $f(y|\xi_k)$ , distinguished through their component-specific parameter vector  $\xi_k$ .

If variables are present which do not influence the components  $f_k(\cdot)$  themselves but rather the prior class membership probabilities  $\pi_k$ , they can be incorporated in the model as so-called *concomitant variables* (Dayton and Macready 1988). In the psychometric literature, such covariates predicting latent information are also employed, e.g., by Tay *et al.* (2011) who advocate a unifying IRT framework that also optionally encompasses concomitant information (labeled *MM-IRT-C* for mixed-measurement IRT with covariates). To embed such concomitant variables  $x_i$  into the general mixture model notation, a model for the component membership probability  $\pi(k|x_i, \alpha)$  with parameters  $\alpha$  is employed:

$$f(y_i|x_i, \alpha, \xi_1, \dots, \xi_K) = \sum_{k=1}^K \pi(k|x_i, \alpha) f(y_i|\xi_k) \quad (6)$$

where commonly a multinomial logit model is chosen to parametrize  $\pi(k|x_i, \alpha)$  (see e.g., Grün and Leisch 2008; Tay *et al.* 2011). Note that the multinomial model collapses to separate  $\pi_k$  ( $k = 1, \dots, K$ ) if there is only an intercept and not real concomitants in  $x_i$ .

### 2.3. Flavors of Rasch mixture models

When combining the general mixture model framework from Equation 6 with the Rasch model based on Equation 1, several options are conceivable for two of the building blocks. First, the component weights can be estimated via a separate parameter  $\pi_k$  for each component or via a concomitant variable model  $\pi(k|x_i, \alpha)$  with parameters  $\alpha$ . Second, the full likelihood function  $f(y_i|\xi_k)$  of the components needs to be defined. If a conditional ML approach is adopted, it is clear that the conditional likelihood  $h(y_i|r_i, \beta)$  from Equation 5 should be one part, but various choices for modeling the score probabilities are available. One option is to model each score probability with its own parameter  $g(r_i) = \psi_{r_i}$ , while another (more parsimonious) option would be to adopt a parametric distribution with fewer parameters (Rost and von Davier 1995). Note that while for a single-component model, the estimates of the item parameters  $\hat{\beta}$  are invariant to the choice of the score probabilities (as long as it is independent from  $\beta$ ), this is no longer the case for a mixture model with  $K \geq 2$ .

#### *Rost's original parametrization*

One of these possible mixtures – the so-called “mixed Rasch model” introduced by Rost (1990) – is already well-established in the psychometric literature. It models the score probabilities through separate parameters  $g(r_i) = \psi_{r_i}$  (under the restriction that they sum to unity) and does not employ concomitant variables. The likelihood of Rost's mixture model can thus

be written as

$$f(y|\pi, \psi, \beta) = \prod_{i=1}^n \sum_{k=1}^K \pi_k h(y_i|r_i, \beta_k) \psi_{r_i, k}. \quad (7)$$

This particular parametrization is implemented in the R package **mRm** (Preinerstorfer 2011). Since subjects who solve either none or all items (i.e.,  $r_i = 0$  or  $m$ , respectively) do not contribute to the conditional likelihood of the item parameters they cannot be allocated to any of the components in this parametrization. Hence, Rost (1990) proposed to remove those “extreme scorers” from the analysis entirely and fix the corresponding score probabilities  $\psi_0$  and  $\psi_m$  at 0. However, if one wishes to include these extreme scorers in the analysis, the corresponding score probabilities can be estimated through their relative frequency (across all components) and the remaining score probabilities within each component are rescaled to sum to unity together with those extreme score probabilities. Nevertheless, the extreme scorers still do not contribute to the estimation of the mixture itself.

#### *Other score distributions*

As noted by Rost and von Davier (1995), the disadvantage of this saturated model for the raw score probabilities is that many parameters need to be estimated ( $K \times (m - 2)$ , not counting potential extreme scorers) that are typically not of interest. To check for DIF, the item parameters are of prime importance while the raw score distribution can be regarded as a nuisance term. This problem can be alleviated by embedding the model from Equation 7 into a more general framework that also encompasses more parsimonious parametrizations. More specifically, a conditional logit model can be established

$$g(r|\delta) = \frac{\exp\{z_r^\top \delta\}}{\sum_{j=1}^{m-1} \exp\{z_j^\top \delta\}}, \quad (8)$$

containing some auxiliary regressors  $z_i$  with coefficients  $\delta$ .

The saturated  $g(r_i) = \psi_{r_i}$  model is a special case when constructing the auxiliary regressor from indicator/dummy variables for the raw scores  $2, \dots, m - 1$ :  $z_i = (I_2(r_i), \dots, I_{m-1}(r_i))^\top$ . Then  $\delta = (\log(\psi_2) - \log(\psi_1), \dots, \log(\psi_{m-1}) - \log(\psi_1))^\top$  is a simple logit transformation of  $\psi$ .

As an alternative Rost and von Davier (1995) suggests a specification with only two parameters that link to mean and variance of the score distribution, respectively. More specifically, the auxiliary regressor is  $z_i = (r_i/m, 4r_i(m - r_i)/m^2)^\top$  so that  $\delta$  pertains to the vector of location and dispersion parameters of the score distribution.

#### *General Rasch mixture model*

Combining all elements of the likelihood this yields a more general specification of the Rasch mixture model

$$f(y|\pi, \alpha, \beta, \delta) = \prod_{i=1}^n \sum_{k=1}^K \pi(k|x_i, \alpha) h(y_i|r_i, \beta_k) g(r_i|\delta_k). \quad (9)$$

with (a) the concomitant model  $\pi(k|x_i, \alpha)$  for modeling component membership, (b) the component-specific conditional likelihood of the item parameters given the scores  $h(y_i|r_i, \beta_k)$ , and (c) the component-specific score distribution  $g(r_i|\delta_k)$ .

## 2.4. Parameter estimation

Parameter estimation for mixture models is usually done via the expectation-maximization (EM) algorithm (Dempster, Laird, and Rubin 1977). It treats group membership as unknown and optimizes the full likelihood including the group membership on basis of the observed values only. It iterates between two steps until convergence: estimation of group membership (E-step) and estimation of the components (M-step).

In the E-step, the posterior probabilities of each observation for the  $k$  components is estimated through:

$$\hat{p}_{ik} = \frac{\hat{\pi}_k f(y_i|\hat{\xi}_k)}{\sum_{g=1}^K \hat{\pi}_g f(y_i|\hat{\xi}_g)} \quad (10)$$

using the parameter estimates for  $\pi$  and  $\xi$  from the previous iteration. In the case of concomitant variables, the component weights are  $\hat{\pi}_{ik} = \pi(k|x_i, \hat{\alpha})$ .

In the M-step, the parameters of the mixture are re-estimated with the posterior probabilities as weights. Thus, observations deemed unlikely to belong to a certain component have little influence on estimation within this component. For each component, the weighted ML estimation can be written as

$$\begin{aligned} \hat{\xi}_k &= \operatorname{argmax}_{\xi_k} \sum_{i=1}^n \hat{p}_{ik} \log f(y_i|\xi_k) \quad (k = 1, \dots, K) \\ &= \left\{ \operatorname{argmax}_{\beta_k} \sum_{i=1}^n \hat{p}_{ik} \log h(y_i|r_i, \beta_k); \operatorname{argmax}_{\delta_k} \sum_{i=1}^n \hat{p}_{ik} \log g(r_i|\delta_k) \right\} \end{aligned} \quad (11)$$

which for the Rasch model amounts to separately maximizing the weighted conditional log-likelihood for the item parameters and the weighted score log-likelihood.

The concomitant model can be estimated separately from the posterior probabilities, e.g., for a multinomial model:

$$\hat{\alpha} = \operatorname{argmax}_{\alpha} \sum_{i=1}^n \sum_{k=1}^K \hat{p}_{ik} \log(\pi(k|x_i, \alpha)). \quad (12)$$

Finally, note that the number of components  $K$  is not a standard model parameter (because the likelihood regularity conditions do not apply) and thus it is not estimated through the EM algorithm. Either it needs to be chosen by the practitioner or by model selection techniques such as information criteria, as illustrated in the following examples.

## 3. Implementation in R

### 3.1. User interface

The function `raschmix()` can be used to fit the different flavors of Rasch mixture models described in Section 2.3: with or without concomitant variables in  $\pi(k|x_i, \alpha)$ , and with different score distributions  $g(r_i|\delta_k)$  (saturated vs. mean/variance parametrization). The function's synopsis is



```

raschmix(formula, data, k, subset, weights, scores = "saturated",
  nrep = 3, cluster = NULL, control = NULL,
  verbose = TRUE, drop = TRUE, unique = FALSE, which = NULL,
  gradtol = 1e-6, deriv = "sum", hessian = FALSE, ...)

```

where the lines of arguments pertain to (1) data/model specification processed within `raschmix()`, (2) control arguments for fitting a single mixture model, (3) control arguments for iterating across mixtures over a range of numbers of components  $K$ , all passed to `stepFlexmix()`, and (4) control arguments for fitting each model component within a mixture (i.e., the M-step) passed to `RaschModel.fit()`. Details are provided below, focusing on usage in practice first.

A formula interface with the usual `formula`, `data`, `subset`, and `weights` arguments is used: The left-hand side of the formula sets up the response matrix  $y$  and the right-hand side the concomitant variables  $x$  (if any). The response may be provided by a single matrix or a set of individual dummy vectors, both of which may be contained in an optional data frame. Example usages are `raschmix(resp ~ 1, ...)` if the matrix `resp` is an object in the working environment or `raschmix(item1 + item2 + item3 ~ 1, data = d, ...)` if the `item*` vectors are in the data frame `d`. In both cases, `~ 1` signals that there are no concomitant variables – if there were, they could be specified as `raschmix(resp ~ conc1 + conc2, ...)`. As an additional convenience, the formula may be omitted entirely if there are no concomitant variables, i.e., `raschmix(data = resp, ...)` or alternatively `raschmix(resp, ...)`.

The `scores` of the model can be set to either `"saturated"` (see Equation 7) or `"meanvar"` for the mean/variance specification of Rost and von Davier (1995). Finally, the number of components  $K$  of the mixture is specified through `k`, which may be a vector resulting in a mixture model being fitted for each element.

To control the EM algorithm for fitting the specified mixture models, `cluster` may optionally specify starting probabilities  $\hat{p}_{ik}$  and `control` can set certain control arguments through a named list or an object of class `"FLXcontrol"`. One of these control arguments named `minprior` sets the minimum prior probability for every component. If in an iteration of the EM algorithm, any component has a prior probability smaller than `minprior`, it is removed from the mixture in the next iteration. The default is 0, i.e., avoiding such shrinkage of the model. If `cluster` is not provided, `nrep` different random initializations are employed, keeping only the best solution (to avoid local optima). Finally, `cluster` can be set to `"mrm"` in which case the fast C++ implementation from `mRm` (Preinerstorfer 2011) can be leveraged to generate optimized starting values. Again, the best solution of `nrep` runs of `mrm()` is used. Note that as of version 1.0 of `mRm` only the model from Equation 7 is supported in `mrm()`, resulting in suboptimal – but potentially still useful – posterior probabilities  $\hat{p}_{ik}$  for any other model flavor.

Internally, `stepFlexmix()` is called to fit all individual mixture models and takes control arguments `verbose`, `drop`, and `unique`. If `k` is a vector, the whole set of models is returned by default but one may choose to select only the best model according to an information criterion. For example, `raschmix(resp, k = 1:3, which = "AIC", ...)` or `raschmix(resp ~ 1, data = d, k = 1:4, which = "BIC", ...)`.

The arguments `gradtol`, `deriv` and `hessian` are used to control the estimation of the item parameters in each M-step (Equation 11) carried out via `RaschModel.fit()`.

Function	Class	Description
<code>summary()</code>	“ <code>raschmix</code> ”	display information about the posterior probabilities and item parameters; returns an object of class “ <code>summary.raschmix</code> ” containing the relevant summary statistics (which has a <code>print()</code> method)
<code>parameters()</code>	“ <code>raschmix</code> ”	extract estimated parameters of the model for all or specified components, extract either parameters $\alpha$ of concomitant model or item parameters $\beta$ and/or score parameters $\delta$
<code>worth()</code>	“ <code>raschmix</code> ”	extract the item parameters $\beta$ under the restriction $\sum_{j=1}^m \beta_j = 0$
<code>scoreProbs()</code>	“ <code>raschmix</code> ”	extract the score probabilities $g(r \delta)$
<code>plot()</code>	“ <code>raschmix</code> ”	base graph of item parameter profiles in all or specified components
<code>xyplot()</code>	“ <code>raschmix</code> ”	lattice graph of item parameter profiles of all or specified components in a single or multiple panels
<code>histogram()</code>	“ <code>raschmix</code> ”	lattice rootogram or histogram of posterior probabilities
<code>print()</code>	“ <code>stepFlexmix</code> ”	simple printed display of number of components, log-likelihoods, and information criteria
<code>plot()</code>	“ <code>stepFlexmix</code> ”	plot information criteria against number of components
<code>getModel()</code>	“ <code>stepFlexmix</code> ”	select model according to either an information criterion or the number of components
<code>print()</code>	“ <code>flexmix</code> ”	simple printed display with cluster sizes and convergence
<code>clusters()</code>	“ <code>flexmix</code> ”	extract predicted class memberships
<code>posterior()</code>	“ <code>flexmix</code> ”	extract posterior class probabilities
<code>logLik()</code>	“ <code>flexmix</code> ”	extract fitted log-likelihood
<code>AIC(); BIC()</code>	“ <code>flexmix</code> ”	compute information criteria AIC, BIC

Table 1: Methods for objects of class “`raschmix`”.

Function `raschmix()` returns objects of class “`raschmix`” or “`stepRaschmix`”, respectively, depending on whether a single or multiple mixture models are fitted. These classes extend “`flexmix`” and “`stepFlexmix`”, respectively, for more technical details see the next section. For standard methods for extracting or displaying information, either for “`raschmix`” directly or by inheritance, see Table 1 for an overview.

### 3.2. Internal structure

As briefly mentioned above, `raschmix()` leverages the **flexmix** package (Leisch 2004; Grün and Leisch 2008) and particularly its `stepFlexmix()` function for the estimation of (sets of) mixture models.

The **flexmix** package is designed specifically to provide the infrastructure for flexible mixture modelling via the EM algorithm, where the type of a mixture model is determined through

the model employed in the components. In the estimation process, this component model definition corresponds to the definition of the M-step (Equation 11). Consequently, the **flexmix** package provides the framework for fitting mixture models by leveraging the modular structure of the EM algorithm. Provided with the right M-step, **flexmix** takes care of the data handling and iterating estimation through both E-step and M-step.

The M-step needs to be provided in the form of a **flexmix** driver inheriting from class “FLXM” (see Grün and Leisch 2008, for details). The **psychomix** package includes such a driver function: `FLXMCrasch()` relies on the function `RaschModel.fit()` from the **psychotools** package for estimation of the item parameters (i.e., maximization of the conditional likelihood from Equation 5) and adds different estimates of raw score probabilities depending on their parameterization.

The reason for employing `RaschModel.fit()` rather than one of the more established Rasch model packages such as **eRm** or **ltm** is speed: `RaschModel.fit()` has been designed with reduced flexibility in order to save time when refitted multiple times as in Rasch mixture models or also Rasch trees in the **psychotree** package (Strobl *et al.* 2011).

In the **flexmix** package, two fitting functions are provided. `flexmix()` is designed for fitting one model once and returns an object of class “flexmix”. `stepFlexmix()` extends this so that either a single model or several models can be fitted. It also provides the functionality to fit each model repeatedly to avoid local optima.

When fitting models repeatedly, only the solution with the highest likelihood is returned. Thus, if `stepFlexmix()` is used to repeatedly fit a single model, it returns an object of class “flexmix”. If `stepFlexmix()` is used to fit several models (repeatedly or just once), it returns an object of class “stepFlexmix”.

This principle extends to `raschmix()`: If it is used to fit a single model, the returned object is of class “raschmix”. If used for fitting multiple models, `raschmix()` returns an object of class “stepRaschmix”. Both classes extend their **flexmix** counterparts.

### 3.3. Illustrations

For illustrating the flexible usage of `raschmix()`, we employ an artificial data set drawn from one of the three data generating processes (DGPs) suggested by Rost (1990) for the introduction of Rasch mixture models. All three DGPs are provided in the function `simRaschmix()` setting the `design` to “rost1”, “rost2”, or “rost3”, respectively. The DGPs contain mixtures of  $K = 1$ , and 2, and 3 components, respectively, all with  $m = 10$  items.

The DGP “rost1” is intended to illustrate the model’s capacity to correctly determine when no DIF is present. Thus, it includes only one latent class with item parameters  $\beta^{(1)} = (2.7, 2.1, 1.5, 0.9, 0.3, -0.3, -0.9, -1.5, -2.1, -2.7)^\top$ . (Rost originally used opposite signs to reflect item easiness parameters but since difficulty parameters are estimated by `raschmix()` the signs have been reversed.) The DGP “rost2” draws observations from two latent classes of equal sizes with item parameters of opposite signs:  $\beta^{(1)}$  and  $\beta^{(2)} = -\beta^{(1)}$ , respectively (see Figure 2 for an example). Finally, the DGP “rost3” adds a third component of smaller size with item parameters  $\beta^{(3)} = (-0.5, 0.5, -0.5, 0.5, -0.5, 0.5, -0.5, 0.5, -0.5, 0.5)^\top$ . In all three DGPs, the person parameters  $\theta$  are drawn from a discrete uniform distribution on  $\{2.7, 0.9, -0.9, -2.7\}$ , except for the third class of DGP “rost3” which uses only one level of ability,  $\{0\}$ . In all DGPs, response vectors for 1800 subjects are initially drawn but the extreme scorers who solved either none or all items are excluded.

Here, a dataset from the second DGP is generated along with two artificial covariates  $x_1$  and  $x_2$ . Covariate  $x_1$  is an informative binary variable (i.e., correlated with the true group membership) while  $x_2$  is an uninformative continuous variable.

```
R> set.seed(1)
R> r2 <- simRaschmix(design = "rost2")
R> d <- data.frame(
+   x1 = rbinom(nrow(r2), prob = c(0.4, 0.6)[attr(r2, "group")], size = 1),
+   x2 = rnorm(nrow(r2))
+ )
R> d$resp <- r2
```

The [Rost \(1990\)](#) version of the Rasch mixture model – i.e., with a saturated score model and without concomitant variables – is fitted for one to three components. As no concomitants are employed in this model flavor, the matrix  $r_2$  can be passed to `raschmix()` without formula:

```
R> m1 <- raschmix(r2, k = 1:3)
R> m1
```

Call:

```
raschmix(formula = r2, k = 1:3)
```

	iter	converged	k	k0	logLik	AIC	BIC	ICL
1	2	TRUE	1	1	-10409.177	20852.35	20944.07	20944.07
2	8	TRUE	2	2	-8738.606	17547.21	17736.04	17809.87
3	105	TRUE	3	3	-8724.237	17554.47	17840.42	18247.37

To inspect the results, the returned object can either be printed, as illustrated above, or plotted yielding a visualization of information criteria (see [Figure 1](#)). Both printed display and visualization show a big difference in information criteria across number of components  $K$ , with the minimum always being assumed for  $K = 2$ , thus correctly recovering the two latent classes constructed in the underlying DGP.

The values of the information criteria can also be accessed directly via the functions of the corresponding names. To select a certain model from a “`stepRaschmix`” object, the `getModel()` function from the `flexmix` package can be employed. The specification of `which` model is to be selected can either be an information criterion, or the number of components as a string, or the index of the model in the original vector  $k$ . In this particular case, `which = "BIC"`, `which = "2"`, and `which = 2` would all return the model with  $K = 2$  components.

```
R> BIC(m1)
```

```
      1      2      3
20944.07 17736.04 17840.42
```

```
R> m1b <- getModel(m1, which = "BIC")
R> summary(m1b)
```




Figure 1: Information criteria for Rost's model with  $K = 1, 2, 3$  components for the artificial scenario 2 data.

Call:

```
raschmix(formula = r2, k = 2)
```

	prior	size	post>0	ratio
Comp.1	0.498	808	1262	0.640
Comp.2	0.502	820	1264	0.649

Item Parameters:

	Comp.1	Comp.2
Item01	-2.5574826	2.4538561
Item02	-2.1844001	2.1573277
Item03	-1.4524543	1.6382597
Item04	-1.0042438	0.9254200
Item05	-0.3639643	0.2675059
Item06	0.3074967	-0.3868124
Item07	0.8828234	-0.8563867
Item08	1.5048074	-1.4900729
Item09	2.1186345	-2.0921240
Item10	2.7487831	-2.6169734

```
'log Lik.' -8738.606 (df=35)
AIC: 17547.21 BIC: 17736.04
```

To inspect the main properties of the model, `summary()` can be called. The information about the components of the mixture includes a priori component weights  $\pi_k$  and sizes as well as the estimated item parameters  $\hat{\beta}$  per component. Additionally, the fitted log-likelihood and

the information criteria AIC and BIC are reported. As one of the item parameters in the Rasch model is not identified, a restriction needs to be applied to the item parameters. In the output of the `summary()` function, the item parameters of each component are scaled to sum to zero.

Two other functions, `worth()` and `parameters()`, can be used to access the item parameters. The sum restriction employed in the `summary()` output is also applied by `worth()`. Additionally, `worth()` provides the possibilities to select several or just one specific component and to transform item difficulty parameters to item easiness parameters. The function `parameters()` also offers these two options but restricts the first item parameter to be zero (rather than the sum of item parameters), as this restriction is used in the internal computations. Thus, for the illustrative dataset with 10 items, `parameters()` returns 9 item parameters, leaving out the first item parameter restricted to zero while `worth()` returns 10 item parameters summing to zero. The latter corresponds to the parametrization employed by Rost (1990) and `simRaschmix()`. For convenience reasons, the true parameters are attached to the simulated dataset as an attribute named "item". These are printed below and visualized in Figure 2 (left), showing that all item parameters are recovered rather well. Note that the ordering of the components in mixture models is generally arbitrary.

```
R> parameters(m1b, "item")
```

	Comp.1	Comp.2
item.Item02	0.3730825	-0.2965283
item.Item03	1.1050282	-0.8155963
item.Item04	1.5532388	-1.5284360
item.Item05	2.1935182	-2.1863502
item.Item06	2.8649793	-2.8406685
item.Item07	3.4403060	-3.3102427
item.Item08	4.0622900	-3.9439290
item.Item09	4.6761171	-4.5459800
item.Item10	5.3062656	-5.0708294

```
R> worth(m1b)
```

	Comp.1	Comp.2
Item01	-2.5574826	2.4538561
Item02	-2.1844001	2.1573277
Item03	-1.4524543	1.6382597
Item04	-1.0042438	0.9254200
Item05	-0.3639643	0.2675059
Item06	0.3074967	-0.3868124
Item07	0.8828234	-0.8563867
Item08	1.5048074	-1.4900729
Item09	2.1186345	-2.0921240
Item10	2.7487831	-2.6169734

```
R> attr(r2, "item")
```

```

      beta1 beta2
1      2.7  -2.7
2      2.1  -2.1
3      1.5  -1.5
4      0.9  -0.9
5      0.3  -0.3
6     -0.3   0.3
7     -0.9   0.9
8     -1.5   1.5
9     -2.1   2.1
10    -2.7   2.7

```

In addition to the item parameters, the `parameters()` function can also return the parameters of the "score" model and the "concomitant" model (if any). The type of parameters can be set via the `which` argument. Per default `parameters()` returns both item and score parameters.

A comparison between estimated and true class membership can be conducted using the `clusters()` function and the corresponding attribute of the data, respectively. As already noticeable from the item parameters, the first component of the mixture matches the second true group of the data and vice versa. This label-switching property of mixture models in general can also be seen in the cross-table of class memberships. We thus have 38 misclassifications among the 1628 observations.

```
R> table(model = clusters(m1b), true = attr(r2, "group"))
```

```

      true
model  1  2
      1 16 792
      2 798  22

```

For comparison, a Rasch mixture model with mean/variance parametrization for the score probabilities, as introduced in Section 2.3, is fitted with one to three components and the best BIC model is selected.

```
R> m2 <- raschmix(data = r2, k = 1:3, scores = "meanvar")
```

```
R> m2
```

Call:

```
raschmix(data = r2, k = 1:3, scores = "meanvar")
```

```

      iter converged k k0      logLik      AIC      BIC      ICL
1       2      TRUE 1  1 -10416.199 20854.40 20913.74 20913.74
2       8      TRUE 2  2  -8747.084 17540.17 17664.25 17737.65
3      60      TRUE 3  3  -8738.314 17546.63 17735.46 17939.80

```

```
R> m2b <- getModel(m2, which = "BIC")
```




Figure 2: True (black) and estimated (blue/red) item parameters for the two model specifications, "saturated" (left) and "meanvar" (right), for the artificial scenario 2 data.

As in the saturated version of the Rasch mixture model, all three information criteria prefer the two-component model. Thus, this version of a Rasch mixture model is also capable of recognizing the two latent classes in the data while using a more parsimonious parametrization with 23 instead of 35 parameters.

```
R> logLik(m2b)
```

```
'log Lik.' -8747.084 (df=23)
```

```
R> logLik(m1b)
```

```
'log Lik.' -8738.606 (df=35)
```

The estimated parameters of the distribution of the score probabilities can be accessed through `parameters()` while the full set of score probabilities is returned by `scoreProbs()`. The estimated score probabilities of the illustrative model are approximately equal across components and roughly uniform.

```
R> parameters(m2b, which = "score")
```

	Comp.1	Comp.2
score.location	-0.10292905	-0.04187548
score.dispersion	-0.09456975	-0.07905495

```
R> scoreProbs(m2b)
```



	Comp.1	Comp.2
0	0.0000000	0.0000000
1	0.1198694	0.1163454
2	0.1155415	0.1133228
3	0.1122157	0.1110791
4	0.1098133	0.1095705
5	0.1082785	0.1087681
6	0.1075758	0.1086567
7	0.1076895	0.1092340
8	0.1086219	0.1105110
9	0.1103944	0.1125124
10	0.0000000	0.0000000

The resulting item parameters for this particular data set are virtually identical to those from the saturated version, as can be seen in Figure 2.

To demonstrate the use of a concomitant variable model for the weights of the mixture, the two artificial variables `x1` and `x2` are employed. They are added on the right-hand side of the formula, yielding a multinomial logit model for the weights (only if `k = 2` or more components are specified).

```
R> cm2 <- raschmix(resp ~ x1 + x2, data = d, k = 2:3, scores = "meanvar")
```

The BIC is used to compare the models with and without concomitant variables. In both cases, the two true groups are recognized correctly, while the model with concomitants manages to employ the additional information and reaches a somewhat improved model fit.

```
R> rbind(m2 = BIC(m2), cm2 = c(NA, BIC(cm2)))
```

	1	2	3
m2	20913.74	17664.25	17735.46
cm2	NA	17611.05	17691.59

As mentioned above, the parameters of the concomitant model can be accessed via the `parameters()` function, setting `which = "concomitant"`. The influence of the informative covariate `x1` is reflected in the large absolute coefficient while the estimated coefficient for the noninformative covariate `x2` is close to zero.

```
R> cm2b <- getModel(cm2, which = "BIC")
R> parameters(cm2b, which = "concomitant")
```

	1	2
(Intercept)	0	0.43688740
x1	0	-0.85033332
x2	0	0.03043432

The corresponding estimated item parameters `parameters(cm2b, "item")` are not very different from the previous models (and are hence not shown here). This illustrative application

shows that the inclusion of concomitant variables can provide additional information, e.g., that *x1* but not *x2* is associated with the class membership. Note also that this is picked up although a rather weak association was simulated here.

```
R> table(x1 = d$x1, clusters = clusters(cm2b))
```

```

  clusters
x1   1   2
  0 320 494
  1 486 328
```

#### 4. Empirical application: Verbal aggression

The verbal aggression dataset (De Boeck and Wilson 2004) contains item response data from 316 first-year psychology students along with gender and trait anger (assessed by the Dutch adaptation of the state-trait anger scale) as covariates (Smits, De Boeck, and Vansteelandt 2004). The 243 women and 73 men responded to 24 items constructed the following way: Following the description of a frustrating situation, subjects are asked to agree or disagree with a possible reaction. The situations are described by the following four sentences:

- S1: A bus fails to stop for me.
- S2: I miss a train because a clerk gave me faulty information.
- S3: The grocery store closes just as I am about to enter.
- S4: The operator disconnects me when I had used up my last 10 cents for a call.

Each reaction begins with either “I want to” or “I do” and is followed by one of the three verbally aggressive reactions “curse”, “scold”, or “shout”, e.g., “I want to curse”, “I do curse”, “I want to scold”, or “I do scold”.

For our illustration, we use only the first two sentences which describe situations in which the others are to blame. Extreme-scoring subjects agreeing with either none or all responses are removed.

```
R> data("VerbalAggression", package = "psychotools")
R> VerbalAggression$resp2 <- VerbalAggression$resp2[, 1:12]
R> va12 <- subset(VerbalAggression,
+   rowSums(resp2) > 0 & rowSums(resp2) < 12)
R> colnames(va12$resp2)
```

```
[1] "S1WantCurse" "S1DoCurse"   "S1WantScold" "S1DoScold"
[5] "S1WantShout" "S1DoShout"   "S2WantCurse" "S2DoCurse"
[9] "S2WantScold" "S2DoScold"   "S2WantShout" "S2DoShout"
```

We fit Rasch mixture models with the mean/variance score model, one to four components, and with and without the two concomitant variables, respectively (the single component model being only fitted without covariates).




Figure 3: Rootogram of posterior probabilities in the 3-component Rasch mixture model on verbal aggression data.

```
R> set.seed(1)
R> va12_mix1 <- raschmix(resp2 ~ 1, data = va12, k = 1:4, scores = "meanvar")
R> va12_mix2 <- raschmix(resp2 ~ gender + anger, data = va12, k = 2:4,
+   scores = "meanvar")
```

The corresponding BIC for all considered models can be computed by

```
R> rbind(BIC(va12_mix1), c(NA, BIC(va12_mix2)))
```

	1	2	3	4
[1,]	3874.632	3857.567	3854.350	3887.361
[2,]	NA	3859.119	3854.822	3880.484

```
R> va12_mix3 <- getModel(va12_mix1, which = "3")
```

showing that three components are preferred regardless of whether or not concomitant variables are used. In this case, they do not lead to a better model fit, thus the 3-class model without concomitant variables is chosen.

The posterior probabilities for the three components can be visualized via `histogram(va12_mix3)` – by default using a square-root scale, yielding a so-called rootogram – as shown in Figure 3. In the ideal case, posterior probabilities of the observations for each component are either high or low, yielding a U-shape in all panels. In this case here, the components are separated acceptably well.

The item profiles in three components can be visualized via `plot(va12_mix3)` or `xyplot(va12_mix3)` with the output of the latter being shown in Figure 4. The first six items are responses to the first sentence (bus), the remaining six refer to the second sentence (train). The six reactions are grouped in “want”/“do” pairs: first for “curse”, then “scold”, and finally “shout”.




Figure 4: Item profiles for the 3-component Rasch mixture model on verbal aggression data. Items 1–6 pertain to situation S1 (bus), items 7–12 to situation S2 (train), each in the following order: want to curse, do curse, want to scold, do scold, want to shout, do shout.

The third component displays a zigzag pattern which indicates that subjects in this component always find it easier or less extreme to “want to” react a certain way rather than to actually “do” react that way. In the other two components this want/do relationship is reversed, except for the shouting response (to either situation).

In the first component, there are no big differences in the estimated item parameters. Neither the situation (S1 or S2) nor the type of verbal response (curse, scold, or shout) is particularly hard to agree to for subjects in this component. In components 2 and 3, the situation is also not very relevant but subjects differentiate between the three verbal responses. This is best visible in component 2 where item difficulty is clearly increasing from response “curse” to response “shout”. Thus, shouting is perceived as the most extreme verbal response while cursing is considered a comparably moderate response. In component 3 this pattern is also visible albeit not as prominently as in component 2.

One could also consider the 3-component model *with* concomitant variables as its BIC was almost equivalent to that of the model *without* concomitant variables. The estimated item parameters are virtually identical between both models and are hence not shown here. Nevertheless, the link between the concomitant variables and the latent classes may still be of interest:

```
R> parameters(getModel(va12_mix2, which = "3"), which = "concomitant")
```

	1	2	3
(Intercept)	0	-0.76040110	-3.6721134
gendermale	0	1.66471685	1.4177908
anger	0	0.01155322	0.1268023

The absolute sizes of the coefficients reflect that there may be some association with gender but less with the anger score. However, as there is a slight increase in BIC compared to the model

without concomitants, the association with the covariates appears to be relatively weak. In comparison to other approaches exploring the association of class membership with covariates *ex post* (e.g., as in Cohen and Bolt 2005), the main advantage of the concomitant variables model lies in the *simultaneous* estimation of the mixture and the influence of covariates.

## 5. Summary

Mixtures of Rasch models are a flexible means of checking measurement invariance and testing for differential item functioning. Here, we establish a rather general unifying conceptual framework for Rasch mixture models along with the corresponding computational tools in the R package **psychomix**. In particular, this includes the original model specification of Rost (1990) as well as more parsimonious parameterizations (Rost and von Davier 1995), along with the possibility to incorporate concomitant variables predicting the latent classes (as in Tay *et al.* 2011).

The R implementation is based on the infrastructure provided by the **flexmix** package, allowing for convenient model specification and selection. The rich set of methods for **flexmix** objects is complemented by additional functions specifically designed for Rasch models, e.g., extracting different types of parameters in different transformations and visualizing the estimated component-specific item parameters in various ways. Optionally, speed gains can be obtained from utilizing the C++ implementation in the **mRm** package for selecting optimal starting values. Thus, **psychomix** provides a comprehensive and convenient toolbox for the application of Rasch mixture models in psychometric research practice.

## Acknowledgments

We are thankful to the participants of the “Psychoco 2011” workshop at Universität Tübingen for helpful feedback and discussions.

## References

- Ackerman TA (1992). “A Didactic Explanation of Item Bias, Item Impact, and Item Validity From a Multidimensional Perspective.” *Journal of Educational Measurement*, **29**(1), 67–91.
- Andersen EB (1972). “A Goodness of Fit Test for the Rasch Model.” *Psychometrika*, **38**(1), 123–140.
- Cohen AS, Bolt DM (2005). “A Mixture Model Analysis of Differential Item Functioning.” *Journal of Educational Measurement*, **42**(2), 133–148.
- Dayton CM, Macready G (1988). “Concomitant-Variable Latent-Class Models.” *Journal of the American Statistical Association*, **83**(401), 173–178.
- De Boeck P, Wilson M (eds.) (2004). *Explanatory Item Response Models: A Generalized Linear and Nonlinear Approach*. Springer-Verlag, New York.
- Dempster A, Laird N, Rubin D (1977). “Maximum Likelihood from Incomplete Data via the EM Algorithm.” *Journal of the Royal Statistical Society B*, **39**(1), 1–38.

- Fischer GH, Molenaar IW (eds.) (1995). *Rasch Models: Foundations, Recent Developments, and Applications*. Springer-Verlag, New York.
- Grün B, Leisch F (2008). “FlexMix Version 2: Finite Mixtures with Concomitant Variables and Varying and Constant Parameters.” *Journal of Statistical Software*, **28**(4), 1–35. URL <http://www.jstatsoft.org/v28/i04/>.
- Leisch F (2004). “FlexMix: A General Framework for Finite Mixture Models and Latent Class Regression in R.” *Journal of Statistical Software*, **11**(8), 1–18. URL <http://www.jstatsoft.org/v11/i08/>.
- Mair P, Hatzinger R (2007). “Extended Rasch Modeling: The **eRm** Package for the Application of IRT Models in R.” *Journal of Statistical Software*, **20**(9), 1–20. URL <http://www.jstatsoft.org/v20/i09/>.
- Preinerstorfer D (2011). *mRm: An R Package for Conditional Maximum Likelihood Estimation in Mixed Rasch Models*. R package version 1.0, URL <http://CRAN.R-project.org/package=mRm>.
- Rasch G (1960). *Probabilistic Models for Some Intelligence and Attainment Tests*. The University of Chicago Press.
- R Development Core Team (2011). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
- Rizopoulos D (2006). “**ltm**: An R Package for Latent Variable Modeling and Item Response Theory Analyses.” *Journal of Statistical Software*, **17**(5), 1–25. URL <http://www.jstatsoft.org/v17/i05/>.
- Rost J (1990). “Rasch Models in Latent Classes: An Integration of Two Approaches to Item Analysis.” *Applied Psychological Measurement*, **14**(3), 271–282.
- Rost J, von Davier M (1995). “Mixture Distribution Rasch Models.” In GH Fischer, IW Molenaar (eds.), *Rasch Models: Foundations, Recent Developments, and Applications*, chapter 14, pp. 257–268. Springer-Verlag, New York.
- Smits DJM, De Boeck P, Vansteelandt K (2004). “The Inhibition of Verbally Aggressive Behaviour.” *European Journal of Personality*, **18**(7), 537–555.
- Strobl C, Kopf J, Zeileis A (2011). “A New Method for Detecting Differential Item Functioning in the Rasch Model.” *Working Paper 2011-01*, Working Papers in Economics and Statistics, Research Platform Empirical and Experimental Economics, Universität Innsbruck. URL <http://EconPapers.RePEc.org/RePEc:inn:wpaper:2011-01>.
- Tay L, Newman DA, Vermunt JK (2011). “Using Mixed-Measurement Item Response Theory with Covariates (MM-IRT-C) to Ascertain Observed and Unobserved Measurement Equivalence.” *Organizational Research Methods*, **14**(1), 147–176.
- Van den Noortgate W, De Boeck P (2005). “Assessing and Explaining Differential Item Functioning Using Logistic Mixed Models.” *Journal of Educational and Behavioral Statistics*, **30**(4), 443–464.

Zeileis A, Strobl C, Wickelmaier F (2011). *psychotools: Infrastructure for Psychometric Modeling*. R package version 0.1-1, URL <http://CRAN.R-project.org/package=psychotools>.

**Affiliation:**

Hannah Frick, Achim Zeileis

Department of Statistics

Universität Innsbruck

Universitätsstr. 15

6020 Innsbruck, Austria

E-mail: [Hannah.Frick@uibk.ac.at](mailto:Hannah.Frick@uibk.ac.at), [Achim.Zeileis@R-project.org](mailto:Achim.Zeileis@R-project.org)

URL: <http://eeecon.uibk.ac.at/~frick/>, <http://eeecon.uibk.ac.at/~zeileis/>

Carolin Strobl

Department of Psychology

Universität Zürich

Binzmühlestr. 14

8050 Zürich, Switzerland

E-mail: [Carolin.Strobl@psychologie.uzh.ch](mailto:Carolin.Strobl@psychologie.uzh.ch)

URL: <http://www.psychologie.uzh.ch/fachrichtungen/methoden.html>

Friedrich Leisch

Institute of Applied Statistics and Computing

Universität für Bodenkultur Wien

Peter Jordan-Str. 82

1180 Wien, Austria

E-mail: [Friedrich.Leisch@R-project.org](mailto:Friedrich.Leisch@R-project.org)

URL: <http://www.rali.boku.ac.at/friedrichleisch.html>

University of Innsbruck - Working Papers in Economics and Statistics  
Recent Papers can be accessed on the following webpage:

<http://eeecon.uibk.ac.at/wopec/>

- 2011-21 **Hannah Frick, Carolin Strobl, Friedrich Leisch, Achim Zeileis:** Flexible Rasch mixture models with package psychomix
- 2011-20 **Thomas Grubinger, Achim Zeileis, Karl-Peter Pfeiffer:** evtree: Evolutionary learning of globally optimal classification and regression trees in R
- 2011-19 **Wolfgang Rinnergschwentner, Gottfried Tappeiner, Janette Walde:** Multivariate stochastic volatility via wishart processes - A continuation
- 2011-18 **Jan Verbesselt, Achim Zeileis, Martin Herold:** Near Real-Time Disturbance Detection in Terrestrial Ecosystems Using Satellite Image Time Series: Drought Detection in Somalia
- 2011-17 **Stefan Borsky, Andrea Leiter, Michael Pfaffermayr:** Does going green pay off? The effect of an international environmental agreement on tropical timber trade
- 2011-16 **Pavlo Blavatskyk:** Stronger Utility
- 2011-15 **Anita Gantner, Wolfgang Höchtel, Rupert Sausgruber:** The pivotal mechanism revisited: Some evidence on group manipulation
- 2011-14 **David J. Cooper, Matthias Sutter:** Role selection and team performance
- 2011-13 **Wolfgang Höchtel, Rupert Sausgruber, Jean-Robert Tyran:** Inequality aversion and voting on redistribution
- 2011-12 **Thomas Windberger, Achim Zeileis:** Structural breaks in inflation dynamics within the European Monetary Union
- 2011-11 **Loukas Balafoutas, Adrian Beck, Rudolf Kerschbamer, Matthias Sutter:** What drives taxi drivers? A field experiment on fraud in a market for credence goods
- 2011-10 **Stefan Borsky, Paul A. Raschky:** A spatial econometric analysis of compliance with an international environmental agreement on open access resources
- 2011-09 **Edgar C. Merkle, Achim Zeileis:** Generalized measurement invariance tests with application to factor analysis



- 2011-08 **Michael Kirchler, Jürgen Huber, Thomas Stöckl:** Thar she bursts - reducing confusion reduces bubbles *modified version forthcoming in American Economic Review*
- 2011-07 **Ernst Fehr, Daniela Rützler, Matthias Sutter:** The development of egalitarianism, altruism, spite and parochialism in childhood and adolescence
- 2011-06 **Octavio Fernández-Amador, Martin Gächter, Martin Larch, Georg Peter:** Monetary policy and its impact on stock market liquidity: Evidence from the euro zone
- 2011-05 **Martin Gächter, Peter Schwazer, Engelbert Theurl:** Entry and exit of physicians in a two-tiered public/private health care system
- 2011-04 **Loukas Balafoutas, Rudolf Kerschbamer, Matthias Sutter:** Distributional preferences and competitive behavior *forthcoming in Journal of Economic Behavior and Organization*
- 2011-03 **Francesco Feri, Alessandro Innocenti, Paolo Pin:** Psychological pressure in competitive environments: Evidence from a randomized natural experiment: Comment
- 2011-02 **Christian Kleiber, Achim Zeileis:** Reproducible Econometric Simulations
- 2011-01 **Carolin Strobl, Julia Kopf, Achim Zeileis:** A new method for detecting differential item functioning in the Rasch model
- 2010-29 **Matthias Sutter, Martin G. Kocher, Daniela Rützler and Stefan T. Trautmann:** Impatience and uncertainty: Experimental decisions predict adolescents' field behavior
- 2010-28 **Peter Martinsson, Katarina Nordblom, Daniela Rützler and Matthias Sutter:** Social preferences during childhood and the role of gender and age - An experiment in Austria and Sweden *Revised version forthcoming in Economics Letters*
- 2010-27 **Francesco Feri and Anita Gantner:** Baragining or searching for a better price? - An experimental study. *Revised version accepted for publication in Games and Economic Behavior*
- 2010-26 **Loukas Balafoutas, Martin G. Kocher, Louis Putterman and Matthias Sutter:** Equality, equity and incentives: An experiment
- 2010-25 **Jesús Crespo-Cuaresma and Octavio Fernández Amador:** Business cycle convergence in EMU: A second look at the second moment
- 2010-24 **Lorenz Goette, David Huffman, Stephan Meier and Matthias Sutter:** Group membership, competition and altruistic versus antisocial punishment: Evidence from randomly assigned army groups

- 2010-23 **Martin Gächter and Engelbert Theurl:** Health status convergence at the local level: Empirical evidence from Austria (*revised Version March 2011*)
- 2010-22 **Jesús Crespo-Cuaresma and Octavio Fernández Amador:** Buiness cycle convergence in the EMU: A first look at the second moment
- 2010-21 **Octavio Fernández-Amador, Josef Baumgartner and Jesús Crespo-Cuaresma:** Milking the prices: The role of asymmetries in the price transmission mechanism for milk products in Austria
- 2010-20 **Fredrik Carlsson, Haoran He, Peter Martinsson, Ping Qin and Matthias Sutter:** Household decision making in rural China: Using experiments to estimate the influences of spouses
- 2010-19 **Wolfgang Brunauer, Stefan Lang and Nikolaus Umlauf:** Modeling house prices using multilevel structured additive regression
- 2010-18 **Martin Gächter and Engelbert Theurl:** Socioeconomic environment and mortality: A two-level decomposition by sex and cause of death
- 2010-17 **Boris Maciejovsky, Matthias Sutter, David V. Budescu and Patrick Bernau:** Teams make you smarter: Learning and knowledge transfer in auctions and markets by teams and individuals
- 2010-16 **Martin Gächter, Peter Schwazer and Engelbert Theurl:** Stronger sex but earlier death: A multi-level socioeconomic analysis of gender differences in mortality in Austria
- 2010-15 **Simon Czermak, Francesco Feri, Daniela Rützler and Matthias Sutter:** Strategic sophistication of adolescents - Evidence from experimental normal-form games
- 2010-14 **Matthias Sutter and Daniela Rützler:** Gender differences in competition emerge early in live
- 2010-13 **Matthias Sutter, Francesco Feri, Martin G. Kocher, Peter Martinsson, Katarina Nordblom and Daniela Rützler:** Social preferences in childhood and adolescence - A large-scale experiment
- 2010-12 **Loukas Balafoutas and Matthias Sutter:** Gender, competition and the efficiency of policy interventions
- 2010-11 **Alexander Strasak, Nikolaus Umlauf, Ruth Pfeifer and Stefan Lang:** Comparing penalized splines and fractional polynomials for flexible modeling of the effects of continuous predictor variables
- 2010-10 **Wolfgang A. Brunauer, Sebastian Keiler and Stefan Lang:** Trading strategies and trading profits in experimental asset markets with cumulative information

- 2010-09 **Thomas Stöckl and Michael Kirchler:** Trading strategies and trading profits in experimental asset markets with cumulative information
- 2010-08 **Martin G. Kocher, Marc V. Lenz and Matthias Sutter:** Psychological pressure in competitive environments: Evidence from a randomized natural experiment: Comment
- 2010-07 **Michael Hanke and Michael Kirchler:** Football Championships and Jersey sponsors' stock prices: An empirical investigation
- 2010-06 **Adrian Beck, Rudolf Kerschbamer, Jianying Qiu and Matthias Sutter:** Guilt from promise-breaking and trust in markets for expert services - Theory and experiment
- 2010-05 **Martin Gächter, David A. Savage and Benno Torgler:** Retaining the thin blue line: What shapes workers' intentions not to quit the current work environment
- 2010-04 **Martin Gächter, David A. Savage and Benno Torgler:** The relationship between stress, strain and social capital
- 2010-03 **Paul A. Raschky, Reimund Schwarze, Manijeh Schwindt and Ferdinand Zahn:** Uncertainty of governmental relief and the crowding out of insurance
- 2010-02 **Matthias Sutter, Simon Czermak and Francesco Feri:** Strategic sophistication of individuals and teams in experimental normal-form games
- 2010-01 **Stefan Lang and Nikolaus Umlauf:** Applications of multilevel structured additive regression models to insurance data

University of Innsbruck

Working Papers in Economics and Statistics

2011-21

Hannah Frick, Carolin Strobl, Friedrich Leisch, Achim Zeileis

Flexible Rasch mixture models with package psychomix

**Abstract**

Measurement invariance is an important assumption in the Rasch model and mixture models constitute a flexible way of checking for a violation of this assumption by detecting unobserved heterogeneity in item response data. Here, a general class of Rasch mixture models is established and implemented in R, using conditional maximum likelihood estimation of the item parameters (given the raw scores) along with flexible specification of two model building blocks: (1) Mixture weights for the unobserved classes can be treated as model parameters or based on covariates in a concomitant variable model. (2) The distribution of raw score probabilities can be parametrized in two possible ways, either using a saturated model or a specification through mean and variance. The function `raschmix()` in the R package "psychomix" provides these models, leveraging the general infrastructure for fitting mixture models in the "flexmix" package. Usage of the function and its associated methods is illustrated on artificial data as well as empirical data from a study of verbally aggressive behavior.

ISSN 1993-4378 (Print)  
ISSN 1993-6885 (Online)