



Dipartimento

Scienze

Economiche

Working Paper

Department
of Economics

Ca' Foscari University of
Venice

Jean-Michel Grandmont

Nonlinear Difference Equations,
Bifurcations and Chaos: an
Introduction



Nonlinear Difference Equations, Bifurcations and Chaos: An Introduction

Jean-Michel Grandmont

ICEF, University of Venice and CREST-GRECSTA (UMR CNRS 2773), Paris

Abstract

The aim of these lecture notes is to present a few mathematical facts about the bifurcations of nonlinear difference equations, in a concise and simple form that might be useable by economic theorists.

Keywords

Nonlinear dynamics, Bifurcations, Chaos, Business cycles

JEL Codes

C02, C61, E32

Address for correspondence.

Jean-Michel Grandmont
Department of Economics
Ca' Foscari University of Venice
Cannaregio 873, Fondamenta S.Giobbe
30121 Venezia - Italy
Phone: (+39) 041 2349173
Fax: (+39) 041 2349176
Jean-Michel.Grandmont@ensae.fr

This Working Paper is published under the auspices of the Department of Economics of the Ca' Foscari University of Venice. Opinions expressed herein are those of the authors and not those of the Department. The Working Paper series is designed to divulge preliminary or incomplete work, circulated to favour discussion and comments. Citation of this paper should consider its provisional character.

A. PRELIMINARIES

We state first a few more or less elementary facts about matrices and differentiable maps, that are used repeatedly in the sequel.

We recall that \mathbb{R}^m is the set of all m -tuples of real numbers. A “point” or a “vector” of \mathbb{R}^m is $x = (x_1, \dots, x_m)$; the number x_i is the i -th *coordinate* of the vector. Vectors x, y are added coordinatewise

$$x + y = (x_1, \dots, x_m) + (y_1, \dots, y_m) = (x_1 + y_1, \dots, x_m + y_m).$$

If α is a real number, the product αx is the vector $(\alpha x_1, \dots, \alpha x_m)$. \mathbb{R}^m is then an m -dimensional real vector space. Its *standard basis* is the collection of vectors (e_1, \dots, e_m) , in which for each $i = 1, \dots, m$, e_i is the vector of coordinates $e_{ij} = \delta_{ij}$, $j = 1, \dots, m$, where δ_{ij} is the Kronecker function, that is $\delta_{ij} = 0$ if $i \neq j$ and 1 if $i = j$. Any vector $x = (x_1, \dots, x_m)$ has then a unique representation as a linear combination of the vectors e_i of the standard basis, that is $x = \sum_i x_i e_i$. A *norm* is a real valued function $\|\cdot\|$ defined on \mathbb{R}^m , with $\|x\| \geq 0$, such that $\|\alpha x\| = |\alpha| \|x\|$, $\|x + y\| \leq \|x\| + \|y\|$, and $\|x\| = 0$ if and only if $x = 0$. The *Euclidean norm* will be denoted $|x| = (\sum_i x_i^2)^{1/2}$.

A.1. Matrix algebra

A square *matrix* of dimension m , i.e. a collection of m^2 real numbers $A = [a_{ij}]$, where $i = 1, \dots, m$ stands for the index of the i -th row of the matrix, and $j = 1, \dots, m$ stands for its j -th column, defines a *linear transformation* (or *map*) T from \mathbb{R}^m into itself, that associates to every vector $x = (x_1, \dots, x_m)$ a new vector $x' = Tx$ of coordinates $x'_i = \sum_j a_{ij} x_j$, for $i = 1, \dots, m$, or in matrix notation, $x' = Ax$. Then if e_1, \dots, e_m is the standard basis of \mathbb{R}^m , the vector represented by the j -th column of A , i.e. $a^j = (a_{1j}, \dots, a_{mj})$, is the image of e_j by T (or A), that is $a^j = Te_j = Ae_j$. The image $x' = Ax$ of any vector $x = (x_1, \dots, x_m)$ is the linear combination of the vectors a^j , with weights x_j :

$$x' = A(\sum_j x_j e_j) = \sum_j x_j a^j$$

Conversely, any linear transformation T from \mathbb{R}^m into itself can be (uniquely) represented by the matrix $A = [a_{ij}]$, in the standard basis, where the j -th column $a^j = (a_{1j}, \dots, a_{mj})$ of the matrix A is the image by T of the vector e_j . It follows from these remarks that a matrix A is *invertible* if and only if the corresponding linear transformation T is onto (i.e. the image of \mathbb{R}^m by T is \mathbb{R}^m itself) or equivalently, if and only if the m vectors a^j are *linearly independent* (i.e. $\sum_j \alpha_j a^j = 0$ implies $\alpha_j = 0$ for all j).

A given linear transformation T of \mathbb{R}^m into itself has different equivalent matrix representations, according to which basis of \mathbb{R}^m is chosen. Consider a *new basis* of \mathbb{R}^m , i.e. a collection of m vectors $\bar{e}_1, \dots, \bar{e}_m$, that are linearly independent. Let (p_{1j}, \dots, p_{mj}) be the coordinates of \bar{e}_j in the standard basis, and P stand for the matrix of which the j -th columns is \bar{e}_j , i.e. $P = [p_{ij}]$. We know from the previous paragraph that P has an inverse P^{-1} . A vector of \mathbb{R}^m of which the coordinates in the old (standard) basis are $x = (x_1, \dots, x_m)$, has coordinates $y = (y_1, \dots, y_m)$ in the new basis. That is, this vector can be (uniquely) expressed as a linear combination of the vectors \bar{e}_j of the new basis, with weights y_j , i.e. $\sum_j y_j \bar{e}_j$. The relationship between new and old coordinates is obtained from the vector equalities

$$\sum_i x_i e_i = \sum_j y_j \bar{e}_j = \sum_j y_j \left(\sum_i p_{ij} e_i \right),$$

which imply $x_i = \sum_j p_{ij} y_j$ for all i , or in matrix notation, $x = Py$, $y = P^{-1}x$.

A given linear transformation T is represented, in matrix notation, by the map $x \rightarrow x' = Ax$ in the standard basis, and by $y \rightarrow y' = By$ in the new basis. Analytically, the matrix B is obtained from A by making the change of variables $x = Py$, which yields $B = P^{-1}AP$. Here again, the j -column of B represents the coordinates, in the new basis, of the image of \bar{e}_j by T .

A linear transformation T of \mathbb{R}^m into itself may thus be given a convenient matrix representation, by choosing an appropriate basis. The remainder of this section is devoted to such a matrix representation, the *real canonical (or Jordan) form* of T .

We look first at the circumstances ensuring that T has a *block diagonal* matrix representation. Let E_1, \dots, E_r be a collection of (linear) subspaces of \mathbb{R}^m , i.e. each E_h is a subset of \mathbb{R}^m that is closed under the operations of addition and scalar multiplication : if x, y are vectors of E_h and α a real number, then $x + y$ and αx belong also to E_h . Assume that any vector x of \mathbb{R}^m has a unique representation of the form $x = x_1 + \dots + x_r$, in which x_h is in E_h for each h . We say then that \mathbb{R}^m is the *direct sum* of the linear subspaces. Assume further that each subspace E_h is *invariant* by T , i.e. if x belongs to E_h , then Tx is also in E_h . Choose now a basis for each E_h , and take the union of the basis elements of the E_h to obtain a basis for \mathbb{R}^m . In that basis, T has the block diagonal form

$$B = \text{diag} \{B_1, \dots, B_r\} = \begin{bmatrix} B_1 & & \\ & \ddots & \\ & & B_r \end{bmatrix}$$

This means that the matrix B_h are put together corner-to-corner diagonally as indicated, all other entries in B being zero (we adopt the convention that the blank entries in a matrix are zeros). Each matrix B_h represents in fact the restriction T_h of T to the invariant subspace E_h .

Conversely, assume that \mathbb{R}^m has a basis in which T has a matrix representation of the above block diagonal form. Let E_h be the linear subspace spanned by the vectors of the basis, the images of which are the columns of the matrix B associated to the submatrix B_h . Then E_h is invariant by T , and \mathbb{R}^m is the direct sum of the E_h . To sum up,

Proposition A.1.1. *Let T be a linear transformation of \mathbb{R}^m into itself. T has a block diagonal matrix representation if and only if there exists a collection of linear subspaces E_h of \mathbb{R}^m , $h = 1, \dots, r$, such that 1) each E_h is invariant by T , and 2) \mathbb{R}^m is the direct sum of the E_h .*

The real canonical matrix representation of T is obtained when the invariant subspaces E_h are taken to be the “real generalized eigenspaces” of T (to be defined shortly), with an appropriate basis.

Distinct eigenvalues

Let A be the matrix representation of T in the standard basis. We recall that an *eigenvalue* λ of T is a (possibly complex) number such that $A\xi = \lambda\xi$ has a nonzero solution ξ , where ξ is an m -dimensional vector, the coordinates of which may be complex numbers. Such a nonzero solution ξ is called an *eigenvector* (of T) corresponding to λ . The set of (zero and nonzero) solutions ξ of $A\xi = \lambda\xi$ is the *eigenspace* corresponding to λ , and is noted $E(\lambda)$.

$E(\lambda)$ is a complex vector space, i.e. if ξ, η are in $E(\lambda)$ and z is a complex number, then $\xi + \eta$ and $z\xi$ are also in $E(\lambda)$. If, however, λ is real, the equation $A\xi = \lambda\xi$ has a nonzero solution x in \mathbb{R}^m , a *real eigenvector*. The *real eigenspace* $F(\lambda)$ is then the set of all real solutions of $A\xi = \lambda\xi$. $F(\lambda)$ is a linear subspace of \mathbb{R}^m , and $E(\lambda)$ is its “complexification” : every ξ in $E(\lambda)$ is of the form $u + iv$, where u, v belong to $F(\lambda)$ and $i = \sqrt{-1}$.

If the eigenvalue is nonreal, i.e. $\lambda = a + ib$ with $b \neq 0$, then every eigenvector ξ corresponding to λ must be nonreal, i.e. have the form $u + iv$ where u, v are vectors of \mathbb{R}^m and $v \neq 0$. In that case, the complex conjugate of λ , i.e. $\bar{\lambda} = a - ib$, is also an eigenvalue, and $\xi = u + iv$ belongs to $E(\lambda)$ if and only if its complex conjugate $\bar{\xi} = u - iv$ belongs to $E(\bar{\lambda})$. This follows from the fact that $A\xi = \bar{\lambda}\bar{\xi}$ is equivalent, through complex conjugation, to $A\bar{\xi} = \lambda\xi$. Consider next the set $F(\lambda)$ of all vectors of \mathbb{R}^m that lie in

in $E(\mu_k)$. The whole basis is actually $\{x_1, \dots, x_r, v_1, u_1, \dots, v_s, u_s\}$. For details, see Hirsch and Smale (1974), Chapter 4, Section 2.

Multiple eigenvalues

We look now at the case in which T may have multiple eigenvalues. The characteristic polynomial can always be factorized as

$$p(\lambda) \equiv (\lambda_1 - \lambda)^{n_1} \dots (\lambda_q - \lambda)^{n_q}$$

where the λ_i are the distinct eigenvalues of T , and n_i is the *multiplicity* of λ_i . One has $n_1 + \dots + n_q = m$, and T has multiple eigenvalues if some $n_i \geq 2$.

In such case, one has to introduce the notions of “generalized” eigenvectors and eigenspaces. Let λ be an eigenvalue of T , and n its multiplicity. The *generalized eigenspace* $E^*(\lambda)$ corresponding to λ is the space of m -dimensional complex vectors ξ that solve $(A - \lambda I)^n \xi = 0$. If ξ is in $E^*(\lambda)$ and $\xi \neq 0$, then ξ is a *generalized eigenvector*. Of course if $n = 1$, these notions reduce to the usual concepts of an eigenspace and an eigenvector.

The arguments employed above to define real eigenvectors or eigenspaces apply equally well here. If λ is real, the space of all vectors in $E^*(\lambda)$ that are real, is the *generalized real eigenspace* $F^*(\lambda)$ corresponding to λ . If x is in $F^*(\lambda)$ and $x \neq 0$, it is a *generalized real eigenvector*. If λ is nonreal, the *generalized real eigenspace* $F^*(\lambda)$ (corresponding to λ) is the set of all real vectors of $E^*(\lambda) + E^*(\bar{\lambda})$. Here again, there is a slight abuse of language in such a definition, since there are no real generalized eigenvectors when λ is nonreal.

One can verify that the dimension of $E^*(\lambda)$ is n ; the dimension of $F^*(\lambda)$ is thus n when λ is real, $2n$ otherwise. It is easy to see that $E^*(\lambda)$ and $F^*(\lambda)$ are invariant by T .

Theorem A.1.4. *Let T be a linear transformation of \mathbb{R}^m into itself. Let $\lambda_1, \dots, \lambda_r$ be its distinct real eigenvalues, with λ_h having multiplicity n_h , and $\mu_1, \bar{\mu}_1, \dots, \mu_s, \bar{\mu}_s$ its distinct nonreal eigenvalues, with $\mu_k = a_k + ib_k$, $b_k > 0$, having multiplicity n_k . Then \mathbb{R}^m is the direct sum of the generalized real eigenspaces $F^*(\lambda_1), \dots, F^*(\lambda_r), F^*(\mu_1), \dots, F^*(\mu_s)$. The dimension of $F^*(\lambda_h)$ is n_h , while the dimension of $F^*(\mu_k)$ is $2n_k$. Each of these eigenspaces is invariant by T .*

Moreover, \mathbb{R}^m has a basis in which T has the following block diagonal matrix representation

Remark A.1.5. In the Theorem, the basis is the union of bases for each generalized real eigenspace $F^*(\lambda_1), \dots, F^*(\lambda_r), F^*(\mu_1), \dots, F^*(\mu_s)$, in that order. For $F^*(\mu_k)$, the basis is of the form $\{v_1, u_1, \dots, v_s, u_s\}$, where $\{u_1 + iv_1, \dots, u_s + iv_s\}$ is a basis for $E^*(\mu_k)$. For details, see Hirsch and Smale (1974, Chapter 6, Section 4, and Appendix III).

A.2. The implicit function theorem

Let G be a map from an open subset W of \mathbb{R}^m into \mathbb{R}^m . We say that G is C^r with $r \geq 1$, if G has continuous partial derivatives of every order $h = 1, \dots, r$ at each point of W . G is a *homeomorphism* if it is continuous and one to one, and if its inverse G^{-1} (a map from $G(W)$ onto W) is also continuous. G is a C^r *diffeomorphism* if it is C^r and one to one, and if its inverse is also C^r .

If G is C^1 , the matrix of partial first derivatives at a point x of W is called the *Jacobian matrix* of G at x , and is noted $DG(x)$.

The following result gives conditions ensuring that a C^r map is locally a diffeomorphism.

Theorem A.2.1. (Inverse function theorem). *Let G be a C^r map from an open set W into \mathbb{R}^m , with $r \geq 1$. Let x be a point of W and assume that the Jacobian matrix $DG(x)$ is invertible. Then x has an open neighbourhood U such that the restriction of G to U is a C^r diffeomorphism onto the open set $G(U)$.*

Under the conditions of the theorem, we shall say that $G^{-1} : G(U) \rightarrow U$ is a *local inverse* of G . Application of the inverse function theorem permits then to see when an implicit equation $F(x, y) = c$ can be solved and generate, at least locally, a functional relation of the form $y = G(x)$.

Theorem A.2.2. (Implicit function theorem). *Let W be an open set in $\mathbb{R}^m \times \mathbb{R}^p$, and F a C^r map from W into \mathbb{R}^p , that is $(x, y) \rightarrow F(x, y)$ where x and y are vectors of \mathbb{R}^m and \mathbb{R}^p , respectively. Let (x_o, y_o) in W be such that $F(x_o, y_o) = c$ and suppose that the Jacobian matrix of the map $F(x_o, \cdot)$ is invertible at $y = y_o$.*

Then there are open sets U and V in \mathbb{R}^m and \mathbb{R}^p , respectively, with x_o in U , y_o in V and $U \times V$ contained in W , and a unique C^r map $G : U \rightarrow V$, such that

$$F(x, G(x)) = c$$

for all x in U , and moreover, $F(x, y) \neq c$ if x is in U , y in V and $y \neq G(x)$.

Notes on the literature

The material of Section A.1 can be found in any textbook on linear algebra. The presentation here is adapted from Hirsch and Smale (1974), see also Palis and De Melo (1982, chap. 2.2). For a proof of the inverse and the implicit function theorems, see Hirsch and Smale (1974, Appendix IV).

B. LINEAR AND NONLINEAR DIFFERENCE EQUATIONS

We shall be concerned in this section, and in the next two ones, with the qualitative behaviour of the trajectories generated by a difference equation of the form

$$(B.1) \quad x_{n+1} = G(x_n)$$

in which G is a map from an open subset U of \mathbb{R}^m into \mathbb{R}^m . Our first specific task, which will be carried out presently, is to proceed to such a study in the immediate vicinity of a *fixed point* of G (also called a *stationary state*, an *equilibrium point*), i.e. an element \bar{x} of U such that $\bar{x} = G(\bar{x})$. It will be seen in particular that if G is continuously differentiable, the trajectories generated by (B.1) that are close enough to \bar{x} , are generally similar to those that are associated to the “linearized” version

$$(B.2) \quad x_{n+1} - \bar{x} = DG(\bar{x}) (x_n - \bar{x})$$

in which $DG(\bar{x})$ stands for the Jacobian matrix of G , evaluated at the fixed point.

A frequently encountered situation giving rise to a formulation such as (B.1) is the following one. Suppose that the successive states y_n of a given (physical, social) system must satisfy

$$(B.3) \quad F(y_{n+1}, y_n, \dots, y_{n-N}) = 0$$

in which each y is a vector of \mathbb{R}^p , and F is a C^r map from an open set W of $\mathbb{R}^{p(N+2)}$ into \mathbb{R}^p . Let \bar{y} be a stationary state, i.e. a vector of \mathbb{R}^p such that $F(\bar{y}, \dots, \bar{y}) = 0$, and assume that the Jacobian matrix of the map $y \rightarrow F(y, \bar{y}, \dots, \bar{y})$, evaluated at \bar{y} , is invertible. The Implicit Function Theorem A.2.2 implies that (B.3) can then be solved in y_{n+1} in a neighbourhood of \bar{y} . That is, there exist open sets U in $\mathbb{R}^{p(N+1)}$ and V in \mathbb{R}^p , containing $(\bar{y}, \dots, \bar{y})$ and \bar{y} respectively, and a unique C^r map $H : U \rightarrow V$ such that $(y_{n+1}, y_n, \dots, y_{n-N})$ satisfies (B.3), with (y_n, \dots, y_{n-N}) in U and y_{n+1} in V , if and only if

$$(B.4) \quad y_{n+1} = H(y_n, \dots, y_{n-N}).$$

The resulting “delayed” difference equation can now be cast in the form (B.1) by considering the variable $x_n = (y_n, \dots, y_{n-N})$, and the map G that associates to every such x_n in U the vector $x_{n+1} = (y_{n+1}, \dots, y_{n-N+1})$, with

y_{n+1} given by (B.4). One may note that $\bar{x} = (\bar{y}, \dots, \bar{y})$ is a fixed point of G , and that the Jacobian matrix of G at \bar{x} has the form

$$DG(\bar{x}) = \begin{bmatrix} D_0H & D_1H & \cdots & D_NH \\ I_p & 0 & & \\ & \ddots & \ddots & \\ & & I_p & 0 \end{bmatrix}$$

in which D_kH , $k = 0, \dots, N$, stands for the matrix of partial derivatives of H with respect to the components y_{n-k} , evaluated at $(\bar{y}, \dots, \bar{y})$, and I_p is the p -dimensional identity matrix. It is easy to see that $DG(\bar{x})$ is invertible if and only if D_NH is itself invertible. In that case, by the Inverse Function Theorem A.2.1, one can choose U small enough to ensure that G is actually a C^r diffeomorphism onto the open set $G(U)$.

B.1. Stability

Consider the difference equation (B.1) in which G maps the open subset U of \mathbb{R}^m into \mathbb{R}^m . For any x in U , one constructs the *trajectory* (or the *orbit*) of x by using repeatedly (B.1) for $n \geq 0 : x_0 = x, x_1 = G(x), \dots, x_n = G^n(x)$, in which $G^n(x)$ is the n -th *iterate* of x and is defined recursively through $G^n(x) = G(G^{n-1}(x))$. Of course, it may occur that for some n , $G^n(x)$ no longer belongs to U ; in that case, the orbit leaves the domain of definition of G and is defined for only finitely many $n \geq 0$. If x belongs to an *invariant set* K , i.e. a subset of U such that $G(K)$ is contained in K , then the orbit of x is defined for all $n \geq 0$. This will be the case for all x , in particular, if $U = \mathbb{R}^m$.

A fixed point \bar{x} of G is *stable* if for every open subset V of U containing \bar{x} , there is an open subset V_1 of V , with \bar{x} also in V_1 , such that for every x in V_1 , the iterates $G^n(x)$ are well defined and belong to V for all $n \geq 0$. If in addition, V can be chosen so that the distance of $G^n(x)$ to \bar{x} tends to zero as $n \rightarrow +\infty$ for each x in V_1 , then \bar{x} is *asymptotically stable*. A fixed point is *unstable* if it is not stable.

Remark B.1.1. The foregoing notions of (asymptotic) stability apply not only to fixed points, but also to any closed set K , without any change.

Remark B.1.2. The analysis of the present section applies immediately to the study of what happens near a *periodic orbit*. Suppose that $G^n(\bar{y}) = \bar{y}$ for some \bar{y} in U and $n > 1$, with $G^j(\bar{y}) \neq \bar{y}$ for $j = 1, \dots, n-1$. Then \bar{y} is

a *periodic point*, of period n , and the corresponding periodic orbit is defined as $\bar{y}, G(\bar{y}), \dots, G^{n-1}(\bar{y})$. The map $F = G^n$ is defined at least on a small open neighbourhood V of \bar{y} . Stability or unstability of the periodic orbit in the dynamics generated by G is equivalent to stability or unstability of the fixed point \bar{y} in the dynamics generated by F .

B.2. Change of variables

There is nothing intrinsic in the formulation embodied in (B.1). Indeed, one may always make a change of variables of the form $y = h(x)$, in which y is a vector of \mathbb{R}^m and h an homeomorphism from \mathbb{R}^m onto itself. The difference equation obtained after the change of variable is $y_{n+1} = F(y_n)$, in which $F = h \circ G \circ h^{-1}$ is a map from $h(U)$ into \mathbb{R}^m (actually, h need be defined only on an open set containing both U and $G(U)$). The two maps F and G are then said to be *topologically conjugate*. If h is a C^r diffeomorphism, they are *C^r conjugate*. Clearly, two difference equations (or two maps) deduced from each other through conjugation generate the same trajectories (up to the change of variable) ; they must therefore be viewed as *equivalent*.

Such an equivalence relation may be only local. Let \bar{x} be a fixed point of G , and consider another map $y \rightarrow F(y)$, with fixed point \bar{y} . Then G is *locally topologically* (or *C^r*) *conjugate* fo F if there is an open neighbourhood V of \bar{x} , and an open neighbourhood W of \bar{y} , such that the restriction of G to V is topologically (or *C^r*) conjugate to the restriction of F to W . In that case, only the orbits generated by G that are sufficiently near \bar{x} , are the same, up to the change of variable, to those of F that are near \bar{y} .

This notion of equivalence between maps or difference equations is important, since it allows one transforming a given difference equation, which may be difficult to analyse, and bringing it through an appropriate change of variables into a more tractable form, where the qualitative features of the trajectories may be more clearly seen. This equivalence notion leads also, quite naturally, to the idea of structural stability.

If the equation (B.1) is meant to describe the evolution of some physical or social phenomenon, one cannot be exactly sure about all the details of the specification. To get reliable results, one should accordingly work with maps G that yield a qualitative picture that remains unaltered, up to a change of variable, when they are slightly perturbed. To be precise, assume that $G : U \rightarrow \mathbb{R}^m$ is C^r , and let us endow the space of all C^r maps $F : U \rightarrow \mathbb{R}^m$ with the so-called *C^r - topology*, i.e. the topology of uniform convergence, on compacta, of the values of the functions and of its derivatives up to the

order r . The map G is then C^r - *structural stable* if there is a neighbourhood V of G , in the C^r topology, such that all C^r perturbations F of G that lie in V , are conjugate to G . The qualitative features of the trajectories generated by a structurally stable map are unchanged after small perturbations.

Here again, structural stability may be required only locally. The C^r map $G : U \rightarrow \mathbb{R}^m$ is C^r *locally structurally stable* (near a fixed point \bar{x}) if there exists an open neighbourhood V of \bar{x} such that the restriction of G to V is C^r - structurally stable.

The foregoing notion of structural stability involves *arbitrary* small perturbations of the map G under consideration. If particular features of the phenomenon to be represented impose restrictions (e.g. symmetry) on the class of maps to be considered, then of course, perturbations and structural stability should be defined *relatively* to this particular class of maps.

Remark B.2.1. Difference equations may be defined more generally on manifolds (a subset M of \mathbb{R}^m is a C^r *manifold of dimension* $m \leq p$, if each element of M has a neighbourhood V for which there is a C^r diffeomorphism that maps V onto \mathbb{R}^m). We shall need only exceptionally to consider explicitly such a more general formulation. But it may be worth noting at this stage that the study of the qualitative properties of trajectories near a fixed point in such a case, may be reduced to the analysis of the text, through an appropriate change of variable. Specifically, let $y_{n+1} = F(y_n)$ be the difference equation to be studied, where F is a map from an open subset of a C^r m -dimensional manifold M into M . If \bar{y} is a fixed point of F , there is a neighbourhood V of \bar{y} , and a C^r diffeomorphism h mapping V onto \mathbb{R}^m . If F is continuous, there is an open neighbourhood W of \bar{y} such that both W and $F(W)$ are subsets of V . Making the change of variable $x = h(y)$, with y in W , shows that the restriction of F to W is C^r conjugate to the map $G : U \rightarrow \mathbb{R}^m$, where $U = h(W)$ and $G = h \circ F \circ h^{-1}$.

B.3. Linear difference equations

Linear difference equations of the form

$$(B.5) \quad x_{n+1} = Ax_n$$

in which x is a point of \mathbb{R}^m and A is an m -dimensional square matrix, are important to study on their own right. In addition, as mentioned earlier, the trajectories associated to a nonlinear difference equation like (B.1) that are

near enough a fixed point \bar{x} , generally behave qualitatively as those associated to the linearized version (B.2).

An obvious fixed point of (B.5) is the origin $x = 0$, and an important issue is to determine its stability. The issue is not difficult to analyze, since the orbit of a point x_0 of \mathbb{R}^m through the repeated action of (B.5) is given by $x_n = A^n x_0$, in which A^n is the n -th power of A , for all $n \geq 0$. A^n may not be easy to compute, however, so we appeal to Theorem A.1.4, to assert the existence of a linear change of variable $x = Py$, in which P is invertible, that transforms (B.5) in

$$(B.6) \quad y_{n+1} = By_n$$

where $B = P^{-1}AP$ is the real canonical form of A . The matrix B is much easier to work with, since it is block diagonal, i.e. $B = \text{diag}\{B_1, \dots, B_p\}$, in which each B_i is the matrix corresponding to a generalized real eigenspace F_i^* associated to an eigenvalue (and its conjugate if it is nonreal), equipped with a canonical basis. Since each eigenspace is invariant under the action of (B.6), one can solve separately each equation $\xi_{n+1} = B_i \xi_n$, where ξ is a vector in the eigenspace F_i^* , which leads to $\xi_n = B_i^n \xi_0$. Complete solutions of (B.6) are obtained by piecing together the solutions associated to each eigenspace, and one gets the solution of the original problem by making the change of variable $x = Py$.

Distinct eigenvalues

When A has distinct eigenvalues, the form of B was given in Theorem A.1.2. Restricting attention to the real eigenspace $F(\lambda)$ corresponding to the real eigenvalue λ leads to the onedimensional difference equation $\xi_{n+1} = \lambda \xi_n$, or equivalently to the geometric sequence $\xi_n = \lambda^n \xi_0$, in which the real number ξ_n is the coordinate of y_n in $F(\lambda)$. Thus if $|\lambda| > 1$, the sequence $|\xi_n|$ diverges monotonically to $+\infty$ for every nonzero initial value ξ_0 . If $|\lambda| < 1$, the sequence converges to 0, all points going to 0 on first iterate when $\lambda = 0$ (the map $\xi \rightarrow \lambda \xi$ is then noninvertible). If $\lambda = 1$, then all points of $F(\lambda)$ are fixed points. If $\lambda = -1$, all points of the eigenspace that differ from 0 are periodic, with period 2. The restriction of (B.6) to $F(\lambda)$ is orientation preserving when $\lambda > 0$, and orientation reversing when $\lambda < 0$.

Considering on the other hand the real eigenspace $F(\mu)$ corresponding to the nonreal eigenvalue $\mu = a + ib$, with $b > 0$, leads to the twodimensional difference equation

$$\xi_{n+1} = D\xi_n, \quad \text{with} \quad D = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$$

or equivalently $\xi_n = D^n \xi_0$, where the twodimensional vector ξ_n represents the coordinates of y_n in the canonical basis of $F(\mu)$. Setting $a = \rho \cos \theta$, $b = \rho \sin \theta$ with $\rho = |\mu| \neq 0$ and $0 < \theta < \pi$, one verifies easily that the action of D in \mathbb{R}^2 is a rotation of angle θ around the origin, followed by an homothety, the center of which is the origin, with ratio ρ , see Fig. B.1. Thus

$$D^n = \rho^n \begin{bmatrix} \cos n\theta & -\sin n\theta \\ \sin n\theta & \cos n\theta \end{bmatrix}$$

If $|\mu| > 1$, the sequence $|\xi_n|$ diverges monotonically to $+\infty$ whenever $\xi_0 \neq 0$. It converges monotonically to 0 when $|\mu| < 1$ (the map $\xi \rightarrow D\xi$ is then a contraction). If $|\mu| = 1$, D is a rotation of angle θ around the origin. Thus $|y_n| = |y_0|$ for all n , and every circle with center the origin is left invariant by D . If θ is of the form $2\pi p/q$, where p and q are positive integers that are relatively prime, every point $y \neq 0$ of the plane is periodic, with period q , since then $D^q y = y$. If $\theta/2\pi$ is irrational, no point of the plane is periodic; the trajectory generated by $y_0 \neq 0$ is in fact dense in the circle centered at the origin of radius $|y_0|$. In all cases, the restriction of (B.6) to the real eigenspace $F(\mu)$ is orientation preserving.

Fig. B.1

An important remark is in order here. The foregoing contracting (or expanding) properties of the application $\xi \rightarrow D\xi$ are valid when one uses the Euclidean norm $|\xi|$ of \mathbb{R}^2 . These properties do not hold any longer, in general, when the application is expressed in another coordinates system, e.g. through a change of variable $x = Q\xi$, where Q is an invertible 2-matrix, if one sticks to the Euclidean norm $|x|$. Clearly, however, the map $x \rightarrow QDQ^{-1}x$ has the same contracting or expanding properties as $\xi \rightarrow D\xi$, if one chooses the norm $\|x\| = |Q^{-1}x|$.

Multiple eigenvalues

When A has multiple eigenvalues, the real canonical form B is more complicated, and a little more care is needed. Consider first the case of a multiple real eigenvalue λ . As stated in Theorem A.1.4, the matrix Λ associated to the corresponding real generalized eigenspace $F^*(\lambda)$ in the real canonical form, is block diagonal and composed of blocks of the form




Fig. B.1

$$(\xi_n)_p = \lambda^n (\xi_0)_p + \binom{1}{n} \lambda^{n-1} (\xi_0)_{p-1} + \dots + \binom{s}{n} \lambda^{n-s} (\xi_0)_{p-s}$$

with $s = \text{Min}(n, p - 1)$. It follows that $|\xi_n \rightarrow 0|$ when $|\lambda| < 1$, and that $|\xi_n| \rightarrow +\infty$ when $\xi_0 \neq 0$ and $|\lambda| > 1$, as n goes to $+\infty$. The convergence (or divergence) needs not be monotonic here, however. But one may show that there exists a norm $\|\xi\|$ such that the map $\xi \rightarrow L\xi$ is a contraction of modulus $k < 1$ when $|\lambda| < 1$ (such that the same property is true for the inverse mapping $\xi \rightarrow L^{-1}\xi$ when $|\lambda| > 1$). If $|\lambda| = 1$, the origin is unstable. One may verify that the restriction of (B.6) to $F^*(\lambda)$ is orientation preserving if $\lambda \geq 0$, or if $\lambda < 0$ and the multiplicity of λ is even, and that it is orientation reversing if $\lambda < 0$ and the multiplicity of λ is odd.

The case of a multiple nonreal eigenvalue $\mu = a + ib$, $b > 0$, is handled by similar methods. According to Theorem A.2.4, the matrix Δ associated in the real canonical form to the real generalized eigenspace $F^*(\mu)$, is block diagonal and composed of blocks of the form

$$M = \begin{bmatrix} D & & & & \\ I & \ddots & & & \\ & \ddots & \ddots & & \\ & & & I & D \end{bmatrix} \quad \text{with } D = \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \quad I = \begin{bmatrix} 1 & \\ & 1 \end{bmatrix}.$$

Consider one of these matrices M and let $2q$, with $q \geq 2$, be its dimension (again, the case $q = 1$ was analysed for distinct eigenvalues). Restricting attention to the subspace spanned by the elements of the canonical basis corresponding to M , leads to the study of the equation $\xi_{n+1} = M\xi_n$, or of the sequence $\xi_n = M^n \xi_0$, $n \geq 0$, where ξ_n stands for the coordinates of y_n in (B.6) in the subspace under consideration.

One has here $M = S + N$, in which

$$S = \begin{bmatrix} D & & & & \\ & \ddots & & & \\ & & & & \\ & & & & D \end{bmatrix}, \quad N = \begin{bmatrix} 0 & & & & \\ I & \ddots & & & \\ & \ddots & \ddots & & \\ & & & I & 0 \end{bmatrix}.$$

The matrices S and N commute, and thus

$$M^n = \sum_{h=0}^n \binom{h}{n} D^{n-h} N^h.$$

The form of D^{n-h} was given when considering distinct eigenvalues. If $a = \rho \cos \theta$ and $b = \rho \sin \theta$ with $\rho = |\mu|$ and $0 < \theta < \pi$, then

$$D^{n-h} = \rho^{n-h} \begin{bmatrix} \cos(n-h)\theta & -\sin(n-h)\theta \\ \sin(n-h)\theta & \cos(n-h)\theta \end{bmatrix}.$$

As for N^h , it is easily verified that

$$N^2 = \begin{bmatrix} 0 & & & & & \\ 0 & \ddots & & & & \\ I & \ddots & \ddots & & & \\ & \ddots & \ddots & \ddots & & \\ & & & I & 0 & 0 \end{bmatrix}, \dots, N^{q-1} = \begin{bmatrix} I \end{bmatrix} \text{ and } N^q = 0.$$

The components of ξ_n can then be easily computed from $\xi_n = M^n \xi_0$, in much the same way as for real eigenvalues. It will suffice here to remark that each component of ξ_n is a linear combination of terms of the form

$$\binom{h}{n} \rho^{n-h} \cos(n-h)\theta \quad , \quad \binom{h}{n} \rho^{n-h} \sin(n-h)\theta$$

with $h = 0, \dots, \text{Min}(n, q-1)$, the coefficients being determined by the initial vector ξ_0 . It follows here again that $|\xi_n| \rightarrow 0$ when $\rho = |\mu| < 1$, and $|\xi_n| \rightarrow +\infty$ when $\xi_0 \neq 0$ and $|\mu| > 1$, as n diverges to $+\infty$. The convergence or divergence needs not be monotonic. But there exists a norm $\|\xi\|$ such that $\xi \rightarrow M\xi$ is a contraction of modulus $k < 1$ when $|\mu| < 1$ (the same being true of the inverse mapping $\xi \rightarrow M^{-1}\xi$ if $|\mu| > 1$). If $|\mu| = 1$, the origin is unstable. In all cases, the restriction of (B.6) to $F^*(\mu)$ is orientation preserving.

This completes the analysis of the solutions of (B.6). One important feature of the solutions is the following one.

Theorem B.3.1. *Every component of $y_n = B^n y_0$, $n > 0$, is a linear combination of terms of the form*

$$\binom{h}{n} \rho^{n-h} \cos(n-h)\theta \quad , \quad \binom{h}{n} \rho^{n-h} \sin(n-h)\theta$$

where $\lambda = \rho(\cos \theta + i \sin \theta)$ is an eigenvalue with multiplicity $k \geq 1$, and $h = 0, \dots, \text{Min}(n, k-1)$ with weights determined by the initial vector y_0 (real eigenvalues correspond to $\theta = 0, \pi$). The solutions to (B.5), being obtained

through the linear transformation $x_n = Py_n$, have the same qualitative feature.

The other important finding concerned the stability of the origin in (B.6), and thus in (B.5). To state the results neatly, it is convenient to introduce the following notation. Let $\lambda_1, \dots, \lambda_r$ be the distinct real eigenvalues of A , and $\mu_1, \bar{\mu}_1, \dots, \mu_s, \bar{\mu}_s$ its distinct nonreal eigenvalues. The *stable space* F^s of the origin is defined as the space spanned by all the vectors that lie in the real generalized eigenspaces $F^*(\lambda_h)$ or $F^*(\mu_k)$ such that $|\lambda_h|$ or $|\mu_k| < 1$. The *unstable space* F^u is the space spanned by the eigenspaces $F^*(\lambda_h)$ or $F^*(\mu_k)$ such that $|\lambda_h|$ or $|\mu_k| > 1$. The *center space* F^c is the space spanned by all the real generalized eigenspaces corresponding to an eigenvalue of modulus 1. Each of these spaces F^s, F^u, F^c is of course invariant under (B.5). Moreover

Theorem B.3.2. *There exist a norm $\|x\|$ in \mathbb{R}^m and two constants $0 \leq k_s < 1 < k_u$ such that every solution $x_n = A^n x_0$ of (B.5) with $x_0 \neq 0$ satisfies*

1. $\|x_n\| \leq k_s^n \|x_0\|$ when $x_0 \in F^s$, for all $n \geq 1$.
2. $\|x_n\| > k_u^n \|x_0\|$ when $x_0 \in F^u$, for all $n \geq 1$.
3. The sequence x_n does not converge to 0 when $x_0 \in F^c$.

If all eigenvalues of A are such that $|\lambda| < 1$, the origin is a *sink*; if $|\lambda| < 1$ for some eigenvalues and $|\lambda| > 1$ for the others: a *saddle point*, and if $|\lambda| > 1$ for all eigenvalues: a *source*.

Structural stability

It is clear that the linear map $x \rightarrow Ax$ cannot be structurally stable (even in the class of linear applications of \mathbb{R}^m) if one of the eigenvalues (or more) has modulus one. For the trajectories lying in the center space F^c can then be removed by a slight perturbation of the matrix A . Nor can it be structurally stable if one of the eigenvalues (or more) is 0. One may expect, however, the linear map to be structurally stable if these two "borderline" cases are assumed away. This conjecture is confirmed by the following fact.

Theorem B.3.3. *Assume that the m -dimensional matrix A has no eigenvalue of modulus equal to 0 or 1. Then the linear map $x \rightarrow Ax$ is structurally stable in the class of all linear maps of \mathbb{R}^m .*

B.4. Hyperbolic fixed points

Let us go back to the nonlinear difference equation

$$(B.7) \quad x_{n+1} = G(x_n)$$

where the map $G : U \rightarrow \mathbb{R}$ is C^r , and consider a fixed point \bar{x} of G . A common procedure for studying the solutions of this equation near \bar{x} , is to linearize it and to analyse the solutions of the associated linear difference equation

$$(B.8) \quad y_{n+1} = DG(\bar{x})y_n$$

in which $DG(\bar{x})$ is the Jacobian matrix of G at \bar{x} . The following fact implies that the procedure is valid, provided that $DG(\bar{x})$ is invertible (has no zero eigenvalue), and that \bar{x} is a *hyperbolic fixed point*, i.e. $DG(\bar{x})$ has no eigenvalue of modulus 1.

Theorem B.4.1. *Let G be a C^r map, $r \geq 1$, from the open subset U of \mathbb{R}^m into \mathbb{R}^m . Let \bar{x} be a hyperbolic fixed point such that the Jacobian matrix $DG(\bar{x})$ is invertible. Then G is locally topologically conjugate to the linear map $y \rightarrow DG(\bar{x})y$.*

It follows from the analysis of the previous section that under the conditions of the theorem, the fixed point \bar{x} is asymptotically stable if all eigenvalues λ of $DG(\bar{x})$ satisfy $|\lambda| < 1$, and that it is unstable when one of them is such that $|\lambda| > 1$. As a matter of fact, this conclusion is valid even when some eigenvalues of $DG(\bar{x})$ are 0.

Corollary B.4.2. *Let G be a C^r map, $r \geq 1$, from the open subset U of \mathbb{R}^m into \mathbb{R}^m , and \bar{x} a hyperbolic fixed point. Then \bar{x} is asymptotically stable if all eigenvalues λ of $DG(\bar{x})$ satisfy $|\lambda| < 1$, and unstable if $|\lambda| > 1$ for some λ .*

In view of the result concerning the structural stability of linear maps of the previous section (Theorem B.3.3), one should expect, under the assumptions of Theorem B.4.1, the map G to be locally structurally stable.

Corollary B.4.3. *Under the assumptions of Theorem B.4.1, G is locally structurally stable.*

B.5. Invariant manifolds

It is convenient, when studying the solutions of a linear difference equation $x_{n+1} = Ax_n$, to split the analysis into three parts, by looking separately at

what happens in each of the three invariant subspaces that were introduced in Theorem B.3.2 : the stable space F^s , the unstable space F^u and the center space F^c of the matrix A . It turns out that a similar decomposition is also possible in the case of a nonlinear difference equation $x_{n+1} = G(x_n)$, at least locally, i.e. near a fixed point \bar{x} . The role of the above spaces is then played by local invariant surfaces that are tangent to the stable, unstable and center spaces of the Jacobian matrix $DG(\bar{x})$, respectively.

To be specific, assume that $G : U \rightarrow \mathbb{R}^m$ is C^r , $r \geq 1$, and that \bar{x} is a fixed point. Given a neighbourhood V of \bar{x} , a *local invariant manifold* $W_{(\bar{x})}$ is a C^1 manifold (a continuously differentiable surface) passing through \bar{x} and contained in V , such that $x \in W_{(\bar{x})}$, $x \in V$ and $G(x) \in V$ imply $G(x) \in W_{(\bar{x})}$. A *local stable manifold* $W_{(\bar{x})}^s$ is then a local invariant manifold that is tangent at \bar{x} to (and has the same dimension as) the stable space F^s of the Jacobian matrix $DG(\bar{x})$. Similarly, a *local unstable (or center) manifold* $W_{(\bar{x})}^u$ (or $W_{(\bar{x})}^c$) is a local invariant manifold that is tangent at \bar{x} to (and has the same dimension as) the unstable (or center) space F^u (or F^c) of $DG(\bar{x})$.

Theorem B.5.1. *Let \bar{x} be a fixed point of the C^r map G from the open subset U of \mathbb{R}^m into \mathbb{R}^m ($r \geq 1$). Then there exists a neighbourhood V of \bar{x} such that*

1. *There is a unique local stable manifold $W_{(\bar{x})}^s$ in V . It is C^r .*
2. *If $DG(\bar{x})$ is invertible, there is a unique local unstable manifold $W_{(\bar{x})}^u$ in V (it is actually the local stable manifold of the local inverse G^{-1}). It is C^r .*
3. *Suppose $r \geq 2$. There is a (generally nonunique) local center manifold $W_{(\bar{x})}^c$ in V . It is C^{r-1} , and locally attracting : if x and $G^n(x)$ are in V for all $n \geq 1$, then the distance between $G^n(x)$ and $W_{(\bar{x})}^c$ tends to 0 as $n \rightarrow +\infty$.*

Of course, in the foregoing statement, if the dimension of one of the spaces F is 0 (if the space reduces to the origin), the corresponding local manifold vanishes. Fig. B.2 describes local stable and unstable manifolds in the case of a hyperbolic fixed point in \mathbb{R}^2 .

Fig. B.2

Motion on the local stable and unstable manifolds




Fig. B.2

Since the local stable and unstable manifolds are diffeomorphic, near \bar{x} , to the stable and unstable spaces F^s and F^u of the Jacobian matrix $DG(\bar{x})$, one should expect, by analogy to Theorem B.4.1, the restriction of G to these manifolds to behave near \bar{x} like the restriction of the linearized map $x \rightarrow DG(\bar{x})x$ to F^s or F^u . Indeed,

Theorem B.5.2. *Suppose that G is as in Theorem B.5.1, and that $DG(\bar{x})$ is invertible. Let B^s, B^u be the submatrices in the real canonical form of $DG(\bar{x})$, that correspond to the stable and unstable spaces F^s and F^u , respectively. Then*

1. *The restriction of G to the local stable (resp. unstable) manifold is locally topologically conjugate to the linear map $x \rightarrow B^s x$, $x \in F^s$ (resp. $x \rightarrow B^u x$, $x \in F^u$).*
2. *There is a norm $\|x\|$ of \mathbb{R}^m , and two constants $k^s < 1 < k^u$ such that if $\delta > 0$ is small enough*

- a. $\|G(x) - \bar{x}\| \leq k^s \|x - \bar{x}\|$ for all x in $W_{(\bar{x})}^s$, $\|x - \bar{x}\| < \delta$.
- b. $\|G(x) - \bar{x}\| \geq k^u \|x - \bar{x}\|$ for all x in $W_{(\bar{x})}^u$, $\|x - \bar{x}\| < \delta$.

Motion on a local center manifold

Part 3 of Theorem B.5.1 shows that any recurrent behaviour (e.g. cycles) near the fixed point, must occur in a local center manifold. An explicit representation of such a manifold, that takes into account possible nonlinearities, is therefore useful. It may be obtained through the following procedure (the reader will verify easily that the procedure applies to local stable and unstable manifolds as well).

To simplify notation, assume that the fixed point has been translated to the origin, i.e. $\bar{x} = 0$, and that a linear change of variables has brought the Jacobian matrix $DG(\bar{x})$ in its real canonical form, which we write $\text{diag}\{C, B\}$, in which C corresponds to the center space F^c , and B to the direct sum of the stable and unstable spaces F^s and F^u . In terms of the notation of Theorem B.5.2, $B = \text{diag}\{B^s, B^u\}$. In the new coordinate system (ξ, η) , where ξ belongs to F^c and η to $F^s + F^u$, the difference equation $x_{n+1} = G(x_n)$ reads

$$(B.9) \quad \xi_{n+1} = C\xi_n + f(\xi_n, \eta_n)$$

$$\eta_{n+1} = B\eta_n + g(\xi_n, \eta_n)$$

in which f and g are defined on an appropriate neighbourhood of the origin. By construction, the functions f and g are C^r , $r \geq 2$ and are equal to 0, as well as their first partial derivatives, at $\xi = 0$, $\eta = 0$.

Since a local center manifold $W_{(\bar{x})}^c$ is tangent at the origin to F^c (the $\eta = 0$ space), one can represent it as the graph of a C^{r-1} function $\eta = \gamma(\xi)$, that is defined on a (small enough) open ball V in F^c , centered at $\xi = 0$, such that γ and its first partial derivatives vanish at $\xi = 0$, see Fig. B.3. The function γ , near the origin, is given implicitly by expressing that $W_{(\bar{x})}^c$ is locally invariant, that is

$$(B.10) \quad B\gamma(\xi) + g(\xi, \gamma(\xi)) = \gamma[C\xi + f(\xi, \gamma(\xi))]$$

whenever ξ and $C\xi + f(\xi, \gamma(\xi))$ are in V (which will be surely true if ξ is small enough). Given such a function γ , the restriction of the difference equation to the local center manifold is obtained by setting $\eta_n = \gamma(\xi_n)$ and $\eta_{n+1} = \gamma(\xi_{n+1})$ in (B.9). Clearly, one obtains an equivalent difference equation by projecting it on F^c , i.e. by considering only the equation corresponding to the component ξ

$$(B.11) \quad \xi_{n+1} = C\xi_n + f(\xi_n, \gamma(\xi_n)) \equiv \Gamma(\xi_n), \xi_n \in V.$$

Trajectories in the local center manifold may then be analyzed by using this “reduced difference equation” in F^c , near $\xi = 0$.

Fig. B.3

Equivalent local dynamics

Up to now, we confined ourselves to the trajectories that lie in the local stable, center and unstable manifolds. The following fact shows that these trajectories contain all the information that is necessary to reconstruct the dynamics generated by G in \mathbb{R}^m , near the fixed point : as for the linear case, trajectories in \mathbb{R}^m may be considered as the “Cartesian product” of trajectories lying in each invariant manifold. Since the motion on the local stable and unstable manifolds is locally conjugate to the linear equation $\eta \rightarrow B\eta$, with η in $F^s + F^u$, this result can be formulated as




Fig. B. 3

Theorem B.5.3. *Let G be as in Theorem B.5.1, and assume that $DG(\bar{x})$ is invertible. Then G is locally topologically conjugate to the application defined by*

$$\begin{aligned} \eta &\rightarrow B\eta, \eta \in F^s + F^u, \text{ where } B = \text{diag}\{B^s, B^u\} \\ \xi &\rightarrow C\xi + f(\xi, \gamma(\xi)) = \Gamma(\xi), \xi \in F^c \text{ small.} \end{aligned}$$

Of course, if the Jacobian matrix $DG(\bar{x})$ has no eigenvalue of modulus 1, the foregoing result reduces to Theorem B.4.1.

Remark B.5.4. One cannot get, in general, an exact analytical expression of a solution γ of (B.10), but one can approximate it by a Taylor series at $\xi = 0$. Suppose that a C^1 function T from F^c into $F^s + F^u$ has been found such that, first, T and its first partial derivatives vanish at $\xi = 0$, and second

$$T[C\xi + f(\xi, T(\xi))] - BT(\xi) - g(\xi, T(\xi)) = 0(|\xi|^p)$$

as $\xi \rightarrow 0$, for some $1 < p < r$. Then $\gamma(\xi) = T(\xi) + 0(|x|^p)$ as $\xi \rightarrow 0$, see Carr (1981, Section 2.8). Here the notation $0(|\xi|^p)$ stands for a function $\alpha(\xi)$ such that there is a constant k with $|\alpha(\xi)| < k|\xi|^p$ for ξ small enough. In practice, $T(\xi)$ is chosen as a Taylor series of γ . Specifically, one expands γ , up to the desired order, as well as f and g , as a Taylor series at $\xi = 0$ in (B.10), and one gets the unknown coefficients of the Taylor expansion of γ by identification of the two members. This shows in particular that although the function γ defining a local center manifold need not be unique, its partial derivatives at $\xi = 0$ are nevertheless uniquely determined. Finally, replacing γ by its Taylor approximation T in (B.11) yields an approximate analytical expression of the reduced difference equation in the local center manifold.

Of course, the principle of such a Taylor approximation applies equally well to local stable and unstable manifolds.

Notes on the literature

The treatment of linear difference equations in section B.3 is a transposition of Hirsch and Smale (1974, chap. 6), who deal with differential equations. The material on hyperbolic fixed points of section B.4 can be found in any book on dynamical systems, see e.g. Hartman (1964), Palis and De Melo (1982, chap. 2.4), Guckenheimer and Holmes (1983, chap. 1.4). Section B.5 on invariant manifolds is adapted from Lanford (1983), Iooss (1979, chap. V), Carr (1981, section 2.8), Guckenheimer and Holmes (1983, chap. 1.4 and 3.2), Palis and De Melo (1982, chap. 2.6). Theorem B.5.3 follows from Palis and Takens (1977) : the analogue result for differential equations (vector fields) is stated there p. 341 (see also Guckenheimer and Holmes (1983, p.

130)), and the result for diffeomorphisms follows by a suspending technique, as p. 340 (I owe this reference to A. Chenciner).

C. LOCAL BIFURCATIONS

As we have seen in Section B.3, linear difference equations can produce cyclical trajectories when some eigenvalues have modulus 1, but such cycles are removed by a small change of the matrix defining the system. By contrast, nonlinear maps may generate cyclical behaviour that is preserved under small perturbations, i.e. that is structurally stable. A most powerful tool to analyse the occurrence of cyclical fluctuations, in many cases the only one presently available, is provided by the theory of bifurcations.

C.1. Introduction

Consider a family of difference equations indexed by a real parameter α , say $x_{n+1} = G_\alpha(x_n)$, where each G_α is a map from an open set U of \mathbb{R}^m into \mathbb{R}^m , α is in an open interval I containing 0, and where the dependence on (α, x) is C^r , $r \geq 1$. In many cases, one requires also that $G_\alpha(0) = 0$, for all admissible α : this simply means that there is a family of fixed points that have been translated to the origin, see Remark C.1.1. The family G_α undergoes a *bifurcation*, say at $\alpha = 0$, if the qualitative features of the orbits of G_α change when α moves from negative to positive values. The bifurcation is *local* (at the origin) if the change of the orbit structure can be observed in an arbitrarily small (but independent of α) neighbourhood of $x = 0$; the bifurcation is *global* otherwise. We focus here on local bifurcations. A brief introduction to global bifurcations, which are far more difficult to understand, will be given in Appendix D.

If a local bifurcation occurs, then in view of Theorem B.5.2, there must be an eigenvalue λ_α of the Jacobian matrix $DG_\alpha(0)$ that either goes through 0 ($\lambda_\alpha = 0$), or crosses the unit circle in the complex plane ($|\lambda_\alpha| = 1$), at $\alpha = 0$. The case $\lambda_\alpha = 0$ is uninteresting, however, since it may change locally a map that is orientation reversing to an orientation preserving one, but it cannot generate cycles near the origin (think of the one dimensional difference equation $x_{n+1} = \alpha x_n + 0(|x_n^2|)$ for α, x small). We shall assume accordingly that the Jacobian matrix $DG_\alpha(0)$ is invertible for each admissible α , and focus on the case where only one eigenvalue λ_α (and of course its conjugate $\bar{\lambda}_\alpha$ when it is nonreal) having multiplicity one, crosses the unit circle at $\alpha = 0$: this is called a *codimension one local bifurcation*¹. There are thus three cases: if $\lambda_0 = +1$, one gets a *saddle node* (or *fold*) *bifurcation*; if $\lambda_0 = -1$, a *flip bifurcation*, and if λ_0 is nonreal, a *Hopf bifurcation*.

It turns out that although the dimension of the ambient space \mathbb{R}^m may be large, all the recurrent behaviour associated with a local bifurcation occurs

necessarily in an invariant surface (manifold) that has a dimension equal to the number of eigenvalues that cross simultaneously the unit circle at $\alpha = 0$. The principle of such a drastic reduction of the dimension of the problem will be shown to be a consequence of the theory of local center manifolds in section C.5. This remarkable result enables us to study the saddle node and the flip bifurcations directly for difference equations on a onedimensional curve, or equivalently on the real line. On the other hand, a Hopf bifurcation may be analysed directly in \mathbb{R}^2 , since then the (nonreal) eigenvalue λ_α and its conjugate $\bar{\lambda}_\alpha$ have modulus 1 at the point of bifurcation $\alpha = 0$.

Remark C.1.1. We describe now a frequently encountered situation, and show how it can generate the formulation described in the text. Assume, as in the introduction of section B, that the successive states y of a given physical or social system must satisfy

$$(C.1) \quad F(y_{n+1}, y_n, \dots, y_{n-N}, \alpha) = 0$$

where y is a vector of \mathbb{R}^p , α is a real number indexing some characteristics of the system, and F is a C^r map from an open set of $\mathbb{R}^{p(N+2)+1}$ into \mathbb{R}^p . Let \bar{y} be a stationary state for $\alpha = 0$, i.e. a vector of \mathbb{R}^p such that $F(\bar{y}, \dots, \bar{y}, 0) = 0$, and assume that the Jacobian matrix of the map $y \rightarrow F(y, \bar{y}, \dots, \bar{y}, 0)$ is invertible at $y = \bar{y}$. From the Implicit Function Theorem, there are open neighbourhoods V of $(\bar{y}, \dots, \bar{y}, 0)$ in $\mathbb{R}^{p(N+1)+1}$, W of y in \mathbb{R}^p , and a unique C^r map $H : V \rightarrow W$ such that $(y_{n+1}, \dots, y_{n-N}, \alpha)$ satisfies (C.1), with $(y_n, \dots, y_{n-N}, \alpha)$ in V and y_{n+1} in W , if and only if

$$(C.2) \quad y_{n+1} = H(y_n, \dots, y_{n-N}, \alpha).$$

To get the formulation of the text, one considers the variable $x_n = (y_n, \dots, y_{n-N})$, and the map G that associates to every (x_n, α) in V the vector $x_{n+1} = (y_{n+1}, y_n, \dots, y_{n-N+1})$ with y_{n+1} given by (C.2). One may restrict V to be the product of an open neighbourhood U of $(\bar{y}, \dots, \bar{y})$ in $\mathbb{R}^{p(N+1)}$, and of an open interval I containing 0 in \mathbb{R} . Then $G(x, \alpha)$ is defined on $U \times I$; in other words, each map $G_\alpha = G(\cdot, \alpha)$ is well defined for each admissible α , on the open set U , which is independent of the parameter α .

We remark that $\bar{x} = (\bar{y}, \dots, \bar{y})$ is a fixed point for $\alpha = 0$, i.e. $G(\bar{x}, 0) - \bar{x} = 0$. This fixed point may of course be translated to the origin through a linear change of variable. In fact, as noted in the text, one often requires that $G(0, \alpha) = G_\alpha(0) = 0$ for all admissible α . We describe now a procedure that leads to such a formulation.

Assume that the Jacobian matrix $D_x G(\bar{x}, 0)$ has no eigenvalue equal to 1. Then, again from the Implicit Function Theorem, the fixed point is bound to persist for small values of α . That is, the equation

$$(C.3) \quad x - G(x, \alpha) = 0$$

can be solved uniquely in x near $(\bar{x}, 0)$. Specifically, there are open neighbourhoods of $\alpha = 0$ in \mathbb{R} and of \bar{x} in $\mathbb{R}^{p(N+1)}$ –which we may take, since we are interested in local bifurcations, to be I and U respectively– and a C^r map from I to U , i.e. $\alpha \rightarrow \bar{x}_\alpha$, such that (C.3) is satisfied on $U \times I$ if and only if $x = \bar{x}_\alpha$. Translating \bar{x}_α to the origin of $\mathbb{R}^{p(N+1)}$ is achieved by making the change of variable $z = x - \bar{x}_\alpha$, for each admissible α , which yields the equivalent family $G^*(z, \alpha) = G(z + \bar{x}_\alpha, \alpha) - \bar{x}_\alpha$. One has then indeed $G^*(0, \alpha) = 0$ for all α . By continuity of the map $\alpha \rightarrow \bar{x}_\alpha$, we are sure that each map $G^* \alpha = G^*(\cdot, \alpha)$ is well defined on a small open ball U^* centered at 0 in $\mathbb{R}^{p(N+1)}$, for $|\alpha| < \delta$, if δ is small enough. Restricting attention to the family $G^* : U^* \rightarrow \mathbb{R}^{p(N+1)}$, $|\alpha| < \delta$, yields the desired formulation.

Remark C.1.2. It should be emphasised that the results of this section apply also to the study of the local bifurcations of a family of difference equations, near a *periodic orbit*. Consider a family of difference equations in \mathbb{R}^m given by the maps F_α , each being defined on a given open set, and suppose that the origin $x = 0$ is a fixed point of the n -th iterate of F_α , i.e. $F_\alpha^n(0) = 0$ for some $n > 1$, with $F_\alpha^j(0) \neq 0$ for $j = 1, \dots, n-1$, for all admissible α . Then $x = 0$ is a periodic point, with period n , the corresponding periodic orbit being $0, F_\alpha(0), F_\alpha^2(0), \dots, F_\alpha^{n-1}(0)$, for each α (again this simply means that there is a family of periodic orbits, and that one of the periodic points on the orbits has been translated to the origin). All the results of this section can then be transposed to the case at hand by considering the family $G_\alpha = F_\alpha^n$.

C.2. The saddle node bifurcation

We look first at the case of a one-parameter family of difference equations on the real line

$$x_{n+1} = G(x_n, \alpha) \equiv G_\alpha(x_n)$$

defined by the C^r map $G : U \times I \rightarrow \mathbb{R}$, where U and I are open intervals of the real line containing 0. In that case, for each value of the parameter α , the trajectories generated by the map G_α are easily visualised with the help of its graph, in the plane (x_n, x_{n+1}) , as shown in Fig. C.1. A fixed point \bar{x}

is then described as a point of intersection of the graph of G_α with the 45° line, and the trajectory associated with an initial condition x_0 is generated by following the arrows as in the Figure. A fixed point \bar{x} of G_α is then asymptotically stable if $\left| \frac{\partial G}{\partial x}(\bar{x}, \alpha) \right| < 1$, unstable if $\left| \frac{\partial G}{\partial x}(\bar{x}, \alpha) \right| > 1$.

Fig. C.1

Assume that the origin is a fixed point for $\alpha = 0$, i.e. $G(0, 0) = 0$, and that $\frac{\partial G}{\partial x}(0, 0) = +1$. A *saddle node bifurcation* will obtain if we suppose the “generic” conditions $\frac{\partial G}{\partial \alpha}(0, 0) \neq 0$ and $\frac{\partial^2 G}{\partial x^2}(0, 0) \neq 0$ (we need accordingly $r \geq 2$).

Fig. C.2.a pictures what happens to the graphs of the maps G_α for α and x small, in the case where $\frac{\partial G}{\partial \alpha}(0, 0) > 0$ and $\frac{\partial^2 G}{\partial x^2}(0, 0) > 0$. If $\alpha < 0$, G_α has two fixed points near 0. The one on the left is asymptotically stable, the other is unstable. For $\alpha = 0$, the origin is stable from the left, unstable from the right. If $\alpha > 0$, G_α has no fixed point near 0. Thus in a saddle node bifurcation, two fixed points having opposite stability properties coalesce to the origin, and disappear. Fig. C.2.b, which represents the qualitative features of this bifurcation for α and x small, is called a *bifurcation diagram*. The curve on the left describes the two branches of fixed points. Arrows show where the discrete orbits generated by G_α go, for a given α .

Fig. C.2.a

Fig.C.2.b

Proposition C.2.1. (Saddle node bifurcation). *Let $G : U \times I \rightarrow \mathbb{R}$ define a one-parameter family of maps, where G is C^r with $r \geq 2$, and U, I are open intervals of the real line containing 0. Assume (1) $G(0, 0) = 0$; (2) $\frac{\partial G}{\partial x}(0, 0) = 1$; (3) $\frac{\partial^2 G}{\partial x^2}(0, 0) > 0$; (4) $\frac{\partial G}{\partial \alpha}(0, 0) > 0$. Then there are $\alpha_1 < 0 < \alpha_2$ and $\varepsilon > 0$ such that*

(i) *If $\alpha_1 < \alpha < 0$, then $G_\alpha = G(\cdot, \alpha)$ has two fixed points $x_{1\alpha} < 0 < x_{2\alpha}$ in $(-\varepsilon, \varepsilon)$. The fixed point $x_{1\alpha}$ is asymptotically stable, the other is unstable.*

(ii) *If $0 < \alpha < \alpha_2$, then G has no fixed point in $(-\varepsilon, \varepsilon)$.*




Fig. C.1




Fig. C.2.a




Fig. C.2.b

It can be shown further that the curve on Fig. C.2.b describing the two branches of fixed points, i.e. the union of the sets $\{x_{1\alpha}, x_{2\alpha}\}$ for $\alpha_1 < \alpha \leq 0$, is indeed a C^{r-1} onedimensional manifold that is tangent to the vertical axis at $x = 0, \alpha = 0$.

Of course, the case $\frac{\partial G}{\partial \alpha}(0, 0) < 0$ is identical to the above after the change of parameter $\alpha \rightarrow -\alpha$. The reader will easily figure out what is the bifurcation diagram when inequality (3) is reversed : it amounts to changing the roles of positive and negative values of α , and reversing the arrows in Fig. C.2.b.

Transcritical bifurcation

The foregoing bifurcation described the “generic” situation. In some cases, however, the class of maps under consideration may be restricted, and a different bifurcation may occur. A common restriction is that the fixed point should persist, or equivalently, after a suitable α -dependent change of variables that translates the fixed point to the origin, that 0 should be a fixed point for all values of the parameter. In that case, one obtains a *transcritical bifurcation*.

Consider a family of maps defined by $G(x, \alpha)$ as above, with $\frac{\partial G}{\partial x}(0, 0) = +1$. We require here that $G(0, \alpha) = 0$ for all α , and not only for $\alpha = 0$ (this implies $\frac{\partial G}{\partial \alpha}(0, 0) = 0$, a violation of condition (4) of Proposition C.2.1), but keep the “generic” assumption $\frac{\partial^2 G}{\partial x^2}(0, 0) \neq 0$. In order to get actually a bifurcation, we must specify that $\frac{\partial G}{\partial x}(0, \alpha)$ goes through +1 when α passes 0, that is $\frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) \neq 0$.

Fig. C.3.a describes what happens to the graphs of G_α near 0 for α small, when $\frac{\partial^2 G}{\partial x^2}(0, 0) > 0$ and $\frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) > 0$. If $\alpha < 0$, the origin is asymptotically stable, but there is another fixed point $x_{1\alpha} > 0$ near 0, which is unstable. If $\alpha > 0$, the origin is unstable, and there exists another fixed point $x_{1\alpha} < 0$ near 0, which is asymptotically stable. Thus in a transcritical bifurcation, there are two fixed points that exchange stability at the point of bifurcation. The corresponding bifurcation diagram is given in Fig. C.3.b.

Fig. C.3.a

Fig. C.3.b




Fig. C.3.a




Fig. C.3.b

Proposition C.2.2. (Transcritical bifurcation). *Let $G : U \times I \rightarrow \mathbb{R}$ define a one-parameter family of maps, where G is C^r with $r \geq 2$, and U, I are open intervals of the real line containing 0. Assume*

(1) $G(0, \alpha) = 0$ for all α ; (2) $\frac{\partial G}{\partial x}(0, 0) = 1$; (3) $\frac{\partial^2 G}{\partial x^2}(0, 0) > 0$; (4) $\frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) > 0$. Then there are $\alpha_1 < 0 < \alpha_2$ and $\varepsilon > 0$ such that

(i) If $\alpha_1 < \alpha < 0$, G_α has two fixed points, 0 and $x_{1\alpha} > 0$ in $(-\varepsilon, \varepsilon)$. The origin is asymptotically stable, the other fixed point is unstable.

(ii) If $0 < \alpha < \alpha_2$, G_α has two fixed points, 0 and $x_{1\alpha} < 0$ in $(-\varepsilon, \varepsilon)$. The origin is unstable, the other fixed point is asymptotically stable.

It can be shown furthermore that each branch of fixed points in Fig. C.3.b, i.e. the union of all fixed points $x_{1\alpha}$ for $\alpha_1 < \alpha < 0$, or for $0 < \alpha < \alpha_2$, is a C^{r-1} onedimensional manifold. Here again, the case $\frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) < 0$ is handled by making the change of parameter $\alpha \rightarrow -\alpha$. The reader will also figure out easily the bifurcation diagram when the inequality (3) is reversed: it amounts, here also, to changing the roles of positive and negative values of α , and reversing the arrows in Fig. C.3.b.

Pitchfork bifurcation

Another interesting, “nongeneric” case arises when one assumes $\frac{\partial^2 G}{\partial x^2}(0, 0) = 0$ in the foregoing proposition. Let G be C^3 and suppose that the third derivative, i.e. $\frac{\partial^3 G}{\partial x^3}(0, 0)$, differs from 0. One gets then a *pitchfork bifurcation*.

Fig. C.4.a describes what happens to the graphs of G_α near the origin for α small, when $\frac{\partial^3 G}{\partial x^3}(0, 0) < 0$. For $\alpha = 0$, the graph has an inflexion point at $x = 0$, and the origin is asymptotically stable. If $\alpha < 0$, G_α has a unique fixed point, at $x = 0$, near the origin, and it is asymptotically stable. If $\alpha > 0$, G_α has three fixed points near 0. The origin is an unstable fixed point, but the two others are asymptotically stable. One says then that the family G_α undergoes a *supercritical* pitchfork bifurcation : the fixed point $x = 0$ loses its stability as α goes through 0, to give rise to a pair of stable fixed points. The corresponding bifurcation diagram is described in Fig. C.4.b.

Fig. C.4.a

Fig. C.4.b

Proposition C.2.3. (Supercritical pitchfork bifurcation). *Let $G : U \times I \rightarrow \mathbb{R}$ be as in Proposition C.2.2, except that G is C^r with $r \geq 3$, $\frac{\partial^2 G}{\partial x^2}(0, 0) = 0$*




Fig. C.4.a




Fig. C.4.b

and $\frac{\partial^3 G}{\partial x^3}(0,0) < 0$. Then there are $\alpha_1 < 0 < \alpha_2$ and $\varepsilon > 0$ such that

(i) If $\alpha_1 < \alpha \leq 0$, then G_α has a unique fixed point, $x = 0$, in $(-\varepsilon, \varepsilon)$. It is asymptotically stable.

(ii) If $0 < \alpha < \alpha_2$, then G has three fixed points in $(-\varepsilon, \varepsilon)$. The origin is an unstable fixed point, the two others, $x_{1\alpha} < 0 < x_{2\alpha}$, are asymptotically stable.

Here again, the two branches of fixed points in Fig. C.4.b, i.e. the union of the sets $\{x_{1\alpha}, x_{2\alpha}\}$ for $0 \leq \alpha < \alpha_2$, is a C^{r-1} one-dimensional manifold that is tangent to the x -axis at $x = 0$, $\alpha = 0$. The case $\frac{\partial^2 G}{\partial x \partial \alpha}(0,0) < 0$ is identical to the above after the change of parameter $\alpha \rightarrow -\alpha$. A more interesting modification arises when considering the case $\frac{\partial^3 G}{\partial x^3}(0,0) > 0$, which yields a so-called *subcritical pitchfork bifurcation*. What happens qualitatively is described in Fig. C.5.a, and the corresponding bifurcation diagram is given in Fig. C.5.b. At the bifurcation point $\alpha = 0$, the graph of G_α has an inflexion point at $x = 0$ as before, but the origin is now unstable (whereas it was stable for $\alpha = 0$ in the case of a supercritical bifurcation). Then for $\alpha < 0$, there are three fixed points near the origin, but only $x = 0$ is asymptotically stable. For $\alpha \geq 0$, the origin is the unique fixed point near $x = 0$, and it is unstable.

Fig. C.5.a

Fig. C.5.b

Remark C.2.4. The reader will easily verify that if the family G_α undergoes a subcritical pitchfork bifurcation when α goes through 0 from below, then the family of its local inverses G_α^{-1} undergoes a supercritical bifurcation when α goes through 0 from above. The family of local inverses is defined as follows. Since $\frac{\partial G}{\partial x}(0,0) \neq 0$, from the implicit Function Theorem, the equation $G(x, \alpha) - y = 0$ can be solved uniquely in x for x, α, y small, to yield $x = F(y, \alpha)$ where F may assumed to be defined on $V \times J$, with V, J being two open intervals containing 0. Then for every α in J , $F_\alpha = F(\cdot, \alpha)$ is indeed a local inverse of $G_\alpha = G(\cdot, \alpha)$, and the family F_α undergoes a supercritical pitchfork bifurcation when α goes through 0 from above.

Remark C.2.5. It is clear that the qualitative features of the local bifurcations discussed here are preserved if one makes a change of parameter $\alpha \rightarrow \varphi(\alpha)$ and an α -dependent change of variable $x \rightarrow h(x, \alpha)$, where φ and $h(\cdot, \alpha)$ are homeomorphisms. It can be shown that the converse result holds. For instance, two bifurcating families satisfying the assumptions of Proposition C.2.1 are locally topologically conjugate : they are the same, up to a




Fig. C.5.a




Fig. C.5.b

change of parameter and a change of variable as above, at least sufficiently near $\alpha = 0$, $x = 0$. Similar statements hold for transcritical or pitchfork bifurcations.

Finally, it should be noted that the bifurcations presented here are (locally) *structurally stable*, i.e. the qualitative features of the bifurcation are preserved, up to a change of variable, if the maps in the family are slightly perturbed (in a way that depends sufficiently smoothly on the parameter). By contrast, as noted earlier, such structural stability could not obtain, had we assumed the maps in the family to be *linear*.

C.3. The flip bifurcation

The previous section was devoted to the case where the one-parameter family of maps of the real line defined by $G(x, \alpha)$, was such that $\frac{\partial G}{\partial x}(0, 0) = 1$. We investigate now the case where $\frac{\partial G}{\partial x}(0, 0) = -1$. To be specific, let the family be defined by the map $G : U \times I \rightarrow \mathbb{R}$ as before, where G is C^3 (the need for continuous third derivatives will become apparent soon). Assume $G(0, \alpha) = 0$ for all α , $\frac{\partial G}{\partial x}(0, 0) = -1$ and $\frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) < 0$.² The origin $x = 0$ is a fixed point of every G_α , and the slope of the graph of G_α , at the origin, decreases and passes through -1 as α goes through 0 from below. The family G_α undergoes then a *flip bifurcation*.

The orbits of G_α near the origin oscillate more and more around the fixed point $x = 0$, when α varies from negative to positive values. If all the maps in the family G_α were linear, then these oscillating trajectories would converge to 0 for $\alpha < 0$, and diverge to infinity for $\alpha > 0$, while every $x \neq 0$ would be a periodic point of period two for $\alpha = 0$, since one would have then $G_0(x) = -x$ for all x . One should thus expect a cycle of period two to be part of the story in the more general case of nonlinear maps, as considered here.

The most efficient way to discover an orbit of period two of G_α is to look at its second iterate, $G_\alpha^2 = G_\alpha \circ G_\alpha$, which is well defined near $x = 0$. Indeed, a cycle of period two for G_α is characterized by a fixed point of G_α^2 that differs from 0.

Let us write $G_\alpha^2(x)$ as $G^2(x, \alpha)$. Since

$$G^2(x, \alpha) = G(G(x, \alpha), \alpha)$$

one has clearly $G^2(0, \alpha) = 0$ for all α , and one gets from the chain rule of differentiation

$$\frac{\partial G^2}{\partial x}(0, 0) = +1 ; \quad \frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) > 0 ; \quad \frac{\partial^2 G^2}{\partial x^2}(0, 0) = 0.$$

If we suppose the “generic” condition $\frac{\partial^3 G^2}{\partial x^3}(0, 0) \neq 0$, we see that *the family G_α^2 undergoes a pitchfork bifurcation at $\alpha = 0$* .³ If $\frac{\partial^3 G^2}{\partial x^3}(0, 0) < 0$, the bifurcation is supercritical, and what happens to the family G_α^2 is qualitatively depicted in Fig. C.4. For $\alpha \leq 0$, there is no cycle of period two near $x = 0$, and the origin is a stable fixed point. When $\alpha > 0$, the origin becomes unstable, there is an orbit of period two near $x = 0$, and this orbit is asymptotically stable (since a cycle of period two is characterized by a fixed point \bar{x} of G_α^2 , asymptotic stability is defined by using G_α^2 , and is thus guaranteed when $\left| \frac{\partial G_\alpha^2}{\partial x}(\bar{x}) \right| < 1$). Therefore, in a supercritical flip bifurcation, a stable fixed point becomes unstable, and gives rise to a stable orbit of period 2. For this reason, the flip bifurcation is sometimes called a *period doubling* (or *subharmonic*) *bifurcation*.

The corresponding bifurcation diagram for the family G_α itself is represented in Fig. C.6. The curve there describes the evolution of the two points on the orbit of period 2 when the parameter α varies. In Fig. C.7.a and b, the graphs of G_α are represented, as well as the trajectories they generate near the origin, for $\alpha < 0$ and $\alpha > 0$ respectively.

Fig. C.6

Fig. C.7.a

Fig. C.7.b

The analogue of Proposition C.2.3 becomes here :

Proposition C.3.1. (Supercritical flip bifurcation). *Let $G : U \times I \rightarrow \mathbb{R}$ define a one-parameter family of maps, where G is C^r , with $r \geq 3$, and U, I are open intervals containing 0. Assume*

- (1) $G(0, \alpha) = 0$ for all α ; (2) $\frac{\partial G}{\partial x}(0, 0) = -1$;
- (3) $\frac{\partial^2 G}{\partial x \partial \alpha}(0, 0) < 0$; (4) $\frac{\partial^3 G^2}{\partial x^3}(0, 0) < 0$.

Then, there are $\alpha_1 < 0 < \alpha_2$ and $\varepsilon > 0$ such that




Fig. C.6





Fig. C.7.a



$$\alpha > 0$$

Fig. C.7.b

(i) If $\alpha_1 < \alpha \leq 0$, then G_α has a unique fixed point at the origin, and no orbit of period two in $(-\varepsilon, \varepsilon)$. The fixed point is asymptotically stable.

(ii) If $0 < \alpha < \alpha_2$, then G_α has a unique fixed point at the origin, and a unique orbit of period two in $(-\varepsilon, \varepsilon)$. The fixed point is unstable, and the orbit of period two is asymptotically stable.

In the same way as for the pitchfork bifurcation, the union of the period two points for $0 \leq \alpha < \alpha_2$ in Fig. C.6, is a C^{r-1} onedimensional manifold that is tangent to the x -axis at $x = 0$, $\alpha = 0$. When $\frac{\partial^3 G^2}{\partial x^3}(0, 0) > 0$, the bifurcation is *subcritical*, and what happens to the family G_α^2 is qualitatively described in Fig. C.5. The basic difference with the supercritical case is that the origin $x = 0$ is now unstable when $\alpha = 0$, while it was stable in the case of supercritical bifurcation. Thus in a subcritical flip bifurcation, an unstable orbit of period two coalesces with a stable fixed point at $\alpha = 0$, to leave an unstable fixed point. The corresponding bifurcation diagram for the family G_α is given in Fig. C.8. Here again, the two branches of the curve represent the evolution of the two points of the orbit of period two. What happens to the graphs of the family G_α and to the trajectories they generate near the fixed point $x = 0$ is described in Fig. C.9.a and b, for $\alpha < 0$ and $\alpha > 0$.

Fig. C.8

Fig. C.9.a

Fig. C.9.b

Proposition C.3.2 (Subcritical flip bifurcation). *Replace the inequality (4) in Proposition C.3.1 by $\frac{\partial^3 G^2}{\partial x^3}(0, 0) > 0$. Then there exist $\alpha_1 < 0 < \alpha_2$ and $\varepsilon > 0$ such that*

(i) *If $\alpha_1 < \alpha < 0$, then G_α has a unique fixed point at the origin, and a unique orbit of period two in $(-\varepsilon, \varepsilon)$. The fixed point is asymptotically stable, the orbit of period two is unstable.*

(ii) *If $0 \leq \alpha < \alpha_2$, then G has a unique fixed point at the origin, and no orbit of period two in $(-\varepsilon, \varepsilon)$. The fixed point is unstable.*

Remark C.3.3. The *Schwarzian derivative* SG of a C^3 map $G : U \rightarrow \mathbb{R}$ where U is an open interval of the real line, is defined as

$$SG(x) = \frac{G'''(x)}{G'(x)} - \frac{3}{2} \left[\frac{G''(x)}{G'(x)} \right]^2$$





Fig. C.8



$$\alpha < 0$$

Fig. C.9.a




Fig. C.9.b

for every x such that $G'(x) \neq 0$. It is not difficult to verify that for the above family G_α , one has

$$\frac{\partial^3 G^2}{\partial x^3}(0, 0) = -2 \frac{\partial^3 G}{\partial x^3}(0, 0) - 3 \left[\frac{\partial^2 G}{\partial x^2}(0, 0) \right]^2$$

and since $\frac{\partial G}{\partial x}(0, 0) = -1$, $\frac{\partial^3 G^2}{\partial x^3}(0, 0) = 2 SG_0(0)$. It follows that condition (4) in Proposition C.3.1 may be expressed as : *the Schwarzian derivative $SG_\alpha(x)$ is negative at $x = 0$ at the point of bifurcation $\alpha = 0$* . Maps of the real line having globally a negative Schwarzian derivative have nice global properties, as we shall see in section D, and their global bifurcations are reasonably well understood. The reason for this should be apparent from the preceding analysis : for such maps, a flip bifurcation is necessarily supercritical.

Remark C.3.4. As for the saddle mode bifurcation, two families undergoing a supercritical (or subcritical) flip bifurcation as in Proposition C.3.1 (or C.3.2) can be deduced from each other through a change of parameter $\alpha \rightarrow \varphi(\alpha)$, and an α -dependent change of variable $x \rightarrow h(x, \alpha)$ where φ and $h(\cdot, \alpha)$ are homeomorphisms. Similarly, the qualitative features of a flip bifurcation are (locally) structurally stable, up to such a change of variable. Here again, one could not get such a structural stability if the maps were linear.

C.4. The Hopf bifurcation

We consider now a one-parameter family of difference equations of the form

$$x_{n+1} = G(x_n, \alpha) \equiv G_\alpha(x_n)$$

defined by the C^r map $G : U \times I \rightarrow \mathbb{R}^2$, where U is an open neighbourhood of the origin of \mathbb{R}^2 , and I is an open interval of the real line containing 0. We assume that the origin is a fixed point for all admissible values of the parameter, i.e. $G(0, \alpha) = 0$ for all α . We suppose further that for each α , the Jacobian matrix $D_x G(0, \alpha)$ has a pair of nonreal, conjugate eigenvalues $\lambda(\alpha)$ and $\bar{\lambda}(\alpha)$, with $\lambda(\alpha) = \rho(\alpha)[\cos \theta(\alpha) + i \sin \theta(\alpha)]$, $\rho(\alpha) \neq 0$ and $\theta(\alpha)$ in $(0, \pi)$. The modulus $\rho(\alpha)$ and the argument $\theta(\alpha)$ depend then in a C^{r-1} fashion on the parameter. In order to get a bifurcation, we assume that $r \geq 2$ and that the eigenvalues pass out of the unit circle in the complex plane as α goes through 0, i.e. $\rho(0) = 1$ and $\frac{d\rho}{d\alpha}(0) > 0$. Then the origin $x = 0$ is asymptotically stable for $\alpha < 0$, unstable for $\alpha > 0$: there is a so-called *Hopf bifurcation* at $\alpha = 0$.⁴

From Section A.1, there is for each α a change of variable $x \rightarrow h(x, \alpha)$ which brings the Jacobian matrix $D_x G(0, \alpha)$ in its real canonical form

$$B_\alpha = \rho(\alpha) \begin{bmatrix} \cos \theta(\alpha) & -\sin \theta(\alpha) \\ \sin \theta(\alpha) & \cos \theta(\alpha) \end{bmatrix},$$

the map h being linear in x and C^{r-1} in (x, α) . As noted in Section B.3, the action on \mathbb{R}^2 of the linear map $y \rightarrow B_\alpha y$ is a rotation of angle $\theta(\alpha)$, followed by an homothety of centre $y = 0$ and ratio $\rho(\alpha)$, see Fig. B.1. Thus the family of linear maps B_α can be represented in polar coordinates by $(\rho, \theta) \rightarrow (\rho(\alpha)\rho, \theta + \theta(\alpha))$. If $\alpha < 0$, all trajectories generated by $y_n = B_\alpha^n y_0$ converge to the origin, while all such trajectories diverge when $y_0 \neq 0$ for $\alpha > 0$. By contrast, for $\alpha = 0$, the map $y \rightarrow B_0 y$ leaves invariant every circle of the plane centered at the origin (of course, every such circle is generally an ellipse in the original coordinates).

Now if the maps in the family G_α are actually nonlinear, they are in fact small perturbations of the linear maps B_α near the origin. Thus one should expect the appearance of an *invariant closed curve*, i.e. a one-dimensional manifold that is homeomorphic to a circle of the plane, near $x = 0$ and for α small. The following set of results shows that this is indeed in general the case.

We first look at the case where the bifurcation is *supercritical*. This case arises when the origin is attracting for $\alpha = 0$. Then an attracting invariant closed curve appears for $\alpha > 0$.

Theorem C.4.1. (Supercritical Hopf bifurcation). *Let $G : U \times I \rightarrow \mathbb{R}^2$ define a one-parameter family of maps of the plane, where G is C^r with $r \geq 6$, U is an open set in \mathbb{R}^2 containing the origin, and I an open interval of the real line containing 0. Assume*

- (1) $G(0, \alpha) = 0$ for all α ;
- (2) for each α , $D_x G(0, \alpha)$ has a pair of nonreal, conjugate eigenvalues $\lambda(\alpha)$ and $\bar{\lambda}(\alpha)$, with $\lambda(\alpha) = \rho(\alpha)[\cos \theta(\alpha) + i \sin \theta(\alpha)]$, $\rho(\alpha) \neq 0$ and $\theta(\alpha)$ in $(0, \pi)$;
- (3) $\rho(0) = 1$ and $\frac{d\rho}{d\alpha}(0) > 0$;
- (4) $\theta(0) \neq 2\pi/q$ for $q = 1, 2, 3$ or 4.

Then there is a C^{r-3} α -dependent change of coordinates that brings the family $G_\alpha = G(\cdot, \alpha)$ into the polar coordinates form

$$(\rho, \theta) \rightarrow (\rho(\alpha)\rho - a(\alpha)\rho^3 + \rho^4 R(\rho, \theta, \alpha), \theta + \theta(\alpha) + b(\alpha)\rho^2 + \rho^3 S(\rho, \theta, \alpha))$$

where $R(\rho, \theta, \alpha)$ is C^{r-4} , $S(\rho, \theta, \alpha)$ is C^{r-3} , while $a(\alpha)$ and $b(\alpha)$ are C^{r-3} and C^{r-2} respectively. If in addition,

$$(5) \quad a(0) > 0$$

then the origin is asymptotically stable for $\alpha = 0$. Furthermore, there exist $\alpha_1 < 0 < \alpha_2$ and an open neighbourhood V of the origin in \mathbb{R}^2 such that

- (i) If $\alpha_1 < \alpha \leq 0$, then G_α has a unique fixed point, at $x = 0$, and no invariant closed curve in V . The origin is asymptotically stable.
- (ii) If $0 < \alpha < \alpha_2$, then G_α has a unique fixed point, at $x = 0$, and a unique closed invariant curve in V . The origin is unstable, while the invariant closed curve is asymptotically stable.

The first part of the theorem states that, if one puts aside the “strong resonance” cases $\theta = 2\pi/q$, $q = 1, 2, 3$ or 4 (these cases are exceptional anyway), each map G_α may be approximated for x small, after a suitable change of variables, by the transformation T_α given in polar coordinates by

$$(\rho, \theta) \rightarrow (\rho(\alpha)\rho - a(\alpha)\rho^3, \theta + \theta(\alpha) + b(\alpha)\rho^2)$$

If the maps G_α are indeed nonlinear, one should have in general $a(0) \neq 0$ and $b(0) \neq 0$. The transformation T_α is the composition of a (ρ -dependent) rotation and of an homothecy. The contracting or expanding properties of the transformation are thus most easily analysed by looking directly at the associated onedimensional difference equation

$$(C.4) \quad \rho_{n+1} = \rho(\alpha)\rho_n - a(\alpha)\rho_n^3 \equiv \gamma(\rho_n, \alpha),$$

for $\rho_n, \rho_{n+1} > 0$. In particular, ρ_α^* is a fixed point of $\gamma_\alpha = \gamma(\cdot, \alpha)$ if and only if T_α leaves invariant the circle of the plane centered at $x = 0$ of radius ρ_α^* . Moreover, ρ_α^* is asymptotically stable in the dynamics generated by γ_α if and only if the corresponding invariant circle is asymptotically stable in the dynamics generated by T_α .

It is easily seen that, if $a(0) \neq 0$ and if one allows negative values of ρ , the family γ_α undergoes a *pitchfork bifurcation* at $\alpha = 0$. Indeed γ is C^3 , and

for all α , $\gamma(0, \alpha) = 0$, $\frac{\partial \gamma}{\partial \rho}(0, \alpha) = \rho(\alpha)$, $\frac{\partial^2 \gamma}{\partial \rho^2}(0, \alpha) = 0$, $\frac{\partial^3 \gamma}{\partial \rho^3}(0, \alpha) = -6a(\alpha)$.

Thus if $a(0) > 0$, the origin $\rho = 0$ is asymptotically stable for $\alpha = 0$, and the bifurcation is supercritical (Proposition C.2.3). What happens in that case to the graphs of γ_α near $\rho = 0$, for α small, is represented in Fig. C.10.a. In terms of the dynamics associated with the transformation T_α , for small $\alpha \leq 0$, the origin of \mathbb{R}^2 attracts all nearby points. For small $\alpha > 0$, the origin becomes unstable and all small $x \neq 0$ in \mathbb{R}^2 are attracted to an invariant circle centered at $x = 0$, of radius $\rho_\alpha^* = \sqrt{(\rho(\alpha) - 1)/a(\alpha)}$. The last part of the theorem states that this picture is qualitatively unchanged when one goes back to the original family G_α . The corresponding bifurcation diagram is given in Fig. C.11.

Fig. C.10.a

Fig. C.10.b

Fig. C.11

The same line of reasoning permits to see easily what happens when $a(0) < 0$. In that case the origin is unstable at $\alpha = 0$, and the family γ_α undergoes a *subcritical* pitchfork bifurcation (Fig. C.10.b). In terms of the dynamics associated with the transformation T_α , for small $\alpha < 0$, the origin is asymptotically stable, but there is a repelling invariant circle centered at $x = 0$, of radius $\rho_\alpha^* = \sqrt{(\rho(\alpha) - 1)/a(\alpha)}$. When α crosses 0 from below, the repelling invariant circle coalesces to the origin and for small $\alpha > 0$, the fixed point $x = 0$ is unstable. The following result states that the picture remains qualitatively the same when going back to the original family G_α . The corresponding bifurcation diagram is given in Fig. C.12. †

Theorem C.4.2. (Subcritical Hopf bifurcation). *Assume in the preceding theorem that (5) is replaced by*

$$(5') \quad a(0) < 0.$$

Then the origin is unstable for $\alpha = 0$. Furthermore, there exist $\alpha_1 < 0 < \alpha_2$, and an open neighbourhood V of the origin in \mathbb{R}^2 such that

- (i) *If $\alpha_1 < \alpha < 0$, then G_α has a unique fixed point, at $x = 0$, and a unique invariant closed curve in V . The origin is asymptotically stable, the closed curve is unstable.*




Fig. C.10.a




Fig. C.10.b




Fig. C.11

(ii) If $0 \leq \alpha < \alpha_2$, then G_α has a unique fixed point, at $x = 0$, and no invariant closed curve in V . The origin is unstable.

Fig. C.12

The sign of the coefficient $a(0)$ determines whether the bifurcation is supercritical or subcritical. The following proposition shows how to obtain in principle this coefficient - although the practical computations involved are sometimes tedious.

Proposition C.4.3. *Under the assumptions of Theorem C.4.1, let $\lambda, \bar{\lambda}$ be the conjugate nonreal eigenvalues at $\alpha = 0$, and assume that coordinates in the plane have been chosen to put G_0 in the form*

$$\begin{pmatrix} x \\ y \end{pmatrix} \rightarrow \begin{bmatrix} \cos \theta(0) & -\sin \theta(0) \\ \sin \theta(0) & \cos \theta(0) \end{bmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix}.$$

Then one has ⁵

$$a(0) = \operatorname{Re} \left[\frac{(1 - 2\lambda)\bar{\lambda}^2}{1 - \lambda} c_{11} c_{20} \right] + \frac{1}{2} |c_{11}|^2 + |c_{02}|^2 - \operatorname{Re}(\bar{\lambda}c_{21}),$$

where

$$\begin{aligned} 8c_{20} &= (f_{xx} - f_{yy} + 2g_{xy}) + i(g_{xx} - g_{yy} - 2f_{xy}) \\ 4c_{11} &= (f_{xx} + f_{yy}) + i(g_{xx} + g_{yy}) \\ 8c_{02} &= (f_{xx} - f_{yy} - 2g_{xy}) + i(g_{xx} - g_{yy} + 2f_{xy}) \\ 16c_{21} &= (f_{xxx} - f_{xyy} + g_{xxy} + g_{yyy}) + i(g_{xxx} + g_{xyy} - f_{xxy} - f_{yyy}) \end{aligned}$$

in which the partial derivatives of f and g are evaluated at $(x, y) = (0, 0)$.

Remark C.4.4. The invariant closed curve C_α , of which the existence is asserted for $\alpha > 0$ in Theorem C.4.1, may be represented as the locus of all points of the plane that have the polar coordinates $(\rho(\theta, \alpha), \theta)$, where θ is an arbitrary real number, $\rho(\theta, \alpha) \neq 0$ for all θ and $\alpha > 0$, and ρ is periodic of period 2π in θ , i.e. $\rho(\theta + 2\pi, \alpha) = \rho(\theta, \alpha)$. Under the assumptions of the Theorem, α_2 can be chosen small enough so that the map ρ is C^{r-5} for $0 < \alpha < \alpha_2$. The union of all invariant closed curves C_α for α in $(0, \alpha_2)$, together with the origin of the plane for $\alpha = 0$, form in the bifurcation diagram of Fig. C.11, a two dimensional C^{r-5} manifold Σ that is actually




Fig. C.12

tangent to the x -plane at $\alpha = 0$. This does not mean that Σ is C^∞ (is C^k for all $k \geq 1$) when G is itself C^∞ , however. In such a case, one can find $\alpha_k > 0$ for each $k \geq 1$ such that ρ and thus Σ is C^k on $(0, \alpha_k)$. Yet the sequence α_k will typically tend to 0 as k tends $+\infty$. The same regularity result holds for $\alpha < 0$ in the case of a subcritical bifurcation.

Remark C.4.5. This remark is the analogue of Remark C.2.4 for the case of a pitchfork bifurcation. One can verify that if the family G_α undergoes a subcritical Hopf bifurcation when α goes through 0 from below, then the family of its local inverses G_α^{-1} undergoes a supercritical bifurcation when α goes through 0 from above. The family of local inverses is defined as follows. Since the eigenvalues of $D_x G(0, 0)$ have modulus 1, this matrix is invertible. From the Implicit Function Theorem, the equation $G(x, \alpha) - y = 0$ can then be solved uniquely in x for x, α, y small, to yield $x = F(y, \alpha)$ where F may be assumed to be defined on $V \times J$, where V is an open set of the plane containing the origin, and J is an open interval containing 0. For every α in J , $F_\alpha = F(\cdot, \alpha)$ is indeed a local inverse of $G_\alpha = G(\cdot, \alpha)$, and the family F_α undergoes a supercritical bifurcation when α goes through 0 from above.

Remark C.4.6. The strong resonance cases $\theta(0) = 2\pi/q$ with $q = 1, 2, 3, 4$, are “nongeneric” in the class of all nonlinear maps. The study of what happens in these cases involves some difficult problems. For more information, see Whitley (1983, Section 2.5), and Ioos (1979, Chapter IV).

Remark C.4.7 (*Motion on the invariant closed curve*).⁶ Once we have obtained an asymptotically stable invariant closed curve C_α in a supercritical Hopf bifurcation (for $\alpha > 0$ in Theorem C.4.1), it remains to understand the behaviour of the trajectories generated by the difference equation $x_{n+1} = G_\alpha(x_n)$, when it is restricted to C_α (the results concerning a supercritical bifurcation can then be transposed to the subcritical case, for instance by using Remark C.4.5).

Under the assumptions of Theorem C.4.1, for small α , the map G_α is only a small perturbation, after a suitable change of variables, of the transformation T_α given in polar coordinates by

$$(\rho, \theta) \rightarrow (\rho(\alpha)\rho - a(\alpha)\rho^3, \theta + \theta(\alpha) + b(\alpha)\rho^2)$$

Thus for small $\alpha > 0$, in this system of polar coordinates, the restriction of G_α to the invariant closed curve C_α is a small perturbation of the restriction to the circle centered at the origin of radius $\rho_\alpha^* = \sqrt{(\rho(\alpha) - 1)/a(\alpha)}$, of a rotation R_α of angle $\theta_\alpha^* = \theta(\alpha) + b(\alpha)(\rho_\alpha^*)^2$.

Let us first look at the dynamics generated by R_α . If $\theta_\alpha^*/2\pi$ is rational, i.e. of the form p/q where p and q are positive integers that are relatively prime, every point of the circle is periodic with period q (i.e. one goes back to it when applying q -times the rotation R_α). If $\theta_\alpha^*/2\pi$ is irrational, then no point of the circle is periodic : the trajectory generated by an arbitrary point of the circle is dense in that circle.

Knowledge of the properties of the restriction of the approximating rotation R_α to the invariant circle of radius ρ_α^* is not enough, unfortunately, to state the properties of the original family G_α restricted to the invariant closed curves C_α . The reason is that such rotations R_α are *not* structurally stable : the fact that $\theta_\alpha^*/2\pi$ is rational or irrational can be altered by a small perturbation of the map. It can be shown nevertheless that, as far as the behaviour of the restriction of G_α to C_α is concerned, the typical (i.e. “generic”) picture is the following one. Assuming that G is C^∞ , and if the parameter α_2 appearing in Theorem C.4.1 is chosen small enough

1. *there is a set A that is open and dense in $(0, \alpha_2)$ such that for every α in A , the map G_α has two families of periodic orbits, all having the same (α -dependent) period, on the invariant closed curve C_α . One family is stable, the other unstable.*

The best way to visualise what happens is to look at the path followed in the complex plane by the eigenvalue $\lambda(\alpha)$ of the Jacobian matrix $D_x G(0, \alpha)$, as shown in Fig. C.13.a. Consider a complex number of the form $\cos \theta + i \sin \theta$ represented by the point M on the unit circle, with $\theta = 2\pi p/q$, where p and q are positive integers that are relatively prime, and $q \neq 1, 2, 3, 4$ (again, we put aside the strong resonance cases). Then there is a narrow cusped open region, near the unit circle, that lies between two curves passing through M and having a common tangent there as in the grey region of Fig. C.13.a, with the following property. Suppose that the point representing the eigenvalue $\lambda(\alpha)$ lies in this narrow “tongue” for $\alpha > 0$ in some interval (α', α'') . Then for every α in that interval, G_α has two families of periodic orbits, of period q , on the invariant closed curve C_α . The periodic orbits of one family are unstable, while the periodic orbits of the other family are asymptotically stable (in the dynamics generated by the restriction of G_α to C_α). The unstable and stable periodic orbits “alternate” on the invariant closed curve ; an example of such a pattern is given in Fig. C.13.b, with $q = 6$ (the arrows indicate stability or unstability ; they are drawn as if the trajectories were continuous, the interpretation in discrete time should be apparent).

Fig. C.13.a

Fig. C.13.b




Fig. C.13.a




Fig. C.13.b

Since rational numbers are dense in the real line, the path followed by the eigenvalue $\lambda(\alpha)$ in the complex plane will cross infinitely many such distinct little tongues. The union A of the corresponding intervals (α', α'') of the parameter will be open and dense in $(0, \alpha_2)$ if α_2 is small enough. Thus for most values of α in $(0, \alpha_2)$, the asymptotic behaviour of the trajectories generated by G_α on the invariant closed curve will be periodic. It should be remarked that many of the periods will be large, i.e. will correspond to a large q .

2. *Although the above mentioned set A is open and dense in $(0, \alpha_2)$, its complement in that interval is not negligible from a measure theoretic viewpoint.* Indeed, the set B of parameters in $(0, \alpha_2)$ such that the restriction of G_α to the invariant closed curve C_α is topologically conjugate to a rotation of angle θ of the unit circle of the plane, with $\theta/2\pi$ irrational, has positive Lebesgue measure $m(B) > 0$. In that case, any orbit generated by the restriction of G_α to C_α is aperiodic : it is dense in C_α . In fact, the relative frequency of such aperiodic behaviour is high for $\alpha > 0$ small : the ratio $m(B)/\alpha_2$ tends to 1 when α_2 goes to 0.

Remark C.4.8. It should be clear from the previous remark that the qualitative features of a Hopf bifurcation cannot be structurally stable. After a perturbation, the path followed by the eigenvalue $\lambda(\alpha)$ would cross the unit circle at a different point, and thus meet different little tongues as in Fig. C.13.a. The two families could not be conjugate.

Remark C.4.9. (*Degenerate Hopf (Chenciner) bifurcation*). The condition $a(0) \neq 0$ is generally satisfied (is "generic") in the class of all families of nonlinear maps. Analogues of Theorems C.4.1 and C.4.2 are actually available in the case $a(0) = 0$, that has been thoroughly studied by Chenciner (1985a, b, 1988), see also Iooss (1979, Chapter III, Section 2) or Kuznetsov (1998). We give here a summary heuristic account of that degenerate bifurcation for $a(0) = 0$ and $a(0)$ very small.

Existence and stability of an invariant closed curve in the local dynamics induced by G_α near the steady state $x = 0$, amounts essentially to existence and stability of invariant circles in the dynamics generated by the Taylor approximation T_α of the family G_α in polar coordinates, or equivalently to existence and stability of fixed points $\rho_\alpha^* > 0$ of the associated onedimensional difference equation $\rho \rightarrow \gamma_\alpha(\rho)$ as in (C.4). Under appropriate (higher order) differentiability conditions, taking into account higher order terms when $a(0)$ is 0 or is very small, leads to the higher order difference equation :

$$(C.5) \quad \rho_{n+1} = \rho(\alpha)\rho_n - a(\alpha)\rho_n^3 - c(\alpha)\rho_n^5 = \tilde{\gamma}_\alpha(\rho_n).$$

For given α , an invariant circle in the dynamics generated by the Taylor approximation T_α is thus identified with a solution $\rho_\alpha^* > 0$ of the equation $\tilde{\gamma}_\alpha(\rho) = \rho$, or equivalently with a positive solution $z_\alpha^* = (\rho_\alpha^*)^2 > 0$ given by

$$(C.6) \quad \left(z_\alpha^* + \frac{a(\alpha)}{2c(\alpha)}\right)^2 = \frac{a^2(\alpha)}{4c(\alpha)} + \frac{\rho(\alpha) - 1}{c(\alpha)}.$$

We focus on the *supercritical case*, i.e. $c(0) > 0$ and thus $c(\alpha) > 0$ for α small. In the benchmark configuration $a(0) = 0$, the graph of $\tilde{\gamma}_0(\rho) = \rho - c(0)\rho^5$ for $\alpha = 0$ looks like that of $\gamma_0(\rho)$ in Fig. C.10.a, which implies that the graphs of $\tilde{\gamma}_\alpha(\rho)$ for α small, near $\rho = 0$, are also similar to those of $\gamma_\alpha(\rho)$ in that same Fig. C.10.a. For $\alpha \leq 0$, the origin $\rho = 0$ is stable in the dynamics induced by $\tilde{\gamma}_\alpha$ or T_α , while there is no invariant circle nearby. For $\alpha > 0$, the origin $\rho = 0$ becomes unstable and gives rise to a stable invariant circle.

The case where $a(0) \neq 0$ is small, is a slight perturbation of the above benchmark configuration. The picture is in fact qualitatively the same as when $a(0) > 0$ and thus $a(\alpha) > 0$ for α small. For $\alpha < 0$, the two solutions $z_{\alpha+}^*$ and $z_{\alpha-}^*$ of (C.6)

$$z_{\alpha\pm}^* = -\frac{a(\alpha)}{2c(\alpha)} \pm \left(\frac{a^2(\alpha)}{4c^2(\alpha)} + \frac{\rho(\alpha) - 1}{c(\alpha)}\right)^{1/2}$$

are both negative. The origin $\rho = 0$ is stable and there is no invariant circle nearby. For $\alpha > 0$, the origin $\rho = 0$ loses its stability and is surrounded by a stable invariant circle corresponding to the unique solution of (C.6) that is positive, i.e. $z_{\alpha+}^* > 0$. This configuration conforms to what is predicted for a standard non-degenerate Hopf bifurcation when $a(0) > 0$.

The picture is slightly different when $a(0) < 0$ and thus $a(\alpha) < 0$ for α small. Since $a(0)$ is very small, there is $\alpha_0 < 0$ such that the right hand side of (C.6) is 0 when $\alpha = \alpha_0$, and positive if and only if $\alpha > \alpha_0$. Therefore (C.6) has no real solution when $\alpha < \alpha_0$: the origin $\rho = 0$ is then stable and there is no invariant circle nearby. When $\alpha = \alpha_0$, (C.6) has two positive identical real solutions $z_{\alpha+}^* = z_{\alpha-}^* = -a(\alpha)/2c(\alpha) > 0$, that separate and remain positive, $0 < z_{\alpha-}^* < z_{\alpha+}^*$, when α goes up. Thus when $\alpha_0 < \alpha < 0$, the origin $\rho = 0$ is still stable, but two invariant circles appear nearby through a *saddle node bifurcation* when a goes up through α_0 , the invariant circle with a smaller (resp. larger) radius being unstable (resp. stable). When α goes up through 0, the origin $\rho = 0$ undergoes a subcritical Hopf bifurcation as

predicted by the general theory since $a(0) < 0$, where the unstable invariant circle of smaller radius coalesces with the origin, to disappear and leave the origin $\rho = 0$ unstable (the smaller solution $z_{\alpha-}^*$ becomes negative). The novel feature here is the presence all along of the stable invariant circle of larger radius corresponding to the larger solution $z_{\alpha+}^* > 0$, for $\alpha > \alpha_0$ and in particular for $\alpha > 0$.

The outcome one gets for the original two-dimensional dynamics generated by the family of maps G_α is essentially a small perturbation of the above picture. In a nutshell, for a *supercritical degenerate Hopf (Chenciner) bifurcation* ($a(0)$ small, $c(0) > 0$):

- for $a(0) \geq 0$ small, the picture is qualitatively identical to what one gets for a *supercritical Hopf bifurcation*. For $\alpha \leq 0$, the stationary state $x = 0$ is stable, while there is no nearby invariant closed curve. For $\alpha > 0$, the stationary state becomes unstable and gives rise to a nearby attracting invariant closed curve.

- For $a(0) < 0$, small, there exists $\alpha_0 < 0$ small such that :
 - for $\alpha < \alpha_0$, the stationary state $x = 0$ is stable, while there is no nearby invariant closed curve.
 - When α goes through α_0 from below, there is a saddle node bifurcation of invariant closed curves : two identical invariant closed curves appear near the stable stationary state $x = 0$ for $\alpha = \alpha_0$ and separate for $\alpha_0 < \alpha < 0$, the smaller invariant closed curve being unstable and the larger one being stable.
 - When α goes through 0 from below, the smaller unstable invariant closed curve coalesces with the stable stationary state $x = 0$ through a *subcritical Hopf bifurcation*.
 - For $\alpha > 0$ small, the stationary state $x = 0$ becomes unstable and coexists with the remaining larger stable invariant closed curve.

Symmetric statements can be obtained for the case of a *subcritical degenerate Hopf (Chenciner) bifurcation*, when $a(0)$ is zero or is very small and $c(0) < 0$. When $a(0) \leq 0$, the outcome is qualitatively what one gets in a subcritical Hopf bifurcation. For $\alpha < 0$, the stationary state $x = 0$ is stable but is surrounded by an unstable invariant closed curve (corresponding to the unique positive root of (C.6), $z_{\alpha+}^* > 0$). When α goes up through 0, the unstable invariant closed curve coalesces with the stationary state $x = 0$, to disappear and leave it unstable. The picture is slightly different when $a(0)$ is positive, so that $a(\alpha) > 0$ for α small, with a saddle node bifurcation of invariant closed curves occurring for $\alpha_0 > 0$ small in such a case. For $\alpha < \alpha_0$,

there is an invariant closed curve (corresponding to the largest root of (C.6), $z_{\alpha+}^* > 0$) near the stationary state $x = 0$, that is unstable, and stays away from $x = 0$. When $\alpha < 0$, the stationary state $x = 0$ is unstable, and there is no other invariant closed curve nearby. When α goes up through 0, the stationary state undergoes a supercritical Hopf bifurcation as predicted by the general theory when $a(0) > 0$. It becomes unstable, and a small stable invariant closed curve appears near $x = 0$ (corresponding to the small root of (C.6), that becomes positive, $z_{\alpha-}^* > 0$) and inside the larger unstable invariant closed curve. When α goes up through α_0 , the smaller stable and larger unstable invariant closed curves coalesce together and disappear through a saddle node bifurcation of invariant closed curves, to leave alone the unstable stationary state.

C.5. The center manifold reduction

Local bifurcations appear to be hard to analyse in \mathbb{R}^m , if m is large. Fortunately, as mentioned in Section C.1, an application of the theory of center manifolds permits to reduce the dimension of the problem to the number of eigenvalues the modulus of which goes through 1 in the bifurcating family. We exploited that fact in the previous sections by looking at the saddle node or the flip bifurcations directly for maps of the real line, and by analysing the Hopf bifurcation directly in the plane. We show now precisely how to implement such a reduction of the dimension of the problem.

Consider a family of difference equations defined by the map $G : U \times I \rightarrow \mathbb{R}^m$, where U is an open set of \mathbb{R}^m containing the origin, I is an open interval of the real line containing 0, and G is C^r , with $r \geq 2$. We assume that the origin of \mathbb{R}^m is a fixed point of each map $G_\alpha = G(\cdot, \alpha)$ of the family, i.e. $G(0, \alpha) = 0$ for all α .⁷ We let A_α be the Jacobian matrix $D_x G(0, \alpha)$, and assume that A_0 is invertible and has some eigenvalues of modulus 1. The family will thus generally undergo a local bifurcation (near the origin) at $\alpha = 0$. We assume furthermore that a linear change of variables has brought A_0 in its real canonical form, i.e. $A_0 = \text{diag}\{C, B\}$, in which C correspond to the center space F^c of A_0 , and B to the direct sum of the stable and unstable spaces F^s and F^u of A . We shall write accordingly $x = (\xi, \eta)$, where ξ belongs to F^c and η to $F^s + F^u$.

It is convenient here to consider that the map G defines a single difference equation in \mathbb{R}^{m+1} by

$$a_{n+1} = \alpha_n, \quad x_{n+1} = G(x_n, \alpha_n)$$

The trajectory generated by this difference equation, that corresponds to the initial condition (α, x_0) , or more precisely its projection on the x -space,

coincides then to the orbit of x_0 generated by the map G_α . Analysing the local bifurcation of the family G_α near $x = 0$, for α small, is thus equivalent to studying the local properties of the map $H : I \times U \rightarrow \mathbb{R}^{m+1}$ defined by

$$(\alpha, x) \rightarrow (\alpha, G(x, \alpha))$$

near its fixed point $\alpha = 0, x = 0$. One can then apply the local analysis developed in Section B.5.

Since $G(0, \alpha) = 0$ for all α , the partial derivatives of G with respect to α , evaluated at $x = 0, \alpha = 0$, are equal to zero. Thus the Jacobian matrix $DH(0, 0)$ is block diagonal and given by

$$DH(0, 0) = \begin{bmatrix} 1 & & \\ & C & \\ & & B \end{bmatrix}.$$

The stable and unstable spaces of $DH(0, 0)$ are thus equal to $\{0\} \times F^s$ and $\{0\} \times F^u$, respectively, while its center space is $\mathbb{R} \times F^c$.

According to Theorem B.5.1, there is a neighbourhood V of $(0, 0)$ in \mathbb{R}^{m+1} such that H has a local center manifold $W_{(0,0)}^c$ in V . Such a center manifold is C^{r-1} and has the same dimension as the center space of $DH(0, 0)$, i.e. $\mathbb{R} \times F^c$, to which it is tangent at $\alpha = 0, \xi = 0, \eta = 0$. More importantly, it is locally attracting : if (α, x) and $H^n(\alpha, x)$ are in V for all $n \geq 1$, then the distance between $H^n(\alpha, x)$ and $W_{(0,0)}^c$ tends to 0 as $n \rightarrow +\infty$. This fact implies that *any recurrent behaviour that may appear, or disappear, in the bifurcating family, such as fixed points, periodic orbits, invariant closed curves, must lie in a center manifold $W_{(0,0)}^c$* . This result is the key to the reduction of the dimension of the problem.

To be specific, assume that the difference equation associated with H , has the following representation

$$\begin{aligned} \alpha_{n+1} &= \alpha_n \\ \xi_{n+1} &= C\xi_n + f(\xi_n, \eta_n, \alpha_n) \\ \eta_{n+1} &= B\eta_n + g(\xi_n, \eta_n, \alpha_n) \end{aligned}$$

in which f and g are defined for $(\xi, \eta) \in U, \alpha \in I$. Of course the last two lines stand for $x_{n+1} = G(x_n, \alpha_n)$. So, by construction, the functions f and g are $C^r, r \geq 2, f(0, 0, \alpha) = 0$ and $g(0, 0, \alpha) = 0$ for all α , and the first partial derivatives of f and g with respect to ξ and η , vanish at $\xi = 0, \eta = 0, \alpha = 0$.

In this system of coordinates, a local center manifold $W_{(0,0)}^c$ can be represented as the graph of a C^{r-1} function $\eta = \gamma(\alpha, \xi)$, defined on a small open

neighbourhood of $\alpha = 0$, $\xi = 0$ which we may take of the form $J \times V$, where J is an open interval of the real line containing $\alpha = 0$, and V an open set in F^c containing $\xi = 0$. By definition, γ and its first partial derivatives vanish at $(0, 0)$. According to the analysis of Section B.5, the motion on the local center manifold near $\alpha = 0$, $x = 0$, is governed by the difference equation, for (α_n, ξ_n) in $J \times V$,

$$\alpha_{n+1} = \alpha_n$$

$$\xi_{n+1} = C \xi_n + f(\xi_n, \gamma(\alpha_n, \xi_n), \alpha_n) \equiv \Gamma(\xi_n, \alpha_n),$$

with $\eta_n = \gamma(\alpha_n, \xi_n)$ for all $n \geq 0$. In fact (Theorem B.5.3), *the map H is locally topologically conjugate to*

$$\alpha \rightarrow \alpha, \xi \rightarrow \Gamma(\xi, \alpha), \eta \rightarrow B\eta, (\alpha, \xi) \text{ in } J \times V, \eta \text{ in } F^s + F^u.$$

Studying the local bifurcation of the family G_α in \mathbb{R}^m amounts to analysing the local bifurcation of the family $\Gamma_\alpha = \Gamma(\cdot, \alpha) : V \rightarrow F^c$. It is possible to show that $\gamma(\alpha, 0) = 0$, in which case $\Gamma(0, \alpha) = 0$, for all α in J , see Iooss (1979, Chapter V, Section 3). One has moreover $D_\xi \Gamma(0, 0) = C$, and C is a matrix the eigenvalues of which have all modulus one. The dimension of the bifurcation problem is indeed reduced to the number of eigenvalues of the Jacobian matrix $D_x G(0, \alpha)$ that have modulus one at $\alpha = 0$. In particular, if only a simple real eigenvalue, or only a pair of simple nonreal conjugate eigenvalues, have modulus one at $\alpha = 0$, one may try and apply to the family Γ_α , the bifurcation theorems presented in the preceding three sections, on the real line or in the plane. A precise representation in the ambient space \mathbb{R}^m , of the bifurcating fixed points, periodic orbits or invariant closed curves, will then be obtained through the transformation $\xi \rightarrow (\xi, \gamma(\alpha, \xi))$, for each α . Stability or unstability in \mathbb{R}^m , in the dynamics generated by G_α , will then be deduced from the fact, already mentioned, that H is locally topologically conjugate to $\alpha \rightarrow \alpha, \xi \rightarrow \Gamma(\xi, \alpha), \eta \rightarrow B\eta$. In particular, if all eigenvalues λ of the matrix B satisfy $|\lambda| < 1$, and if an object K in F^c (fixed point, periodic orbit, etc ...) has been found to be asymptotically stable in the dynamics generated in F^c by Γ_α , the image of K in \mathbb{R}^m after the transformation $\xi \rightarrow (\xi, \gamma(\alpha, \xi))$, is also asymptotically stable in the dynamics generated in \mathbb{R}^m by G_α .

Remark C.5.1. In practice, in order to apply the bifurcation theorems of the preceding sections, one needs only to know the partial derivatives of $\Gamma(\xi, \alpha)$, and thus of $\gamma(\alpha, \xi)$, at $\alpha = 0$, $\xi = 0$. One can compute them by using the identification technique described in Remark B.5.4.

Remark C.5.2. We already noted that $\gamma(\alpha, 0) = 0$ for all α in J . For every $\bar{\alpha}$, the graph in \mathbb{R}^m of $\gamma(\bar{\alpha}, \cdot)$ represents the section of $W_{(0,0)}^c$ in \mathbb{R}^{m+1} by the hyperplane of equation $\alpha = \bar{\alpha}$; it is therefore a local invariant manifold $W_{\bar{\alpha}}$ of $G_{\bar{\alpha}}$ near $(\xi, \eta) = 0$, having the same dimension as F^c . Since $W_{(0,0)}^c$ is locally attracting in the dynamics generated in \mathbb{R}^{m+1} by the map H , the manifold W_{α} is also locally attracting in the dynamics generated in \mathbb{R}^m by G_{α} , for each α . Since by construction $D_{\xi}\gamma(0, 0) = 0$, W_0 is actually a local center manifold of G_0 . Under some regularity conditions, it can be shown that for $\alpha \neq 0$, the manifold W_{α} is tangent to the generalized eigenspace of $DG_{\alpha}(0)$ corresponding to the eigenvalues that cross the unit circle at $\alpha = 0$, see Iooss (1979, Chapter V.3, Lemma 3).

The fact that local center manifolds need not be unique is not troublesome here. Any invariant closed set that bifurcates from the origin in \mathbb{R}^m for the family G_{α} , such as fixed points, periodic orbits or invariant closed curves, must belong to W_{α} for each α , and this no matter which specific choice of $W_{(0,0)}^c$ was made. For a precise statement, see Iooss (1979, Chapter V.3, Lemma 4).

Notes on the literature

The presentation of the saddle node and flip bifurcations in Sections C.2 and C.3 is a direct transposition of Whitley (1983, Section 1.1). The material of Section C.4 on the Hopf bifurcation is adapted from Iooss (1979, chap. III), Whitley (1983, Sections 2.2, 2.3 and 2.4), Guckenheimer and Holmes (1983, chap. 3.5). Other useful references are Carr (1981, chap. 3.4), Chenciner (1983), Marsden and McCracken (1976, Section 6). Remark C.4.9 on the degenerate Hopf (Chenciner) bifurcation is adapted from the presentation made in Gaunersdorfer, Hommes and Wagener (2006). The material on the center manifold reduction is taken from Iooss (1979, chap. V), Marsden and McCracken (1976, Section 2), Carr (1981, chap. 2.8), Guckenheimer and Holmes (1983, chap. 3.2).

FOOTNOTES TO SECTION C

- (1) Local bifurcations associated with several or multiple eigenvalues crossing the unit circle are far more difficult to handle, and at any rate are “rare” (nongeneric).
- (2) We could have assumed $G(0, 0) = 0$ instead of $G(0, \alpha) = 0$ for all α . The origin is then a fixed point of $G_0 = G(\cdot, 0)$, but since $\frac{\partial G}{\partial x}(0, 0) \neq 1$, the fixed point would persist for small α , and one would go back to the formulation of the text by translating it to the origin, according to the procedure outlined in the last part of Remark C.1.1.
- (3) By continuity, we are sure that all G_α^2 are well defined on a small common open interval containing 0 for α small enough. Thus all conditions of Proposition C.2.3 are satisfied.
- (4) We could have assumed only $G(0, 0) = 0$ instead of $G(0, \alpha) = 0$ for all α . For then, since $D_x G(0, 0)$ has no eigenvalue equal to 1, the fixed point would persist for small α , and one would translate it to the origin according to the procedure outlined in Remark C.1.1.
- (5) For any complex number $z = x + iy$, the notation $\operatorname{Re}(z)$ stands for the real part of z , i.e. $\operatorname{Re}(z) = x$.
- (6) What follows is adapted from Whitley (1983, Sections 2.3 and 2.4). See also Iooss (1979, chap. III.2).
- (7) Here again, this simply means that there is a persistent fixed point of G_α , that has been translated to the origin by a suitable α -dependent change of coordinates.

D. GLOBAL BIFURCATIONS AND CHAOS

We were interested in the preceding section in *local* bifurcations of a family of difference equations $x_{n+1} = G_\alpha(x_n)$, i.e. qualitative changes of the corresponding trajectories that can be observed in *every*, however small, neighbourhood of a fixed point of the family, when the indexing parameter varies. Such qualitative changes may of course occur only in *some* neighbourhood of the stationary state, sometimes quite far away from it. In that case, the bifurcation is *global*. Quite complex phenomena may then obtain: periodic orbits or even “chaotic” trajectories that look random although they are generated by a purely deterministic system. Many of these “exotic” phenomena are still poorly understood. We shall content ourselves here to give a brief account of what happens in the case of one-dimensional nonlinear difference equations, and of so-called “homoclinic” bifurcations of diffeomorphisms of the plane. Some notes at the end of the section will permit the interested reader to pursue her/his own inquiry.

D.1. Maps of an interval

We consider the difference equation

$$(D.1) \quad x_{n+1} = G(x_n)$$

where x is a real number. We are interested in trajectories generated by (D.1) that do not diverge to infinity, so we shall assume that G leaves invariant an interval $[a, b]$, and consider only the restriction of G to that interval, i.e. $G : [a, b] \rightarrow [a, b]$.

If G is continuous, it has a fixed point in the interval. More generally, (D.1) may have solutions that are periodic, or display an irregular, “turbulent” behaviour. We say that x is a *periodic point* of period $k \geq 1$, if x is a fixed point of the k -th iterate of G , i.e. $x = G^k(x)$, and if k is the smallest integer having this property. As usual, the iterates of G are defined recursively through $G^j(x) = G(G^{j-1}(x))$. The corresponding *periodic orbit*, or *cycle*, is then the set $\{x, G(x), \dots, G^{k-1}(x)\}$.

Existence of cycles

Global conditions ensuring the existence of cycles are relatively easy to find and to verify in this simple context, when the period is small. We give an example for cycles of period two and three.

Lemma D.1.1. *Assume that $G : [a, b] \rightarrow [a, b]$ is continuous. Let \bar{x} in (a, b) be a fixed point, with $G(x) > x$ for all x in (a, \bar{x}) . If there exists x_0 in (a, \bar{x}) such that $G^2(x_0) \leq x_0$ (resp. $G^3(x_0) \leq x_0$), then G has a cycle of period two (resp. three).*

We give a short proof in the period two case. Let x_0 in (a, \bar{x}) be such that $G^2(x_0) \leq x_0$. Assume first that the end point a is not a fixed point of G . Since $G^2(a) \geq a$, by continuity, G^2 has a fixed point x in $[a, x_0]$ that is not a fixed point of G : x is a periodic point with period 2. If on the other hand $G(a) = a$, one has $G^2(x_1) > x_1$ for x_1 close enough to a . Then, again by continuity, G^2 has a fixed point in $(x_1, x_0]$, which is a periodic point with period 2. The proof for the period three case follows the same line. The reader will easily verify that the argument works for any period k that is prime.

Fig. D.1 represents the graph of a map G that satisfies the conditions of the Lemma, and that has accordingly a cycle of period two *and* a cycle of period three. The example shows that we should generally expect cycles of different periods to coexist. The following result characterizes the way in which such a coexistence may obtain. It implies in particular that a continuous map from an interval into itself that has a cycle of period three, as in Fig. D.1, has in fact cycles of *every* period, and can thus generate quite complex trajectories.

Fig. D.1

Theorem D.1.2 (Sarkovskii). *Consider the following ordering of the integers*

$$\begin{array}{l}
 3 \succ 5 \succ 7 \succ \dots\dots\dots \\
 \succ 2.3 \succ 2.5 \succ 2.7 \succ \dots\dots\dots \\
 \dots\dots\dots \\
 \succ 2^m.3 \succ 2^m.5 \succ 2^m.7 \succ \dots\dots\dots \\
 \dots\dots\dots \\
 \succ \dots \succ 2^p \succ \dots \succ 8 \succ 4 \succ 2 \succ 1.
 \end{array}$$

If the continuous map $G : [a, b] \rightarrow [a, b]$ has a cycle of period k , then it has also a cycle of period k' for all $k' \prec k$.

Stability

Finding which cycles are stable, if there are any, is an important task when one is faced with so many possible equilibria of the system. Stability and asymptotic stability were defined in Section B for a fixed point or for any closed set (Remark B.1.1). The definition applies therefore to a periodic orbit $(x_0, x_1, \dots, x_{k-1})$, where $G^k(x_0) = x_0$ and $G^n(x_0) = x_n \neq x_0$ for $n = 1, \dots, k-1$. As a matter of fact, since x_0 is a fixed point of G^k , (asymptotic) stability of the periodic orbit is equivalent to the (asymptotic) stability of x_0 in the difference equation associated to G^k (Remark B.1.2). If G is continuously differentiable, we have from the chain rule of differentiation

$$(D.2) \quad (G^k)'(x_0) = G'(x_{k-1}) \dots G'(x_0).$$

Thus if $d = (G^k)'(x_0)$, the cycle is asymptotically stable when $|d| < 1$, unstable when $|d| > 1$. In the sequel it will be useful to say that the periodic orbit is *weakly stable* if $|d| \leq 1$. This terminology is a slight abuse of language, since when $|d| = 1$, the cycle may be stable or unstable depending upon the sign of higher order derivatives of G^k at x_0 , but it will turn out to be consistent in the cases under consideration. The cycle will be said *superstable* if $d = 0$; in view of (D.2), this means that a *critical point* of G , i.e. a point x such that $G'(x) = 0$, belongs to the periodic orbit. Finally, one may remark that none of the foregoing definitions depends upon the particular point chosen on the period orbit. Indeed, we get, again from the chain rule of differentiation, for any element x_n of the cycle

$$\begin{aligned} (G^k)'(x_n) &= G'(x_{n-1}) \dots G'(x_0)G'(x_{k-1}) \dots G'(x_n) \\ &= (G^k)'(x_0). \end{aligned}$$

There is little hope to be able to say something on the stable periodic solutions of (D.1) if one does not make simplifying regularity assumptions about the map G , and its iterates. The first regularity assumption we shall make is that G is *single peaked*.

(D.3) G is C^1 and single peaked, i.e. there exists x^* in (a, b) such that $G'(x^*) = 0$, with $G'(x) > 0$ for $x < x^*$ and $G'(x) < 0$ for $x > x^*$. Moreover $G(x^*) = b$.

The assumption $G(x^*) = b$ involves no loss of generality whenever $G(x^*) > x^*$, since one can then restrict attention to the invariant interval $[a, G(x^*)]$. The assumption implies that G has a unique fixed point \bar{x} in (x^*, b) . The next

condition states that G has no fixed point in the interval (a, x^*) , and that if the end point a is a fixed point, it is unstable.

$$(D.4) \quad G(x) > x \quad \text{for all } x \text{ in } (a, x^*), \text{ and } G'(a) > 1 \text{ if } G(a) = a.$$

The assumption that G is single peaked rules out multiple ups and downs in the graph of G . The curve representing this graph may still make a lot of “little waves” within each interval of monotony of G , and that is something we wish to avoid, not only for G but also for the iterates G^n .

It turns out that the phenomenon is indeed prevented to occur if one assumes that G has a *negative Schwarzian derivative*. This is a global version of a condition that we already met when studying the local bifurcations of maps of the real line : we saw that a flip bifurcation is supercritical if, and in general only if, the value of the Schwarzian derivative of the map, evaluated at the fixed point, was negative (Remark C.3.3). If we note the Schwarzian derivative (assuming that G is of class C^3)

$$SG(x) = \frac{G'''(x)}{G'(x)} - \frac{3}{2} \left[\frac{G''(x)}{G'(x)} \right]^2,$$

the condition reads

$$(D.5) \quad G \text{ is } C^3 \text{ and } SG(x) < 0 \text{ for every } x \text{ such that } G'(x) \neq 0.$$

One can verify by direct computation that

$$SG = -2 |G'|^{1/2} [|G'|^{-1/2}]''.$$

So the assumption (D.5) –which we will write for short $SG < 0$ – means that $|G'|^{-1/2}$ is convex on every interval of monotony of G . It will be satisfied in particular when $|G'|$ or $\text{Log } |G'|$ is concave on such intervals. One may note, incidentally, that the concavity of G is neither necessary nor sufficient to guarantee $SG < 0$.

The next fact implies that assumption (D.5) rules indeed out “little waves” not only in the graph of G , but also in the graphs of *all* its iterates.

Lemma D.1.3. 1) *If $SG < 0$, then $|G'|$ has no positive local minimum.*
 2) *For any two C^3 maps G and H from the real line into itself, one has*

$$S(G \circ H)(x) = SG(H(x))[H'(x)]^2 + SH(x).$$

Thus $SG < 0$ and $SH < 0$ imply $S(G \circ H) < 0$. In particular, $SG < 0$ implies $SG^n < 0$ for all $n \geq 1$.

It turns out that the assumption $SG < 0$ achieves a lot more. Indeed, when G is single peaked, i.e. satisfies assumptions (D.3) and (D.4), there exists *at most* one weakly stable cycle, as the following theorem shows. To state the results most conveniently, it will be useful to say that a periodic orbit P *attracts* a point x if the set of limit points of the sequence of all iterates $G^n(x)$, as n tends to $+\infty$, coincides with P .

Theorem D.1.4. *Assume that $G : [a, b] \rightarrow [a, b]$ satisfies (D.3), (D.4) and (D.5). Then*

- 1) *the map G has at most one weakly stable periodic orbit. This periodic orbit lies in the interval $[G(b), b]$.*
- 2) *If G has a weakly stable periodic orbit, it attracts the critical point x^* . If in addition*

$$(D.6) \quad G''(x^*) < 0,$$

the weakly stable periodic orbit attracts all points of the interval $[a, b]$, except for a set of Lebesgue measure 0.

This result is remarkable for it shows that if we restrict attention to those maps that satisfy the regularity conditions (D.3) through (D.6), we are sure that there exists at most one weakly stable periodic orbit, even though there may be an infinity of cycles. When such a stable periodic orbit exists, it attracts actually most points of the interval. The last feature yields a nice practical method to discover the stable cycle when it exists. It suffices indeed to choose at random a point in the interval, to iterate it by using a computer, and to check whether the iterates converge to a periodic trajectory. If they do, we are sure that the limit cycle is weakly stable (if it were not, one would not have converged to it, if only as a result of rounding off errors). Since iterations have to be stopped after a finite time in practice, this experimental method will enable us to discover cycles that have not too large a period, and that are attracting enough. Of course, this way of proceeding will not allow us distinguishing between the presence of a weakly stable cycle that has a very long period or that is very weakly attracting, and the absence of any stable cycle.

The foregoing theorem can also be used to construct maps satisfying (D.3) through (D.5), that have *no* stable cycle : it suffices that the iterates of the

critical point x^* do not converge, or that they fall upon an unstable cycle. Fig. D.2.a and b give an example of such a pattern, in which the iterates of the critical point hit an unstable fixed point. The maps pictured in these figures satisfy the conditions of Lemma D.1.1 ; they have thus a cycle of period three, and therefore from Sarkovskii's theorem, cycles of every period. None of these cycles is stable, however, when the maps have a negative Schwarzian derivative. In such a case, for any initial condition that does not belong to a periodic orbit, the corresponding trajectory will be aperiodic no matter how long one waits. Maps in this category are good candidates to portray "turbulent" or "chaotic" behaviour. We shall go back to such maps later on when dealing specifically with that issue.

Fig. D.2.a

Fig. D.2.b

Bifurcations

Consider a family of maps G_α indexed by the real number α . When the parameter varies, the map G_α , or one of its iterates G_α^n , may undergo a bifurcation. Since we are considering here maps of the real line, the bifurcation may be of the saddle node type (Section C.2), or a flip bifurcation (Section C.3). Moreover, if each element G_α of the family is assumed to have a negative Schwarzian derivative, any iterate G_α^n has the same property, and every such flip bifurcation should be *supercritical*, a stable cycle of period k becoming unstable and giving rise to a stable cycle of period $2k$ (Remark C.3.3). When dealing with such families, one should expect to observe a cascade of period doubling bifurcations, involving stable cycles, as the parameter varies. This is indeed the pattern one observes in numerical experiments. We review here some mathematical results that explain the phenomenon.

Consider a family of maps G_α from $[a_\alpha, b_\alpha]$ into itself, in which the parameter α belongs to $[0, 1]$. We assume that each G_α is single peaked, i.e. satisfies assumption (D.3) with a unique critical point x_α^* , and moreover, that a_α and b_α depend continuously on α , as well as G_α and its first derivative. We shall say that the family is *full* if

- 1) for $\alpha = 0$, one has $G_0^2(x_0^*) = G_0(b_0) > x_0^*$. In that case all iterates of the critical point, i.e. $G_0^n(x_0^*)$, belong to the interval $[G_0(b_0), b_0]$;
- 2) for $\alpha = 1$, one has $G_1^2(x_1^*) < x_1^*$ and $G_1^3(x_1^*) < x_1^*$.

Thus in a full family, for α near 0, the only periodic orbits that G_α may have, are fixed points or cycles of period 2, as one can easily verify. By contrast, for




Fig. D.2.a




Fig. D.2.b

α near 1, the map G_α has a cycle of period three, in view of Lemma D.1.1, and thus from Sarkovskii's theorem, cycles of every period.

Although Sarkovskii's theorem is not a statement about stable cycles, one should expect nevertheless that, in a full family, for any period k , a stable cycle of that period should appear for some open interval of the parameter, somewhere in $[0, 1]$. Furthermore, in view of the ordering of the integers appearing in Theorem D.1.2, stable cycles of a period equal to a power of 2 should appear first, i.e. for smaller values of α . The next fact makes precise this intuition.

Theorem D.1.5. *Consider a continuous, full, one-parameter family of single peaked maps G_α indexed by α in $[0, 1]$, as above. Then*

- 1) *given any $k \geq 2$, the set of parameters α for which G_α has a superstable cycle of period k is non-empty and closed. Given such an α , there is an open interval containing α such that G_β has a stable cycle of period k for every β in this interval.*
- 2) *Let α_p^* be the first value of α for which there exists a superstable cycle of period 2^p for $p \geq 1$. Then the sequence α_p^* increases with p and converges to some value $\alpha_\infty^* < 1$ as p tends to $+\infty$. For each α in $[0, \alpha_\infty^*)$, all cycles of the map G_α have a period that is a power of 2 or are fixed points.*
- 3) *If superstable cycles of periods 2^p and 2^q with $q > p + 1$ occur respectively for the values α and β in $[0, \alpha_\infty^*)$, then a superstable cycle of period 2^k , with $q > k > p$, must appear for some value in the open interval defined by α and β .*

The foregoing result is valid even for maps that have not a negative Schwarzian derivative. If we make that additional assumption, we get

Theorem D.1.6. *Consider a continuous, full, one-parameter family of maps G_α that satisfy each (D.3), (D.4), (D.5) and (D.6). Then*

- 1) *for any α in $[0, \alpha_\infty^*)$, the map G_α has a unique weakly stable cycle ;*
- 2) *there is an uncountable set of values of α in $(\alpha_\infty^*, 1]$, for which G_α has no weakly stable periodic orbit.*

These results can be translated into a (global) bifurcation diagram, in the spirit of the (local) bifurcation diagrams that were used in the previous section by putting on an horizontal axis the parameter α , and by representing on a vertical axis, for each value of α , the corresponding stable periodic orbit(s), whenever it exists. If we focus on the case where each member of the family has a negative Schwarzian derivative, we obtain a cascade of global period doubling bifurcations, as in Fig. D.3, on the interval $[0, \alpha_\infty^*)$. There, stable periods equal to a power of 2 are “visited” consecutively as in the last line of Sarkovskii’s ordering of the integers (but not necessarily monotonically: think of a nonmonotonic reparametrization of the family). Stable cycles with a period different from a power of 2 must (and will) occur in open intervals of $(\alpha_\infty^*, 1]$. Once such a stable period obtain, say an odd number, one should observe subsequently a cascade of global period doubling bifurcations. For an uncountable number of values in $(\alpha_\infty^*, 1]$, the map is “aperiodic”: there is no stable cycle.

Fig. D.3

Such a pattern can be produced “experimentally”, by using Theorem D.1.4, and simulating the difference equations on a computer. One may employ a grid of values of the parameter, and plot on a vertical axis above each α in the grid, the values of the iterates $G_\alpha^n(x)$, say for $n = 100$ to $n = 200$, of a point chosen at random in the corresponding interval $[a_\alpha, b_\alpha]$. If each G_α has a negative Schwarzian derivative, this procedure will display clearly stable cycles that have a low period and are attracting enough. The experimental procedure will thus reproduce neatly the cascade of period doubling bifurcations in $[0, \alpha_\infty^*)$. Beyond α_∞^* , we shall observe typically in some places, little windows with small stable periods such as 3 or 5, together with the beginning of the corresponding period doubling cascade ; in other places, we will observe a “mess” : there is then either no stable cycle, or a stable cycle with a long period or that is only little attracting.

Fig. D.4 describes such an experimental bifurcation diagram for the *logistic family*

$$G_\alpha(x) = 1 - \alpha x^2 \quad , \quad x \text{ in } [-1, 1],$$

for $0 < \alpha \leq 2$. For each value of the parameter α in that interval, G_α is single peaked, with a unique critical point at $x^* = 0$ and $G_\alpha(x^*) = 1 > x^*$; it has a negative Schwarzian derivative, since $G_\alpha'''(x) \equiv 0$. It is easy to verify that $G_\alpha^2(x^*) > x^*$, for α close to 0. For $\alpha = 2$, one has $G_\alpha^2(x^*) = G_\alpha^3(x^*) = -1 < x^*$; the graph of G_α is then as in Fig. D.2.b : there is a cycle of period three,




Fig. D.3

hence a cycle of every period, and none of these cycles is stable. The logistic family, for α in $(0, 2]$, is full.

Fig. D.4

Topological chaos

Nonlinear difference equations on the real line can generate periodic orbits, as we have seen. They can also yield much more complex trajectories, that look “random” and rather unpredictable, although they are generated by a purely deterministic dynamical system. We have already alluded to the possibility of such a behaviour when giving examples of maps that have no stable cycle (Fig. D.2). The phenomenon has attracted a lot of attention, for it seems to open the possibility to “explain” phenomena that look apparently random by deterministic nonlinear dynamical systems, and thus to reconcile 19th century determinism and unpredictability.

The notion of chaos that we wish to discuss first is due to Li and Yorke. We say that the map G from $[a, b]$ into itself displays *topological chaos* if there is an uncountable invariant subset S of the interval, containing no periodic orbits, such that

- 1) for every x, y in S , $x \neq y$,

$$\limsup_n |G^n(x) - G^n(y)| > 0,$$

$$\liminf_n |G^n(x) - G^n(y)| = 0 ;$$

- 2) for every x in S and y periodic,

$$\limsup_n |G^n(x) - G^n(y)| > 0.$$

This definition means that for any two (different) initial conditions x and y in the invariant chaotic set S , the corresponding trajectories become infinitely often extremely close to each other and also infinitely often noticeably separated. The notion seems to account for one essential feature of “turbulence” : small perturbations of initial conditions lead to trajectories that are notably different. This property is called *sensitive dependence on initial conditions*.

Li and Yorke (1975) proved that if the continuous map G has a cycle of period three, then G is indeed chaotic in the above sense. On the other hand, it is not difficult to see that if an iterate G^n is chaotic in the above sense




Fig. D.4

for some $n \geq 1$, then G displays also topological chaos. If one puts together these facts with Sarkovskii's Theorem, one gets

Theorem D.1.7. *If the continuous map $G : [a, b] \rightarrow [a, b]$ has a cycle of a period that is not a power of two, then G displays topological chaos.*

Another popular notion of topological chaos, which is in fact essentially equivalent to that of Li and Yorke, can be found in Devaney (2003, Section 1.8, Definition 8.5). The map G from $[a, b]$ into itself is said to be *chaotic in the sense of Devaney* if there is an uncountable invariant subset S of the interval such that :

- (1') G has sensitive dependence on initial conditions on S , i.e. there exists $\delta > 0$ such that for any x in S and any neighborhood U of x in S , there is y in U such that $|G^n(y) - G^n(x)| > \delta$ for some $n > 1$.
- (2') G is "topologically transitive" on S , i.e. G has an orbit that is dense in S .
- (3') S contains a dense set of periodic orbits.

Theorem D.1.7 is equally valid when topological chaos is defined in the sense of Devaney instead of Li and Yorke. It may be noted that quite generally (i.e. when $G : X \rightarrow X$ acts continuously on a compact metric space), the sensitive dependence on initial conditions property (1') is implied by the two others (2') and (3'), see Banks et al. (1992).

The above notion of "topological chaos" is not quite satisfactory from a "physical" point of view, however. It happens very often that although a given map G is chaotic in the above sense, there is a stable periodic orbit that attracts most points of the interval with respect to the Lebesgue measure. In view of Theorem D.1.4, this will happen if G is single peaked, has a negative Schwarzian derivative, and has a stable cycle of a period that is not a power of two. In a bifurcating family of maps G_α that are single peaked and have a negative Schwarzian derivative as in Theorems D.1.5 and D.1.6, maps G_α will often be chaotic in the sense of Li and Yorke or Devaney in the whole interval $(\alpha_\infty^*, 1]$, yet for many open subintervals, there will be a stable cycle attracting most points of $[a_\alpha, b_\alpha]$. In such circumstances, the invariant chaotic set S has Lebesgue measure 0, and its presence may affect the trajectories for some time, but it will not influence the asymptotic behaviour of the system: one talks sometimes of "*transient chaos*" in such a case.

Ergodic chaos

The above discussion shows that in this context, *a satisfactory notion of chaos should involve maps that have no stable cycle*. In view of Theorem D.1.4, if G is single peaked and satisfies $SG < 0$, this means that the iterates of the critical point either do not converge, or hit an unstable periodic orbit.

Among the class of such aperiodic maps, of special interest are those which have an “invariant probability measure” that has an integrable density with respect to the Lebesgue measure. The probability measure ν on $[a, b]$ (endowed with its Borel σ -algebra) is said to be *invariant* with respect to G if $\nu(G^{-1}(A)) = \nu(A)$ for any Borel set. An example of such an invariant probability measure is provided by considering a periodic orbit $\{x_0, x_1, \dots, x_{k-1}\}$ and the probability measure ν that assigns the weight $1/k$ to each x_j , $j = 0, \dots, k - 1$. More generally, the support of an invariant probability measure, i.e. the smallest closed set Λ that has probability one, is an invariant set : $G(x)$ is in Λ whenever x belongs to Λ . If ν has a density (we shall say then that ν is *continuous*), the invariant set contains an uncountable set of points, and has positive Lebesgue measure. Finally, ν is said to be *ergodic* if for every ν -integrable real function f ,

$$\frac{1}{n} \sum_{j=1}^n f(G^{j-1}(x)) \rightarrow \int f d\nu$$

as n tends to $+\infty$, for ν -almost every x . This implies in particular that if one considers for each x and every n , the empirical distribution $\nu_n(x)$ that is generated by the iterates $G^j(x)$ for $j = 0, \dots, n - 1$, which assigns the weight $1/n$ to each $G^j(x)$, then the sequence $\nu_n(x)$ converges weakly to ν for ν -almost every x (see, e.g. Parthasarathy (1967, Theorem 9.1) for a proof of that fact, and for a definition of the weak convergence of probability measures). Thus if G has an invariant probability measure ν that is continuous and ergodic, most of the trajectories starting in the support Λ of ν will stay there and will look quite complex since they will eventually fill up the entire support of the limit distribution ν . In fact, most of these trajectories will be *dense* in Λ . Yet these trajectories display strong statistical regularities, since empirical distributions and time averages become asymptotically fairly stable for ν -almost every initial point.

It turns out that maps having a graph like those of Fig. D.2. do have an invariant, continuous, ergodic probability measure, as the next fact shows.

Theorem D.1.8. *Assume that the map G from $[a, b]$ into itself satisfies (D.3), (D.4), (D.5) and (D.6), that it has no weakly stable periodic orbit, and that there exists an open neighbourhood V of the critical point x^* such that the iterates $G^n(x^*)$ do not belong to V , for $n \geq 1$. Then G has a unique continuous invariant probability measure. It is ergodic.*

The foregoing conditions are met if G satisfies (D.3), (D.4), (D.5) and (D.6), and if the iterates of the critical point hit an unstable periodic orbit.

One may note that when G has an invariant probability measure ν that is continuous and ergodic, the trajectories in the support Λ of ν exhibit sensitive dependence on initial conditions, as defined for the notion of topological chaos in the sense of Li and Yorke (condition (1)) or of Devaney (condition (1')). The basic (and essential!) difference with topological chaos is that *here the invariant "chaotic" set Λ has positive Lebesgue measure.*

The idea of sensitive dependence on initial conditions can also be captured by using the notion of so-called *Lyapounov exponents*. Let x be a point in $[a, b]$, and let $x_j = G^j(x)$ be its iterates for $j \geq 0$, with the convention that $x_0 = x$. The Lyapounov exponent of G at x is then defined as

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \text{Log} |G'(x_{j-1})| = \lim_{n \rightarrow \infty} \frac{1}{n} \text{Log} |(G^{n-1})'(x)|$$

whenever the limit exists. When the definition is specialised to the case of a fixed point, or more generally a periodic point of period k , the corresponding Lyapounov exponent is simply the logarithm of the modulus of $(G^k)'(x)$. Intuitively, the Lyapounov exponent measures the average (exponential) rate of separation of the orbits of points near x from the trajectory generated by x . Thus the idea that trajectories are sensitive to small changes of the initial condition x , may be expressed by the property that the Lyapounov exponent of G at x exists and is positive.

Consider now a map G that has an ergodic, continuous invariant probability measure ν . Then by definition of ergodicity, Lyapounov exponents exist for ν -almost every x in the support Λ of ν ; they are in fact equal to $\int \text{Log} |G'(x)| d\nu$, and are thus independent of x . We may then speak of *the* Lyapounov exponent of G on Λ . If we assume that it is positive, then the set Λ has positive Lebesgue measure, and almost every initial condition x in Λ gives rise to a trajectory that is dense in Λ and displays sensitive dependence on initial conditions. Although the law of motion of the system is deterministic, trajectories in Λ will look random, and will be in fact unpredictable after a significant number of iterations if one makes even small errors of measurement of the initial condition. The set Λ is then an instance of what is called a *wild attractor*, and gives rise to *ergodic chaos*.

Abundance of asymptotically stable periodic behaviour versus ergodic chaos

A natural question arises in this context : what are the relative frequencies of stable periodic behaviour and of chaos in one-dimensional difference equations ? The issue involves in fact a subtle conflict between two notions of “genericity”.

A convenient framework to analyze the issue is, again, to consider a one-parameter family of single peaked maps having a negative Schwarzian derivative, as in Theorem D.1.6. First, one should expect asymptotic stability of a periodic orbit to be *structurally stable*, i.e. to persist after a small C^r -perturbation of the map G_α (this follows from the structural stability of maps satisfying the so-called "Axiom A", see Nusse (1986) or De Melo and van Strien (1993, p. 227)). In fact, in the class of S -unimodal maps satisfying conditions (D.3), (D.4), (D.5) and (D.6) as in Theorem D.1.6, asymptotic stable periodic behaviour turns out to be *dense* (this follows from the early work of Jakobson (in Russian), and that of Kozlovski (1998, Ch. 7); see Jakobson and Swiatek (2002, section 4.2)). So under the assumptions of Theorem D.1.6, the set of values of the parameter α for which a stable period obtains is *open and dense* in $[0, 1]$. According to this topological viewpoint, maps G_α that have no stable cycles in the family are “rare”, and moreover, *structurally unstable*, i.e. a small perturbation of the parameter α would lead back to a stable period. Examples of such structural instability are most easily visualized by looking at Fig. D.2. According to Theorem D.1.8, the maps there have an invariant continuous probability measure, because the iterates of the critical point happen to hit upon an unstable fixed point. It is clear, however, that such a feature can be destroyed by a small smooth perturbation of the map under consideration.

By contrast, it was shown by Jakobson (1981) that for the logistic family of quadratic maps as in Fig. D.4, the set of values of the parameter α for which G_α has an invariant, continuous ergodic probability measure, and displays sensitive dependence on initial conditions (has positive Lyapounov exponent) has *positive Lebesgue measure*. That the same result holds for a one-parameter family of S -unimodal maps as in Theorem D.1.6, is a consequence of the works of Misiurewicz (1981, 1983) and of Thieullen, Tresser and Young (1994). According to this measure theoretic viewpoint, chaotic behaviour *cannot* be considered as rare. If one chooses at random the parameter α in the family, there is a positive probability to obtain a map that leads to ergodic chaos.

Theorem D.1.9. *Consider a continuous, full, one-parameter family of maps G_α satisfying (D.3), (D.4), (D.5) and (D.6) as in Theorem D.1.6. Then*

- 1) *the set of parameters α such that G_α has a unique stable cycle, is open*

and dense,

- 2) *the set of parameters α such that G_α has a continuous, ergodic invariant probability measure ν_α , with a support Λ_α having a positive Lyapounov exponent, has positive Lebesgue measure.*

The conclusion of this discussion is that one should be cautious when evaluating claims of the existence of chaos. Topological chaos in the sense of Li and Yorke or Devaney is a quite unsatisfactory notion since the invariant chaotic set may (and will for an open and dense set of the parameter α in Theorem D.1.9) be of Lebesgue measure zero and coexist with a stable cycle that attracts almost all initial conditions in the sense of the Lebesgue measure. Focusing on the stronger and more satisfactory notion of ergodic chaos as in 2) of Theorem D.1.9, one gets that such ergodic chaos occurs with a positive probability. Yet such a phenomenon is structurally unstable, i.e. it goes away, and one gets back to the existence of a stable cycle that attracts almost all initial conditions, after a small smooth perturbation. As a consequence, one should be very cautious when facing claims of the existence of “chaos” based upon the result of numerical simulations as in the bifurcation diagram reproduced in Fig. D.4 for the logistic family of quadratic maps. This family satisfies the assumptions of Theorem D.1.9. The apparently “messy” trajectories reproduced in Fig. D.4 cannot therefore be interpreted, strictly speaking, as “chaotic” since for an open and dense set of the bifurcating parameter α , stable asymptotic behaviour prevails. These complicated orbits may in most cases be only viewed as the result of the transient influence of topological chaos, or as reflecting the presence of a stable cycle with a very long period together with some small amount of noise due to rounding off errors in the simulation.

Remark D.1.10. It was noted in the text that ergodic chaos, i.e. the existence of a continuous ergodic invariant probability measure ν , implied sensitive dependence on initial conditions in the support Λ of ν , this support having positive Lebesgue measure. Conversely, topological chaos in the sense of Li and Yorke or of Devaney on an invariant “chaotic” set S having positive Lebesgue measure may occur without implying ergodic chaos, i.e. the existence of a continuous invariant probability measure. That this could occur for S -unimodal maps satisfying the assumptions of Theorems D.1.6 and D.1.9 (even for the logistic family of quadratic maps of Fig. D.4) was shown by Johnson (1985) and by Bruin, Keller, Nowicki and van Strien (1996). Be it as it may, “thick” topological chaos (on an invariant chaotic set S with positive Lebesgue measure) can only occur, within a one-parameter family

of S -unimodal maps as in Theorem D.1.9, outside an open and dense set of parameters α .

Remark D.1.11. It is not difficult to verify that assumptions (D.3), (D.4) and (D.6) are preserved when making a change of variable of the form $x \rightarrow h(x)$, where h is a C^r diffeomorphism that preserves orientation, i.e. such that $h'(x) > 0$ for all x . By contrast, the condition $SG < 0$, as any global convexity assumption, is *not* preserved through a nonlinear change of variable. This situation is mathematically awkward and makes one wonder about the actual status of the negative Schwarzian derivative condition (D.5). Recent results appear to claim that such a condition would be in fact redundant in the case of C^3 unimodal maps (as in (D.3), (D.4)) with a nonflat critical point (satisfying (D.6)). The central argument is that there is then an interval around the critical point such the first entry map to this interval has a negative Schwarzian derivative, so that condition (D.5) might not be actually needed (De Melo and van Strien (1988) ; Kozlovski (1998, section 3.4), (2000)).

D.2. Homoclinic bifurcations and horseshoes

The example of complex periodic orbits and of chaos that were just described involved single peaked maps of the real line with a significant “hump”. Such maps are admittedly special, but they contain a mechanism that seems to show up in many other cases of deterministic chaos that have been found in higher dimensions : if we look at the maps represented in Figure D.2, for instance, the image of the interval is obtained by stretching it, and folding it back into itself. We present now a diffeomorphism of the plane, the so-called *horseshoe* map, that is due to Smale, and leads to a complex invariant set through a similar mechanism. As we shall see horseshoes arise for diffeomorphisms of the plane when there is a transverse crossing of the stable and unstable manifolds of a hyperbolic fixed point. The phenomenon was known, as least qualitatively, to Poincaré and Birkhoff.

The Horseshoe

Consider a C^1 diffeomorphism G of the plane, that sends the unit square S onto the horseshoe shaped set $G(S)$, as in Figure D.5, with A' being the image of A , B' that of B , etc. (the map may actually be defined only on a neighborhood of the region that interests us). The transformation involves first a uniform vertical contraction by a factor $a < 1$, second a uniform horizontal expansion by a factor $b > 1$, and finally, a folding of the resulting rectangle in its middle. The only nonlinear transformation is the third one,

and it is assumed that it concerns only points of the horseshoe that fall outside the unit square.

Figure D.5

We are interested in characterizing the set Λ of points x of the unit square such that $G^n(x)$ is in S for all n , positive and negative. In this respect, it is useful to let Δ_0 and Δ_1 stand for the pre-images of the intersection of $G(S)$ and S , as in Figure D.6.a. By construction, the set $G(S) \cap S$ is composed of two horizontal rectangles of height a , while Δ_0 and Δ_1 are vertical rectangles of width b^{-1} .

The set of points x of the unit square such that $G(x)$ belongs to S , is thus equal to the union of Δ_0 and Δ_1 , or equivalently to $G^{-1}(G(S) \cap S)$. By iterating the map twice, we see that the image by G of $G(S) \cap S$ is composed of two thinner horseshoes that are contained in $G(S)$. Their intersections with the unit square yield four horizontal rectangles of height a^2 , the union of which form the set

$$G(G(S) \cap S) \cap S = G^2(S) \cap G(S) \cap S,$$

see Figure D.6.b. The set of points x of the unit square such that both $G(x)$ and $G^2(x)$ belong to S is the union of the images by G^{-2} of these four horizontal rectangles. This set is therefore composed of four vertical rectangles of width b^{-2} , as shown in Figure D.6.c. Proceeding by induction on n , one sees that the set of points x of S such that $G^j(x)$ is in S for every $j = 1, \dots, n$, is the union of 2^n vertical rectangles of width b^{-n} , given by

$$G^{-n}(G^n(S) \cap \dots \cap G(S) \cap S).$$

In the limit for $n \rightarrow +\infty$, the iterates $G^j(x)$ of x lie in the unit square for all $j \geq 0$ if and only if x belongs to a family of vertical segments of S . The foregoing construction makes apparent that the projection of this family of vertical segments on the horizontal basis of S is a *Cantor set* (also called a *perfect set*) Λ_+ , i.e. a closed set such that 1) the largest connected subset is a point, and 2) every point of Λ_+ is an accumulation point of Λ_+ .

Figure D.6.a

Figure D.6.b

Figure D.6.c

One can proceed in a similar fashion for backward iterates. The set of points x of S such that $G^{-1}(x)$ is also on S is equal to the intersection of




Fig. D.5




Fig. D.6.a




Fig. D.6.b




Fig. D.6.c

$G(x)$ and S , i.e. to the two horizontal rectangles of Figure D.6.a. Similarly, the set of points x of the unit square having the property that both $G^{-1}(x)$ and $G^{-2}(x)$ are in S , is composed of the four horizontal rectangles of Figure D.6.b. Proceeding by induction on n and going to the limit, yields that the set of points x of S such that $G^{-n}(x)$ is in S for every $n \geq 0$ is a family of horizontal segments. Its projection on the vertical side of the unit square is, again, a Cantor set Λ_- .

Therefore $G^n(x)$ belongs to S for all n , positive and negative, if and only if x belongs to the product of these two Cantor sets, i.e. $\Lambda = \Lambda_+ \times \Lambda_-$.

To describe the structure of the orbits that lie in the invariant set Λ that we just obtained, it is convenient to employ a method called *symbolic dynamics*. The principle of the approach is to substitute to the complex task of tracking down where the forward and backward orbits of a given point lie exactly, the much simpler task of identifying *regions* in which the points of the orbit may fall.

In the case at hand, if x is in Λ , every iterate $G^n(x)$ belongs either to Δ_0 or Δ_1 , for each positive or negative n . For any x in Λ , we can therefore define a bi-infinite sequence $I(x)$, the n -th element of the sequence, i.e. $I_n(x)$, being equal to 0 if $G^n(x)$ is in Δ_0 , and to 1 if $G^n(x)$ belongs to Δ_1 , for $-\infty < n < +\infty$.

It can be shown that the map $x \rightarrow I(x)$ from Λ to the set Σ of bi-infinite sequences of 0's and 1's, is onto and one-to-one. In fact, a much stronger statement is true. Suppose that the set Σ is endowed with the metric defined by

$$(D.7) \quad d(I, J) = \sum_{-\infty}^{+\infty} \delta_n 2^{-|n|},$$

with $\delta_n = 0$ if $I_n = J_n$, and $\delta_n = 1$ otherwise. Two sequences I and J are thus close in the topology induced by the metric (D.7), if and only if they agree on a long enough "central block", i.e. $I_n = J_n$ for all $|n| \leq N$ for some large enough N . Then the map $x \rightarrow I(x)$ from Λ onto Σ , endowed with this metric, is actually an *homeomorphism*. Since for any element x of Λ , the sequence corresponding to $G(x)$, i.e. $I(G(x))$, is obtained from the sequence $I(x)$ by shifting indices one place to the left, the restriction of G to the invariant set Λ is *topologically conjugate to the (left) shift automorphism* $\sigma : \Sigma \rightarrow \Sigma$ defined by $[\sigma(I)]_n = I_{n+1}$, for $-\infty < n < +\infty$ (see Guckenheimer and Holmes (1983, Theorem 5.1.1)).

This result enables us to substitute to the study of the orbits of points x of Λ , the analysis of the corresponding sequences $I(x)$. For instance, an

element x of Λ is a periodic point, of period k , if and only if $I(x)$ is periodic, of period k . It follows that G has two fixed points in S , one in Δ_0 and the other in Δ_1 . More generally, G has periodic orbits of every period in Λ . It is not difficult to see that the set of these periodic orbits is actually dense in Λ . Indeed, let x be in Λ such that the corresponding sequence $I(x)$ is not periodic. Let J be a sequence that agrees with $I(x)$ for all $|n| \leq N$, and that has period $k > N$. By taking N (and thus k) large enough, the sequence J can be made as close as one wishes to $I(x)$ in Σ , endowed with the metric (D.7). Since the map $x \rightarrow I(x)$ is an homeomorphism, the element y of Λ such that $J = I(y)$, is a periodic point, with period k , and is also as close as one wishes to x .

The same type of argument shows that Λ contains an uncountable set of nonperiodic orbits, that is dense in Λ . It is also quite easy to show that Λ contains a dense orbit. Indeed, let J be a sequence in Σ that has as a subsequence every finite sequence of 0's and 1's one can think of. By adapting the above density argument, the orbit generated by the element y of Λ such that $I(y) = J$ is actually dense in Λ . To sum up this discussion,

Theorem D.2.1. (Smale) : *The horseshoe map G has an invariant Cantor set Λ such that*

- 1) Λ has periodic orbits of every period. The set of such periodic orbits is dense.
- 2) Λ has an uncountable set of nonperiodic orbits. This set is also dense in Λ .
- 3) Λ has a dense orbit.

It is intuitively clear from the preceding discussion (but nontrivial to prove), that the qualitative features of the example should be structurally stable, i.e. should remain valid after a small enough C^1 perturbation. Indeed,

Theorem D.2.2. (Smale) : *Let G^* be a diffeomorphism of the plane that is C^1 close enough to the horseshoe map G (at least in the region of interest). Then G^* has an invariant Cantor set Λ^* such that the restriction of G^* to Λ^* is topologically conjugate to the restriction of G to Λ .*

Remark D.2.3 : Despite its many attractive features, the invariant set Λ of the horseshoe map is *not* an attractor : if one perturbrates slightly the initial condition x so as to make it to fall outside Λ , the forward or backward

orbit of x will eventually leave the set S . Yet, the presence of the complicated invariant set Λ will affect the behaviour of nearby trajectories, often for quite some time. For an illuminating example of a complicated invariant set that can be analyzed through the above symbolic dynamics methods, and that is indeed an attractor (the so-called *solenoid*), see Lanford (1983). The example is three dimensional, and involves stretching and folding a *solid torus*.

Homoclinic Bifurcations

The qualitative features of the horseshoe map can be generated in the case of diffeomorphisms of the plane, when there is transverse crossing of the stable and unstable manifolds of a hyperbolic fixed point.

To see this point, consider a C^r diffeomorphism G of the plane, with $r \geq 1$, and let \bar{x} be fixed point of G (here again G may be defined only on a neighborhood of \bar{x} that is sufficiently large to make the following arguments to go through). We assume \bar{x} to be hyperbolic, i.e. the Jacobian matrix $DG(\bar{x})$ has no eigenvalue of modulus one. We know that the *local* stable and unstable manifolds $W_{(\bar{x})}^s$ and $W_{(\bar{x})}^u$ are then well defined and unique in a sufficiently small neighborhood of the fixed point (Theorem B.5.1). The *global stable manifold* of \bar{x} is then defined by taking the union of the backward iterates of $W_{(\bar{x})}^s$,

$$W^s = \bigcup_{n \geq 0} G^{-n}(W_{(\bar{x})}^s).$$

Similarly, the *global unstable manifold* of \bar{x} is obtained by taking the union of the forward iterates of $W_{(\bar{x})}^u$,

$$W^u = \bigcup_{n \geq 0} G^n(W_{(\bar{x})}^u).$$

We consider the case where the global stable and unstable manifolds of \bar{x} intersect transversely, as in Figure D.7.a. The point of intersection, say y , is then an *homoclinic point*. There must be actually infinitely many such homoclinic points : all forward and backward iterates of y , i.e. $G^n(y)$ for $-\infty < n < +\infty$, belong to both W^s and W^u . The (forward and backward) orbit of y is then called a *transversal homoclinic orbit*. Since $G^n(y) \rightarrow \bar{x}$ as $n \rightarrow +\infty$, the unstable manifold W^u must cross more and more often the stable manifold W^s , and oscillate more and more wildly as one gets closer to the fixed point \bar{x} . A similar statement holds for the stable manifold.

Figure D.7.a

Figure D.7.b




Fig. D.7.a




Fig. D.7.b

One should expect complex dynamics to occur in such a case. If one considers a small tubular neighborhood V of the stable manifold, the image $G^n(V)$ should be, for $n > 0$ large enough, a tubular neighborhood of the unstable manifold, as in Figure D.7.b. Intuitively, all the qualitative features of the previous horseshoe example should also be present here. Indeed

Theorem D.2.4. (Smale homoclinic theorem) : *Let G be a C^1 diffeomorphism of the plane and \bar{x} a hyperbolic fixed point. If the stable and unstable manifolds of \bar{x} intersect transversely at $y \neq \bar{x}$, there exists a neighborhood V of the fixed point and an integer $n > 0$ such that the restriction of G^n to V is topologically conjugate to the horseshoe map.*

The same phenomenon will occur in particular when a family of diffeomorphisms G_α of the plane, undergoes a (global) *homoclinic bifurcation*, i.e. when the unstable manifold of a hyperbolic fixed point becomes tangent (in fact at infinitely many points) to the stable manifold, say when $\alpha = 0$, to lead to a transverse crossing for $\alpha > 0$. The qualitative features of such a bifurcation are described in Figure D.8.a, b, and c.

Figure D.8.a

Figure D.8.b

Figure D.8.c

Remark D.2.5. (*Hénon-like diffeomorphisms*). The above horseshoe map displays a feature that is likely to be interesting for generating complex dynamics in multidimensional settings, where a small ball is stretched significantly in one direction and shrunk in all other directions. The dynamics of such a multidimensional diffeomorphism should be close to the dynamics generated by a one-dimensional map if the contraction is strong enough. One would expect then the multidimensional diffeomorphism to exhibit the same kind of dynamic complexities (topological, ergodic chaos) as those which we described in section D.1, if the one-dimensional map is unimodal with a significant hump, as considered there.

An example of this type of multidimensional dynamics, that has been extensively studied in the literature, is the (twodimensional, two parameters) Hénon family of maps $G_{\alpha,\beta}$:

$$x_{n+1} = 1 - \alpha x_n^2 + \beta y_n,$$

$$y_{n+1} = x_n,$$

that was introduced by Hénon in 1978. The map $G_{\alpha,\beta}$ is indeed a diffeomorphism when $\beta \neq 0$. But it is no longer one when β is set equal to 0, as it is in that case the composition of a projection with the logistic family. One would expect then for β small, the diffeomorphism $G_{\alpha,\beta}$ to display




Fig. D.8.a




Fig. D.8.b




Fig. D.8.c

complex dynamics as the logistic family, e.g. to exhibit ergodic chaos with positive probability as stated by Jakobson’s theorem (1981) (see 2) of Theorem D.1.9). Indeed, it can be shown that for any given $\beta > 0$ small, there is a set A_β of parameters α , with positive Lebesgue measure, such that $G_{\alpha,\beta}$ has a continuous invariant probability measure $\nu_{\alpha,\beta}$, with a support $\Lambda_{\alpha,\beta}$ having a positive Lyapounov exponent, for α in A_β (Benedicks and Carleson (1991), Benedicks and Young (1992, 1993)). For further information, in particular on extensions and open issues for “Hénon-like” maps, the reader may consult the very clear recent exposition by Benedicks (2003).

D.3. Backward and forward dynamics : the “Inverse Limit” approach

A basic characteristic of socioeconomic models is that the current state of the system depends not only on past states, but also on the agents’ expectations about the future, since those influence current decisions. Perfect foresight dynamics are thus often given in implicit form by a map that can be solved “backward”, $x_t = g(x_{t+1})$, where x belongs to some open subset U of \mathbb{R}^p , and where g maps U , but that may not be one to one (see e.g. Benhabib and Day (1982), Grandmont (1985)). If one restricts attention to a compact invariant subset X of U , i.e. with $g(X) = X$, one can apply the results surveyed in these lecture notes to the trajectories generated by the iterates of g , i.e. $g^j(x)$ for $j \geq 0$, but unfortunately these trajectories go backward in time. Getting information on trajectories that go forward in time may be problematic when g is not a diffeomorphism (as when g is a unimodal map on an interval), since reversing the arrow of time may generate a difference equation $x_{t+1} \in \varphi(x_t)$ where φ , the “inverse” of g , is a multivalued correspondence. The “inverse limit” approach is a method that has been designed to cope with that sort of issue. The main difficulty originates from the fact that, while a backward trajectory $(\dots, g^2(x_0), g(x_0), x_0)$ is uniquely determined by its initial condition x_0 in X , there may be many (even infinitely many) forward trajectories corresponding to x_0 . The formal “trick” employed in the inverse limit approach is to consider the space of forward trajectories as the state space itself.

Specifically, let X (the *factor space*) be compact and $g : X \rightarrow X$ (the *bonding map*) with $g(X) = X$ be (at least) continuous. The pair (X, g) is called an *inverse system*. A forward trajectory is an element $\mathbf{x} = (x_0, x_1, x_2, \dots)$ in the product space X^∞ (endowed with a metric that is compatible with the product topology), such that $x_t = g(x_{t+1})$ or $x_{t+1} \in \varphi(x_t)$ for all $t \geq 0$. This space of forward trajectories is called the *inverse limit* of the inverse system (X, g) and is noted $\varprojlim(X, g) = Z$. The action of the bonding map

on X induces a *natural map* G acting on the inverse limit space Z . For any $\mathbf{x} = (x_0, x_1, x_2, \dots)$ in Z ,

$$G(\mathbf{x}) = (g(x_0), g(x_1), g(x_2), \dots) = (g(x_0), x_0, x_1, \dots).$$

It turns out that G is indeed an homeomorphism on Z . Its inverse G^{-1} is in fact the (*left*) *shift homeomorphism* σ defined by

$$\sigma(\mathbf{x}) = \sigma(x_0, x_1, x_2) = (x_1, x_2, \dots)$$

for any (x_0, x_1, x_2, \dots) in Z . Therefore by considering the space of solutions of the implicit difference equation $x_t = g(x_{t+1})$ or $x_{t+1} \in \varphi(x_t)$, one transforms the non-invertible map g acting on X into an invertible map G acting on Z , to get finally its inverse $\sigma = G^{-1}$ moving forward in time.

The inverse limit approach aims accordingly at studying the forward dynamics induced by g by looking at the stability properties or complex dynamics generated by the corresponding shift map σ on the inverse limit space Z . An important ingredient of the analysis is of course the topology employed on that space. The route usually taken is to endow Z with the metric structure generated by the following distance $d(\mathbf{x}, \mathbf{y})$ between two elements $\mathbf{x} = (x_0, x_1, x_2, \dots)$ and $\mathbf{y} = (y_0, y_1, y_2, \dots)$ in that space :

$$(D.8) \quad d(\mathbf{x}, \mathbf{y}) = \sum_{t=0}^{\infty} \|x_t, y_t\| 2^{-t}$$

where $\|x_t, y_t\|$ is the Euclidean distance between x_t and y_t in X . Two infinite sequences \mathbf{x} and \mathbf{y} are close to each other in this metric if and only if they are very close component-wise for a very long but finite time, but they can differ significantly asymptotically as time goes to infinity. The metric defined by (D.8) appears to be directly inspired from the fact that complex dynamics are often analyzed in the field by representing orbits as an infinite sequence of indices taking values e.g. in $\{0, 1\}$, depending on which “region” the state x_t lies in, and by measuring the distance between two such index sequences by a formula like (D.8) : see e.g. (D.7) for the horseshoe map.

Topological chaos

Since the backward map g is well defined on the compact space X , the standard definition of *topological chaos in the sense of Devaney* can be transposed here, as in conditions (1') (2') (3') in section D.1 (sensitive dependence on initial conditions, topological transitivity, existence of a dense set of periodic orbits). It is easy to see that the same definitions apply to the inverse forward multivalued difference equation, $x_{t+1} \in \varphi(x_t)$. Intuitively, one should

expect topological chaos in this sense to prevail for φ whenever it occurs for g . Indeed (see Kennedy and Stockman (2008, Theorem 6), that is actually valid when X is a general compact metric space) :

Theorem D.3.1. *Suppose that X is a compact subset of \mathbb{R}^p and that $g : X \rightarrow X$ is continuous and onto. The backward dynamics on X generated by $x_t = g(x_{t+1})$ displays topological chaos in the sense of Devaney if and only if the forward dynamics generated by the multivalued inverse forward difference equation $x_{t+1} \in \varphi(x_t)$ also exhibits topological chaos in the sense of Devaney.*

Remarkably, the above property is valid for g and the inverse correspondence φ quite independently of the inverse limit formalism. On the other hand, one can also transpose directly the definition of topological chaos in the sense of Devaney for the left shift map σ acting on the inverse limit space Z . Here we get (Li (1992, Theorem C, p. 97, which is also valid when X is compact metric space) :

Corollary D.3.2. *Under the assumptions of Theorem D.3.1, assume that the inverse limit space $Z = \varprojlim (X, g)$ is endowed with the metric (D.8). Then the backward map g displays topological chaos in the sense of Devaney on X if and only if the corresponding shift map σ displays topological chaos in the sense of Devaney on the inverse limit space Z .*

Fig. D.9.a illustrates the mechanisms underlying the emergence of topological chaos for the forward multivalued difference equation $x_{t+1} \in \varphi(x_t)$, when the backward map g is acting on an interval $[a, b]$, as for instance in Fig. D.1. Here, g is assumed to be single peaked, with a unique critical point at x^* and to satisfy the equivalent of (D.3) and (D.4), so that $b = g(x^*) > x^* > g^2(x^*) \geq a$. The backward map g has then a unique fixed point \bar{x} in (a, b) , and the key feature of Fig. D.9.a is that $g^3(x^*) = g^2(b) < \bar{x}$. The forward dynamics generated by $x_{t+1} \in \varphi(x_t)$ on the invariant interval $X = [g^2(x^*), b]$ involve the correspondence φ , that has two “branches” in Fig. D.9.a, one corresponding to the part of the graph of g that lies in $[x^*, b]$ and the other associated with $[g^2(x^*), x^*]$. As usual, the interpretation of the economic mechanism that drives the forward dynamics is the agents’ choice at each date of a (deterministic) self-fulfilling expectation. A particular forward trajectory may then be viewed as driven by a sequence of *deterministic “sunspots”* that determine at each date which “branch” of the correspondence φ is selected and thus the corresponding self-fulfilling expectation (“sunspots” were formally introduced in economics by Shell (1977), Cass and Shell (1983)). Fig. D.9.a makes intuitively clear how such deterministic sunspots driven forward dynamics can generate chaotic trajectories, in

particular sensitive dependence on initial conditions, by generating switches between the two branches at irregular, “chaotic” dates. The stationary state \bar{x} is unstable in the backward dynamics g and is actually stable in the forward dynamics whenever the latter is constrained to stay in $[x^*, b]$. Forward trajectories can stay on this “right” branch of φ for long time, get very close to \bar{x} , suddenly jump to the “left” branch at irregular and unpredictable times and go back to the stable “right” branch immediately after. In particular, two trajectories can stay very close in X (or even be identical) on the “right” branch for a long time and separate significantly thereafter with one of the trajectories jumping to the other branch while the other does not. One may note that such sensitive dependence on initial conditions for the forward dynamics is going to occur for a “large” set of initial conditions in X (having a nonempty interior), whereas this will not typically be the case for the backward dynamics whenever the map g is S -unimodal as in Theorem D.1.4. Early economic studies of such deterministic sunspot driven sensitive dependence on initial conditions can be found in De Vilder (1995), Hommes and De Vilder (1995). The same deterministic sunspot driven branch switching mechanism in Fig. D.9.a helps in explaining intuitively how the shift map σ acting on the inverse limit space Z with the metric (D.8), can display sensitive dependence on initial conditions, as in Corollary D.3.2. One may note, however, that in such a case, it is unclear whether the set of elements z with sensitive dependence on initial conditions, is “large” or “small” in the infinite dimensional space Z .

Figure D.9.a

Figure D.9.b

Figure D.9.c

Forward stability in the inverse limit space

The above deterministic branch switching mechanism, that appears to be responsible for the emergence of sensitive dependence on initial conditions for the shift map σ acting on the inverse limit space $Z = \varprojlim (X, g)$, seems to suggest that in a situation such as described in Fig. D.9.a, the stationary sequence $(\bar{x}, \bar{x}, \bar{x}, \dots)$ might not be stable in the forward dynamics $\sigma : Z \rightarrow Z$, despite the fact that \bar{x} is a repelling fixed point in the backward dynamics $g : X \rightarrow X$. For simplicity, we state a general “instability” result of this kind for unimodal maps of the interval. Similar results of this sort are in fact valid also for quite general metric spaces X , see Medio and Raines (2007, Theorems 1 and 2).

Theorem D.3.3. *Let the backward map g act from a closed interval $[a, b]$ of the real line into itself, with $a < b$, and be unimodal, i.e. satisfy the analogue of (D.3) and (D.4). Let K be a (backward) periodic orbit for g and \tilde{K} be*




Fig. D.9.a




Fig. D.9.b




Fig. D.9.c

corresponding (forward) periodic orbit in the limit inverse space $Z = \varprojlim ([a, b], g)$.

Let Z be endowed with the metric (D.8). If the image of K by the inverse multivalued correspondence φ differs from K , i.e. $\varphi(K) \neq K$, then \tilde{K} is not asymptotically stable in the forward dynamics generated by the shift map σ on Z .

The same sort of negative result holds not only for periodic orbits but also for closed sets that are g and σ -invariant. The implications for asymptotic stability of the forward dynamics σ on the inverse limit space Z in situations such as those represented in Fig. D.9.a, b, c are then rather neat. Assume that in all three cases, the backward map g is S -unimodal, i.e. satisfies the analogue of assumptions (D.3), (D.4), (D.5) and (D.6), so that it is single peaked and has a negative Schwarzian derivative.

In the case of Fig. D.9.a, where $g^3(x^*) < \bar{x}$, one can focus on the invariant interval $X = [g^2(x^*), b]$. Assume further that g has infinitely many cycles (because for instance $g^3(x^*) < x^*$), in which case g has a cycle of period 3), and that one of these is g -stable (the generic case from a topological viewpoint, according to 1) of Theorem D.1.9). From Theorem D.1.4, all the others are then g -unstable. Theorem D.3.3 above implies nevertheless that *none of these g -unstable periodic orbits will be asymptotically stable in the forward dynamics generated by the shift map σ on the inverse limit space $\varprojlim (X, g)$* . The reason is again branch switching : any periodic orbit must have an element on the right branch of the graph of g in Fig. D.9.a, so that the image by the inverse correspondence φ of that periodic orbit is multivalued and is thus bound to differ from P .

In the case of Fig. D.9.b, where $g^3(x^*) > \bar{x}$, this branch switching along forward trajectories generated by $x_{t+1} \in \varphi(x_t)$ no longer occurs for a significant part of the right branch of the graph of g for x in $[x^*, b]$, that includes the g -unstable stationary state \bar{x} . One might expect then the stationary state $(\bar{x}, \bar{x}, \dots)$ to be stable in the forward dynamics associated with the shift map σ acting on the inverse limit space $Z = \varprojlim (X, g)$, where X is again taken to be the invariant interval $[g^2(x^*), b]$. In fact, the *stationary state can then be shown to be the only asymptotically stable, “topological attractor” (i.e. that attracts “most” initial conditions from a topological viewpoint) for the shift map σ in the inverse limit space $Z = \varprojlim ([g^2(x^*), b], g)$ (Medio and Raines (2007, Theorem 4))*.

Finally, the third case described in Fig. D.9.c differs from that of Fig. D.9.b by specifying $g(x)$ for x in $[a, g^2(x^*)]$ with $g(a) = a$ and $g'(a) > 1$: the

end point a is a fixed point of g , that is unstable in the backward dynamics g (of course if one restricts forward orbits to lie in $[g^2(x^*), b]$, one gets back to the case represented in Fig. D.9.b). The main observation here is that there is no longer any possibility of switching between the two branches of the graph of g along forward trajectories when they get close to the g -unstable fixed point a . Indeed $\varphi(x)$ is single valued when x lies in $[a, g^2(x^*))$ and forward trajectories that hit that interval get trapped there and must converge to a . It can be in fact shown that this is the “typical” fate of forward trajectories, so that (a, a, a, \dots) is the only asymptotically stable, “topological attractor” for the forward shift map σ acting on the inverse limit space $Z = \varprojlim ([a, b], g)$ (see Medio and Raines (2007, Theorem 3)).

Forward local stability

The above results intend to provide a neat formal picture of the asymptotic properties of forward trajectories. Apart from Theorem D.3.1, they use the inverse limit formalism, by focusing on the topological or metric properties of the orbits generated by the shift map σ on the inverse limit space $Z = \varprojlim (X, g)$. A crucial ingredient of the analysis is the metric introduced by the distance (D.8). Such a metric exhibits a special feature, namely that two forward trajectories (x_0, x_1, x_2, \dots) and (y_0, y_1, y_2, \dots) are considered as close to each other even if they differ significantly asymptotically as time goes to infinity, provided that they stay very close to each other for a sufficiently long but finite time. This type of metric seems to be well adapted to account for the potential instability of forward trajectories associated with *global switching* among different branches of the multivalued correspondence φ governing the forward dynamics, through $x_{t+1} \in \varphi(x_t)$. The induced topological properties of the shift map σ , such as sensitive dependence on initial conditions (as in Corollary D.3.2), or the forward σ -instability of a backward g -unstable periodic orbit (as in Theorem D.3.3 and the stationary state \bar{x} in Fig. D.9.a and c) seem to reflect well this potential global branch switching instability mechanism.

Care should nevertheless be exercised by looking at the robustness of the outcomes of the inverse limit formalism with respect to the topology or the metric employed, especially when dealing with infinite dimensional spaces as here (for an April 1st joke on this issue, see Grandmont, Kirman and Neufind (1974)). Indeed very different results would be obtained if the inverse space Z was endowed with the alternative distance defined by, for two forward trajectories $\mathbf{x} = (x_0, x_1, x_2, \dots)$ and $\mathbf{y} = (y_0, y_1, y_2, \dots)$ in Z :

$$(D.9) \quad d^*(\mathbf{x}, \mathbf{y}) = \text{Sup}_{t \geq 0} \|x_t, y_t\|$$

where $\|x_t, y_t\|$ is the Euclidean distance in \mathbb{R}^p . Two forward trajectories in Z are considered close in that case if and only if they stay close to each other *at all dates* $t \geq 0$. As a consequence, while the multivalued forward difference equation $x_{t+1} \in \varphi(x_t)$ can exhibit sensitive dependence on initial conditions (as in Theorem D.3.1), the forward shift map σ cannot when the inverse limit space Z is equipped with the “sup-metric” (D.9) (two forward trajectories that are close must stay close after the repeated application of the shift map), while it does when Z is endowed with the distance (D.8). So on this count, the distance (D.8) appears to be more adapted than (D.9) to deal with the presence of topological chaos due to global switching among different branches of the forward correspondence φ .

On the other hand, with the distance (D.9), the focus is on uniform closeness of forward trajectories, so the local forward dynamics generated by the shift map σ on the inverse limit space Z near a (hyperbolic) periodic orbit of period k , are then essentially equivalent to the local dynamics generated by the local inverse g^{-1} near that periodic orbit, or the local inverse of the appropriate iterate g^k near a corresponding periodic point. So with this distance (D.9), a hyperbolic g -unstable periodic orbit in the factor space X will be σ -asymptotically stable in the inverse limit space. Similar statements are valid more generally for hyperbolic g -unstable and σ -asymptotically stable closed invariant sets. Here, while the distance (D.8) appears to be more adapted to the detection of instability mechanisms associated with *global* switching among different branches of the forward multivalued correspondence φ (as documented in the discussion above of Fig. D.9.a and c), the distance (D.9) seems to be better suited to the detection of multiple asymptotically stable *local* attractors and the corresponding dynamic complexities. For instance, let the backward map g act on a closed interval $[a, b]$, be S -unimodal as in Theorem D.1.4, and have a cycle of period 3, hence cycles of all periods by Sarkovskii’s Theorem. From 1) of Theorem D.1.9, we know that the generic situation is that all but one of these cycles are unstable in the backward dynamics g . The argument above shows that with the distance (D.9) and the implied emphasis on *local* forward dynamics, all these g -unstable cycles in $[a, b]$ should be σ -asymptotically stable in Z . This is indeed a non-trivial multiplicity that involves interesting dynamic complexities as well, and that would be absent from the picture one would get from the exclusive use of the distance (D.8), as our discussion of Fig. D.9.a shows.

At any rate, the information that a hyperbolic periodic orbit is g -unstable in the factor space X and thus asymptotically stable in the local forward dynamics obtained by inverting locally the backward map g near the periodic orbit, has significant implications for economic modeling, beyond nonlinear

dynamics technicalities *stricto sensu*. Such a configuration is indeed a particular case of what is known in economics as *local indeterminacy*, i.e. a situation in which *the number of stable local eigenvalues in the local forward dynamics near the periodic orbit under consideration exceeds the number of predetermined variables inherited at each date from the past*. In such a case, economic agents have some degrees of freedom (the number of stable local eigenvalues of the local forward dynamics in X , in excess of the number of predetermined variables) in formulating their expectations while preserving perfect foresight and local stability conditions. It can be shown that *such local indeterminacy implies the existence of infinitely many stochastic stationary “sunspot” equilibria arbitrarily near the periodic orbit, along which expectations are driven by random shocks while being still self-fulfilling* (Azariadis (1981), Woodford (1986), Grandmont, Pintus and De Vilder (1998)). Such multiple equilibria display indeed dynamic complexities of very high magnitude by comparison even to chaotic deterministic systems, as each of these involve truly random processes.

Notes on the literature

The presentation of the global bifurcations of maps of the real line, in section D.1, is adapted from a previous survey, Grandmont (1986), which was itself derived from Collet and Eckmann (1980). See also Guckenheimer and Holmes (1983, Chaps. 5.6 and 6.3), May (1976), Misiurewicz (1983), De Melo and van Strien (1993). The discussion of the abundance of stable periodic behaviour versus ergodic chaos leading to Theorem D.1.9 is adapted from Jakobson and Swiatek (2002).

Section D.2 on the horseshoe map and the homoclinic theorem is adapted from the very clear exposition of Lanford (1983), and from Guckenheimer and Holmes (1983, Chap. 5.1 and Theorem 5.3.5) ; see also Smale (1980).

The use of symbolic dynamics to describe complex orbits is pervasive in this context. For further information and references on this question, the reader may consult Guckenheimer and Holmes (1983, Chap. 5).

The discussion of the abundance of ergodic chaos for families of Hénon-like diffeomorphisms in Remark D.2.5 is adapted from Wilkinson (2008, section 8) and Benedicks (2003).

The presentation of the “Inverse Limit” approach in section D.3 is adapted from Kennedy and Stockman (2008) and from Medio and Raines (2007), which the reader may usefully consult for further information.

ACKNOWLEDGEMENTS

The material of these lecture notes was presented initially as a short graduate course at the Summer Workshop in Economic Theory at Stanford University in the summer of 1987. I am greatly indebted to Mordecai Kurz and Kenneth Arrow who made this possible. Comments and suggestions from the participants are gratefully acknowledged. This work was supported by a grant of the National Science Foundation at the Institute for Mathematical Studies in the Social Sciences (IMSSS) of Stanford University. Financial support from the Suntory Toyota International Centre for Economics and Related Disciplines (STICERD) at the London School of Economics in 1987 is also gratefully acknowledged.

I greatly benefited from the kind advice and outstanding expertise of Alain Chenciner, in particular from a two days course he gave on the Hopf bifurcation at the Groupement de Recherche en Economie Quantitative et Econométrie (GREQAM), Ecole des Hautes Etudes en Sciences Sociales (EHESS) in Marseille in the fall of 1986. I wish to thank him very much here.

The lecture notes have been circulated first as a Stanford Technical Report and as a CEPREMAP Working Paper in 1988. I have taught this material over the years as part of Topics Ph.D. courses in various places, in particular at the Universities of Yale, Bonn, Louvain la Neuve (CORE), Venice, Taipei (Academia Sinica), Kyoto (KIER), Alicante, Marseille (GREQAM), Tokyo (Keio University), Kobe (RIEB). The current revision of the lecture notes, made with the prospect of a publication in *Research in Economics*, owes much to the comments and suggestions of many students and colleagues : they are gratefully acknowledged. I had in particular useful exchanges with Michael Jakobson on the abundance of ergodic chaos in one-parameter families of S -unimodal maps. Finally, I am greatly indebted to the efficient typing of Ms. Josselyne BITAN and Nadine GUEDJ and of secretaries of the Printing Office of Ecole Polytechnique. I am also very grateful to Nicolas Dromel, who very ably put my handwritten Figures into electronic form.

The english version of the lecture notes will be translated into japanese with in view a publication by Chisenshokan co., Ltd.

REFERENCES

- Azariadis, C. (1981), "Self-Fulfilling Prophecies", *Journal of Economic Theory*, 25, 380-396.
- Banks, J., Brooks, J., Cairns, G., Davis, G. and P. Stacey (1992), "On Devaney's Definition of Chaos", *Trans. of the Amer. Math. Soc.* , 99, 332-334.
- Benedicks, M. (2003), "Non Uniformly Hyperbolic Dynamics : Hénon Maps and Related Dynamical Systems", in Li, Ta-tsien (ed.), *Proceedings of the International Congress of Mathematicians, Beijing 2002*, Higher Education Press, Beijing, Vol. III, 265-278.
- Benedicks, M. and L. Carleson (1991), "The Dynamics of the Hénon Map", *Ann. of Math.*, 133, 73-169.
- Benedicks, M. and L.-S. Young (1992), "Absolutely Continuous Invariant Measures and Random Perturbations for Certain One-Dimensional Maps", *Ergodic Theory and Dynamical Systems*, 12, 13-37.
- Benedicks, M. and L.-S. Young (1993), "Sinai-Bowen-Ruelle Measures for Certain Hénon Maps", *Invent. Math.*, 112, 541-576.
- Benhabib, J. and R. Day (1982), "A Characterization of Erratic Dynamics in the Overlapping Generations Model", *Journal of Economic Dynamics and Control*, 4, 37-55.
- Bruin, H., Keller, G., Nowicki, T. and S. van Strien (1996), "Wild Cantor Attractors Exist", *Ann. of Math.*, 143, 97-130.
- Carr, J. (1981), *Applications of Center Manifold Theory*, Springer Verlag, New York.
- Cass, D. and K. Shell (1983), "Do Sunspots Matter ?", *Journal of Political Economy*, 91, 193-227.

- Chenciner, A. (1983), “Bifurcations de Diffeomorphismes de R^2 au Voisinage d’un Point Fixe Elliptique”, in Iooss, G., Helleman, R.H.G. and R. Stora (eds.), 1983.
- Chenciner, A. (1985a), “Bifurcations de Points Fixes Elliptiques. I. Courbes Invariantes”, *IHES-Publications Mathématiques*, 61, 67-127.
- Chenciner, A. (1985b), “Bifurcations de Points Fixes Elliptiques. II. Orbites Périodiques et Ensembles de Cantor Invariants”, *Inventiones Mathematicae*, 80, 81-106.
- Chenciner, A. (1988), “Bifurcations de Points Fixes Elliptiques. III. Orbites Périodiques de ”Petites” Périodes et Elimination Résonnante des Couples de Courbes Invariantes”, *IHES-Publications Mathématiques*, 66, 5-91.
- Collet, P. and J.P. Eckmann (1980), *Iterated Maps on the Interval as Dynamical Systems*, Birkhäuser, Boston.
- Devaney, R.L. (2003), *An Introduction to Chaotic Dynamical Systems*, 2nd ed., Westview Press, Boulder.
- Gaunersdorfer, A., Hommes, C.H. and F.O.O. Wagener (2006), “Bifurcation Routes to Volatility Clustering under Evolutionary Learning”, mimeo, CNDEF, University of Amsterdam, forthcoming in *Journal of Economic Behavior and Organization*.
- Grandmont, J.-M. (1985), “On Endogenous Competitive Business Cycles”, *Econometrica*, 53, 995-1045.
- Grandmont, J.-M. (1986), “Periodic and Aperiodic Behaviour in Discrete One-Dimensional Dynamical Systems”, in Hildenbrand, W. and A. Mas-Colell (eds.), 1986.
- Grandmont, J.-M., Kirman, A.P. and W. Neuefind (1974), “A New Approach to the Uniqueness of Equilibrium”, *Review of Economic Studies*, 41, 289-291.
- Grandmont, J.-M., Pintus, P. and R. de Vilder (1998), “Capital-Labor Sub-

- stitution and Competitive Nonlinear Endogenous Business Cycles”, *Journal of Economic Theory*, 80, 14-59.
- Guckenheimer, J. and P. Holmes (1983), *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer Verlag, New York.
- Hartman, P. (1964), *Ordinary Differential Equations*, Wiley, New York.
- Hildenbrand, W. and A. Mas-Colell (1986), *Contributions to Mathematical Economics, In Honour of Gerard Debreu*, North-Holland, New York.
- Hirsch, M.W. and S. Smale (1974), *Differential Equations, Dynamical Systems, and Linear Algebra*, Academic Press, New York.
- Hommel, C. and R. de Vilder (1995), “Sunspot Equilibria in an Implicitly Defined Overlapping Generations Model”, mimeo, University of Amsterdam.
- Iooss, G. (1979), *Bifurcations of Maps and Applications*, North-Holland, New York.
- Iooss, G., Helleman, R.H.G. and R. Stora (eds.) (1983), *Chaotic Behaviour of Deterministic Systems*, North-Holland, New York.
- Jakobson, M.V. (1981), “Absolutely Continuous Invariant Measures for One-Parameter Families of One-Dimensional Maps”, *Comm. Math. Phys.*, 81, 39-88.
- Jakobson, M.V. and G. Swiatek (2002), “One-Dimensional Maps”, in *Handbook of Dynamical Systems*, Hasselblatt, B. and A. Katok (eds.), Elsevier, Amsterdam, Vol. 1, Part 1, Chap. 10, 599-664.
- Johnson, S. (1985), “Singular Measures without Restrictive Intervals”, *Commun. Math. Phys.*, 110, 185-190.
- Kennedy, J.A. and D.R. Stockman (2008), “Chaotic Equilibria in Models with Backward Dynamics”, *Journal of Economic Dynamics and Control*, 32, 939-955.
- Kozlovski, O. S. (1998), *Structural Stability in One-Dimensional Dynamics*,

- Ph. D. Thesis, University of Amsterdam.
- Kozlovski, O. S. (2000), “Getting Rid of the Negative Schwarzian Derivative Condition”, *Annals of Math.*, *152*, 743-762.
- Kuznetsov, Y.A. (1998), *Elements of Applied Bifurcation Theory*, 2nd ed., Springer Verlag, New York.
- Lanford III, O.E. (1983), “Introduction to the Mathematical Theory of Dynamical Systems”, in Iooss, G., Helleman, R.H.G. and R. Stora (eds.), 1983.
- Li, S.H. (1992), “Dynamical Properties of the Shift Maps on the Inverse Limit Spaces”, *Ergodic Theory and Dynamical Systems*, *12*, 95-108.
- Li, T. and J.A. Yorke (1975), “Period Three Implies Chaos”, *American Mathematical Monthly*, *82*, 985-992.
- Marsden, J.E. and M. McCracken (1976), *The Hopf Bifurcation and its Applications*, Springer Verlag, New York.
- May, R.B. (1976), “Simple Mathematical Models with Very Complicated Dynamics”, *Nature*, *261*, 459-467.
- Medio, A. and B. Raines (2007), “Backward Dynamics in Economics. The Inverse Limit Approach”, *Journal of Economic Dynamics and Control*, *31*, 1633-1671.
- De Melo, W. and S. van Strien (1988), “One-Dimensional Dynamics : the Schwarzian Derivative and Beyond”, *Bull. of the American Math. Society*, *18*, 159-162.
- De Melo, W. and S. van Strien (1993), *One-Dimensional Dynamics*, Springer Verlag, New York.
- Misiurewicz, M. (1981), “Absolutely Continuous Measures for Certain Maps of an Interval”, *Publications Mathématiques de l’IHES*, *53*, 17-51.
- Misiurewicz, M. (1983), “Maps of an Interval”, in Iooss, G., Helleman, R.H.G. and R. Stora (eds.), 1983.

- Nusse, H.E. (1886), “Persistence of Order and Structure in Chaos”, *Physica D*, 20, 374-386.
- Palis, Jr., J. and W. De Melo (1982), *Geometric Theory of Dynamical Systems*, Springer Verlag, New York.
- Palis, Jr., J. and F. Takens (1977), “Topological Equivalence of Normally Hyperbolic Dynamical Systems”, *Topology*, 16, 335-345.
- Parthasarathy, K.R. (1967), *Probability Measures on Metric Spaces*, Academic Press, New York.
- Sarkovskii, A.N. (1964), “Coexistence of Cycles of a Continuous Map of the Line into Itself”, *Ukr. Mat. Z.*, 16, 61-71.
- Shell, K. (1977), “Monnaie et Allocation Intertemporelle”, mimeo, Séminaire d’Econométrie Roy-Malinvaud, CNRS, Paris. Title and abstract in French, text in English.
- Singer, D. (1978), “Stable Orbits and Bifurcations of Maps of the Interval”, *SIAM J. Applied Math.*, 35, 260.
- Smale, S. (1980), *The Mathematics of Time*, Springer Verlag, New York.
- Thieullen, Ph. Tresser, C. and L. S. Young (1994), “Positive Lyapounov Exponent for Generic One-Parameter Families of Unimodal Maps”, *J. Anal. Math.*, 64, 121-172.
- De Vilder, R. (1995), *Endogenous Business Cycles*, Ph. D. dissertation, Tinbergen Institute Research Series N° 96, University of Amsterdam.
- Whitley, D. (1983), “Discrete Dynamical Systems in Dimensions One and Two”, *Bull. London Math. Soc.*, 15, 177-217.
- Wilkinson, A. (2008), “Smooth Ergodic Theory”, mimeo, Northwestern University, forthcoming in *Encyclopedia of Complexity and Systems Science*, R. Meyers (ed.), Springer Verlag, New York.
- Woodford, M. (1986), “Stationary Sunspot Equilibria in a Finance Constrained

Economy”, *Journal of Economic Theory*, 40, 128-137.