

2009/61



On marginal likelihood computation
in change-point models

Luc Bauwens and Jeroen V.K. Rombouts



CORE

The logo for CORE (Center for Operations Research and Econometrics) features the word "CORE" in a bold, black, sans-serif font. A thin, light blue arc curves over the letters, starting from the top left of the 'C' and ending at the bottom right of the 'E'.

DISCUSSION PAPER

Center for Operations Research
and Econometrics

Voie du Roman Pays, 34
B-1348 Louvain-la-Neuve
Belgium

<http://www.uclouvain.be/core>

CORE DISCUSSION PAPER
2009/61

**On marginal likelihood computation
in change-point models**

Luc BAUWENS¹ and Jeroen V.K. ROMBOUTS²

October 2009

Abstract

Change-point models are useful for modeling time series subject to structural breaks. For interpretation and forecasting, it is essential to estimate correctly the number of change points in this class of models. In Bayesian inference, the number of change points is typically chosen by the marginal likelihood criterion, computed by Chib's method. This method requires to select a value in the parameter space at which the computation is done. We explain in detail how to perform Bayesian inference for a change-point dynamic regression model and how to compute its marginal likelihood. Motivated by our results from three empirical illustrations, a simulation study shows that Chib's method is robust with respect to the choice of the parameter value used in the computations, among posterior mean, mode and quartiles. Furthermore, the performance of the Bayesian information criterion, which is based on maximum likelihood estimates, in selecting the correct model is comparable to that of the marginal likelihood.

Keywords: BIC, change-point model, Chib's method, marginal likelihood.

JEL Classification: C11, C22, C53

¹ Université catholique de Louvain, CORE, B-1348 Louvain-la-Neuve, Belgium.

E-mail: luc.bauwens@uclouvain.be. This author is also member of ECORE, the association between CORE and ECARES.

² Institute of Applied Economics at HEC Montréal, CIRANO, CIRPEE, and CORE (Université catholique de Louvain).

Research supported by the contract "Projet d'Actions de Recherche Concertées" 07/12-002 of the "Communauté française de Belgique", granted by the "Académie universitaire Louvain".

This paper presents research results of the Belgian Program on Interuniversity Poles of Attraction initiated by the Belgian State, Prime Minister's Office, Science Policy Programming. The scientific responsibility is assumed by the authors.

1 Introduction

Economic and financial time series are subject to changes in their pattern over long periods, see e.g. Stock and Watson (1996) for US macroeconomic data, Pastor and Stambaugh (2001) and Liu and Maheu (2008) for financial series. It is therefore interesting to take into account the possibility of structural change in time series models, both for interpreting historical data and for forecasting future values. Researchers do not usually assume that break dates are known. Models allowing for the possibility of changing structure or parameters have thus been developed over the last twenty years. In particular, the change in model parameters can be modelled by a Markov-switching discrete process, following the impetus given by Hamilton (1989). The states of this process, which correspond to the parameter values, are recurrent as the process can move from one state to any other state at any date. A particular case of this model is the change-point model that has non-recurrent states: the process can only stay in the same state or move to the next one. An important and difficult issue is the choice of the number of states of the hidden Markov chain, since this determines the number of structural breaks in the time series. For this issue, Bayesian inference is useful: one can estimate the model for a range of values of the number of states and choose the model that delivers the highest marginal likelihood. When maximum likelihood estimation is feasible, one can likewise choose the model according to the Bayesian (or Schwarz) information criterion (BIC).

The computation of the marginal likelihood of a given Markov-switching model can be performed by the method of Chib (1995). This method only uses the Gibbs output, while other methods like those Newton and Raftery (1994) and Gelfand and Dey (1994) are either unstable or need an additional tuning function. Chib's method is based on the marginal likelihood identity, namely that the marginal likelihood is equal to the product of the likelihood and the prior divided by the posterior. Since each element on the right hand side of this identity depends on the model parameters, a particular value must be chosen, even if the result does not depend in principle on this value. Chib recommends to choose a high posterior density value for numerical efficiency reasons. Many researchers use the posterior mean, see e.g. Kim and Nelson (1999), Elerian, Chib, and Shephard (2001), Kim and Piger (2002), Kim, Morley, and Nelson (2005), Pesaran, Pettenuzzo, and Timmermann (2006), Johnson and Sakoulis (2008), Liu and Maheu (2008), Maheu and Gordon (2008), Fruhwirth-Schnatter and Wagner

(2008), Nakajima and Omori (2009), and few, such as Paroli and Spezia (2008), use the mode. To the best of our knowledge, no study has clearly documented the sensitivity of the marginal likelihood value obtained by Chib's method to the value of the parameter chosen for its computation in the case of Markov-switching models.

The first goal in this paper is precisely to assess this sensitivity in the case of a change-point regression model. As mentioned above, change-point models are a particular case of Markov-switching models. In the general case of recurrent states, the number of parameters of the transition matrix of the Markov chain is equal to $K^2 - K$, where K is the number of states. A change-point model is much more parsimonious in this dimension since it requires only $K - 1$ parameters. Even if a change-point model may require a higher K than a general Markov-switching one, it is not likely that the number of change points will be as high as $K^2 - K$. Another advantage of change-point models is the fact that they are not subject to the label switching problem of general Markov-switching models, which renders estimation much more difficult.

Based on a simulation study, our conclusion is that Chib's method is not much sensitive to the chosen value of the parameter for computing the marginal likelihood, and we recommend to use indifferently the posterior mode, mean or median rather than other values. Our simulation study is based on three change-point regression models whose specification and parameter values are closely inspired by models that we estimated using real time series.

Moreover, for models of the type we consider, Chib's method uses, as always, the output of the Gibbs sampler, but requires additional simulations due to the presence of the latent variables. This explains, at least partially, why computations are typically heavy. The second objective of this paper is to compare the model choices resulting from the application of the BIC and of the marginal likelihood criterion. Computing the BIC requires of course to maximize the log-likelihood function, which is not an easy task when the number of change points is large, the reason being the existence of multiple local maxima. The simulation results show that the BIC leads to choose the correct model in a higher proportion of replications than the marginal likelihood criterion, and that a correct computation of the BIC is not necessarily less heavy than for the marginal likelihood.

The paper is structured as follows: in Section 2, we present change-point models and how Bayesian inference is done. This is done in detail so as to make the presentation, done in

Section 3, of the computation of the marginal likelihood and BIC understandable. Section 4 contains the empirical examples and Section 5 the simulation results. Section 6 concludes.

2 Model and inference

In Section 2.1 we explain how breaks can be introduced in a time series model, drawing on the work of Chib (1998) and Pesaran, Pettenuzzo, and Timmermann (2006). In Section 2.2, we define the prior densities, and in Section 2.3, we explain how to compute the posterior density of the parameters of this type of model by Gibbs sampling.

Concerning notations, we adopt some conventions: by default vectors are columns and y' denotes the transpose of y . Data densities (including predictive) for observable or latent variables, whether discrete or continuous, are denoted by $f(\cdot)$. Prior and posterior densities are denoted by $\varphi(\cdot)$. Parameters of prior densities are identified by underscore bars (e.g. \underline{a}), those of posterior densities by overscore bars (e.g. \bar{a}).

2.1 Model

Let y_t be the time series we want to model over the sample period $\{1, 2, \dots, T\}$. Let s_t be an integer latent variable (also called state variable) taking its value in the set $\{1, 2, \dots, K\}$, K being assumed known. The state variable s_t indicates the active regime generating y_t at period t in the sense that y_t is generated from the data density

$$f(y_t | Y_{t-1}, \theta_{s_t}), \tag{1}$$

where $Y_{t-1} = (Y_0' \ y_1 \ \dots \ y_{t-1})'$, Y_0 being a vector of known initial conditions (observations prior to date $t = 1$), and θ_{s_t} is a vector of parameters indexing the density. There are potentially K regimes, hence K parameter vectors $\theta_1, \theta_2, \dots, \theta_K$.

We assume that the active regime at t is selected by a discrete first-order Markov process for the s_t process. As in Chib (1998), the transition probability matrix P_K (more simply denoted by P when indicating the number of regimes is not important) allows either to stay in the regime operating at $t - 1$ or to switch to the next regime and has therefore the following

structure:

$$P_K = \begin{pmatrix} p_{11} & 1 - p_{11} & 0 & \dots & 0 & 0 \\ 0 & p_{22} & 1 - p_{22} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & p_{K-1,K-1} & 1 - p_{K-1,K-1} \\ 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}. \quad (2)$$

Notice that the last regime is an absorbing state over the sample period. Given the zero entries in P , the discrete Markov chain generates potentially $K - 1$ breaks, at random dates τ_k ($k = 1, 2, \dots, K - 1$) defined by τ_k being the integer in $\{1, 2, \dots, T\}$ such that $s_{\tau_k} = k$ and $s_{\tau_{k+1}} = k + 1$. A posterior density on these dates is therefore a direct by-product of the inference on the state variables. A convenient prior density, based on the assumption of independence between the parameters of the matrix P , takes the form of a product of identical beta densities with parameters \underline{a} and \underline{b} :

$$\varphi(p_{11}, p_{22}, \dots, p_{K-1,K-1}) \propto \prod_{i=1}^{K-1} p_{ii}^{\underline{a}-1} (1 - p_{ii})^{\underline{b}-1}. \quad (3)$$

The assumption that the beta densities are identical can be easily relaxed. This prior implies that the (strictly positive integer) duration of regime k , $d_k = \tau_k - \tau_{k-1}$ (setting $\tau_0 = 0$ and $\tau_K = T$) is approximately geometrically distributed with parameter p_{ii} and expected value $(\underline{a} + \underline{b})/\underline{a}$. For example, by fixing $\underline{a} = \underline{b} = 1$, the prior is uniform for each probability. Actually, fixing $\underline{a} = \underline{b}$ implies that $E(d_k) = 2$ and a priori, on average, a lot of regimes. To choose a prior on the probabilities consistent with K , T/K should be the expected number of observations per regime. Hence one should fix $E(p_{ii}) = 1 - (K/T)$, which implies that $\underline{a}/\underline{b} = K/(T - K)$ and fixes one of the parameters given the other. The other parameter can be deduced by fixing the variance of the beta distribution and solving.¹

Following Pesaran, Pettenuzzo, and Timmermann (2006), an essential ingredient of the model specification is the prior assumption that the parameter vectors θ_i are drawn independently from a common distribution, i.e. $\theta_i \sim \varphi(\theta_i|\theta_0)$ where θ_0 itself is a parameter vector

¹If one substitutes $\underline{b}K/(T - K)$ for \underline{a} in $v = \underline{a}\underline{b}/[(\underline{a} + \underline{b})^2(\underline{a} + \underline{b} + 1)]$ where v is the assigned prior variance, one obtains a second degree equation in \underline{b} and a positive solution can be found under some conditions.

endowed with a prior density $\varphi(\theta_0|\underline{A})$, \underline{A} denoting the prior hyper-parameters. This is called a hierarchical prior or a meta-distribution. For example, if θ_i contains location parameters and a scale one, the prior can be a normal density on the location parameters and a gamma density on the scale one. Generally, the joint prior on the θ parameters is

$$\varphi(\theta_0, \theta_1, \theta_2, \dots, \theta_K) = \varphi(\theta_0|\underline{A}) \prod_{i=1}^K \varphi(\theta_i|\theta_0). \quad (4)$$

Behind the common prior (4) lies the belief that the regime parameters differ and evolve independently of each other. Another possible prior belief is that the regime parameters evolve in a more structured way. For example the conditional mean of y_t could be increasing ($\mu_{k-1} < \mu_k$). This idea can be formalized through a joint normal prior on $(\mu_2 - \mu_1, \mu_3 - \mu_2, \dots, \mu_K - \mu_{K-1})'$ with mean vector m_0 and covariance matrix V_0 , where m_0 and V_0 are the hyper-parameters to be endowed with a prior density implying that m_0 is positive with high probability.

The model is fully specified by defining the conditional densities $f(y_t|Y_{t-1}, \theta_{s_t})$. In this study, we use the following assumption:

$$y_t|Y_{t-1}, \theta_{s_t} \sim N(x_t' \beta_{s_t}, \sigma_{s_t}^2), \quad (5)$$

where x_t is a vector of m exogenous variables and β_{s_t} a vector of coefficients, so that $\theta_{s_t} = (\beta_{s_t}' \sigma_{s_t}^2)'$. In our applications, the model within a regime is autoregressive, i.e. x_t includes the constant 1 and p lags of y_t , hence $m = p + 1$.

2.2 Prior densities for location and scale parameters

In the model developed in the previous sub-section, the hierarchical prior (4) is defined by a prior for each β_j ($j = 1, 2, \dots, K$) and an independent prior for each σ_j^2 , given additional random parameters (forming θ_0) defined below, to which prior densities are assigned. The hierarchy can be divided into two independent pieces, one for the regression coefficients and one for the variances. We define the two pieces, provide the formulas of some moments, and discuss how to define weakly informative (proper) prior densities. Improper prior densities are excluded as we want to compute the marginal likelihood.

Regression coefficients

The hierarchical prior is

$$\beta_j | b_0, B_0 \sim N_m(b_0, B_0), \quad (6)$$

$$b_0 \sim N_m(\underline{\mu}_\beta, \underline{\Sigma}_\beta), \quad (7)$$

$$B_0^{-1} \sim W_m(\underline{\nu}_\beta, \underline{V}_\beta^{-1}), \quad (8)$$

with b_0 independent of B_0 . $W_m(\nu, S)$ denotes a Wishart density of dimension m with scale parameter S , a positive-definite symmetric matrix, and ν degrees of freedom ($\nu > m - 1$). If $m = 1$, this reduces to a $\text{Gamma}(\nu, s)$ defined below, see (14). Equivalently, B_0 has an inverted Wishart density, $B_0 \sim IW_m(\nu, S)$, see Appendix A of Bauwens, Lubrano, and Richard (1999), with $E(B_0) = \underline{V}_\beta^{-1} / (\underline{\nu}_\beta - m - 1)$ if $\underline{\nu}_\beta > m + 1$. Although the prior marginal density of β_j is not known analytically, its moments can be obtained by applying the law of iterated expectations:

$$E(\beta_j) = \underline{\mu}_\beta, \quad (9)$$

$$\text{Var}(\beta_j) = E(B_0) + \text{Var}(b_0) = \frac{\underline{V}_\beta^{-1}}{\underline{\nu}_\beta - m - 1} + \underline{\Sigma}_\beta \text{ if } \underline{\nu}_\beta > m + 1. \quad (10)$$

To be weakly informative on the β_j parameters, one usually sets B_0^{-1} close to 0 and diagonal in (6), meaning the variances are large and the elements are independent, and $b_0 = 0$, though taking another value does not matter if B_0^{-1} close to 0. In our hierarchical setup one can set $\underline{\mu}_\beta = 0$ instead of $b_0 = 0$. In view of (10), one can achieve large variances through $\text{Var}(b_0)$ or $E(B_0)$. It is more direct and easier for understanding the prior to make $\text{Var}(b_0)$ large, rather than $E(B_0)$. For $\text{Var}(b_0)$, this means setting $\underline{\Sigma}_\beta = c_1 I_m$ with a large value of c_1 , but what "large" means has to be considered relatively to the order of magnitude of the β_j parameters. We set $\underline{\nu}_\beta = m + 2$ and $\underline{V}_\beta = I_m$, such that $E(B_0) = I_m$.

Variances

The hierarchical prior is defined as

$$\sigma_j^{-2} | v_0, d_0 \sim \text{Gamma}(v_0, d_0), \quad (11)$$

$$v_0 \sim \text{Gamma}(\lambda_0, \rho_0), \quad (12)$$

$$d_0 \sim \text{Gamma}(\underline{c}_0, \underline{d}_0), \quad (13)$$

with v_0 independent of d_0 . We use the following definition of a Gamma(ν, s) density for x where ν is the degrees of freedom and s is the scale parameter:

$$f_G(x|\nu, s) = \left(\frac{s}{2}\right)^{\frac{\nu}{2}} \left[\Gamma\left(\frac{\nu}{2}\right)\right]^{-1} x^{\frac{\nu}{2}-1} \exp\left(-\frac{s}{2}x\right) \text{ for } x > 0. \quad (14)$$

Its expected value is ν/s and its variance is $2\nu/s^2$. Equivalently to (11), and using the terminology of Bauwens, Lubrano, and Richard (1999), σ_j^2 has an inverted gamma-2 density, $\sigma_j^2|v_0, d_0 \sim IG(v_0, d_0)$, with

$$E(\sigma_j^2|v_0, d_0) = d_0/(v_0 - 2) \text{ if } v_0 > 2, \quad (15)$$

$$\text{Var}(\sigma_j^2|v_0, d_0) = \frac{4d_0^2}{(v_0 - 2)^2(v_0 - 4)} \text{ if } v_0 > 4. \quad (16)$$

The prior marginal density of σ_j^{-2} (or its inverse) cannot be computed analytically, but its expectation can:

$$E(\sigma_j^{-2}) = E(v_0)E\left(\frac{1}{d_0}\right) = \frac{\lambda_0}{\rho_0} \frac{\underline{d}_0}{\underline{c}_0 - 2} \text{ if } \underline{c}_0 > 2. \quad (17)$$

Note however that $E(\sigma_j^2)$ does not exist since $v_0 < 2$ has a non-zero probability if $v_0 \sim \text{Gamma}(\underline{\lambda}_0, \underline{\rho}_0)$.

The non-informative version of a prior density on a variance σ_j^2 is usually taken as $\varphi(\sigma_j^2) \propto 1/\sigma_j^2$ or equivalently $\varphi(\sigma_j^{-2}) \propto 1/\sigma_j^{-2}$. This corresponds to setting $v_0 = d_0 = 0$. This approach is not feasible when v_0 and d_0 are random variables. The same type of non-informative prior for v_0 and d_0 is obtained by setting their hyper-parameters at 0. This implies that $E(\sigma_j^{-2})$ does not exist. The full conditional posterior density of d_0 is still a proper density even if $\underline{c}_0 = \underline{d}_0 = 0$ since it is a Gamma($Kv_0 + \underline{c}_0, \sum_{j=1}^K \sigma_j^{-2} + \underline{d}_0$), as shown in the next sub-section. The latter result suggests that \underline{d}_0 should not dominate the sum of the inverted variances if one wishes to be weakly informative on d_0 . Setting the prior degrees of freedom \underline{c}_0 is more difficult since it is added to Kv_0 which is random. The prior mean of Kv_0 , equal to $K\underline{\lambda}_0/\underline{\rho}_0$, can serve as reference value to fix \underline{c}_0 . As PPT, we set $\underline{c}_0 = 1$ and $\underline{d}_0 = 0.01$, hence $E(d_0) = 100$ and $\text{Var}(d_0) = 20000$.

In the next sub-section where the Gibbs algorithm is described, it is shown that the full conditional posterior density of v_0 does not belong to a known class of density functions. Hence it is not possible to compare the hyper-parameters $\underline{\lambda}_0$ and $\underline{\rho}_0$ to their counterparts from the information set. We set $\underline{\lambda}_0 = 1$ and $\underline{\rho}_0 = 0.01$, so that the prior mean and standard deviation of v_0 are both equal to 100, such that our prior densities are weakly informative

2.3 Inference

The joint posterior density of $S_T = (s_1 \ s_2 \ \dots, s_T)'$ and the parameters is proportional to

$$\prod_{t=1}^T f(y_t|Y_{t-1}, \theta_{s_t}) f(s_t|s_{t-1}, P) \prod_{i=1}^{K-1} p_{ii}^{\underline{a}-1} (1 - p_{ii})^{\underline{b}-1} \prod_{i=1}^K \varphi(\theta_i|\theta_0) \varphi(\theta_0|\underline{A}), \quad (18)$$

where $f(s_t|s_{t-1}, P)$ is the transition probability from state $t - 1$ to state t and is one of the non null elements of P . The parameters are θ_i , $i = 0, 1, \dots, K$, jointly denoted by Θ (or Θ_K to emphasize that K regimes are assumed), and the diagonal elements of the matrix P . This density lends itself to simulation by Gibbs sampling in three blocks corresponding to the full conditional densities:

1. $\varphi(S_T|\Theta, P, Y_T) \propto \prod_{t=1}^T f(y_t|Y_{t-1}, \theta_{s_t}) f(s_t|s_{t-1}, P)$,
2. $\varphi(P|S_T) \propto \prod_{t=1}^T f(s_t|s_{t-1}, P) \prod_{i=1}^{K-1} p_{ii}^{\underline{a}-1} (1 - p_{ii})^{\underline{b}-1}$, which does not depend on Θ and Y_T , and
3. $\varphi(\Theta|S_T, Y_T) \propto \prod_{t=1}^T f(y_t|Y_{t-1}, \theta_{s_t}) \prod_{i=1}^K \varphi(\theta_i|\theta_0) \varphi(\theta_0|\underline{A})$, which does not depend on P .

Sampling S_T is done as Chib (1998) and detailed below (notice that $s_T = K$ by assumption). Sampling P is done by simulating each p_{ii} from a beta density with parameters $\underline{a} + T_i$ and $\underline{b} + 1$ where T_i is the number of states equal to i in the sampled S_T vector. Sampling Θ implies usually to break it into sub-blocks and to sample each sub-block given the other, plus S_T . We give all the necessary details below for the case when $f(y_t|Y_{t-1}, \theta_{s_t})$ is the normal density defined in (5).

Chib's algorithm for sampling the states

The algorithm is explained in Chib (1996) for the case when the matrix P is not restricted and in Chib (1998) for the case when P is restricted like in (2), but the differences between the two cases are minor. The state vector S_T is sampled from $\varphi(S_T|\Theta, P, Y_T)$, by sampling sequentially from T to 1 using the backward sequential factorization

$$\varphi(s_{T-1}|s_T, Y_T, \Theta, P) \dots \varphi(s_t|s_{t+1}, Y_T, \Theta, P) \dots \varphi(s_2|s_3, Y_T, \Theta, P) \quad (19)$$

and the fact that the model structure implies that $s_T = K$ and $s_1 = 1$ with probability 1. In the above factorization, $\varphi(s_t|s_{t+1}, Y_T, \Theta, P)$ should in principle be replaced by

$\varphi(s_t|s_{t+1}, s_{t+2}, \dots, s_T, Y_T, \Theta, P)$, but the added conditions s_{t+2}, \dots, s_T are not needed since

$$\varphi(s_t|s_{t+1}, s_{t+2}, \dots, s_T, Y_T, \Theta, P) \propto f(s_t|Y_t, \Theta, P)f(s_{t+1}|s_t, P), \quad (20)$$

which also shows that $\varphi(s_t|s_{t+1}, Y_T, \Theta, P) = \varphi(s_t|s_{t+1}, Y_t, \Theta, P)$. Indeed, as in equation (7) in Chib (1996), and using the notation $S^{t+1} = (s_{t+1} s_{t+2} \dots s_T)$ and similarly for Y^{t+1} ,

$$\varphi(s_t|S^{t+1}, Y_T, \Theta, P) \propto f(s_t|Y_t, \Theta, P)f(Y^{t+1}, S^{t+1}|Y_t, s_t, \Theta, P) \quad (21)$$

$$\propto f(s_t|Y_t, \Theta, P)f(s_{t+1}|s_t, P)f(Y^{t+1}, S^{t+2}|Y_t, s_t, s_{t+1}, \Theta, P) \quad (22)$$

$$\propto f(s_t|Y_t, \Theta, P)f(s_{t+1}|s_t, P). \quad (23)$$

The first line (21) follows from

$$\begin{aligned} \varphi(s_t|S^{t+1}, Y_T, \Theta, P) &\propto f(s_t, S^{t+1}|Y_t, Y^{t+1}, \Theta, P) \\ &\propto f(s_t, S^{t+1}, Y^{t+1}|Y_t, \Theta, P) \\ &\propto f(s_t|Y_t, \Theta, P)f(S^{t+1}, Y^{t+1}|s_t, Y_t, \Theta, P). \end{aligned}$$

For (22),

$$f(Y^{t+1}, S^{t+1}|Y_t, s_t, \Theta, P) = f(s_{t+1}|Y_t, s_t, \Theta, P)f(Y^{t+1}, S^{t+2}|s_{t+1}, Y_t, s_t, \Theta, P)$$

and $f(s_{t+1}|Y_t, s_t, \Theta, P) = f(s_{t+1}|s_t, P)$ from the model assumption about the Markov chain for the states. The third line (23) follows from the fact that the third factor of the rhs of the second line does not depend on s_t .

One starts with generating a random draw from $\varphi(s_{T-1}|s_T = K, Y_{T-1}, \Theta, P)$, then continues with $\varphi(s_{T-2}|s_{T-1}, Y_{T-2}, \Theta, P)$ where s_{T-1} is set equal to the random draw generated from the previous factor, and so on until s_2 is generated. Actually, $\varphi(s_t|s_{t+1}, Y_t, \Theta, P)$ can only take two values given the structure of P in (2), therefore one can just compute these two values according to the rhs of (20) and divide by their sum to obtain normalized probabilities.

In the rhs of (20), $f(s_{t+1}|s_t, P)$ is the same as in (18), and $f(s_t|Y_t, \Theta, P)$ is computed by recurrence from $f(s_{t-1}|Y_{t-1}, \Theta, P)$. One starts at $t = 2$ with initial condition $f(s_1|Y_1, \Theta, P) = 1$ if $s_1 = 1$ and $= 0$ otherwise (at $t = 1$, only state 1 can occur), and one ends at $T - 1$ since $f(s_T|Y_T, \Theta, P) = 1$ if $s_T = K$ and $= 0$ otherwise. The passage from $f(s_{t-1}|Y_{t-1}, \Theta, P)$ to $f(s_t|Y_t, \Theta, P)$ is done in two steps (prediction and updating):

Prediction step: by the law of total probability,

$$f(s_t|Y_{t-1}, \Theta, P) = \sum_{j=1}^K f(s_t|s_{t-1} = j, P) f(s_{t-1} = j|Y_{t-1}, \Theta, P), \quad (24)$$

given that $f(s_t|s_{t-1} = j, P) = f(s_t|s_{t-1} = j, Y_{t-1}, \Theta, P)$ from the assumptions on the model.

Update step: by Bayes theorem,

$$f(s_t|Y_t, \Theta, P) \propto f(s_t|Y_{t-1}, \Theta, P) f(y_t|Y_{t-1}, \Theta_{s_t}), \quad (25)$$

which is easily normalized by dividing by the sum of the rhs over all values of s_t (even if actually the rhs is strictly positive only for two values of s_t).

Gibbs algorithm for Θ

Given (5), $y_t|Y_{t-1} \sim N(x_t'\beta_j, \sigma_j^2)$ when $s_t = j$. The prior densities are given in (6)-(8) and (11)-(13). The parameters are divided into $4 + 2K$ blocks: b_0, B_0, v_0, d_0 (altogether corresponding to θ_0 in (4)), β_j , and σ_j^2 (for each $j = 1, \dots, K$). We provide the conditional posterior density of each block (given the relevant other parameters and the state vector S_T when it is needed). Each of these is obtained by keeping all parts of $\prod_{t=1}^T f(y_t|Y_{t-1}, \theta_{s_t}) \prod_{i=1}^K \varphi(\theta_i|\theta_0) \varphi(\theta_0|\underline{A})$ that depend on the relevant elements. The hyperparameter \underline{A} corresponds to the parameters of (7), (8), (12), and (13). Given S_T , we assign each observation to one of the K regimes and thus we form K vectors y_j and matrices X_j that contain the y_t and x_t' observations assigned to regime j . We denote by T_j the number of observations assigned to regime j . Below we state the result for each block and sketch how we obtain it.

- For each $j = 1, \dots, K$: $\beta_j|\sigma_j^2, b_0, B_0, S_T, Y_T \sim N_m(\bar{\beta}_j, \bar{V}_j)$, where $\bar{V}_j = (\sigma_j^{-2} X_j' X_j + B_0^{-1})^{-1}$ and $\bar{\beta}_j = \bar{V}_j (\sigma_j^{-2} X_j' y_j + B_0^{-1} b_0)$. Indeed, denoting $\hat{\beta}_j = (X_j' X_j)^{-1} X_j' y_j$,

$$\begin{aligned} p(\beta_j|\sigma_j^2, b_0, B_0, S_T, Y_T) &\propto \exp -0.5[(\beta_j - b_0)' B_0^{-1} (\beta_j - b_0) + \sigma_j^{-2} (y_j - X_j \beta_j)' (y_j - X_j \beta_j)] \\ &\propto \exp -0.5[(\beta_j - b_0)' B_0^{-1} (\beta_j - b_0) + \sigma_j^{-2} (\beta_j - \hat{\beta}_j)' X_j' X_j (\beta_j - \hat{\beta}_j)] \\ &\propto \exp -0.5[(\beta_j - \bar{\beta}_j)' \bar{V}_j^{-1} (\beta_j - \bar{\beta}_j)]. \end{aligned}$$

- For each $j = 1, \dots, K$: $\sigma_j^{-2}|\beta_j, v_0, d_0, S_T, Y_T \sim \text{Gamma}(v_0 + T_j, d_0 + (y_j - X_j \beta_j)' (y_j - X_j \beta_j))$, since

$$\begin{aligned} p(\sigma_j^{-2}|\beta_j, v_0, d_0, S_T, Y_T) &\propto (\sigma_j^{-2})^{v_0/2-1} \exp[-0.5\sigma_j^{-2}d_0] \\ &\quad (\sigma_j^{-2})^{T_j/2} \exp[-0.5\sigma_j^{-2}(y_j - X_j\beta_j)'(y_j - X_j\beta_j)]. \end{aligned}$$

- $b_0|\beta_1, \dots, \beta_K, B_0, Y_T \sim N_m(\bar{\mu}_\beta, \bar{\Sigma}_\beta)$, where $\bar{\Sigma}_\beta = (\underline{\Sigma}_\beta^{-1} + KB_0^{-1})^{-1}$ and $\bar{\mu}_\beta = \bar{\Sigma}_\beta(\underline{\Sigma}_\beta^{-1}\underline{\mu}_\beta + B_0^{-1}\sum_{j=1}^K\beta_j)$. This comes from

$$p(b_0|\beta_1, \dots, \beta_K, B_0, Y_T) \propto \exp[-0.5(b_0 - \underline{\mu}_\beta)'\underline{\Sigma}_\beta^{-1}(b_0 - \underline{\mu}_\beta)] \prod_{j=1}^K \exp[-0.5(\beta_j - b_0)'B_0^{-1}(\beta_j - b_0)].$$

- $B_0^{-1}|\beta_1, \dots, \beta_K, b_0, Y_T \sim W_m(\bar{\nu}_\beta, \bar{V}_\beta^{-1})$, where $\bar{\nu}_\beta = \underline{\nu}_\beta + K$ and $\bar{V}_\beta^{-1} = \underline{V}_\beta^{-1} + \sum_{j=1}^K(\beta_j - b_0)(\beta_j - b_0)'$. Indeed,

$$\begin{aligned} p(B_0^{-1}|\beta_1, \dots, \beta_K, b_0, Y_T) &\propto |B_0^{-1}|^{\frac{\underline{\nu}_\beta+m+1}{2}} \exp[-0.5\text{tr}B_0^{-1}\underline{V}_\beta^{-1}] \\ &\quad \prod_{j=1}^K |B_0|^{-1/2} \exp[-0.5(\beta_j - b_0)'B_0^{-1}(\beta_j - b_0)] \\ &\propto |B_0^{-1}|^{\frac{\underline{\nu}_\beta+K+m+1}{2}} \exp[-0.5\text{tr}B_0^{-1}(\underline{V}_\beta^{-1} + \sum_{j=1}^K(\beta_j - b_0)(\beta_j - b_0)')]. \end{aligned}$$

- $d_0|\sigma_1, \dots, \sigma_K, v_0, Y_T \sim \text{Gamma}(\underline{c}_0 + Kv_0, \underline{d}_0 + \sum_{j=1}^K\sigma_j^{-2})$, since

$$p(d_0|\sigma_1, \dots, \sigma_K, v_0, Y_T) \propto d_0^{\underline{c}_0/2-1} \exp[-0.5d_0\underline{d}_0] \prod_{j=1}^K d_0^{v_0/2} \exp[-0.5d_0\sigma_j^{-2}].$$

- $v_0|\sigma_1, \dots, \sigma_K, d_0, Y_T \propto f_g(v_0|\underline{\lambda}_0, \underline{\rho}_0) \prod_{j=1}^K f_g(\sigma_j^{-2}|v_0, d_0)$. Since this is not belonging to a known class of densities, we simulate it by inverting numerically its cdf computed by Simpson's rule (rather than using a Metropolis algorithm as PPT do).

3 Marginal likelihood and BIC computation

In this section, we give details about the computation of the marginal log-likelihood (MLL) and the Bayesian information criterion (BIC). We are interested by the following questions:

1. Is the value of the MLL computed by Chib's algorithm reliable?
2. If we apply the BIC to choose the number of change points, do we get approximately the same results as if we apply the MLL criterion?

Answers to these questions are provided in Sections 4 and 5. The motivation for using the BIC is that it is well known, and in large samples it usually leads to choose the model also picked

by the MLL criterion, see the discussion in Kass and Raftery (1995). The BIC is generally more quickly programmed and computed than the MLL. For the change-point models we consider, this is especially true in terms of programming time. Once a correct computer program is available, only computing time matters, and in that respect computing the BIC is not necessarily quicker than computing the MLL, because obtaining the global maximum of the log-likelihood function may require many computations (see below).

3.1 Marginal likelihood

The posterior density defined in (18) is conditional on a known value of K , the number of regimes in the sample period. Obviously, K can range from 1, the no break scenario, to T (the sample size). The latter case can be interpreted as a time-varying parameter (TVP) model, for which one often assumes $\theta_i = \theta_0 + \Phi\theta_{i-1} + v_t$ where θ_0 , Φ are parameters (often set equal to 0 and I , respectively) and v_t is a multivariate normal vector with zero mean and unknown covariance matrix.

We can choose the model corresponding to the value of K that maximizes the marginal likelihood (also called predictive density) of the sample. We can also use model averaging, over a range of values of K , say $K \in \{1, 2, \dots, \bar{K}\}$ where \bar{K} is the largest number of regimes that we wish to consider. Model averaging requires posterior model probabilities, which themselves require prior model probabilities and the marginal likelihood value for each model.

To compute the marginal log-likelihood for the data Y_T and the model M_K with parameters $\Theta_K = (\theta_0, \theta_1, \dots, \theta_K)$ and $P_K = (p_{11}, p_{22}, \dots, p_{K-1, K-1})$ defined in the previous section, we use the idea of Chib (1995). The predictive density is related to the prior, posterior and data density by the equality

$$f(Y_T|M_K) = f(Y_T|M_K, \Theta_K, P_K)\varphi(\Theta_K, P_K|M_K)/\varphi(\Theta_K, P_K|M_K, Y_T).$$

Since this holds for any admissible parameter value, we can pick a value (Θ_K^*, P_K^*) of the parameters and compute

$$\log f(Y_T|M_K, \Theta_K^*, P_K^*) + \log \varphi(\Theta_K^*, P_K^*|M_K) - \log \varphi(\Theta_K^*, P_K^*|M_K, Y_T) \quad (26)$$

to approximate $\log f(Y_T|M_K)$. Notice that all densities in the above equation must be proper (i.e. integrate to 1), hence the requirement to use a proper prior. In principle, (Θ_K^*, P_K^*)

can be any value in the parameter space, but the posterior mean or mode is an easy and recommended choice. In Sections 4 and 5, we report results throwing light on the sensitivity of the value of the marginal log-likelihood with respect to the value of Θ_K^* .

The first term in (26) is the log-likelihood function of the model, which can be written as

$$\log f(Y_T|M_K, \Theta_K^*, P_K^*) = \sum_{t=1}^T \log f(y_t|Y_{T-1}, \Theta_K^*, P_K^*) \quad (27)$$

where

$$f(y_t|Y_{T-1}, \Theta_K^*, P_K^*) = \sum_{j=1}^K f(y_t|Y_{t-1}, \Theta_K^*, P_K^*, s_t = j) f(s_t = j|Y_{t-1}, \Theta_K^*, P_K^*). \quad (28)$$

In each term of the above sum, the first factor is equivalent to $f(y_t|Y_{t-1}, \theta_j^*)$ and the second is obtained by the prediction step of Chib's algorithm, see (24). Actually, the log-likelihood function does not depend on θ_0^* .

The prior density is defined analytically and easily computed. The difficult part is the computation of the log-posterior since it must be computed numerically. We use the factorization

$$\varphi(\Theta_K^*, P_K^*|M_K, Y_T) = \varphi(\Theta_K^*|M_K, Y_T)\varphi(P_K^*|\Theta_K^*, M_K, Y_T), \quad (29)$$

where

$$\varphi(P_K^*|\Theta_K^*, M_K, Y_T) = \int \varphi(P_K^*|M_K, Y_T, S_T)\varphi(S_T|\Theta_K^*, M_K, Y_T)dS_T$$

since P_K is independent of Θ_K given S_T . This is estimated by

$$H^{-1} \sum_{h=1}^H \varphi(P_K^*|M_K, Y_T, S_{T,h}),$$

where $\{S_{T,h}\}_{h=1}^H$ are H draws generated from the Gibbs sampler described in sub-section 2.3, *conditioned on* Θ^* , i.e. it iterates between $\varphi(S_T|\Theta^*, P, Y_T)$ and $\varphi(P|S_T)$.

The first density in the right-hand side of (29) can be expressed as

$$\varphi(\Theta_K^*|M_K, Y_T) = \int \varphi(\Theta_K^*|M_K, Y_T, S_T)\varphi(S_T|M_K, Y_T)dS_T.$$

It could be estimated by

$$G^{-1} \sum_{g=1}^G \varphi(\Theta_K^*|M_K, Y_T, S_{T,g}), \quad (30)$$

where $\{S_{T,g}\}_{g=1}^G$ are G draws generated from the posterior density through the Gibbs sampler described above, *if* $\varphi(\Theta_K^*|M_K, Y_T, S_{T,g})$ were known analytically. However this is not typically the case. To solve this problem, we partition Θ_K^* in B blocks $\{\Theta_{K,b}^*\}_{b=1}^B$, such that the full conditional posterior densities $\varphi(\Theta_{K,b}^*|M_K, Y_T, S_T, \{\Theta_{K,c}^*\}_{c \neq b})$ are known analytically or if not can be computed by numerical integration. These blocks are those used for implementing step 3 of the Gibbs sampler (see sub-section 2.3). Then, we factorize $\varphi(\Theta_K^*|M_K, Y_T)$ as

$$\varphi(\Theta_{K,1}^*|M_K, Y_T)\varphi(\Theta_{K,2}^*|M_K, Y_T, \Theta_{K,1}^*) \dots \varphi(\Theta_{K,B}^*|M_K, Y_T, \Theta_{K,1}^*, \dots, \Theta_{K,B-1}^*),$$

and we implement an auxiliary Gibbs sampler to compute each density of this factorization in the same way as (30). For example, $\varphi(\Theta_{K,2}^*|M_K, Y_T, \Theta_{K,1}^*)$ is estimated as

$$G^{-1} \sum_{g=1}^G \varphi(\Theta_{K,2}^*|M_K, Y_T, S_{T,g}, \Theta_{K,1}^*, \Theta_{K,3,g}, \dots, \Theta_{K,B,g})$$

where $\{S_{T,g}, \Theta_{K,3,g}, \dots, \Theta_{K,B,g}\}_{g=1}^G$ are draws from the auxiliary Gibbs sampler. This sampler is the one defined in sub-section 2.3 *conditioned on* $\Theta_{K,1}^*$ and $\Theta_{K,2}^*$.

In the context of the normal regression model (5) for each regime, $B = 6$ and $\Theta_{K,1}$ corresponds to the K vectors β_j , $\Theta_{K,2}$ to the K parameters σ_j^{-2} , $\Theta_{K,3}$ to b_0 , $\Theta_{K,4}$ to B_0^{-1} , $\Theta_{K,5}$ to d_0 , and $\Theta_{K,6}$ to v_0 . This way of partitioning Θ_K minimizes the computational time by exploiting the conditional independence features of the posterior, since the the last four blocks do not require integration with respect to the states. The last block is reserved for v_0 since it requires numerical integration that can be done only once since the other parameters are then fixed.

3.2 BIC

To compute the BIC, we must of course compute the maximum likelihood estimator since we need to evaluate the log-likelihood at its maximum and penalize it by the usual term $0.5m_K \log T$, where m_K is the number of parameters of model M_K . The log-likelihood function of the change-point model is given in (27)-(28). We maximize it by a gradient method (algorithm BFGS of Ox). There may be several local maxima. Thus we maximize the objective function many times, using different starting values. We choose them as follows: for a given number of change points, we draw randomly the break dates, taking care that each regime has at least 30 observations, and given these dates, we estimate the model by

OLS in each regime, thus obtaining starting values for the autoregressive coefficients and the variance of the error. The starting values for the transition probabilities p_{kk} are set to 0.98 or 0.99, since transitions are not frequent. For the applications reported in the next section, we did 100 maximizations and chose as value of the log-likelihood for computing the BIC the maximum of the 100 log-likelihood values. For the simulation results in Section 5, we used 20 maximizations in each experiment.

For running the Gibbs sampler, we use as starting values the MLE values used for computing the BIC.

4 Empirical examples

In this section, we apply the change-point model to three time series. Our empirical results serve to motivate and design the simulation experiments reported in the next section.

4.1 Quarterly growth rate of US GDP

The sample for this series covers the period from the first quarter of 1947 to the last one of 2008 (248 observations). The original series of the real GDP is seasonally adjusted. It was downloaded from the web site of the Federal Reserve Bank of St. Louis (<http://alfred.stlouisfed.org/>, series GDPC1.20090130). The growth rate series is plotted in Figure 6.

We use an AR(p) model in each regime, with $p \leq 4$. We limit the number of regimes to be equal to 3 at most. According to the BIC and MLL criteria, the best model is an AR(1) with two regimes. Table 1 reports, for $p = 1$, the BIC and several MLL values, obtained by applying formula (26) for different parameter values: posterior means, mode, medians, 0.25-quantiles, and 0.75-quantiles, where the quantiles are from each marginal distribution. The choice of two regimes is consistent across the BIC and all values of the MLL. The latter hardly differ for each K . The posterior means, standard deviations, mode, medians and other quantiles of the main parameters of the best model are reported in Table 2. The posterior median of the break date is 1983, second quarter (July) as shown in Figure 6. The break corresponds to a large reduction of the error variance and is called "the great moderation" by economists. The persistence parameter of the AR(1) hardly changes after the break.

4.2 Monthly growth rate of US industrial production

The sample for this series covers the period from January 1950 to January 2009 (709 observations). The original series, downloaded from Datastream (USIPTOT.G series), is a volume index of the industrial production of the USA and is seasonally adjusted, equal to 100 in 2002. We work with the growth rate series, plotted in Figure 6, defined as the first difference of the logarithm of the original series.

We use an AR(1) model in each regime. We limit the number of regimes to be equal to 7 at most. According to the BIC and MLL criteria, the best model has four regimes, see Table 3. The choice of four regimes is consistent across the BIC and the values of the MLL with the exception of the MLL evaluated at the mode which leads to select $K = 5$. One can notice that differences between the criteria values between 3 and 4 regimes are much more important than between 4 and 5.

The MLL values hardly differ for each K . The posterior central values, standard deviations, and 0.25/0.75 quantiles of the main parameters of the best model are reported in Table 4. The posterior medians of the break dates are April 1960, January 1984, and June 2008, as shown in Figure 6. The first break corresponds to a large reduction in the variance of the growth rate: the posterior mean of the residual variance is divided by three. The second break corresponds to a further reduction (division by 2.5). The last break corresponds clearly to the big recession triggered by the subprime crisis of 2007, with a huge increase of the residual variance. The coefficients of the AR(1) equation are very similar in the first two regimes, and change a lot in the last two regimes though the precision of the estimation for the last regime is low due to the very small number of observations.

4.3 Monthly US 3-month T-bill rate

This monthly series was analyzed by Pesaran, Pettenuzzo, and Timmermann (2006) using an AR(1) model for the level. We extend the sample by almost five years, covering the period July 1947 to September 2008 (735 observations). The estimation results for an AR(1) model indicate that most of the regimes are nearly integrated. Therefore, we use an AR(0) model for the first difference. The series (from the CRSP database) is shown in Figure 6, together with the regimes resulting from the detected break dates (posterior medians of change points). The number of regimes is equal to 11 according to the BIC to the different values of the MLL

criterion (see Table 5). The decrease of the criteria values as K increases is marked until $K = 8$, indicating that the choice of the number of regimes after that is not clear. A similar pattern is prevalent in the other applications, suggesting that there is a much bigger risk of picking too many regimes than too few in this type of model.

The posterior means, standard deviations, and modes for 11 regimes are reported in Table 6. The regime changes seem to correspond mainly to changes in the the error term variance. Changes in the mean coefficients of the interest rate changes (β_{j2} , $j=1,2,\dots,11$) seem less important as their posterior means are close to 0 except in regimes 3, 10 and 11.

5 Simulation study

The simulation design is inspired by the empirical results of the previous section. We choose three DGPs, one with a single change point, one with three, and one with seven. In each regime, a sample of size T is generated by an AR(1) process with parameter values and regime durations close to the corresponding estimated results of the previous section. The three DGPs are presented in Tables 7, 10, and 13. For DGPs 1 and 2, 100 repetitions are used, and for the last DGP this number is reduced to 50 because computations are taking a lot of time. A repetition means that a sample of size T is simulated from the DGP and for each sample, the BIC and MLL values are computed for AR(1) models with varying number of regimes. These sets of 100 (or 50) values are then used to produce averages and proportions.

5.1 Single change point ($K = 2$)

The results of the simulations are contained in Tables 8 and 9. Table 8 contains means (across the 100 repetitions) of BIC and MLL values. Table 9 shows the proportions in which the models with 1, 2 (the true value), 3 or 4 regimes are selected by the BIC or the MLL criterion. From these results, the answers to the two questions we raise in the beginning of 3 are rather clear:

1) The MLL values depend very little on the parameter value chosen for its computation, when the values being considered are the means, mode, medians, 0.25-quantiles (q25), and 0.75-quantiles (q75), where the quantiles are from each marginal distribution. For a given K , the largest difference is 2 ($K = 4$, $T = 500$) when comparing mean, mode, and median-based

results (Table 8), which is very small compared to 692. Between mode and median, the difference is always smaller than 1. When comparing mean, mode or median with the quartiles, the differences are a bit larger, with the largest difference occurring for q25 in the cases $K = 2$, $T = 250$ (difference of 3.622) and 500 (difference of 6.19). The sensitivity does not seem to be related to the sample size T and may be slightly more pronounced for $K = 4$ than for smaller values. This is expected as inference for an over-parameterized model is likely to be less precise.

2) Concerning the model selection criteria (Table 9), the performance of the BIC in the selection of the correct model is very good (at least 97 per cent of correct choices) and even slightly better than for the MLL. The MLL performance is the best when its computation is based on central values of the parameters (more than 90 per cent of correct choices). When T is increased from 250 to 500, the performance of the BIC and MLL computed with the mode and median increases.

5.2 Three change points ($K = 4$)

The results of the simulations are contained in Tables 11 and 12. Table 11 contains means (across the 100 repetitions) of BIC and MLL values. Table 12 shows the proportions in which the models with 2 to 6 regimes are selected by the BIC or the MLL criterion. From these results, the answers to the questions raised are rather clear:

1) The MLL values depend very little on the parameter values chosen for its computation. For a given K , the largest difference is 3.95 (for $K = 3$) when comparing mean, mode, and median-based results (Table 11), which is very small compared to 913. Between mode and median, the difference is smaller than 1 except for $K = 3$, where the value at the mode is slightly different. When comparing mean, mode or median with the quartiles, the differences are increasing with K , because inference for an over-parameterized model is likely to be less precise.

2) Concerning the model selection criteria (Table 12), the performance of the BIC and the MLL criterion (whatever the evaluation point) is good, since the right model (4 regimes) is picked in about 89 to 94 per cent of the repetitions. The risk of choosing a wrong number of regimes is entirely concentrated on the over-parameterized models (especially the model with 5 regimes).

5.3 Ten change points ($K = 11$)

The results of the simulations are contained in Tables 14 and 15. Table 14 contains means (across the 50 repetitions) of BIC and MLL values. Table 15 shows the proportions in which the models with 9 up to 13 regimes are selected by the BIC or the MLL criterion. From these results, the answers to the questions we are interested in are rather clear:

1) The MLL values (Table 14) do not depend much on the parameter value chosen for its computation, but they vary more than in the two previous cases. The largest difference occurs for $K = 13$ between mean and mode (difference of the order of 10 per cent). However, between mode and median, the differences are very small.

2) Concerning the model selection criteria (Table 15), the performance is not good. The BIC selects the correct number of breaks only in 24 per cent of the repetitions and puts too much weight (50 per cent) on over-parameterized models ($K = 12$ and 13). This is even much more pronounced with the MLL criterion, since it selects the models with too many regimes in 90 per cent of the repetitions. This bad performance of the criteria may be explained by the following feature of the DGP: there are many regimes, among which some with few observations, and some which are not very different. This confirms our intuition, based on the applications to real data, that there is a tendency to select too many breaks with the type of model we consider.

6 Conclusion

There are two objectives in this paper. The first is to assess the sensitivity of the marginal likelihood value obtained by Chib's method in the case of a change-point regression model. The second objective is to compare the model choices resulting from the application of the BIC and of the marginal likelihood criterion. Our findings are firstly that the value of the marginal likelihood computed by Chib's algorithm is reliable for the class of Markov-switching autoregressive models. Secondly that the MLL criterion and BIC provide a good performance in choosing the right model, with a marginally better performance of the BIC, except apparently in the case where the number of regimes is large and difficult to identify.

References

- BAUWENS, L., M. LUBRANO, AND J. RICHARD (1999): *Bayesian Inference in Dynamic Econometric Models*. Oxford University Press, Oxford.
- CHIB, S. (1995): “Marginal likelihood from the Gibbs output,” *Journal of the American Statistical Association*, 90, 1313–1321.
- (1996): “Calculating posterior distributions and modal estimates in Markov mixture models,” *Journal of Econometrics*, 75, 79–97.
- (1998): “Estimation and comparison of multiple change-point models,” *Journal of Econometrics*, 86, 221–241.
- ELERIAN, O., S. CHIB, AND N. SHEPHARD (2001): “Likelihood inference for discretely observed nonlinear diffusions,” *Econometrica*, 69(4), 959–993.
- FRUHWIRTH-SCHNATTER, S., AND H. WAGNER (2008): “Marginal likelihoods for non-Gaussian models using auxiliary mixture sampling,” *Computational Statistics & Data Analysis*, 52(10), 4608–4624.
- GELFAND, A., AND D. DEY (1994): “Bayesian model choice: Asymptotics and exact calculations,” *Journal of the Royal Statistical Society B*, 56, 501–514.
- HAMILTON, J. (1989): “A New Approach to the Economic Analysis of Nonstationary Time Series and the Business Cycle,” *Econometrica*, 57, 357–384.
- JOHNSON, L. D., AND G. SAKOULIS (2008): “Maximizing equity market sector predictability in a Bayesian time-varying parameter model,” *Computational Statistics & Data Analysis*, 52(6), 3083–3106.
- KASS, R., AND A. RAFTERY (1995): “Bayes Factors,” *Journal of the American Statistical Association*, 90, 773–795.
- KIM, C., J. MORLEY, AND C. NELSON (2005): “The structural break in the equity premium,” *Journal of Business & Economic Statistics*, 23(2), 181–191.

- KIM, C., AND C. NELSON (1999): “Has the US economy become more stable? A Bayesian approach based on a Markov-switching model of the business cycle,” *Review of Economics and Statistics*, 81(4), 608–616.
- KIM, C., AND J. PIGER (2002): “Common stochastic trends, common cycles, and asymmetry in economic fluctuations,” *Journal of Monetary Economics*, 49(6), 1189–1211.
- LIU, C., AND J. M. MAHEU (2008): “Are there structural breaks in realized volatility?,” *Journal of Financial Econometrics*, 6(3), 326–360.
- MAHEU, J. M., AND S. GORDON (2008): “Learning, forecasting and structural breaks,” *Journal of Applied Econometrics*, 23(5), 553–583.
- NAKAJIMA, J., AND Y. OMORI (2009): “Leverage, heavy-tails and correlated jumps in stochastic volatility models,” *Computational Statistics & Data Analysis*, 53(6, Sp. Iss. SI), 2335–2353.
- NEWTON, M., AND A. RAFTERY (1994): “Approximate Bayesian inference by the weighted likelihood bootstrap,” *Journal of the Royal Statistical Society B*, 56, 3–48.
- PAROLI, R., AND L. SPEZIA (2008): “Bayesian inference in non-homogeneous Markov mixtures of periodic autoregressions with state-dependent exogenous variables,” *Computational Statistics & Data Analysis*, 52(5), 2311–2330.
- PASTOR, L., AND R. F. STAMBAUGH (2001): “The Equity Premium and Structural Breaks,” *Journal of Finance*, 56, 1207–1239.
- PESARAN, M. H., D. PETTENUZZO, AND A. TIMMERMANN (2006): “Forecasting Time Series Subject to Multiple Structural Breaks,” *Review of Economic Studies*, 73, 1057–1084.
- STOCK, J. H., AND M. W. WATSON (1996): “Evidence on Structural Instability in Macroeconomic Time Series Relations,” *Journal of Business & Economic Statistics*, 14, 11–30.

Table 1: BIC and MLL of AR(1) models (US GDP growth rate)

K	BIC	MLL mean	MLL mode	MLL median	MLL q25	MLL q75
1	-336.44	-349.03	-349.06	-349.03	-348.96	-349.01
2	-321.04	-332.66	-332.39	-332.66	-333.62	-333.88
3	-322.02	-334.98	-333.37	-334.68	-335.78	-336.86

BIC: Bayesian information criterion. MLL: marginal log-likelihood, computed by formula (26) using different parameter posterior values: mean, mode, median, 0.25-quantile (q25) and 0.75 quantile (q75).

Table 2: Posterior results for 1-change point AR(1) model (US GDP growth rate)

Parameter	Mean	St. dev.	Mode	Median	q25	q75
β_{11}	0.576	0.420	0.592	0.580	0.491	0.672
β_{12}	0.345	0.313	0.334	0.336	0.276	0.396
β_{21}	0.478	0.369	0.554	0.481	0.404	0.558
β_{22}	0.333	0.409	0.246	0.330	0.242	0.417
σ_1^2	1.471	2.889	1.381	1.282	1.178	1.398
σ_2^2	0.344	0.779	0.261	0.278	0.252	0.313
p_{11}	0.985	0.053	0.993	0.991	0.986	0.996

Table 3: BIC and MLL of AR(1) models (US industrial production growth rate)

K	BIC	MLL mean	MLL mode	MLL median	MLL q25	MLL q75
1	-927.28	-945.11	-945.13	-945.11	-944.91	-944.94
2	-885.37	-894.61	-894.35	-894.60	-894.90	-894.79
3	-869.70	-882.18	-882.39	-882.16	-882.96	-882.72
4	-855.70	-859.11	-859.35	-859.13	-859.93	-859.67
5	-859.79	-861.68	-858.98	-862.22	-861.35	-863.64
6	-864.41	-864.02	-862.44	-861.88	-867.37	-865.45
7	-872.63	-865.63	-859.97	-864.84	-883.67	-869.86

See Table 1 for explanations.

Table 4: Posterior results for 3-change point AR(1) model (US industrial production growth rate)

Parameter	Mean	St. dev.	Mode	Median	q25	q75
β_{11}	0.195	0.157	0.307	0.199	0.103	0.291
β_{12}	0.470	0.093	0.441	0.470	0.415	0.526
β_{21}	0.171	0.057	0.200	0.170	0.136	0.207
β_{22}	0.405	0.062	0.395	0.405	0.368	0.443
β_{31}	0.191	0.034	0.171	0.191	0.168	0.213
β_{32}	0.100	0.067	0.144	0.099	0.059	0.139
β_{41}	-0.200	0.520	-0.036	-0.189	-0.523	0.129
β_{42}	0.184	0.354	0.463	0.179	-0.045	0.400
σ_1^2	2.075	0.364	2.065	2.043	1.868	2.240
σ_2^2	0.680	0.066	0.668	0.678	0.639	0.717
σ_3^2	0.262	0.025	0.283	0.260	0.246	0.275
σ_4^2	5.390	4.174	4.857	4.259	2.993	6.523
p_{11}	0.988	0.011	0.996	0.991	0.984	0.995
p_{22}	0.995	0.004	0.998	0.996	0.993	0.998
p_{33}	0.995	0.004	0.999	0.996	0.993	0.998

Table 5: BIC and MLL of AR(0) models (US T-bill rate)

K	BIC	MLL mean	MLL mode	MLL median	MLL q25	MLL q75
1	-512.41	-523.48	-523.48	-523.52	-523.49	-523.49
2	-425.85	-494.43	-438.59	-452.81	-531.50	-527.74
3	-295.82	-307.67	-307.61	-307.67	-308.12	-307.96
4	-269.43	-280.02	-280.08	-280.02	-280.29	-280.32
5	-253.42	-263.06	-263.11	-263.02	-263.53	-263.63
6	-229.82	-235.82	-235.93	-235.86	-236.31	-236.34
7	-226.00	-229.10	-229.47	-229.14	-229.70	-229.76
8	-215.08	-214.96	-215.20	-215.02	-215.56	-215.63
9	-211.38	-208.81	-209.40	-208.90	-209.72	-209.90
10	-212.86	-223.06	-207.33	-210.18	-224.07	-262.45
11	-209.98	-207.65	-198.02	-197.58	-199.03	-208.86
12	-219.39	-213.93	-199.20	-201.70	-221.98	-241.31

See Table 1 for explanations.

Table 6: Posterior results for 10-change point AR(0) model (US T-bill rate)

Parameter	Mean	St. dev.	Mode	Median	q25	q75
β_1	0.023	0.014	0.027	0.023	0.014	0.032
β_2	-0.03	0.083	-0.064	-0.036	-0.091	0.018
β_3	0.036	0.015	0.040	0.036	0.026	0.046
β_4	0.027	0.065	-0.006	0.030	-0.011	0.069
β_5	0.073	0.078	0.116	0.069	0.017	0.125
β_6	-0.04	0.194	-0.045	-0.042	-0.172	0.085
β_7	-0.005	0.046	-0.046	-0.006	-0.036	0.026
β_8	-0.03	0.028	0.039	-0.037	-0.052	-0.019
β_9	-0.08	0.124	-0.357	-0.035	-0.057	-0.022
β_{10}	0.092	0.033	0.032	0.095	0.068	0.114
β_{11}	-0.22	0.122	-0.137	-0.226	-0.307	-0.147
σ_1^2	0.023	0.003	0.026	0.022	0.021	0.024
σ_2^2	0.264	0.073	0.247	0.252	0.212	0.302
σ_3^2	0.015	0.003	0.015	0.014	0.012	0.017
σ_4^2	0.187	0.123	0.351	0.114	0.075	0.303
σ_5^2	0.218	0.210	0.099	0.257	0.090	0.314
σ_6^2	2.541	0.640	2.563	2.439	2.083	2.880
σ_7^2	0.158	0.028	0.103	0.155	0.138	0.175
σ_8^2	0.049	0.009	0.038	0.050	0.044	0.055
σ_9^2	0.028	0.048	0.132	0.007	0.006	0.011
σ_{10}^2	0.019	0.005	0.017	0.018	0.016	0.021
σ_{11}^2	0.258	0.113	0.256	0.232	0.185	0.302
P_{11}	0.988	0.010	0.992	0.991	0.983	0.995
P_{22}	0.959	0.033	0.995	0.967	0.943	0.983
P_{33}	0.979	0.018	0.989	0.983	0.971	0.991
P_{44}	0.969	0.060	0.989	0.983	0.965	0.992
P_{55}	0.973	0.046	0.986	0.985	0.969	0.993
P_{66}	0.962	0.030	0.980	0.970	0.948	0.984
P_{77}	0.982	0.014	0.984	0.986	0.975	0.993
P_{88}	0.990	0.009	0.987	0.992	0.986	0.996
P_{99}	0.938	0.057	0.925	0.954	0.917	0.977
P_{1010}	0.966	0.029	0.989	0.974	0.954	0.987

Table 7: DGP with 1 change point

Regime	Duration	Equation
1	140	$y_t = 0.60 + 0.35y_{t-1} + \sqrt{1.50}z_t$
2	110	$y_t = 0.45 + 0.30y_{t-1} + \sqrt{0.35}z_t$

z_t simulated from $N(0, 1)$ distribution.

Table 8: MLL evaluated in different points for DGP with 1 change point

K	Mean	Mode	Median	q25	q75
$T = 250$ (140 + 110)					
1	-373.45	-373.45	-373.42	-373.34	-373.38
2	-358.73	-358.07	-358.49	-361.69	-359.93
3	-361.00	-360.36	-360.26	-361.72	-362.05
4	-364.08	-363.17	-362.44	-366.09	-365.89
$T = 500$ (280 + 220)					
1	-730.14	-730.14	-730.11	-729.97	-730.02
2	-686.71	-685.95	-686.22	-692.14	-687.30
3	-689.09	-688.59	-688.25	-689.15	-689.67
4	-692.50	-691.29	-690.50	-692.60	-692.99

Reported MLL values are means computed from 100 replications. In each replication, T observations are simulated from the DGP defined in Table 7.

Table 9: Model selection performance for DGP with 1 change point

K	BIC	Mean	Mode	Median	q25	q75
$T = 250$						
1	0.01	0.01	0.01	0.01	0.02	0.02
2	0.97	0.95	0.91	0.95	0.72	0.91
3	0.02	0.04	0.08	0.04	0.26	0.07
4	0.00	0.00	0.00	0.00	0.00	0.00
$T = 500$						
1	0.00	0.00	0.00	0.00	0.00	0.00
2	1.00	0.93	0.96	0.98	0.79	0.90
3	0.00	0.07	0.04	0.02	0.21	0.10
4	0.00	0.00	0.00	0.00	0.00	0.00

Proportions based on same replications as for Table 8.

Table 10: DGP with 3 change points

Regime	Duration	Equation
1	123	$y_t = 0.20 + 0.47y_{t-1} + \sqrt{2.10}z_t$
2	285	$y_t = 0.17 + 0.40y_{t-1} + \sqrt{0.68}z_t$
3	293	$y_t = 0.19 + 0.10y_{t-1} + \sqrt{0.26}z_t$
4	16	$y_t = -0.2 + 0.18y_{t-1} + \sqrt{5.39}z_t$

z_t simulated from $N(0, 1)$ distribution.

Table 11: MLL evaluated in different points for DGP with 3 change points

K	Mean	Mode	Median	q25	q75
2	-937.32	-937.18	-937.32	-937.66	-937.58
3	-913.62	-909.67	-912.69	-916.82	-915.14
4	-878.51	-878.21	-878.41	-882.22	-879.36
5	-882.55	-879.71	-879.81	-886.76	-882.79
6	-885.43	-881.90	-881.93	-892.35	-887.00

Reported MLL values are means computed from 100 replications. In each replication, 707 observations are simulated from the DGP defined in Table 10.

Table 12: Model selection performance for DGP with 3 change points

K	BIC	Mean	Mode	Median	q25	q75
2	0.00	0.00	0.00	0.00	0.00	0.00
3	0.00	0.00	0.00	0.00	0.00	0.00
4	0.91	0.94	0.89	0.96	0.91	0.94
5	0.09	0.06	0.10	0.04	0.08	0.05
6	0.00	0.00	0.01	0.00	0.01	0.01

Proportions based on same replications as for Table 11.

Table 13: DGP with 10 change points

Regime	Duration	Equation
1	122	$y_t = 0.023 + \sqrt{0.02}z_t$
2	33	$y_t = -0.040 + \sqrt{0.26}z_t$
3	72	$y_t = 0.038 + \sqrt{0.02}z_t$
4	103	$y_t = 0.006 + \sqrt{0.28}z_t$
5	52	$y_t = 0.110 + \sqrt{0.12}z_t$
6	37	$y_t = -0.040 + \sqrt{2.51}z_t$
7	91	$y_t = -0.01 + \sqrt{0.15}z_t$
8	129	$y_t = -0.01 + \sqrt{0.04}z_t$
9	20	$y_t = -0.23 + \sqrt{0.08}z_t$
10	59	$y_t = 0.06 + \sqrt{0.02}z_t$
11	20	$y_t = -0.23 + \sqrt{0.26}z_t$

z_t simulated from $N(0, 1)$ distribution.

Table 14: MLL evaluated in different points for DGP with 10 change points

K	Mean	Mode	Median	q25	q75
9	-210.70	-210.76	-210.39	-211.47	-212.44
10	-209.56	-207.17	-207.75	-211.77	-215.59
11	-207.55	-202.33	-202.71	-207.12	-213.84
12	-206.81	-198.47	-199.45	-204.46	-210.51
13	-205.84	-193.95	-194.22	-199.76	-207.00

Reported MLL values are means computed from 50 replications. In each replication, 736 observations are simulated from the DGP defined in Table 13.

Table 15: Model selection performance for DGP with 10 change points

K	BIC	Mean	Mode	Median	q25	q75
9	0.12	0.00	0.00	0.00	0.00	0.00
10	0.16	0.04	0.00	0.00	0.04	0.04
11	0.24	0.08	0.00	0.04	0.04	0.04
12	0.20	0.16	0.20	0.12	0.20	0.20
13	0.28	0.72	0.80	0.84	0.72	0.72

Proportions based on same replications as for Table 14.




Figure 1: Real U.S. GDP growth rate. Sample period: 1947.Q1-2008.Q4 (248 observations)




Figure 2: U.S. Industrial production growth rate. Sample period: January 1950-January 2009 (707 observations)




Figure 3: Three month U.S. T-bill rate change. Sample period: July 1947-September 2008 (735 observations)

Recent titles

CORE Discussion Papers

- 2009/22. Claude D'ASPREMONT and Rodolphe DOS SANTOS FERREIRA. Household behavior and individual autonomy.
- 2009/23. Helmuth CREMER, Philippe DE DONDER and Pierre PESTIEAU. Education and social mobility.
- 2009/24. Maria Eugenia SANIN and Francesco VIOLANTE. Understanding volatility dynamics in the EU-ETS market: lessons *from* the future.
- 2009/25. Marco DI SUMMA and Laurence A. WOLSEY. Lot-sizing with stock upper bounds and fixed charges.
- 2009/26. Johanna M.M. GOERTZ and François MANIQUET. On the informational efficiency of simple scoring rules.
- 2009/27. Jean GABSZEWICZ, Ornella TAROLA and Skerdilajda ZANAJ. On uncertainty when it affects successive markets.
- 2009/28. Jerzy A. FILAR, Jacek B. KRAWCZYK and Manju AGRAWAL. On production and abatement time scales in sustainable development. Can we loosen the *sustainability screw* ?
- 2009/29. Maria Eugenia SANIN and Skerdilajda ZANAJ. Clean technology adoption and its influence on tradeable emission permit prices.
- 2009/30. Antoine BOMMIER, Marie-Louise LEROUX and Jean-Marie LOZACHMEUR. On the public economics of annuities with differential mortality.
- 2009/31. Gilles GRANDJEAN, Ana MAULEON and Vincent VANNETELBOSCH. Connections among farsighted agents.
- 2009/32. Axel GAUTIER and Xavier WAUTHY. On the nature of price competition under universal service obligations: a note.
- 2009/33. Santanu S. DEY and Laurence A. WOLSEY. Constrained infinite group relaxations of MIPs.
- 2009/34. Jean-François MAYSTADT and Philip VERWIMP. Winners and losers among a refugee-hosting population.
- 2009/35. Pierre DEHEZ. Allocation of fixed costs and the weighted Shapley value.
- 2009/36. Sabien DOBBELAERE, Roland Iwan LUTTENS and Bettina PETERS. Starting an R&D project under uncertainty.
- 2009/37. Carlotta BALESTRA and Davide DOTTORI. Aging society, health and the environment.
- 2009/38. Alain PHOLO BALA. Urban concentration and economic growth: checking for specific regional effects.
- 2009/39. Alain PHOLO BALA. Gates, hubs and urban primacy in Sub-Saharan Africa.
- 2009/40. Nicolas BOCCARD. On efficiency, concentration and welfare.
- 2009/41. Taoufik BOUEZMARNI, Jeroen V.K. ROMBOUTS and Abderrahim TAAMOUTI. A nonparametric copula based test for conditional independence with applications to Granger causality.
- 2009/42. Jozsef KONINGS and Hylke VANDENBUSSCHE. Antidumping protection hurts exporters: firm-level evidence from France.
- 2009/43. Pierre PESTIEAU and Uri M. POSSEN. Retirement as a hedge.
- 2009/44. Santanu S. DEY and Laurence A. WOLSEY. Lifting group inequalities and an application to mixing inequalities.
- 2009/45. Jean CAVAILHES, Pierre FRANKHAUSER, Dominique PEETERS and Isabelle THOMAS. Residential equilibrium in a multifractal metropolitan area.
- 2009/46. Daisuke OYAMA, Yasuhiro SATO, Takatoshi TABUCHI and Jacques-François THISSE. On the impact of trade on industrial structures: The role of entry cost heterogeneity.
- 2009/47. Ken-Ichi SHIMOMURA and Jacques-François THISSE. Competition among the big and the small.
- 2009/48. Frédéric BABONNEAU, Yurii NESTEROV and Jean-Philippe VIAL. Design and operations of gas transmission networks.
- 2009/49. Olivier BOS. How lotteries outperform auctions for charity.

Recent titles

CORE Discussion Papers - continued

- 2009/50. Nicolas BOCCARD and Xavier WAUTHY. Entry accommodation under multiple commitment strategies: judo economics revisited.
- 2009/51. Joachim GAHUNGU and Yves SMEERS. Multi-assets real options.
- 2009/52. Nicolas BOCCARD and Xavier WAUTHY. Regulating quality by regulating quantity: a case against minimum quality standards.
- 2009/53. David DE LA CROIX and Frédéric DOCQUIER. An incentive mechanism to break the low-skill immigration deadlock.
- 2009/54. Henry TULKENS and Vincent VAN STEENBERGHE. "Mitigation, adaptation, suffering": In search of the right mix in the face of climate change.
- 2009/55. Santanu S. DEY and Quentin LOUVEAUX. Split rank of triangle and quadrilateral inequalities.
- 2009/56. Claire DUJARDIN, Dominique PEETERS and Isabelle THOMAS. Neighbourhood effects and endogeneity issues.
- 2009/57. Marie-Louise LEROUX, Pierre PESTIEAU and Maria RACIONERO. Voting on pensions: sex and marriage.
- 2009/58. Jean J. GABSZEWICZ. A note on price competition in product differentiation models.
- 2009/59. Olivier BOS and Martin RANGER. All-pay auctions with endogenous rewards.
- 2009/60. Julio DAVILA and Marie-Louise LEROUX. On the fiscal treatment of life expectancy related choices.
- 2009/61. Luc BAUWENS and Jeroen V.K. ROMBOUTS. On marginal likelihood computation in change-point models.

Books

- Public goods, environmental externalities and fiscal competition: 22 selected papers in public economics by Henry Tulkens*, edited and introduced by Parkash Chander, Jacques Drèze, C. Knox Lovell and Jack Mintz, Springer, Boston 2006 (588 pp.).
- V. GINSBURGH and D. THROSBY (eds.) (2006), *Handbook of the economics of art and culture*. Amsterdam, Elsevier.
- J. GABSZEWICZ (ed.) (2006), *La différenciation des produits*. Paris, La découverte.
- L. BAUWENS, W. POHLMEIER and D. VEREDAS (eds.) (2008), *High frequency financial econometrics: recent developments*. Heidelberg, Physica-Verlag.
- P. VAN HENTENRYCKE and L. WOLSEY (eds.) (2007), *Integration of AI and OR techniques in constraint programming for combinatorial optimization problems*. Berlin, Springer.
- P-P. COMBES, Th. MAYER and J-F. THISSE (eds.) (2008), *Economic geography: the integration of regions and nations*. Princeton, Princeton University Press.
- J. HINDRIKS (ed.) (2008), *Au-delà de Copernic: de la confusion au consensus ?* Brussels, Academic and Scientific Publishers.
- J-M. HURIOT and J-F. THISSE (eds) (2009), *Economics of cities*. Cambridge, Cambridge University Press.

CORE Lecture Series

- C. GOURIÉROUX and A. MONFORT (1995), *Simulation Based Econometric Methods*.
- A. RUBINSTEIN (1996), *Lectures on Modeling Bounded Rationality*.
- J. RENEGAR (1999), *A Mathematical View of Interior-Point Methods in Convex Optimization*.
- B.D. BERNHEIM and M.D. WHINSTON (1999), *Anticompetitive Exclusion and Foreclosure Through Vertical Agreements*.
- D. BIENSTOCK (2001), *Potential function methods for approximately solving linear programming problems: theory and practice*.
- R. AMIR (2002), *Supermodularity and complementarity in economics*.
- R. WEISMANTEL (2006), *Lectures on mixed nonlinear programming*.