

**Birkbeck ePrints: an open access repository of the  
research output of Birkbeck College**

<http://eprints.bbk.ac.uk>

---

Avery, P.B.; Queeley, G.L.; Faull, J.; Simmonds, M.S.J.

**Effect of photoperiod and host distribution on the horizontal  
transmission of *Isaria fumosorosea* (Hypocreales: Cordycipitaceae)  
in greenhouse whitefly assessed using a novel model bioassay**

*Biocontrol Science and Technology* - 20 (10),  
(2010), pp.1097-1111

---

This is a peer-reviewed author's draft of an article published in *Biocontrol  
Science and Technology* (ISSN 0958-3157) made available here with kind  
permission of:

© 2010 Taylor & Francis. All rights reserved.

All articles available through Birkbeck ePrints are protected by intellectual  
property law, including copyright law. Any use made of the contents should  
comply with the relevant law.

Citation for this version:

Avery, P.B.; Queeley, G.L.; Faull, J.; Simmonds, M.S.J.

**Effect of photoperiod and host distribution on the horizontal  
transmission of *Isaria fumosorosea* (Hypocreales: Cordycipitaceae) in  
greenhouse whitefly assessed using a novel model bioassay**

London: *Birkbeck ePrints*. Available at: <http://eprints.bbk.ac.uk/1258>

Citation for publisher's version:

Avery, P.B.; Queeley, G.L.; Faull, J.; Simmonds, M.S.J.

**Effect of photoperiod and host distribution on the horizontal  
transmission of *Isaria fumosorosea* (Hypocreales: Cordycipitaceae) in  
greenhouse whitefly assessed using a novel model bioassay**

*Biocontrol Science and Technology* - 20 (10), (2010), pp.1097-1111

---

<http://eprints.bbk.ac.uk>

Contact Birkbeck ePrints at [lib-eprints@bbk.ac.uk](mailto:lib-eprints@bbk.ac.uk)

Target Journal: Biocontrol Science and Technology

**Effect of photoperiod and host distribution on the horizontal transmission of *Isaria fumosorosea* (Hypocreales: Cordycipitaceae) in greenhouse whitefly using a novel model bioassay**

Pasco B. Avery<sup>1,2</sup>, Gilbert L. Queeley<sup>3</sup>, Jane Faull<sup>1</sup> and Monique S.J. Simmonds<sup>4</sup>  
(Received 6 August, 2009; Returned 18 September; Accepted 10 August, 2010)

<sup>1</sup>Department of Biological Sciences, Birkbeck, University of London, Malet Street, Bloomsbury, London, WC1E 7HX, UK

<sup>2</sup>University of Florida, Institute of Food and Agricultural Science, Indian River Research and Education Center, 2199 S. Rock Road, Ft. Pierce, FL, 34945, USA

<sup>3</sup>College of Engineering Sciences, Technology and Agriculture, Cooperative Extension Programs, 215 Perry Paige Bldg., S., Florida A & M University, Tallahassee, FL, 32307, USA

<sup>4</sup>Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK

Proofs should be sent to:

Pasco Avery, Ph.D., UF/IFAS/IRREC, 2199 S. Rock Road, Ft. Pierce, FL, 34945  
e-mail address: [pbavery@ufl.edu](mailto:pbavery@ufl.edu) ; phone: 772-468-3922 x 161, fax: 772-468-5668.

## Abstract

A model bioassay was used to evaluate the epizootic potential and determine the horizontal transmission efficiency of *Isaria fumosorosea* Trinidadian strains against *Trialeurodes vaporariorum* pharate adults under optimum conditions ( $25 \pm 0.5$  °C, ~100% RH) at two different photoperiods. Untreated pharate adults were arranged on laminated graph paper at different distributions to simulate varying infestation levels on a leaf surface. Four potential hosts were located 7, 14, and 21 mm away from a central sporulating cadaver simulating high, medium and low infestation levels, respectively. Percent hosts colonized were recorded 7, 12, 14 and 21 days post-treatment during a 16 and 24 hour photophase. After 21 days, mean percent hosts colonized at the highest, middle and lowest infestation levels were 93 and 100%, 22 and 58%, 25 and 39% under a 16 and 24 hour photophase, respectively. From the results, it was concluded that the longer the photophase, the greater the percentage of hosts colonized, and as host distance increased from the central sporulating cadaver, colonization decreased. The use of this novel model bioassay technique is the first attempt to evaluate the epizootic potential and determine the horizontal transmission efficiency of *I. fumosorosea* Trinidadian strains under optimal environmental conditions at different photoperiods. This bioassay can be used to assess transmission efficiency for the selection of fungi being considered for commercial biopesticide development.

**Keywords:** blastospore, epizootic potential, whitefly, leaf model bioassay, secondary cycling

## Introduction

The greenhouse whitefly, *Trialeurodes vaporariorum* (Westwood) (Hemiptera: Aleyrodidae) is a pest of agricultural crops worldwide that necessitates frequent control using insecticides (Castañé & Albajes 1992; Antonious & Snyder 1995; Wang et al. 2003). Due to the inappropriate use of insecticides, environmental concerns and evolution of resistance to important insecticides, there is need for alternative methods to control whiteflies (Omer et al. 1992; Sanderson & Rousch 1992; Bi & Toscano 2007).

Entomopathogenic fungi are being developed as major components of integrated pest management programs for controlling *T. vaporariorum* and other whiteflies (Fransen 1993; Poprawski & Jones 2000; Meekes et al. 2002; Vidal et al. 2003; Cuthbertson & Walters 2005; Cuthbertson et al. 2008). Most research using fungi for controlling aleyrodids has focused primarily on using *Beauveria bassiana* (Balsamo) Vuillemin, *Lecanicillium muscarium* (Petch) Zare & Gams and *Aschersonia* species; however, *Isaria fumosorosea* (= *Paecilomyces fumosoroseus*) Wize (Hypocreales: Cordycipitaceae), has also been shown to cause epizootics in whitefly populations on various host plants (Lacey et al. 1995; Wraight et al. 2000; Meekes et al. 2000; Gökçe & Er 2005; Saito 2005; Alma et al. 2007).

To determine the efficacy of an entomopathogen used in any control program, both biotic (host density, developmental stage, etc.) and abiotic (humidity, temperature, photoperiod, etc.) factors that regulate epizootics in insect pests need to be evaluated (Shah & Pell 2003; Vidal et al. 2003; Pell 2008; Vidal & Fargues 2008). One of many factors important in epizootics, such as epizootic potential, depends not only on their ability to sporulate on cadavers, but also their ability to spread to healthy insects, a process known as secondary cycling (Thomas et al. 1995, Luz & Fargues 1997, 1999; Luz et al. 1999; de la Rosa et al. 2000; Long et al. 2000a; Arthurs & Thomas 2001a, b). Factors that promote secondary cycling are important for achieving long term control and can increase the horizontal transmission efficiency of the entomopathogen.

Horizontal transmission efficiency is dependent upon several parameters including the number and distribution of individuals in the infected population. The importance of density dependent transmission efficiency has been confirmed in the laboratory (Galani & Almasan 1984; Carruthers & Soper 1987) and field (Brown & Hasibuan 1995; Keller et al. 1997; de la Rosa et al. 2000; Long et al. 2000b; Kreutz et al. 2004).

Epizootic parameters of entomopathogenic fungi against various arthropods have been studied under laboratory conditions using different model systems. Hall (1984) used leaf disc bioassays to study the epizootic potential of *L. muscarium* isolates against aphids, whereas Mier et al. (1991) used glass slides in moisture chambers to conduct *in vitro* pathogenicity tests on whitefly nymphs with Mexican isolates of *L. muscarium*. Landa et al. (1994) used a glass slide bioassay to compare the pathogenicity of conidia from different isolates of *I. fumosorosea*, *L. muscarium* and *B. bassiana* against early fourth-instar whitefly nymphs. Later, Landa and Bohatá (1999) used glass slides to determine the compatibility of *I. fumosorosea* with natural insecticides against whitefly nymphs. Most recently, Avery et al. (2004) used a glass slide bioassay to compare the pathogenicity of blastospores and conidia of *I. fumosorosea* Trinidadian strains against pharate adults of *T. vaporariorum* at two different photoperiods. Using this model, horizontal transmission efficiency of *I. fumosorosea* strains after colonization could not be assessed, therefore; a novel model bioassay simulating a leaf surface was designed.

The purpose of this study was to determine the horizontal transmission efficiency of *I. fumosorosea* Trinidadian fungal strains in colonizing susceptible pharate adults of the greenhouse whitefly at different distances from a sporulating cadaver under different photoperiod regimes. Two photoperiods were chosen to compare the horizontal transmission of the fungal strains under different growing conditions in a greenhouse. This study reports the use of a simulated model bioassay to determine the epizootic potential and horizontal transmission of *I. fumosorosea* Trinidadian strains at two different photophases.

## Materials and methods

### *Test insects*

Greenhouse whitefly pharate adults (17-18 d old; = 5<sup>th</sup> instar) were supplied by British Crop Protection Ltd., England. Prior to the assay, detached leaves (*Nicotiana tabacum* L.) were washed with sterile distilled water to remove other potential fungi contaminating the hosts and allowed to dry in a fume hood. Pharate adults were carefully removed at random from the semi-desiccated leaf surface with a probe made from a flattened hypodermic needle (Landa et al. 1994) for use in the bioassays.

### *Fungal strains*

*Isaria fumosorosea* (*Ifr*) strains T, T10 and T11 used in this study were originally collected in Trinidad and maintained at CABI BioScience (Egham, UK) on potato dextrose agar (PDA; Oxoid™) slants. Strains T and T11 were both originally isolated from *Bemisia tabaci* (Gennadius) in 1990 and 1991, respectively, and strain T10 from an unknown aphid species in 1990. *Ifr* Trinidadian strains were maintained on PDA slants at Birkbeck, University of London and later cultured at  $25 \pm 0.5$  °C on PDA prior to use in the experiments.

### *Liquid culture media for blastospores*

Liquid culture medium modified from Jackson et al. (1997) was used to produce the blastospores of *Ifr* strains. The liquid culture media contained the following ingredients per liter:  $\text{KH}_2\text{PO}_4$ , 2 g;  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ , 0.4 g;  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 0.3 g;  $\text{Fe}_2\text{SO}_4 \cdot 7\text{H}_2\text{O}$ , 50 mg;  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ , 37 mg;  $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ , 16 mg;  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 14 mg; glucose, 80 g and Casamino acids, 13.2 g. To supply the required vitamins for the blastospore growth (Jackson et al. 1997), a 1 mL aliquot of *Aspergillus* vitamin solution was added to the medium which contained the following: biotin stock, 15 mL, nictinomide, 100 mg, pyridoxin, 250 mg, p-aminobenzoic acid, 200 mg, riboflavin, 100 mg, aneuren HCl, 0.5 mg, and pantothenic acid, 200 mg per liter of solution. Lastly, glucose was added to the liquid culture media stock, adjusted to a pH of 5.5 by the addition of HCl, autoclaved and then combined with the sterile *Aspergillus* vitamin stock solution.

### *Ifr conidial inoculum preparation and blastospore production*

A circular plug (0.7 cm diameter) was removed from a conidial culture grown on PDA for 14 d at  $25 \pm 0.5$  °C under constant fluorescent light for each strain, i.e. conditions previously determined to be optimum for conidial production for *Ifr* on PDA solid media (Avery 2002). Each plug was transferred to a PDA slant in a glass tube and incubated at  $25 \pm 0.5$  °C light for 14 d under constant fluorescent light. These new PDA slant conidial cultures served as the inoculum for the liquid-culture media for producing blastospores. Conidial suspensions for

inoculating the submerged cultures were prepared by flooding each PDA slant with 10 mL of sterile, distilled water. Sterile glass beads (20-25 beads; 1 mm in diameter) were then added, and the tube suspension was agitated using a vortex mixer. The concentration of the inoculum for submerged cultures per strain was standardized using a hemocytometer to  $2.2 \pm 0.2 \times 10^6$  conidia  $\text{mL}^{-1}$ .

*Ifr* inoculum suspensions (10 mL) were then poured into three separate flasks containing the liquid-culture medium. Liquid-cultures (100 mL in 250 ml Erlenmeyer flasks) were incubated at 25 °C on an orbital shaker (Gallenkamp™) at 140 rpm for 4 d and the pH was not buffered during culture growth. After incubation, each liquid culture broth was filtered through a double layer of sterile muslin to separate any fungal hyphae from the blastospores.

From each replicate filtered culture, a 30  $\mu\text{L}$  pipette sample was spread with a bent sterile glass rod onto two separate PDA plates to determine the purity of each culture. All PDA plates were cultured at  $25 \pm 0.5$  °C under constant fluorescent light for 2-3 d and observed to determine if contaminated with other fungi. Pure blastospore suspension per *Ifr* strain (based on observed plate culture) was chosen as the source of inoculum for producing the cadavers at both photoperiod regimes. The mean number  $\pm$  SEM of viable blastospores  $\text{mL}^{-1}$  used to inoculate the whitefly pharate adult hosts for all *Ifr* strains produced in each flask was  $1.8 \pm 0.13 \times 10^7$  (Avery et al. 2004).

#### *Cadaver conidial production*

Filtered blastospore droplets ( $\sim 2$   $\mu\text{L}$  per drop) of each *Ifr* strain suspended in 0.01% Triton X-100 were placed on four sterile glass slides using an inoculating loop as described by Avery et al. (2004). Pharate adults of similar size were randomly removed from one of the two semi-desiccated tobacco leaves, placed in the middle of each suspension drop on the slide and allowed to dry in the fume hood. After the drops had dried, each slide containing 10 pharate adults were placed inside a sterile plastic Petri dish (100 x 15 mm) containing PDA and sealed using Parafilm®. The PDA in the Petri dishes maintained a high relative humidity (RH  $\sim 100\%$ ) for the duration of the assay. Each Petri dish was then placed in a growth chamber, and the assays were incubated at  $25 \pm 0.5$  °C under a 24 h photophase. After 8 d of incubation, 5 sporulating cadavers for each strain were selected and scraped from a slide with a sterile razor blade and vortexed in 1 mL of 0.01% Triton X-100 (v/v) for 60 s. Ten individual aliquots (10  $\mu\text{L}$  each) were taken from each suspension and the average conidial concentration per cadaver per *Ifr* strain was determined using a hemocytometer. The mean number  $\pm$  SEM of conidia  $\text{mL}^{-1}$  produced per cadaver for strain T, T10 and T11 was  $11.6 \pm 3.2$ ,  $10.3 \pm 1.4$  and  $3.3 \pm 0.5 \times 10^3$ , respectively.

#### *Model bioassay protocol*

*Ifr*-infected pharate adult cadavers used for these studies were produced as described above. The dried inoculated pharate adults were then carefully transferred to the center of each sterile (alcohol swabbed) laminated graph paper square (5 x 5 mm) using the flattened hypodermic needle and adhered to the surface by a drop of water. Four healthy un-treated pharate adults were gently removed from a semi-desiccated leaf and placed surrounding the central inoculated pharate adult (cadaver) according to the specific grid arrangement (Figure 1). Grid arrangement A, B and C, simulated high, medium and low infestation levels on a leaf surface, respectively.

Each complete square (5 total insect hosts including the central cadaver) was carefully laid onto a sterile V-shaped glass rod previously placed on PDA agar in a plastic Petri dish (100 x 15 mm) and then sealed with Parafilm<sup>®</sup>. The glass rod was used to elevate the plastic square above the PDA so as to minimize contamination of other possible fungal pathogens present on the susceptible insect hosts. The sealed dishes were incubated at  $25 \pm 0.5$  °C in growth chambers under a 16 and 24 h photophase. Each dish which contained the completed grid with insect hosts surrounding a central cadaver resting on the glass rod represented a model bioassay.

At assessment of each bioassay, Petri dish lids were carefully removed and the total number of hosts (including exuviae or eclosed adults) colonized by the fungus was recorded 7, 12, 14 and 21 d post-treatment. Colonized hosts were defined as producing conidia. To enhance observation of the transparent hyphal growth across the laminated surface, a colored plastic Petri dish was placed under the bioassay plate. The number of hosts colonized at both photoperiods was observed and recorded during the same period. All experiments were conducted using random samples of pharate adults from a single batch of whiteflies on two different leaves and tests under different photoperiods were conducted simultaneously. The different densities in the bioassay per *Ifr* strain tested were replicated 3 times and the experiment was repeated under a 16 and 24 h photophase.

#### Statistical analysis

Results were subjected to a three-way RMANOVA ( $\alpha = 0.05$ ) factorial analysis to determine the effect of time, fungal strain, host distribution and photoperiod on host colonization. If no interaction was determined, then the mean percent hosts colonized between strains and infestation levels were separated using a one-way ANOVA + Tukey HSD test ( $\alpha = 0.05$ ) after being arcsine transformed. Where no significant differences in percent colonization were found between the strains at the different densities over time, the data were pooled and re-analyzed. Percent of non-treated hosts colonized at different infestation levels per bioassay over time were arcsine transformed prior to being analyzed using a repeated measures (RM) ANOVA ( $\alpha = 0.05$ ). To determine the relationships between host distance from the central cadaver and percent colonization, data were submitted to regression analyses and compared at both photoperiods. These statistical tests were conducted using PROC GLM procedures of SAS WIN\_PRO platform (SAS Institute Inc., Cary, NC, 1999-2001).

Using Least Squares Dummy Regression Analysis (LSDR) we tested the null hypothesis:  $H_0: B_{11} = B_{12} = 0$ . That is, the regression coefficients for distance predicting colonization would be the same across the two photoperiods. For this analysis, distance was coded 1 for the 16:8 LD photoperiod and 2 for the 24:0 LD photoperiod. The two conceptual regression models were as follows:

Model 1:  $y_1 = \beta_{01} + \beta_{11} x$  for  $x \leq 24:0$  LD .

Model 2:  $y_2 = \beta_{02} + \beta_{12} x$  for 16:8 LD  $x \geq 24:0$  LD .

Where

$y_1$  = photoperiod 1 (16:8 LD)

$y_2$  = photoperiod 2 (24:0 LD)

$x$  = Distance

These statistical tests were conducted using the PROC REG and 'TEST' statements in SAS for Windows version 9.1 (SAS Institute Inc., Cary, NC, 1999-2001).

## Results

### *Effect of photoperiod regimes and fungal strains on colonization of insect hosts*

Percent of hosts colonized at each infestation level for both photoperiod regimes increased significantly over time ( $F = 3.53$ ;  $df = 3, 215$ ;  $P = 0.0170$ ). A significant interaction effect of host distribution on the percent colonization was observed amongst all the levels ( $F = 4.74$ ;  $df = 2, 215$ ;  $P < 0.0001$ ) and, when comparing infestation levels, host colonization was significantly greater at the highest level ( $F = 34.9$ ;  $df = 2, 215$ ;  $P < 0.0001$ ) compared to the medium or the lowest level after 7 days (Table 1). No differences ( $F = 0.27$ ;  $df = 2, 53$ ;  $P = 0.767$ ) were observed between the performance of the fungal strains T, T10 and T11 on the percent colonization at either photoperiod regime for 7, 12, 14 and 21 days post treatment.

The percent of hosts colonized 21 d post-treatment under constant light for the highest infestation level (mean  $\pm$  SEM:  $100 \pm 0.0$ ) was 42% and 61% higher ( $F = 6.21$ ;  $df = 2, 26$ ;  $P = 0.0017$ ) compared to the medium ( $58 \pm 12.5$ ) and lowest ( $39 \pm 11.1$ ) level which were similar, respectively (Table 2). Under a 16 h photophase, the percent hosts colonized 21 d post treatment for the highest infestation level (mean  $\pm$  SEM:  $92 \pm 4.2$ ) was 70% and 67% higher compared to the medium ( $22 \pm 5.7$ ) and lowest ( $25 \pm 8.3$ ) level which were similar, respectively.

### *Effect of time, infestation level and photoperiod regimes on colonization of hosts*

Colonization of the whitefly hosts by all fungal treatments was significantly affected by time (RMANOVA:  $F = 21.78$ ;  $df = 3, 211$ ;  $P < 0.0001$ ), infestation level ( $F = 84.18$ ;  $df = 2, 211$ ;  $P < 0.0001$ ) and photoperiod ( $F = 134.34$ ;  $df = 1, 211$ ;  $P < 0.0001$ ). At all infestation levels, colonization increased over time ( $F = 4.74$ ;  $df = 6, 211$ ;  $P = 0.0002$ ) and the longer photoperiod (24 h) had a significant positive effect ( $F = 3.53$ ;  $df = 1, 211$ ;  $P = 0.0170$ ) compared to the shorter photoperiod (16 h) over the 21 d observation period.

Linear regression slopes of percent colonization of the susceptible pharate hosts were significant at 16 h ( $F = 26.6$ ;  $df = 1, 26$ ;  $P < 0.0001$ ) and 24 h ( $F = 20.1$ ;  $df = 1, 26$ ;  $P < 0.0001$ ) 21 d post treatment (Figure 2). As the distance from the sporulating cadaver and the surrounding susceptible hosts increased, the percentage of hosts colonized significantly decreased at both 16 h ( $R^2 = -0.6297$ ;  $P < 0.001$ ) and 24 h ( $R^2 = -0.8326$ ;  $P < 0.001$ ) photoperiod regimes. Regression estimates suggest that distance is a stronger predictor of colonization for the 24:0 photoperiod (slope =  $-4.32046$ ) than for the 16:8 LD photoperiod (slope =  $-4.71409$ ). Results from the F test confirmed a significant difference between the slopes ( $F=18.50$ ,  $p = 0.0007$ ) with the 24 h photophase having a higher percent of susceptible hosts colonized at all infestation levels compared to the 16 h photoperiod. The null hypothesis that the two slopes were identical was rejected, confirming that the slopes do indeed differ significantly across the 2 photoperiods.

## Discussion

### *Epizootic potential of fungal strains against whitefly hosts*

In this study, potential hosts surrounding the central sporulating cadaver were colonized either as pharate adults or exuviae with eclosed adults. Sometimes *Ifr* hyphae were observed to entangle the eclosing adults by surrounding the nymphal casing (exuviae) with a netted mycelial mass and then sporulate. Wraight et al. (1998) observed postmortem hyphal growth and sporulation of *Ifr*



(20°C and 100% RH) rapidly covered the dead whitefly host and extended several millimeters onto the surrounding leaf surface.

In the distribution studies, with a single pharate adult being the only major nutrient resource for the fungus, the *Ifr* hyphae were able to colonize some of the surrounding hosts or host exuviae with eclosed adults located 21 mm away. However, honeydew transferred with the initially infected nymphs may also have provided a secondary carbon resource for the fungal hyphae. Fokkema et al. (1983) noted that aphid honeydew stimulated the growth of various saprophytic fungi on the wheat leaf phylloplane.

Colonization of healthy whitefly hosts may have been influenced by possible auto-dissemination of spores by the eclosed adults (range of eclosed adults per density: 0-50%) while inside each bioassay chamber. Auto-dissemination of *Ifr* PFR 97 spores by whitefly adults was recorded in greenhouse conditions on cucumber plants (Bohatá & Landa 2004). However, in this study, the auto-dissemination effect would be similar because the percent eclosion of the pharate adults after 21 d exposure to 16 h ( $10.9 \pm 2.90$ ) or 24 h ( $13.9 \pm 7.35$ ) photophases were similar for the different infestation levels. Once whiteflies eclosed, regardless of the photophase, they were observed to be impeded from flying freely and immediately became entangled in the infective fungal mycelium extending away from the central cadaver. Also, after analyzing the percent colonization over time, a consistent increase was found for all infestation levels at both photoperiod regimes.

The greatest distance infective hyphae spread from the central cadaver to susceptible hosts across the artificial surface was 21 mm. Wraight et al. (1998) indicated that the ability of *Ifr* hyphae to grow extensively over a leaf surface under humid conditions is a characteristic that certainly enhances its rapid spread through whitefly populations and its ability to cause epizootics. The increase in photophase from 16 to 24 h resulted in greater colonization of the whitefly hosts with *Ifr* conidia regardless of fungal strain. In contrast, *Ifr* Trinidadian conidial treatments cultured on PDA and applied against pharate whitefly adults on glass slides, indicated that the infection and colonization rates were similar under either photophase (Avery et al. 2004). The increased virulence may have resulted from the fungal strains being initially passaged through the central cadaver; however, this hypothesis was not verified in the present study. In addition to fungal virulence, an increased photophase has also been demonstrated to have a positive effect on the infection rate, conidiation and development of *Ifr* (Sakamoto et al. 1985; de la Torre & Cárdenas-Cota 1996; Avery et al. 2004).

In this model system, by varying distances healthy whitefly hosts were located from the central sporulating cadaver, it was possible to simulate high, medium and low infestation levels found on a leaf surface. However, this model system is limited and cannot account for behavioral parameters of the whitefly or plant characteristics that could affect the horizontal transmission of the fungus. For instance, in this study, whiteflies were held under optimum conditions for fungal development (Avery 2002), which may not reflect greenhouse or field environmental conditions due to the microclimatic conditions present on the leaf surface. In tritrophic interactions, parameters not included were the possible effects trichome type and density or allelochemicals found on the leaf surface on the growth of *Ifr*. In a similar study with greenhouse whitefly hosts, *Ifr* strains were not inhibited by the presence of exudates from the trichomes of *Pelargonium* cultivars (Avery 2002). However, Vega et al. (1997), and Lacey and Mercadier (1998) found that *Ifr* growth was inhibited by selected allelochemicals or secondary compounds commonly found

on the surface of leaves. In addition, Poprawski et al (2000) tested similar allelochemicals against conidial suspensions of both *Ifr* and *B. bassiana*. Perhaps these allelochemicals could be spread on the laminated surface of this model bioassay to determine what effect they may have on the potential infectivity, virulence and horizontal transmission of *Ifr* or other entomopathogenic fungi. Lastly, whitefly nymphs usually feed on the underside of the leaf surface, therefore the honeydew could be removed naturally from the phylloplane which may affect the colonization rate of *Ifr*. The leaf model bioassay was designed simply to determine the effect of photoperiod on *Ifr* strains and whether the hyphae from one infected insect (cadaver) could colonize surrounding hosts at different spatial arrangements simulating different infestation levels under optimum conditions of temperature and humidity.

The choice of a 24 h photophase used in this model bioassay was based on previous data showing that *Ifr* Trinidadian strains grew faster under constant light compared to a shorter photophase (Avery 2002, Avery et al. 2004). In addition, certain plants such as pepper plants grown under continuous lighting (24 h) have resulted in earlier flowering and fruiting giving better yields under greenhouse conditions (Demers and Gosselin 1999). Therefore in this model, the choice of comparing the virulence and horizontal transmission of *Ifr* during either a 24 or 16 h photoperiod may have some practical application. For instance, although the colonization was faster under constant light, the end result after 21 days was not significantly different. Therefore, in greenhouse production, the added expense incurred for increasing the photoperiod to supplement the daylight hours is not warranted according to this model. The same result could be obtained at a shorter photophase over time and theoretically save the grower the added expense of using more electricity for lights in the greenhouse. However, this model is based on laboratory data and needs to be confirmed under greenhouse conditions.

#### *Effect of environmental conditions and host distribution on transmission efficiency*

Many authors have indicated that epizootics are dependent upon various abiotic and biotic factors which include environmental conditions and population distribution. For example, in a laboratory experiment using hypocrealean under optimum growing conditions, conidia transfer between inoculated *B. bassiana* and untreated spruce bark beetles at different ratios resulted in 96% mortality at a ratio of 1:1 and 90% mortality at a ratio of 1:2 (Kreutz et al. 2004). In addition at ratios of 1:5, 1:10 and 1:20 the mortalities after 7 days were 83, 77 and 75%, respectively. In soil experiments, horizontal transmission efficiency of infection by Hypocreales was correlated with the distribution of the pest population (Keller et al. 1997; Long et al. 2000a, b).

In this study, the colonization of whitefly hosts was positively correlated to the distribution of the population under optimum environmental conditions. The number of susceptible hosts colonized after 21 days was greatest when nearest to the sporulating cadaver. It was observed that host colonization occurred 61 and 67 % faster at the high infestation level than with the low level under a 24 and 16 hour photoperiod regime, respectively. By decreasing the distance susceptible hosts are located from the cadaver, the contact time by hyphae from the infected host is potentially shortened and subsequently the transmission efficiency should increase.

Transmission efficiencies of *Ifr* thus appear to be dependent upon host distribution under high humidity. However, this was a model system and it does not take into account all the parameters involved in epizootics. In closed production systems, e.g. greenhouses, where optimum growing conditions for *Ifr* can be provided by short periods of misting, secondary cycling of the fungus

may occur as well. Vega et al. (1999) found that the *Ifr* blastospore formulation can germinate in 6-8 h under high humidity conditions. *Ifr* blastospore products, PFR 97™ 20% WDG (Certis, USA) and PreFeRal® WG, (BioBest, Belgium) are both recommended to be most effective when application is initiated before or at the first sign of insect presence. Also, these products are recommended to be applied when populations of whitefly are low in ‘hot spots’ before high infestations occur. Therefore, based on this model system, with only one cadaver infected with *Ifr*, at the lowest infestation level, the fungal hyphae were still able to infect the other whitefly hosts 21 mm away. This implies that transmission efficiency and secondary cycling can occur at very low infestation levels with *Ifr* under high humidity conditions readily available in a greenhouse. Commercial growers could remediate with *Ifr* at this low infestation level with potential infection and secondary cycling occurring for the next whitefly generation.

#### *Model systems as tools for predicting fungal epizootics*

Most modeling on transmission efficiency has dealt with entomopathorean fungi (Brown and Hasibaun 1995; Oduor et al. 1997; Verghese and Sreedevi 2006); however, similar work for hypocrealean-host relationships is both lacking and needed. By using model systems, such as those used in this study, the likelihood of epizootics developing may be more accurately predicted for a specific fungus, given certain environmental and population parameters. In addition, this model bioassay can be used for simulating and predicting the virulence and horizontal transmission of *Ifr* or other entomopathogenic fungi against aleyrodid species or insects with a sessile developmental stage prior to application in the field. This novel model bioassay technique is the first to evaluate the epizootic potential and determine the horizontal transmission efficiency of blastospores of *I. fumosorosea* Trinidadian strains under optimal environmental conditions at different photoperiods.

#### **Acknowledgements**

We thank the technical assistance of Cindy Avery, Jenny Smale, and Birkbeck, University of London, School of Biological Sciences, Malet Street, London WC1E 7HX, UK for the use of the laboratory space and facilities. Thanks goes to Drs. Penelope Smith and Paul Bridge at CABI BioScience, UK, for making available the Trinidadian fungal strains of *I. fumosorosea*. We thank the Biological Crop Protection Ltd, Acorn Nurseries, Chapel Lane, West Wittering, Chichester, PO20 8QG, UK for supplying the greenhouse whitefly pharate adults. We also thank Drs. S. Wraight, V. Wekesa and S. Arthurs for their suggestions and comments on an early draft of this paper.

## References

- Alma CR, Goettel MS, Roitberg BD, Gillespie DR. 2007. Combined effects of the entomopathogenic fungus, *Paecilomyces fumosoroseus* Apopka-97, and the generalist predator, *Dicyphus hesperus*, on whitefly populations. *BioControl* 52: 669-681.
- Antonious GF, Snyder JC 1995. Pirimiphos-methyl residues and control of greenhouse whitefly (Homoptera: Aleyrodidae) on seven vegetables. *Journal of Entomological Science* 30: 191-201.
- Arthurs S, Thomas MB. 2001a. Effect of dose, pre-mortem host incubation temperature and thermal behaviour on host mortality, mycosis and sporulation of *Metarhizium anisopliae* var. *acridum* in *Schistocerca gregaria*. *Biocontrol Science and Technology* 11: 411-420.
- Arthurs S., Thomas MB. 2001b. Effects of temperature and relative humidity on sporulation of *Metarhizium anisopliae* var. *acridum* in mycosed cadavers of *Schistocerca gregaria*. *Journal of Invertebrate Pathology* 78: 59-65.
- Avery PB. 2002. Tritrophic interactions among *Paecilomyces fumosoroseus*, *Encarsia formosa* and *Trialeurodes vaporariorum* on *Phaseolus vulgaris* and *Pelargonium* spp. Ph.D. Dissertation, Birkbeck, University of London, Department of Biological Sciences.
- Avery PB, Faull J, Simmonds MSJ. 2004. Effect of different photoperiods on the growth, infectivity and colonization of Trinidadian fungal strains and spore types of *Paecilomyces fumosoroseus* on the greenhouse whitefly, *Trialeurodes vaporariorum*, using a glass slide bioassay. *Journal of Insect Science* 4: 38, 10pp. Available online: [insectscience.org/4.38](http://insectscience.org/4.38)
- Bi JL, Toscano NC. 2007. Current status of the greenhouse whitefly, *Trialeurodes vaporariorum*, susceptibility to neonicotinoid and conventional insecticides on strawberries in Southern California. *Pest Management Science* 63: 747-752.
- Bohatá A, Landa Z. 2004. Influence of humidity on efficacy of entomopathogenic fungi-case study “*Paecilomyces fumosoroseus*-*Trialeurodes vaporariorum*”. *Collection of Scientific Papers of Faculty of Agriculture in České Budějovice Series for Crop Sciences* 21: 167-171.
- Brown GC, Hasibuan R. 1995. Conidial discharge and transmission efficiency of *Neozygites floridana*, an entomopathogenic fungus infecting two-spotted spider mites under laboratory conditions. *Journal of Invertebrate Pathology* 65: 10-16.
- Carruthers RI, Soper RS. Fungal diseases. 1987. In: Fuxa JR, Tanada Y, editors. *Epizootiology of insect diseases*. New York: Wiley. pp 357-416.
- Castañé C, Albajes R. 1992. *Pelargonium* cultivar selection by adults of greenhouse whitefly (Homoptera: Aleyrodidae). *Environmental Entomology* 21: 269-275.

Cuthbertson AGS, Walters KFA. 2005. Pathogenicity of the entomopathogenic fungus, *Lecanicillium muscarium*, against the sweet potato whitefly *Bemisia tabaci* under laboratory and glasshouse conditions. *Mycopathologia* 160: 315-319.

Cuthbertson AGS, Blackburn LF, Northing P, Luo W, Cannon RJC, Walters KFA. 2008. Further compatibility tests of the entomopathogenic fungus *Lecanicillium muscarium* with conventional insecticide products for control of sweet potato whitefly, *Bemisia tabaci* on poinsettia plants. *Insect Science* 15: 355-360.

de la Torre M, Cárdenas-Cota HM. 1996. Production of *Paecilomyces fumosoroseus* conidia in submerged culture. *Entomophaga* 41: 443-453.

de la Rosa W, Segura HR, Barrera JF, Williams T. 2000. Laboratory evaluation of the impact of entomopathogenic fungi on *Prorops nasuta* (Hymenoptera: Bethyridae), a parasitoid of the coffee berry borer. *Environmental Entomology* 29: 126-131.

Demers, D-A, Gosselin A. 1999. Supplemental lighting of greenhouse vegetables: limitations and problems related to long photoperiods. *ISHS Acta Horticulturae* 481: 469-473.

Fokkema NJ, Riphagen I, Poot RJ, de Jong, C. 1983. Aphid honeydew, a potential stimulant of *Cochliobolus sativus* and *Septoria nodorum* and the competitive role of saprophytic mycoflora. *Transactions of the British Mycological Society* 81: 355-363.

Fransen JJ. 1993. Integrated pest management in glasshouse ornamentals in the Netherlands: a step by step policy. *IOBC/WPRS Bulletin* 16: 35-38.

Galani G, Almasan L. 1984. Detection of a natural fungal epizootic in the whitefly, *Trialeurodes vaporariorum* Westwood. *Buletinul de Protecția Plantelor* 1: 33-36.

Gökçe A, Er MK. 2005. Pathogenicity of *Paecilomyces* spp. to the glasshouse whitefly, *Trialeurodes vaporariorum*, with some observations on the fungal infection process. *Turkish Journal of Agriculture and Forestry* 29: 331-339.

Hall RA. 1984. Epizootic potential for aphids of different isolates of the fungus, *Verticillium lecanii*. *Entomophaga* 29: 311-321.

Jackson MA, McGuire MR, Lacey LA, Wraight SP. 1997. Liquid culture production of desiccation tolerant blastospores of the bioinsecticidal fungus *Paecilomyces fumosoroseus*. *Mycological Research* 101: 35-41.

Keller S, Schweizer S, Keller E, Brenner H. 1997. Control of white grubs (*Melolontha melolontha* L.) by treating adults with the fungus *Beauveria brongniartii*. *Biocontrol Science and Technology* 7: 105-116.

Kreutz J, Zimmermann G, Vaupel O. 2004. Horizontal transmission of the entomopathogenic fungus *Beauveria bassiana* among the spruce bark beetle, *Ips typographus* (Col., Scolytidae) in the laboratory and under field conditions. *Biocontrol Science and Technology* 14: 837-848.

Lacey LA, Fransen JJ, Carruthers R. 1995. Global distribution of naturally occurring fungi of *Bemisia*, their biologies and use as biological control agents. In: Gerling D, and R. Mayer R, editors. *Bemisia* 1995: Taxonomy, Biology, Damage Control and Management. Hants, UK: Intercept Ltd. pp 401-433.

Lacey, LA, Mercadier G. 1998. The effect of selected allelochemicals on germination of conidia and blastospores and mycelial growth of the entomopathogenic fungus, *Paecilomyces fumosoroseus* (Deuteromycotina: Hyphomycetes). *Mycopathologia* 142: 17-25.

Landa Z, Osborne L, Lopez F, Eyal J. 1994. A bioassay for determining pathogenicity of entomogenous fungi on whiteflies. *Biological Control* 4: 341-350.

Landa Z, Bohatá A. 1999. Compatibility of entomogenous fungus *Paecilomyces fumosoroseus* with natural insecticides based on azadirachtin and Neem oil. *Collection of Scientific Papers Faculty of Agriculture in Ceske Budejovice Series for Crop Sciences* 16: 99-105.

Long DW, Groden E, Drummond FA. 2000a. Horizontal transmission of *Beauveria bassiana* (Bals.) Vuill. *Agricultural and Forest Entomology* 2: 11-17.

Long, D.W., Drummond, F.A, Groden, E., and Donahue, D. W. 2000b. Modeling *Beauveria bassiana* horizontal transmission. *Agriculture and Forest Entomology* 2: 19-32.

Luz C, Fargues J. 1997. Temperature and moisture requirements for conidial germination of an isolate of *Beauveria bassiana*, pathogenic to *Rhodnius prolixus*. *Mycopathologia* 138: 117-125.

Luz C, Fargues J. 1999. Dependence of the entomopathogenic fungus, *Beauveria bassiana*, on high humidity for infection of *Rhodnius prolixus*. *Mycopathologia* 146: 33-41.

Luz C, Silva IG, Cordeiro CM, Tigano MS. 1999. Sporulation of *Beauveria bassiana* on cadavers of *Triatoma infestans* after infection at different temperatures and doses of inoculum. *Journal of Invertebrate Pathology* 73: 223-225.

Meekes ETM, Voorst, SV, Joosten NN, Fransen JJ, van Lenteren JC. 2000. Persistence of the fungal whitefly pathogen, *Aschersonia aleyrodis*, on three different plant species. *Mycological Research* 104: 1234-1240.

Meekes ETM, Fransen JJ, van Lenteren JC. 2002. Pathogenicity of *Aschersonia* spp. against whiteflies *Bemisia argentifolii* and *Trialeurodes vaporariorum*. *Journal of Invertebrate Pathology* 81: 1-11.

Mier T, Rivera F, Bermudez JC, Dominquez Y, Benavides C, Ulloa M. 1991. First report in Mexico on the isolation of *Verticillium lecanii* from whitefly and *in vitro* pathogenicity tests on this insect. *Revista Mexicana de Micologia* 7: 149-156.

Oduor, G. I., Sabelis, M. W., Lingeman, R., De Moraes, G. J., Yaninek, J. S. 1997. Modelling fungal (*Neozygites cf. floridana*) epizootics in local populations of cassava green mites (*Mononychellus tanajoa*). *Experimental and Applied Acarology* 21, 485-506.

Omer AD, Leigh TF, Granett J. 1992. Insecticide resistance in field populations of greenhouse whitefly (Homoptera: Aleyrodidae) in the San Joaquin Valley, California cotton cropping system. *Journal of Economic Entomology* 85: 21-27.

Pell JK. 2008. Ecological approaches to pest management using entomopathogenic fungi; concepts, theory, practice and opportunities. In: Ekesi S, Maniana NK, editors. Use of entomopathogenic fungi in biological pest management. Kerala, India: Research Signpost. pp 145-178.

Poprawski TJ, Greenburg SM, Ciomperlik MA. 2000. Effect of host plant on *Beauveria bassiana*- and *Paecilomyces fumosoroseus*-induced mortality of *Trialeurodes vaporariorum* (Homoptera: Aleyrodidae). *Environmental Entomology* 29: 1048-1053.

Poprawski TJ, Jones WJ. 2000. Host plant effects on activity of the mitosporic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* against two populations of *Bemisia* whiteflies (Homoptera: Aleyrodidae). *Mycopathologia* 151: 11-20.

Saito T. 2005. Preliminary experiments to control the silverleaf whitefly with electrostatic spraying of a mycoinsecticide. *Applied Entomology and Zoology* 40: 289-292.

Sakamoto M, Inoue Y, Aoki J. 1985. Effect of light on the conidiation of *Paecilomyces fumosoroseus*. *Transactions of the Mycological Society of Japan* 26: 499-509.

Sanderson JP, Rousch RT. 1992. Monitoring insecticide resistance in greenhouse whitefly (Homoptera: Aleyrodidae) with yellow sticky cards. *Journal of Economic Entomology* 85: 634-641.

Shah PA, Pell JK. 2003. Entomopathogenic fungi as biocontrol agents. *Applied Microbiology and Biotechnology* 61: 413-423.

Thomas MB, Wood SN, Lomer CJ. 1995. Biological control of locusts and grasshoppers using a fungal pathogen: the importance of secondary cycling. *Proceedings of the Royal Society of London* 259: 265-270.


- Vega FE, Jackson MA, McGuire MR. 1999. Germination of conidia and blastospores of *Paecilomyces fumosoroseus* on the cuticle of the silverleaf whitefly, *Bemisia argentifolii*. *Mycopathologia* 147: 33-35.
- Vega FE, Dowd PF, McGuire MR, Jackson MA, Nelsen TC. 1997. *In vitro* effects of secondary plant compounds on germination of blastospores of the entomopathogenic fungus *Paecilomyces fumosoroseus* (Deuteromycotina: Hyphomycetes). *Journal of Invertebrate Pathology* 70: 209-213.
- Verghese A, Sreedevi K. 2006. Epizootics of an entomopathogenic fungus *Entomophthora aphidis* Hoggman on the pomegranate aphid *Aphis punicae passerine* (Homoptera: Aphididae). *Communications in Agriculture and Applied Biological Science* 71: 47-52.
- Vidal C, Fargues J, Rougier M, Smits N. 2003. Effect of air humidity on the infection potential of hyphomycetous fungi as mycoinsecticides for *Trialeurodes vaporariorum*. *Biocontrol Science and Technology* 13: 183-198.
- Vidal C, Fargues J. 2008. Climatic constraints for fungal biopesticides. In: Ekesi S, Maniana NK, editors. *Use of entomopathogenic fungi in biological pest management*. Kerala, India: Research Signpost. pp 39-56.
- Wang KY, Kong XB, Jiang XY, Yi MQ, Liu TX. 2003. Susceptibility of immature and adult stages of *Trialeurodes vaporariorum* (Hom., Aleyrodidae) to selected insecticides. *Journal of Applied Entomology* 127: 527-533.
- Wraight SP, Carruthers RI, Bradley CA, Jaronski ST, Lacey LA, Wood P, Galaini-Wraight S. 1998. Pathogenicity of the entomopathogenic fungi *Paecilomyces* spp. and *Beauveria bassiana* against the silverleaf whitefly, *Bemisia argentifolii*. *Journal of Invertebrate Pathology* 71: 217-226.
- Wraight SP, Carruthers RI, Jaronski ST, Bradley CA, Garza CJ, Wraight GS. 2000. Evaluation of the entomopathogenic fungi *Beauveria bassiana* and *Paecilomyces fumosoroseus* for microbial control of the silverleaf whitefly, *Bemisia argentifolii*. *Biological Control* 17: 203-217.



## Figures

Figure 1. Distribution of infected (solid circles) and susceptible (open circles) *Trialeurodes vaporariorum* hosts and characteristics for the various grid arrangements A, B and C observed under 16:8 and 24:0 hour LD photoperiod regimes. Each unit square measures 5 x 5 mm.

Figure 2. Correlation between the distance of surrounding susceptible *Trialeurodes vaporariorum* hosts from the sporulating cadaver and the percentage colonized at different photoperiods when incubated at  $25 \pm 0.5$  °C with ~100 RH. There are 3 data points (1 per strain) for the varying distances from the cadaver represented, although some are obscured due to similar values.



Grid arrangement	A	B	C
Infestation level	High	Medium	Low
Distance from cadaver (mm)	7.1	14.1	21.2

**Figure 1**




Figure 2

Table 1. Effect of photophase (hr), distance from the cadaver and Trinidadian fungal strains on the percent colonization of susceptible *Trialeurodes vaporariorum* (GW) hosts over time under laboratory conditions.

Main Effect	n	Mean % GW hosts colonized <sup>a</sup> /days post-treatment			
		7	12	14	21
Photophase (hr)					
16	27	0.0a	17.6a	19.4a	44.0a
24	27	45.4b	51.9b	57.4b	65.7b
distance from the cadaver <sup>b</sup>					
7	18	30.6a	65.3b	72.2b	95.8b
14	18	20.8a	22.2a	25.0a	36.8a
21	18	16.6a	16.7a	18.1a	31.9a
Fungal strain					
T	18	30.6a	40.3a	44.4a	62.5a
T10	18	19.4a	33.3a	36.1a	54.9a
T11	18	18.1a	30.6a	34.7a	47.2a

<sup>a</sup>Means followed by the same letter are not significantly different (Duncan multiple range test,  $P < 0.05$ )

<sup>b</sup>Four potential hosts were located 7.1mm (high), 14.1mm (medium) and 21.2 mm (low) from a central sporulating *T. vaporariorum* cadaver.(Infestation level)

Table 2. Main effect<sup>a</sup> of host distribution on the colonization of *Trialeurodes vaporariorum* (GW) with *Isaria fumosorosea* Trinidadian strains at different photoperiods (h L:D) incubated at 25 ± 0.5 °C with ~100% RH.

Host distribution (h L:D)	Mean percentage ± SEM of GW colonized / day <sup>bc</sup>			
	7	12	14	21
High (16:8)	0 ± 0.0c	53 ± 2.8b	58 ± 4.2b	92 ± 4.2a
High (24:0)	61 ± 11.1a	78 ± 7.7a	83 ± 7.2a	100 ± 0.0a
Medium (16:8)	0 ± 0.0c	0 ± 0.0c	6 ± 6.0c	22 ± 5.7c
Medium (24:0)	42 ± 11.7ab	44 ± 12.3b	50 ± 12.5b	58 ± 12.5b
Low (16:8)	0 ± 0.0c	0 ± 0.0c	0 ± 0.0c	25 ± 8.3c
Low (24:0)	33 ± 10.2b	33 ± 10.2b	36 ± 9.4b	39 ± 11.1cb

<sup>a</sup>The performance data for all *I. fumosorosea* strains for both photoperiods and distances from the cadaver were found similar (MANOVA, DMRT,  $P > 0.05$ ) and were therefore combined prior to determining the mean percentage of GW colonized per day.

<sup>b</sup>Four potential hosts were located 7.1, 14.1 and 21.2 mm from a central sporulating GW cadaver for the highest, medium and lowest densities, respectively. (n = 9 replicates for all densities).

<sup>c</sup>Actual untransformed mean values in a column followed by different letters among the densities per hour (h) LD photoperiod are significantly different (Tukey HSD test,  $P < 0.05$ ). Mean values were arcsine transformed prior to analysis.