



**University of  
Leicester**

**DEPARTMENT OF ECONOMICS**

**BAYESIAN SEMIPARAMETRIC INFERENCE  
IN MULTIPLE EQUATION MODELS**

**Gary Koop**

**University of Leicester**

**Dale Poirier**

**University of California at Irvine**

**Justin Tobias**

**University of California at Irvine**

**Working Paper No. 04/17**

**June 2003**

# Bayesian Semiparametric Inference in Multiple Equation Models

Gary Koop  
Department of Economics  
University of Glasgow  
G.Koop@socsci.gla.ac.uk

Dale Poirier  
Department of Economics  
University of California at Irvine  
dpoirier@uci.edu

and

Justin Tobias  
Department of Economics  
University of California at Irvine  
jtobias@uci.edu

June 19, 2003

**KEYWORDS:** nonparametric regression, nonparametric instrumental variables, SUR model, endogeneity, nonlinear simultaneous equations

---

**ABSTRACT:** This paper outlines an approach to Bayesian semiparametric regression in multiple equation models which can be used to carry out inference in seemingly unrelated regressions or simultaneous equations models with nonparametric components. The approach treats the points on each nonparametric regression line as unknown parameters and uses a prior on the degree of smoothness of each line to ensure valid posterior inference despite the fact that the number of parameters is greater than the number of observations. We derive an empirical Bayesian approach that allows us to estimate the prior smoothing hyperparameters from the data. An advantage of our semiparametric model is that it is written as a seemingly unrelated regressions model with independent Normal-Wishart prior. Since this model is a common one, textbook results for posterior inference, model comparison, prediction and posterior computation are immediately available. We use this model in an application involving a two-equation structural model drawn from the labor and returns to schooling literatures.

---

# 1 Introduction

Despite the proliferation of theories for semiparametric and nonparametric regression, the use of these techniques remains relatively rare in empirical practice. Increased computational difficulty and mathematical sophistication, and perhaps most importantly, the *curse of dimensionality* - the unfortunate reality wherein the rate of convergence of the nonparametric regression estimator slows with the number of variables treated nonparametrically - all seem to provide barriers which prevent the widespread use of nonparametric techniques.

The rapid increase in computing power and growth in nonparametric routines found in statistical software packages has helped to mitigate computational concerns. To combat the curse of dimensionality problem, many researchers have adopted the use of *partially linear* or *semilinear* regression models. This model, though not fully nonparametric, provides a convenient generalization of the standard linear model which is not as susceptible to the curse of dimensionality since only one, or perhaps a few, variables are treated nonparametrically. Finally, some studies (*e.g.* Blundell and Duncan (1998), Yatchew (1999), and DiNardo and Tobias (2001)) have tried to bridge the gap between theory and practice, and make these techniques accessible to applied researchers.

In this paper we continue in this tradition, and describe and implement simple and intuitive Bayesian methods for semiparametric and nonparametric regression. Importantly, the methods we describe can be used in the context of multiple equation models, thus generalizing the scope of models for which simple nonparametric methods have been described. In our discussion, we focus primarily on the Seemingly Unrelated Regression (SUR) model. This model is of interest in and of itself, but is also of interest as the (possibly restricted) reduced form of a semiparametric simultaneous equations model (or the structural form of a triangular simultaneous equations model).

Before describing the contributions of this paper, it is useful to briefly outline the method we used in related work (*e.g.* Koop and Poirier (2003a)) in the single equation partially linear regression model. This partially linear model divides the explanatory variables into a set which is treated parametrically,  $z$ , and a set which is treated nonparametrically,  $x$ , and relates them to a dependent variable  $y$  as:

$$y_i = z_i'\beta + f(x_i) + \varepsilon_i,$$

for  $i = 1, \dots, N$  where  $f(\cdot)$  is an unknown function. Because of the curse of dimensionality,  $x_i$  must be of low dimension and is often a scalar (see Yatchew, 1998, for an excellent introduction to the partial linear model).

In this model we assumed  $\varepsilon_i \stackrel{iid}{\sim} N(0, \sigma^2)$  for  $i = 1, \dots, N$ , and all explanatory variables were fixed or exogenous. Observations were ordered so that  $x_1 < x_2 < \dots < x_N$ . Define  $y = (y_1, \dots, y_N)'$ ,  $Z = (z_1, \dots, z_N)'$  and  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_N)'$ . Letting  $\gamma = (f(x_1), \dots, f(x_N))'$ ,  $W = (Z : I_N)$  and  $\delta = (\beta', \gamma)'$ , we showed that you

could write the previous equation as:

$$y = W\delta + \varepsilon.$$

Thus, the partially linear model can be written as the standard Normal linear regression model where the unknown points on the nonparametric regression line are treated as unknown parameters. This regression model is characterized by insufficient observations in that the number of explanatory variables is greater than  $N$ . However, Koop and Poirier (2003a) showed that, if a natural conjugate prior is used, the posterior is still well-defined. In fact, we showed that the natural conjugate prior did not even have to be informative in all dimensions and that prior information about the smoothness of the nonparametric regression line was all that was required to ensure valid posterior inference. Thus, for the subjective Bayesian, prior information can be used to surmount the problem of insufficient observations. Furthermore, for the researcher uncomfortable with subjective prior information, the required amount of prior information was quite small, involving the selection of a single prior hyperparameter which we called  $\eta$  that governed the smoothness of the nonparametric regression line. In Koop and Poirier (2003b), we went even further and showed how (under weak conditions) empirical Bayesian methods could be used to estimate  $\eta$  from the data.

The advantages of remaining within the framework of the Normal linear regression model with a natural conjugate prior are clear. This model is very well understood and standard textbook results for estimation, model comparison and prediction are immediately available. Analytical results for posterior moments, marginal likelihoods and predictives exist and, thus, there is no need for posterior simulation. This means methods which search over many values for  $\eta$  (e.g. empirical Bayesian methods or cross-validation) can be implemented at a low computational cost. Furthermore, as shown in our previous work, the partial linear model can serve as a component in numerous other models which do involve posterior simulation (e.g. semiparametric tobit and probit models or the partial linear model with the errors treated flexibly by using mixtures of Normals). The ability to simplify the estimation of the nonparametric component in such a complicated empirical exercise may provide the researcher a great computational benefit.

In this paper we take up the case of Bayesian semiparametric estimation in multiple equation models, and adopt a similar approach for smoothing the regression functions. In particular, we consider the estimation of a semiparametric SUR model of the form:

$$y_{ij} = z'_{ij}\beta_j + f_j(x_{ij}) + \varepsilon_{ij}, \tag{1.1}$$

where  $y_{ij}$  is the  $i^{th}$  observation ( $i = 1, \dots, N$ ) on the  $j^{th}$  endogenous variable ( $j = 1, \dots, m$ ),  $z_{ij}$  is a  $k_j \times 1$  vector of observations on the exogenous variables which enter linearly,  $f_j(x_{ij})$  is an unknown function which depends on a vector of exogenous variables,  $x_{ij}$ , and  $\varepsilon_{ij}$  is the error term. For equations which have nonparametric components,  $z_{ij}$  does not contain an intercept since the first point on a nonparametric regression line plays the role of an intercept.

The approach we describe for the estimation of this model is simple and intuitive, and hopefully will appeal to practitioners seeking to add degrees of flexibility to their multiple equation analyses. As in our previous work,

we employ a prior which serves to smooth the nonparametric regression functions. It is important to recognize that for the (parametric) seemingly unrelated regressions model (and the reduced form of the simultaneous equations model), the natural conjugate prior suffers from well known criticisms (see Rothenberg, 1963, or Dreze and Richard, 1983). On the basis of these, Dreze and Richard (1983, page 541) argue against using the natural conjugate prior (except for certain noninformative limiting cases not relevant for our class of models). Their arguments carry even more force in the present semiparametric context since it can be shown that the natural conjugate prior places some undesirable restrictions on the way smoothing is carried out on nonparametric regression functions in different equations (i.e. the nonparametric component in each equation is smoothed in the same way). Thus, in the present paper we do not adopt a natural conjugate prior, but rather use an independent Normal-Wishart prior.

The basic ideas behind our approach are straightforward extensions of standard textbook Bayesian methods for the SUR model (see, e.g., Koop, 2003, pages 137-142). Thus, textbook results for estimation, model comparison (including comparison of parametric to nonparametric models) and prediction are immediately available. This, we argue, is an advantage relative to the relevant non-Bayesian literature (see, e.g., Pagan and Ullah, 1999, chapter 6) and to other, more complicated, Bayesian approaches to nonparametric seemingly unrelated regression such as Smith and Kohn (2000). Nevertheless, the notation for our most general semiparametric model is, admittedly, a bit messy.

We illustrate the use of our methods by estimating a two-equation simultaneous equations model in parallel with the development of our theory. This application takes data from the National Longitudinal Survey of Youth (NLSY) and involves estimating the returns to schooling, job tenure, and ability for a cross-sectional sample of white males. Our triangular simultaneous equations model has two equations, one for the (log) wage and the other for the quantity of schooling attained. After estimating standard parametric models that have appeared in the literature, we first extend them to allow for nonparametric treatment of an exogenous variable (weeks of tenure on the current job) in the wage equation (Case 1). Subsequently, we consider Case 2 where single explanatory variables enter nonparametrically in each equation. In this model we additionally allow a measure of cognitive ability to enter the schooling equation nonparametrically. We complete our empirical work with Case 3 by giving cognitive ability a nonparametric treatment in both the wage and schooling equations (with tenure on the job also given a nonparametric treatment in the wage equation).

Our results reveal the practicality and usefulness of our approach. In some cases, our semiparametric treatment yields results which are very similar to those from simple parametric nonlinear models (e.g. quadratic). However, one advantage of a semiparametric approach is that a particular functional form such as the quadratic does not have to be chosen, either in an *ad hoc* fashion or through pre-testing. Furthermore, in some cases our semiparametric approach yields empirical results that could not be easily obtained using standard parametric methods. In terms of our application, our results reveal the empirical importance of controlling for nonlinearities in ability in both the wage and schooling equations when trying to estimate the return to education.

The outline of our paper is as follows. In the next section, we outline our basic semiparametric SUR model, describe our data, and obtain parametric results and semiparametric results where job tenure is treated nonparametrically. In section 3, we describe the process of estimating a model with nonparametric components in both equations, and estimate the model in Case 2. Finally, in section 4, we describe how to handle the estimation of additive models and provide estimation results for our most general Case 3. The paper concludes with a summary in section 5.

## 2 Case 1: A Single Nonparametric Component in a Single Equation

We begin by considering a simplified version of (1.1) where a nonparametric component enters a single equation (chosen to be the  $m^{th}$  equation) and the explanatory variable which receives a nonparametric treatment,  $x_{im}$ , is a scalar. In later sections, we consider cases where several equations have nonparametric components each depending on a different explanatory variable (or variables).

We assume that the data is ordered so that  $x_{1m} < \dots < x_{Nm}$  and define  $\gamma_i = f_m(x_{im})$  for  $i = 1, \dots, N$  to be the unknown points on the nonparametric regression line in the  $m^{th}$  equation. We also let  $\gamma = (\gamma_1, \dots, \gamma_N)'$  and  $\zeta_i$  be the  $i^{th}$  row of  $I_N$ . With these definitions, we can write the model as:

$$y_i = W_i \delta + \varepsilon_i, \quad (2.1)$$

where  $y_i = (y_{i1}, \dots, y_{im})'$ ,  $\varepsilon_i = (\varepsilon_{i1}, \dots, \varepsilon_{im})'$ ,

$$W_i = \begin{bmatrix} z'_{i1} & 0 & \cdot & \cdot & \cdot & 0 \\ 0 & z'_{i2} & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 0 & z'_{i,m-1} & 0 & 0 \\ 0 & \cdot & \cdot & 0 & z'_{im} & \zeta_i \end{bmatrix}$$

and  $\delta = (\beta'_1, \dots, \beta'_m, \gamma)'$  is a  $K + N$  vector where  $K = \sum_{j=1}^m k_j$ . For future reference, we define the partition  $W_i = [W_i^{(1)} : W_i^{(2)}]$  where  $W_i^{(1)}$  is an  $m \times (K + 2)$  matrix and  $W_i^{(2)}$  is  $m \times (N - 2)$ . The likelihood for this model is defined by assuming  $\varepsilon_i \stackrel{iid}{\sim} N(0, \Sigma)$ .

We define smoothness according to second differences of points on the nonparametric regression line. In light of this, it proves convenient to transform the model. Define the  $(N - 2) \times N$  second-differencing matrix as:

$$D = \begin{bmatrix} 1 & -2 & 1 & \cdot & \cdot & \cdot & \cdot & 0 \\ 0 & 1 & -2 & 1 & 0 & \cdot & \cdot & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & 0 & \cdot & \cdot & \cdot & 1 & -2 & 1 \end{bmatrix}, \quad (2.2)$$

so that  $D\gamma$  is the vector of second differences,  $\Delta^2 \gamma_i$ . Prior information about smoothness of the nonparametric regression line will be expressed in terms of  $R\delta$ , where the  $(N - 2) \times (K + N)$  matrix  $R = [0_{(N-2) \times K} : D]$ .

For future reference, we partition  $R = [R_1 : R_2]$  where  $R_1$  is an  $(N - 2) \times (K + 2)$  matrix and  $R_2$  is  $(N - 2) \times (N - 2)$  (i.e. the nonsingular matrix  $R_2$  is  $D$  with the first two columns deleted). Note that other degrees of differencing can be handled by re-defining (2.2) as appropriate (see, e.g., Yatchew, 1998, pages 695-698 or Koop and Poirier, 2003a).

Using standard transformations (see, e.g. Poirier, 1995, pages 503-504), (2.3) can be written as:

$$y_i = V_i^{(1)}\lambda_1 + V_i^{(2)}\lambda_2 + \varepsilon_i = V_i\lambda + \varepsilon_i, \quad (2.3)$$

where  $\lambda = (\lambda'_1, \lambda'_2)'$ ,  $\lambda_1 = (\beta'_1, \dots, \beta'_m, \gamma_1, \gamma_2)'$ ,  $\lambda_2 = D\gamma$ ,  $V_i^{(1)} = W_i^{(1)} - W_i^{(2)}R_2^{-1}R_1$  and  $V_i^{(2)} = W_i^{(2)}R_2^{-1}$ . Note that  $\lambda_2$  is the vector of second differences of the points on the nonparametric regression line and it is on this parameter vector that we place our smoothness prior.

We use an independent Normal-Wishart prior for  $\lambda$  and  $\Sigma^{-1}$  which is a common choice for the parametric SUR model (see, e.g., Chib and Greenberg, 1995 or 1996). Thus,

$$\lambda \sim N(\underline{\lambda}, \underline{V}_\lambda) \quad (2.4)$$

and

$$\Sigma^{-1} \sim W(\underline{V}_\Sigma^{-1}, \underline{\nu}), \quad (2.5)$$

where  $W(\underline{V}_\Sigma, \underline{\nu})$  denotes the Wishart distribution (see, e.g., Poirier, 1995, page 136).

Our empirical work is based on a Gibbs sampler involving  $p(\lambda|y, \Sigma^{-1})$  and  $p(\Sigma^{-1}|y, \lambda)$ . Straightforward manipulations show these to be:

$$\lambda|y, \Sigma^{-1} \sim N(\bar{\lambda}, \bar{V}_\lambda) \quad (2.6)$$

and

$$\Sigma^{-1}|y, \lambda \sim W(\bar{V}_\Sigma^{-1}, \bar{\nu}), \quad (2.7)$$

where

$$\bar{\nu} = \underline{\nu} + N, \quad (2.8)$$

$$\bar{V}_\Sigma^{-1} = \left[ \underline{V}_\Sigma + \sum_{i=1}^N (y_i - V_i\lambda)(y_i - V_i\lambda)' \right]^{-1}, \quad (2.9)$$

$$\bar{V}_\lambda = \left( \underline{V}_\lambda^{-1} + \sum_{i=1}^N V_i'\Sigma^{-1}V_i \right)^{-1} \quad (2.10)$$

and

$$\bar{\lambda} = \bar{V}_\lambda \left( \underline{V}_\lambda^{-1}\underline{\lambda} + \sum_{i=1}^N V_i'\Sigma^{-1}y \right). \quad (2.11)$$

Of course, many values may be selected for the prior hyperparameters,  $\underline{\lambda}$ ,  $\underline{V}_\lambda$ ,  $\underline{V}_\Sigma^{-1}$  and  $\underline{\nu}$ . Here we describe a particular prior elicitation strategy that requires a minimal amount of subjective prior information. We assume

$$\underline{V}_\lambda = \begin{bmatrix} \underline{V}_1 & 0 \\ 0 & V(\eta) \end{bmatrix} \quad (2.12)$$

where  $\underline{V}_1$  and  $V(\eta)$  are the prior covariance matrices for  $\lambda_1$  and  $\lambda_2$ , respectively. We set  $\underline{V}_1^{-1} = 0$ , the noninformative choice. Since  $\lambda_2 = D\gamma$  is the vector of second differences of points on the nonparametric regression line,  $V(\eta)$  controls its degree of smoothness. We assume  $\underline{V}(\eta)$  depends on a scalar parameter,  $\eta$ . As discussed in Koop and Poirier (2003a), several sensible forms for  $V(\eta)$  can be chosen. In this paper, we set  $V(\eta) = \eta I_{N-2}$ . We also set  $\underline{\lambda} = 0_{K+N}$ . Note that these assumptions imply we are noninformative about  $\lambda_1 = (\beta'_1, \dots, \beta'_m, \gamma_1, \gamma_2)'$ , but have an informative prior for the remaining parameters which reflect the degree of smoothness in the nonparametric regression line. Our information about this smoothness is of the form:  $\Delta^2\gamma_i \sim N(0, \eta)$  for  $i = 3, \dots, N$ .<sup>1</sup>

In this paper we adopt an empirical Bayesian approach where  $\eta$  is chosen so as to maximize the marginal likelihood. However, it is worth noting that  $\eta$  could be treated either as a prior hyperparameter to be selected by the researcher or as a parameter in a hierarchical prior. If the latter approach were adopted,  $\eta$  could be integrated out of the posterior. Our empirical Bayesian approach is equivalent to this hierarchical prior approach using a noninformative flat prior for  $\eta$  (and plugging in the posterior mode of  $\eta$  instead of integrating out this parameter).<sup>2</sup>

The results of Fernandez, Osiewalski and Steel (1997) imply that an informative prior is required for  $\Sigma^{-1}$  in order to ensure propriety of the posterior. However, in related work with a single equation model (see Koop and Poirier, 2003b), we found that use of a proper, but relatively noninformative prior yielded sensible (and robust) results. Accordingly, we set  $\underline{\nu} = 10$  in our application. Using the properties of the Wishart distribution, we obtain the prior mean  $E(\sigma_{ij}^{-1}) = \underline{\nu} V_{\Sigma ij}^{-1}$ , where  $\sigma_{ij}^{-1}$  and  $V_{\Sigma ij}^{-1}$  are the  $ij^{th}$  elements of  $\Sigma^{-1}$  and  $\underline{V}_{\Sigma}^{-1}$ , respectively. To center the prior correctly, we calculate the OLS estimate of  $\Sigma$  based on a parametric SUR model where all variables (including  $x_{im}$ ) enter linearly. We set  $\underline{\nu} \underline{V}_{\Sigma}^{-1}$  equal to the inverse of this OLS estimate.

In order to compare models or estimate/select  $\eta$  in our empirical Bayesian approach, the marginal likelihood (for a given value of  $\eta$ ) is required. No analytical expression for this exists. However, we can estimate the marginal likelihood using Gibbs sampler output and the Savage-Dickey density ratio (see, e.g., Verdinelli and Wasserman, 1995). Define  $M_1$  to be the semiparametric SUR model given in (2.3) with prior given by (2.4) and (2.5) and a particular value for  $\eta$  selected. Define  $M_2$  to be  $M_1$  with the restriction  $\lambda_2 = 0_{N-2}$  imposed (with the same prior for  $\lambda_1$  and  $\Sigma^{-1}$ ). Using the Savage-Dickey density ratio, the Bayes factor comparing  $M_1$  to  $M_2$  can be written as:

$$BF(\eta) = \frac{p(\lambda_2 = 0 | M_1)}{p(\lambda_2 = 0 | y, M_1)}, \quad (2.13)$$

where the numerator and denominator are the prior and posterior, respectively, of  $\lambda_2$  in the semiparametric SUR model evaluated at the point  $\lambda_2 = 0_{N-2}$ . This Bayes factor may be of interest in and of itself since it

<sup>1</sup>This approach to prior elicitation does not include any information in  $x_{im}$  other than order information (i.e. data is ordered so that  $x_{1m} < \dots < x_{Nm}$ ). If desired, the researcher could include  $x_{im}$  by eliciting a prior, e.g., of the form  $\Delta^2\gamma_i \sim N(0, \eta \Delta^2 x_{im})$ .

<sup>2</sup>Further motivation for our approach can be obtained by noting that our framework is similar to a state space model and, in such a model, a parameter analogous to  $\eta$  would be the error variance in the state equation and be estimated from the data.



compares the semiparametric SUR model to a sensible parametric alternative.<sup>3</sup> However, it can also be used in an empirical Bayesian analysis to select  $\eta$ . That is, since  $\eta$  does not enter the prior for  $M_2$ ,  $BF(\eta)$  will be proportional to the marginal likelihood for the semiparametric SUR model for a given value of  $\eta$ . The empirical Bayes estimate of  $\eta$  can be implemented by running the Gibbs sampler for a grid of values for  $\eta$  and choosing the value which maximizes  $BF(\eta)$ . Alternative methods for selecting  $\eta$  include cross-validation or extensions of the reference prior approach discussed in van der Linde (2000).

Note that  $BF(\eta)$  can be calculated in the Gibbs sampler in a straightforward manner. The quantity  $p(\lambda_2 = 0|M_1)$  can be directly evaluated using the Normal prior given in (2.4), while  $p(\lambda_2 = 0|y, M_1)$  can be evaluated in the Gibbs sampler in the same way as any posterior function of interest. That is, if we define

$$\hat{p}(\lambda_2 = 0|y, M_1) = \frac{1}{S} \sum_{s=1}^S p(\lambda_2 = 0|y, \Sigma^{(s)}),$$

where  $\Sigma^{(s)}$  for  $s = 1, \dots, S$  denotes draws from the Gibbs sampler (after discarding initial burn-in draws), then

$$\hat{p}(\lambda_2 = 0|y, M_1) \rightarrow p(\lambda_2 = 0|y, M_1)$$

as  $S \rightarrow \infty$ . Note that the posterior conditional  $p(\lambda_2 = 0|y, \Sigma^{(s)})$  is simple to evaluate since it is Normal (see equation 2.6).

This semiparametric SUR model can be used as a restricted reduced form of a semiparametric simultaneous equations model and, thus, the methods described above allow for Bayesian inference in the latter model. That is, our Gibbs sampler provides us with draws from the posterior of the reduced form parameters. Provided the model is identified, the structural form parameters will be a transformation of the reduced form parameters and the draws of the latter can be transformed into draws of the former. The triangular structure of the model in our application means we do not have to adopt such an approach. However, it is useful to note that our approach can be used with more general simultaneous equations models.

Before introducing our application, we briefly discuss related (parametric) work on simultaneous equations models. The literature on Bayesian analysis of simultaneous equations models is voluminous. Dreze and Richard (1983) surveys the literature through the early 1980s, while Kleibergen (1997), Kleibergen and van Dijk (1998) and Kleibergen and Zivot (2003) are recent references. The more recent literature focusses on issues of identification and prior elicitation which are of little relevance for our work. For instance, some of this recent work discusses problems with the use of noninformative priors at points in the parameter space which imply non-identification (and show how noninformative priors based on Jeffreys' principle overcome these problems). However, these problems are less empirically relevant if the posterior is located in regions of the parameter space away from the point of non-identification or if informative priors are used. Furthermore, parameters in the reduced form model do not suffer from these problems. Hence, we feel some of the problems

---

<sup>3</sup>Note that  $\lambda_2 = 0$  implies the nonparametric regression line is perfectly smooth (i.e. is a straight line). Thus,  $M_2$  is nearly equivalent to a SUR model with an intercept and  $x_{im}$  entering linearly. It is not exactly equivalent since we are only using ordering information about  $x_{im}$ .

discussed in, e.g., Kleibergen and van Dijk (1998) are not critical for our work. In some sense, these problems all involve prior elicitation and, with moderately large data sets, data information should predominate.<sup>4</sup> In practice, we argue that our approach should be a sensible one for practitioners and that the advantage of being semiparametric outweighs any costs associated with not eliciting priors directly off of structural form parameters.

## 2.1 The Parametric SEM

In this section we provide an empirical example to illustrate how our techniques can be applied in practice. Our specific example, though primarily illustrative in nature, simultaneously addresses several topics of considerable interest in labor economics. Specifically, we will introduce and estimate a two equation structural simultaneous equations model and permit various nonparametric specifications within this system. The two endogenous variables in our system will be the log hourly wage received by individuals in the labor force and the quantity of schooling attained by those individuals. While many studies have recognized the potential endogeneity of schooling in standard log wage equations (see *e.g.* Card, 1999, for a review of recent IV studies on this issue), these studies do not typically estimate the full underlying structural model, and have not allowed for nonparametric specifications within these systems.

To fix ideas, the fully parametric version of our model may be written as:

$$\begin{aligned} s_i &= z_i^{C'} \alpha_1^S + z_i^{S'} \alpha_2^S + u_i^S \\ w_i &= \alpha_0 + \rho s_i + z_i^{C'} \alpha_1^W + z_i^{W'} \alpha_2^W + u_i^W, \end{aligned} \tag{2.14}$$

with

$$\begin{bmatrix} u_i^s \\ u_i^w \end{bmatrix} \stackrel{iid}{\sim} N \left[ \begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_s^2 & \sigma_{sw} \\ \sigma_{sw} & \sigma_w^2 \end{pmatrix} \right] \equiv N(0, \Sigma).$$

In the above equation  $z_i^C$  is a  $k_C$ -vector of exogenous variables common to both equations,  $z_i^S$  is a  $k_S$ -vector of exogenous variables which enter only the schooling equation (i.e. these are the instruments) and  $z_i^W$  is a  $k_W$ -vector of exogenous variables which enter only the wage equation. The parameters in (2.14) are structural, with  $\rho$  being the returns to schooling parameter that is often of central interest. The triangular structure of (2.14) implies that the Jacobian is unity, so that we can directly estimate the structural form using the methods we have developed in the previous section for the semiparametric SUR model.

In our empirical work we generalize this fully parametric structural model by permitting nonparametric specifications for a few variables in this system. We divide our empirical analysis into three models, with each case adding a new nonparametric component. In Case 1 we add a nonparametric specification to our wage equation and treat tenure on the job nonparametrically. Several studies in labor economics (e.g. Altonji and Shakotko 1987, Topel 1991, Light and McGarry, 1998 and Bratsberg and Terrell, 1998) have addressed

---

<sup>4</sup>This statement should be qualified by noting that the case where unbounded uniform priors over the structural parameters should be avoided. In this case, local non-identification issues can imply impropriety of the posterior.

the issue of separating the effects of on-the-job tenure and total labor market experience, with all of these studies specifying parametric (typically quadratic) specifications for each of these variables. In Case 1, we include a linear experience term and a nonparametric tenure term to flexibly investigate the shape of the relationship between job tenure and labor market experience. In Case 2, we add a nonparametric component to the education equation and treat the “ability” variable nonparametrically. In this analysis, “ability” refers to measured cognitive ability, and is proxied by a (continuous) test score that is available in our data set. Finally, in Case 3 we also treat this ability variable nonparametrically in our wage equation.<sup>5</sup> Before discussing our models and results in more detail, we first describe the data used in this analysis.

## 2.2 The Data

To estimate our models we take data from the National Longitudinal Survey of Youth (NLSY). The NLSY is a widely-used panel study containing a wealth of demographic information regarding a young cohort of U.S. men and women. Survey information from the NLSY begins in 1979, at which point the respondents range in age from 14 and 22. Sample participants are re-interviewed annually until 1994, and then additional biennial interviews were conducted.

To illustrate the use of our methods and remain consistent with the models described in section 2, we abstract from the panel aspect of the NLSY and focus only on cross-sectional wage outcomes in 1992. We choose this year since key variables of interest are directly available in that year, and since the NLSY participants range in age from 27-35 in 1992 and thus are likely to have completed their education and possess a reasonable degree of labor market experience. In keeping with this literature and to abstract from selection issues into employment, we also focus exclusively on the outcomes of white males in the NLSY.

Key to identification of this simultaneous equations model is the availability of an instrument or exclusion restriction. In the context of our application we need to find some variable that affects the quantity of schooling attained, yet has no direct structural effect on wages given the other controls we employ. To this end, we depart from the usual supply-side IV literature (*e.g.* Card 1999) and use the quantity of schooling attained by the respondent’s oldest sibling (SIBED) as our instrument.<sup>6</sup> The argument behind the use of this instrument is that sibling’s education should be strongly correlated with one’s own schooling. This correlation could arise, for example, from unobserved family attitudes toward the importance of education, or credit constraints faced by the family. However, we argue that the only channel through which sibling’s education affects one’s own wages is an indirect one (through the quantity of schooling attained), since conditioned on the schooling of the respondent himself and added controls for family background, the education of the

---

<sup>5</sup>In related work, Cawley, Heckman and Vytlačil (1999) argue that ability enters the wage equation nonlinearly, and Blackburn and Neumark (1995), Heckman and Vytlačil (2001) and Tobias (2003) examine if returns to schooling vary with ability. The latter two of these studies obtain results by allowing for flexible specifications of the relationship between ability and log wages.

<sup>6</sup>In the base year of the NLSY survey, participants are asked to report the highest grade completed by the oldest sibling. To ensure that the oldest sibling had completed his/her education, we restrict our attention to those observations where the oldest sibling was at least 24 years of age. Thus, our analysis conditions on the white males in the NLSY with an older sibling who is at least 24 years old.

sibling should play no structural role in the wage equation.<sup>7</sup>

To estimate our models of interest we also need to obtain information about the actual labor market experience of the individual as well as his tenure on the current (CPS) job in 1992. The job tenure (TENURE) variable is readily available, as the NLSY directly provides information on the total tenure (in weeks) with the current employer. As for total labor market experience, in each year of the survey the NLSY constructs the total number of weeks worked since the previous interview date. Since information for some weeks is occasionally missing, the NLSY also has a companion question that provides the percentage of weeks unaccounted in each year in the construction of this weeks of work variable. As such, we confine our attention to only those individuals whose weeks are fully accounted for in each year, and aggregate these experience variables across years to obtain our measure of total labor market experience (EXPERIENCE).<sup>8</sup>

In both the schooling and wage equations, we include the respondent's Armed Forces Qualifying Test (AFQT) score which is standardized by age (denoted ABILITY), highest grade completed by the respondent's mother (MOMED) and father (DADED), and a dummy variable equal to 1 if the respondent lives with both of his parents at age 14 (NON-BROKEN). In the wage equation, we also include weeks of actual labor market experience (EXPERIENCE), weeks of tenure at the current job (TENURE), a dummy for residence in an urban area (URBAN), and a continuous measure of the local unemployment rate (UNEMP). When measured in weeks, both EXPERIENCE and TENURE can be regarded as approximately continuous variables. Our sample restrictions are quite strict, and produce a clean, but relatively small data set. We limit our focus to white men in 1992 with older siblings at least 24 years of age in the base year of the survey and with complete information on the remaining variables. In addition, we exclude several extra observations for those individuals who report to be currently enrolled in school in 1992, who are in the military subsample, whose hourly wage exceeds \$100 or is less than \$1 per hour or who report to have completed less than 9 years of schooling. This leaves us with a total of  $N = 303$  observations, for which an exact finite-sample Bayesian analysis seems particularly useful.

## 2.3 Parametric Results

Before moving on to present results from our elaborated semiparametric models, we briefly present results using two standard parametric approaches. The first of these simply estimates the structural wage equation ignoring potential endogeneity problems. In this model we include an intercept, linear terms in the explanatory variables described above and a quadratic in TENURE. We estimate this model using a fully noninformative prior so that our results can be interpreted as the Bayesian counterpart to using OLS tech-

---

<sup>7</sup>Simple regression analyses that included sibling's education along with the other controls found no significant role for SIBED in the log wage equation.

<sup>8</sup>This definition is not without controversy, with many researchers (*e.g.* Wolpin 1992, and Bratsberg and Terrell 1998) only considering labor market experience after the completion of high school (or looking at only "terminal" high school graduates). Light (1998) investigates this issue and finds sensitivity of results to the definition of the career starting point. In this analysis, we do not make a distinction between pre-high school and post-high school labor market experience.

niques on the structural wage equation, for now, ignoring potential endogeneity issues. The second of our parametric models estimates the two equation structural model in (2.14). The prior for all of the regression coefficients is noninformative, and for  $\Sigma^{-1}$  we use the same prior as for the semiparametric model (see the discussion of the prior in Section 2.1).

Table 1 presents empirical results for the coefficients in these parametric models. The results are mostly sensible. The coefficient on our instrument SIBED is positive, as expected, with a posterior standard deviation considerably smaller than its posterior mean.<sup>9</sup> We also note that results from single equation estimation of the wage equation (which ignores the endogeneity problem) are quite similar to results obtained the two equation system. For instance, in both cases the point estimate of the return to schooling parameter is roughly 8 percent, and the results for the remaining coefficients are highly similar. The main differences in results occur with the posterior standard deviations for the coefficients on the highly correlated variables SCHOOL and ABILITY, which are much larger in the two equation model. This reduction in precision can be explained by the fact that although the point estimate of the correlation between the errors in the two equations is not far from zero (i.e. it is 0.054), it is relatively imprecisely estimated with this modest sample size (i.e. the posterior standard deviation of this coefficient is 0.252). Although the point estimate suggests that endogeneity is not a problem in this data set (and, hence, point estimates of key parameters do not change much when we control for endogeneity), the posterior for  $\Sigma$  is quite dispersed and allocates appreciable probability to regions of the parameter space where endogeneity is a problem. Given this uncertainty regarding the empirical importance of endogeneity, the standard errors associated with these parameters tend to increase relative to the model which ignores endogeneity concerns.

The finding that appreciable posterior probability is allocated to regions where the correlation between the errors in the two equations is near zero or small in magnitude is consistent with some of the other empirical work in this literature.<sup>10</sup> That is, it has often been either assumed or more formally argued that after controlling for a rich set of explanatory variables, endogeneity problems are likely to be mitigated. Our analysis lends some additional credence to this claim, as we “test down” from a structural model that permits endogeneity, and find little evidence that endogeneity is a serious empirical issue for this model. As we show in later sections, however, we find evidence against this basic parametric model, and in our generalized model specifications, there is some indication of a need to control for endogeneity of schooling.

---

<sup>9</sup>In fact, we find a “significant” role for this instrument in all of our model specifications.

<sup>10</sup>Blackburn and Neumark (1995) using NLSY data argue that once ability is controlled for, there is little evidence that schooling remains endogenous.

Table 1: Posterior Results for Parametric Models

Explanatory Variable	Single Equation Model		Two Equation Model			
	Wage Equation		Wage Equation		Schooling Equation	
	Mean	St. Dev.	Mean	St. Dev.	Mean	St. Dev.
INTERCEPT	0.571	0.296	0.566	0.662	9.255	0.714
ABILITY	0.020	0.038	0.022	0.098	1.253	0.136
MOMED	-0.015	0.013	-0.015	0.017	0.107	0.053
DADED	0.019	0.010	0.019	0.011	0.043	0.037
NON-BROKEN	-0.077	0.076	-0.077	0.080	0.190	0.297
SIBED	--	--	--	--	0.116	0.049
EXPERIENCE	$8.7 \times 10^{-4}$	$2.7 \times 10^{-4}$	$8.7 \times 10^{-4}$	$2.7 \times 10^{-4}$	--	--
TENURE	$1.4 \times 10^{-3}$	$4.3 \times 10^{-4}$	$1.4 \times 10^{-3}$	$4.3 \times 10^{-4}$	--	--
TENURE <sup>2</sup>	$-1.3 \times 10^{-4}$	$6.4 \times 10^{-5}$	$-1.3 \times 10^{-4}$	$6.4 \times 10^{-5}$	--	--
URBAN	0.119	0.061	0.119	0.061	--	--
UNEMP	$-6.6 \times 10^{-3}$	0.010	$-6.6 \times 10^{-3}$	0.010	--	--
SCHOOL	0.081	0.015	0.080	0.068	--	--

## 2.4 Case 1: Application

In this model, we elaborate (2.14), and allow the variable TENURE to enter the log wage equation nonparametrically.<sup>11</sup> Formally, in Model 1 we specify:

$$s_i = z_i^{C'} \alpha_1^S + z_i^{S'} \alpha_2^S + u_i^S$$

$$w_i = \rho s_i + z_i^{C'} \alpha_1^W + z_i^{W'} \alpha_2^W + f(x_i) + u_i^W,$$

where  $x_i$  denotes the number of weeks of work on the current job (TENURE) and  $z_i^W$  no longer contains TENURE. In our analysis, we set  $\eta = 5 \times 10^{-9}$ , which is the empirical Bayes estimate that maximizes the marginal likelihood (see Figure 2). Figure 1 plots the fitted nonparametric regression line against the data (after removing the effect of the other explanatory variables). That is, Figure 1 plots the posterior mean of the nonparametric regression line (and +/- two standard deviation bands) and the “data” points have coordinates  $x_i$  and  $w_i - \rho s_i - z_i^{C'} \alpha_1^W - z_i^{W'} \alpha_2^W$  for  $i = 1, \dots, N$  where all parameters are evaluated at their posterior means. Figure 2 plots the log of the Bayes factor in favor of the semiparametric model over the parametric model with a linear TENURE term across alternate choices of  $\eta$  (see equation 2.13). It can be seen that the maximum of the log Bayes factor is 0.026. Thus, there is only slight support for our semiparametric model over the parametric model with a linear tenure term. Note that as  $\eta \rightarrow 0$ , the nonparametric and linear models become equivalent and the log Bayes factor is zero. The value of  $\eta$  that maximizes the marginal likelihood is quite close to this case. However, Figure 2 does indicate an interior maximum, so a model with slight nonlinearities that suggest a concave tenure profile is preferred over the parametric model with a linear TENURE term.

<sup>11</sup>We found little evidence to support nonlinear relationships between total labor market experience and log wages.


Table 2: Posterior Results for the Case 1 Semiparametric Model

Explanatory Variable	Two Equation Model			
	Wage Equation		Schooling Equation	
	Mean	St. Dev.	Mean	St. Dev.
INTERCEPT	---	---	9.271	0.688
ABILITY	0.009	0.093	1.252	0.131
MOMED	-0.017	0.016	0.108	0.054
DADED	0.019	0.010	0.042	0.036
NON-BROKEN	-0.079	0.079	0.195	0.296
SIBED	---	---	0.116	0.050
EXPERIENCE	$8.5 \times 10^{-4}$	$2.6 \times 10^{-4}$	---	---
URBAN	0.118	0.061	---	---
UNEMP	$-6.5 \times 10^{-3}$	0.010	---	---
SCHOOL	0.090	0.064	---	---

The results in Table 2 are very similar to the two equation results in Table 1 which differ in that TENURE and TENURE<sup>2</sup> are included parametrically. The point estimate of returns to schooling, at 9.0%, is slightly higher than with the parametric models. However, relative to its posterior standard deviation this difference is minor. The posterior mean of the correlation between the errors in the two equations is also very similar to that of the parametric model (i.e. its posterior mean 0.031 and standard deviation is 0.231). Overall, we find our nonparametric function of TENURE to be playing a nearly identical role to the quadratic specification of this variable in the parametric model.

A comparison between a parametric SUR with TENURE entering quadratically to the semiparametric SUR can be done by first calculating the Bayes factor in favor of the quadratic SUR model of Table 1 against the parametric SUR with TENURE entering linearly (call this Bayes factor  $BF^*$  to distinguish it from  $BF(\eta)$  defined in equation 2.13).<sup>12</sup> That is,  $BF(\eta)$  compares the semiparametric SUR against a linear SUR (subject to the qualification of footnote 2), and thus the two Bayes factors  $BF^*$  and  $Bf(\eta)$  will be comparing a nonlinear (either quadratic or nonparametric) specification to the linear one. However, Bayes factor calculation requires an informative prior over parameters which are not common to both models. Thus, to calculate  $BF^*$  we require an informative prior for the coefficient on TENURE<sup>2</sup> which we choose to be  $N(0, \underline{v}_q)$ . With this prior,  $BF^*$  can be calculated using the Savage-Dickey density ratio with the strategy discussed above (see the discussion around equation 2.13). The elicitation of prior hyperparameters such as  $\underline{v}_q$  can be difficult (which is a further motivation for our empirical Bayesian analysis of a semiparametric model). In our application, values of  $\underline{v}_q$  greater than  $10^{-10}$  indicate support for the linear model (i.e.  $BF^* < 1$ ). This apparently informative choice of prior variance is actually not that informative relative to the data information (note that the posterior standard deviation of this coefficient in Table 1 is  $6.4 \times 10^{-5}$ ). For  $\underline{v}_q < 10^{-10}$ , the quadratic model is supported (i.e.  $BF^* > 1$ ). However, there is no value for  $\underline{v}_q$  for which the quadratic model receives overwhelming support. The maximum value for  $BF^*$  is 2.77 which occurs when

<sup>12</sup>Of course, given the Bayes factor of the semiparametric SUR against the linear model, and the Bayes factor of the quadratic model against the linear model, one can calculate the Bayes factor of the semiparametric SUR against the quadratic model.




$$\underline{v}_q = 10^{-12}.^{13}$$

This prior sensitivity analysis for the quadratic model can be interpreted in various ways, but regardless of how it is interpreted it is clear that the performance of the semiparametric and quadratic models (as measured by their marginal likelihoods) is similar. This is despite the fact that there is a strong support for parsimony implicit in Bayes factors. In the quadratic parametric model, deviations from linearity are modelled by adding a single extra parameter, but in our semiparametric model deviations from linearity are modelled by adding  $N - 2$  extra parameters. Hence, we would expect our semiparametric model to be penalized relative to the quadratic model. In any empirical exercise, if the researcher knows the correct functional form of the nonlinearity then it is best to work with the parametric model which captures this functional form. However, the advantage of the nonparametric approach is that the data can be used to decide the (here roughly quadratic) functional form. With the parametric model, preliminary estimation and pretesting were required to select “down” to the quadratic functional form. Hence, in our case it is not advisable to compare the quadratic model to the semiparametric model using a Bayes factor which ignores the model selection issues involved with the quadratic model.

<sup>13</sup>The Bayes Factor in favor of the quadratic model becomes larger if the prior mean of the quadratic coefficient is located closer to the posterior mean. However, we do not consider this case since it is common practice to center the prior over the restriction being tested.





### 3 Case 2: A Single Nonparametric Component in Several Equations

In this section, we consider the more general semiparametric SUR model given in (1.1) where a nonparametric component potentially exists in every equation. That is,  $\gamma_{ij} = f_j(x_{ij})$  for  $j = 1, \dots, m$  is the  $i^{th}$  point on the nonparametric regression line in the  $j^{th}$  equation. We maintain the assumption that  $x_{ij}$  is a scalar. Simple Bayesian methods for this model can be developed similarly to those developed for Case 1. We adopt the same strategy of treating unknown points on the nonparametric regression lines as unknown parameters and, hence, augment each equation with  $N$  new explanatory variables (as in equation 2.1). We then use a smoothness prior on each nonparametric regression line (analogous to equations 2.4 and 2.12). The resulting posterior can be handled using a Gibbs sampler (analogous to equations 2.6 and 2.7). Note, however, that we expressed our smoothness prior in terms of the second-differencing matrix  $D$  given in (2.2). This prior required the data to be ordered so that  $x_{1m} < \dots < x_{Nm}$ . However, unless each equation has its nonparametric component depending on the same explanatory variable (i.e.  $x_{ij} = x_{im}$  for  $j = 1, \dots, m-1$ ), the data in the  $j^{th}$  equation (for  $j = 1, \dots, m-1$ ) will not be ordered in such a way that a smoothness prior can be expressed in terms of  $D$ . However, this can be corrected for by redefining the explanatory variables. This requires some new, somewhat messy, notation. Unless otherwise noted, all other assumptions and notation are as for Case 1. For future reference, define  $\gamma_j = (\gamma_{1j}, \dots, \gamma_{Nj})'$

In Case 1, the inclusion of the nonparametric component implied that the identity matrix,  $I_N$ , was included as a matrix of explanatory variables (see equation 2.1 and the surrounding definitions). Here we define  $I_j^*$  which is a rearranged version of the identity matrix to correspond to the ordering of the data in the  $j^{th}$  equation for  $j = 1, \dots, m$ . Thus, since  $x_{1m} < \dots < x_{Nm}$ ,  $I_m^*$  is simply  $I_N$ , but the other equations potentially involving a reordering of the columns of  $I_N$ . Also define  $\zeta_{ij}$  to be the  $i^{th}$  row of  $I_j^*$ .

A concrete example of how this works might help. Suppose we have  $N = 5$  observations and the explanatory variables treated nonparametrically in the  $m = 2$  equations have values in the columns of the following matrix:

$$\begin{bmatrix} 3 & 1 \\ 4 & 2 \\ 1 & 3 \\ 2 & 4 \\ 5 & 5 \end{bmatrix}.$$

The data has been ordered so that the second explanatory variable is in ascending order,  $x_{12} < \dots < x_{52}$  and, hence, a smoothness prior using  $D$  can be directly applied in the second equation. However, the first explanatory variable is not in ascending order. However, we can reorder the identity matrix

$$I_1^* = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

It can be seen that with  $I_1^*$  used to define the nonparametric explanatory variables for the first equation,  $\gamma_{11}$  is the first point on the nonparametric regression line,  $\gamma_{21}$  is the second point, etc.. Thus, the smoothness prior can be expressed as restricting  $D\gamma_1$ .

In Case 1, we noted that the smoothness prior was only an  $N - 2$  dimensional distribution for the  $N$  points on each nonparametric regression line. Implicitly, this prior did not provide any information about the initial conditions (i.e. what we called  $\gamma_1$  and  $\gamma_2$  in Case 1), but only the second differences of points on the nonparametric regression line,  $\gamma_i - 2\gamma_{i-1} + \gamma_{i-2}$ . For the initial conditions, we used a noninformative prior. This need to separate out initial conditions necessitates the introduction of more notation. Define the  $2 \times 1$  vector of initial conditions in every equation as  $\gamma_j^0$  for  $j = 1, \dots, m$ . Let  $\gamma_j^*$  be  $\gamma_j$  with these first two elements deleted. Similarly, let  $I_j^{**}$  be  $I_j^*$  with its first two columns deleted and  $I_j^{***}$  be the two deleted columns. Also define  $\zeta_{ij}^*$  to be  $\zeta_{ij}$  with the first two elements deleted and  $\zeta_{ij}^0$  be the two deleted elements. Analogously, partition  $D = [D^{**} D^*]$  where  $D^{**}$  contains the first two columns of  $D$ .

With all these definitions, we can write the Case 2 semiparametric SUR as (2.1) with

$$W_i = \begin{bmatrix} z'_{i1} & \zeta_{i1}^0 & 0 & \cdot & \cdot & \cdot & \cdot & \cdot & 0 & \zeta_{i1}^* & 0 & \cdot & \cdot & 0 \\ 0 & 0 & z'_{i2} & \zeta_{i2}^0 & \cdot & \cdot & \cdot & \cdot & \cdot & 0 & \zeta_{i2}^* & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 0 & z'_{i,m-1} & \zeta_{i,m-1}^0 & 0 & 0 & \cdot & \cdot & 0 & \zeta_{i,m-1}^* & 0 \\ 0 & \cdot & \cdot & \cdot & \cdot & \cdot & 0 & z'_{im} & \zeta_{im}^0 & 0 & \cdot & \cdot & \cdot & \zeta_{im}^* \end{bmatrix}, \quad (2.15)$$

where  $\delta = (\beta'_1, \gamma_{11}^{0'}, \dots, \beta'_m, \gamma_{m1}^{0'}, \gamma_{11}^{*'}, \dots, \gamma_{m1}^{*'})'$  is a  $K + mN$  vector of coefficients.

Prior information about smoothness of the nonparametric regression lines will be expressed in terms of  $R\delta$ , where the  $m(N - 2) \times (K + mN)$  matrix  $R$  is given by

$$R = \begin{bmatrix} 0 & D^{**} & 0 & \cdot & \cdot & \cdot & \cdot & 0 & D^* & 0 & \cdot & \cdot & 0 \\ \cdot & & 0 & D^{**} & \cdot & \cdot & \cdot & \cdot & 0 & D^* & \cdot & \cdot & \cdot \\ & & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ & & \cdot & \cdot & \cdot & D^{**} & 0 & 0 & \cdot & \cdot & \cdot & D^* & 0 \\ & & \cdot & \cdot & \cdot & 0 & 0 & D^{**} & 0 & \cdot & \cdot & 0 & D^* \end{bmatrix}. \quad (2.16)$$

The remainder of the derivations are essentially the same as for Case 1. Define the partitions  $W_i = [W_i^{(1)} : W_i^{(2)}]$  where  $W_i^{(1)}$  is an  $m \times (K + 2m)$  matrix and  $W_i^{(2)}$  is  $m \times m(N - 2)$  and  $R = [R_1 : R_2]$  where  $R_1$  is an  $m(N - 2) \times (K + 2m)$  matrix and  $R_2$  is  $m(N - 2) \times m(N - 2)$ . Transform the model as:

$$y_i = V_i^{(1)}\lambda_1 + V_i^{(2)}\lambda_2 + \varepsilon_i = V_i\lambda + \varepsilon_i, \quad (2.17)$$

where  $\lambda = (\lambda'_1, \lambda'_2)'$ ,  $\lambda_1 = (\beta'_1, \gamma_{11}^{0'}, \dots, \beta'_m, \gamma_{m1}^{0'})'$ ,  $\lambda_2 = R\delta = [(D\gamma_1)']', \dots, (D\gamma_m)']'$ ,  $V_i^{(1)} = W_i^{(1)} - W_i^{(2)}R_2^{-1}R_1$  and  $V_i^{(2)} = W_i^{(2)}R_2^{-1}$ .

This model is now in the same form as Case 1. Given an independent Normal-Wishart prior as in (2.4) and (2.5), posterior analysis can be carried out using the Gibbs sampler described in (2.6) through (2.11). As in Case 1, we use a noninformative prior for  $\lambda_1$ . The prior for  $\Sigma^{-1}$  uses the same hyperparameter values as in Case 1. The smoothness prior relates to  $\lambda_2$  and, for this, we extend the prior of Case 1 (see equation 2.12) to be:

$$V(\eta_1, \dots, \eta_m) = \begin{bmatrix} \eta_1 I_{N-2} & 0 & \cdot & \cdot & 0 \\ 0 & \eta_1 I_{N-2} & 0 & \cdot & \cdot \\ \cdot & 0 & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 0 \\ 0 & \cdot & \cdot & 0 & \eta_m I_{N-2} \end{bmatrix}. \quad (2.18)$$

Thus, the nonparametric component of each equation can be smoothed to a different degree. An empirical Bayesian analysis can be carried out as described above (see equation 2.13). The computational demands of empirical Bayes in this general case can be quite substantial since a search over  $m$  dimensions of the smoothing parameter vector must be carried out.

### 3.1 Case 2: Application

For Case 2, we extend the Case 1 model to allow for an exogenous variable in the schooling equation to receive a nonparametric treatment. The model we consider here is:

$$\begin{aligned} s_i &= z_i^{C'}\alpha_1^S + z_i^{S'}\alpha_2^S + f_1(x_{i1}) + u_i^S \\ w_i &= \rho s_i + z_i^{C'}\alpha_1^W + z_i^{W'}\alpha_2^W + f_2(x_{i2}) + u_i^W \end{aligned}, \quad (2.19)$$

where all definitions are as for Case 1 except that  $x_{i1}$  is ABILITY and  $x_{i2}$  is TENURE and  $z_i^C$  no longer contains ABILITY in the schooling equation. Empirical Bayesian methods are used to select  $\eta_1$  and  $\eta_2$  which

smooth the nonparametric regression lines in the two equations. This leads us to set  $\eta_1 = 5 \times 10^{-6}$  and  $\eta_2 = 10^{-11}$ .

Table 3 presents posterior results for the parametric coefficients in this semiparametric model. As found in our previous results, the correlation between the errors in the two equations has a point estimate near to, but now farther away from zero (i.e. its posterior mean is 0.102) and remains very imprecisely estimated (i.e. its posterior standard deviation is 0.142). Thus, we have more evidence that endogeneity is an issue in model specification. Other results can be seen to be similar as for Case 1.

Interestingly, this analysis finds rather strong evidence of nonlinearities in the relationship between ability and schooling. The log of the Bayes factor of our semiparametric model against the linear-in-schooling model is 4.645, which indicates significantly more support for departures from linearity than was found in Case 1. Figures 3 and 4 plot the posterior means of the two nonparametric regression lines against the data (after controlling for parametric explanatory variables). That is, the “data” points in Figure 3 plot TENURE against  $w_i - \rho s_i - z_i^{C'} \alpha_1^W - z_i^{W'} \alpha_2^W$  for  $i = 1, \dots, N$  where all parameters are evaluated at their posterior means. The comparable points in Figure 4 plot ABILITY against  $s_i - z_i^{C'} \alpha_1^S - z_i^{S'} \alpha_2^S$  (evaluated at the posterior means for  $\alpha_1^S$  and  $\alpha_2^S$ ). Figure 3 looks very similar to Figure 1 and indicates some slight nonlinearities that appear quadratic. Figure 4 indicates more interesting (and more precisely estimated) nonlinear effects that would not be captured by simple parametric methods (e.g. including ABILITY in a quadratic manner).<sup>14</sup> Specifically, the graph suggests that marginal increments in ability for low ability individuals does little to increase the quantity of schooling attained (i.e. the graph is quite flat to the left of zero). However, for those individuals above the mean of the ability distribution, marginal increments in ability significantly increase the likelihood of acquiring more schooling. The fact that ability is a strong predictor of schooling has been well-documented (e.g. Heckman and Vytlacil, 2000), and here we add to this result by finding that it is relatively high ability individuals whose schooling choices are most affected by changes in ability.

It is also of interest to note that results for the returns for schooling parameter are slightly lower than what we have seen in either the parametric model or Case 1, with a posterior mean of 0.058 and posterior standard deviation of 0.038. We will now try to reconcile why this reduction has taken place. Our semiparametric estimation results found strong evidence of a nonlinear (and convex) relationship between ability and the quantity of schooling attained. To illustrate how this convex relationship may lead to a reduction of the schooling coefficient, let’s suppose for the sake of simplicity that the actual relationship between schooling and ability is quadratic, with a positive coefficient on the squared term. Since the correlation between the errors of the structural equations of Case 2 is non-zero (or at least most of the posterior mass is concentrated away from zero), this implies that the conditional mean of wages given schooling (i.e. the “reduced form” wage equation from our structural model), will now contain the nonlinear ability term that enters the education

<sup>14</sup>If we add ABILITY<sup>2</sup> to the parametric SUR in (2.14) its coefficient has posterior mean which is roughly one posterior standard deviation from zero. Thus, a parametric analysis using a quadratic functional form for ABILITY would likely conclude that no nonlinearities existed in the ABILITY/SCHOOL relationship.

equation. This nonlinear term was, of course, not present in the conditional mean of Case 1 since that model only contained a linear ability term. So, we can regard the differences between the conditional means of Case 2 and Case 1 as essentially an omitted variable problem - in Case 2 we have an added quadratic ability term that is positively correlated with education (see Figure 3) and also positively correlated with log wages (we provide evidence of this in the next section). Using standard omitted variable bias formulas, we would thus predict a reduction in the “reduced form” schooling coefficient upon controlling for this nonlinearity in ability. This result has potentially significant implications for this literature, as it suggests the importance of controlling for potential nonlinearities in ability (in both the schooling or wage equations) in order to extract accurate estimates of the return to education. Despite this result, it is also important to recognize that the shift in the posterior of this key parameter is small relative to its posterior standard deviation.

Table 3: Posterior Results for the Case 2 Semiparametric Model

Explanatory Variable	Wage Equation		Schooling Equation	
	Mean	St. Dev.	Mean	St. Dev.
SIBED	--	--	0.087	0.047
ABILITY	0.051	0.061	--	--
MOMED	-0.012	0.015	0.123	0.052
DADED	0.021	0.010	0.050	0.035
NON-BROKEN	-0.068	0.077	0.045	0.287
EXPERIENCE	$8.0 \times 10^{-4}$	$2.7 \times 10^{-4}$	--	--
URBAN	0.124	0.062	--	--
UNEMP	$-6.4 \times 10^{-3}$	0.010	--	--
SCHOOL	0.058	0.038	--	--

## 4 Case 3: Nonparametric Components Depend on Several Explanatory Variables: Additive Models

To this point we have only considered cases where the nonparametric component in a given equation depended on a single explanatory variable. That is,  $x_{ij}$  was assumed to be a scalar. In this section, we assume  $x_{ij}$  to be a vector of  $p$  explanatory variables.<sup>15</sup> The curse of dimensionality (see, e.g., Yatchew, 1998, pages 675-676) implies that it is difficult to carry our nonparametric inference (whether Bayesian or non-Bayesian) when  $p$  is even moderately large. The intuition underlying our smoothness prior is that values of  $x_{ij}$  which are near one another should have points on the nonparametric regression line which are near one another. When  $x_{ij}$  is a scalar, the definition of “nearby” points is simple and is expressed through our ordering of the data as  $x_{1m} < \dots < x_{Nm}$ . When  $x_{ij}$  is not a scalar, it is possible to order the data in an analogous way using some distance metric. If it is sensible to order the data in this way, then the approach of Case 2 can be applied directly. However, this approach is apt to be sensitive to choice of distance definition and which point to choose as the first on each nonparametric regression line. In the single equation case, Yatchew (1998,

<sup>15</sup>The case where  $p$  varies across equations is a trivial extension of what is done in this section. We do not consider this extension to keep already messy notation from getting even messier.

Figure 3: Fitted Nonparametric Regression Line in Wage Equation





Figure 4: Fitted Nonparametric Regression Line in Schooling Equation



page 697) argues that the classical Differencing Estimator works well provided the dimension of  $x_i$  does not exceed 3. It is likely that such an approach could work well in our Bayesian semiparametric framework when dimensionality is as low as this. Nevertheless, in this section we develop a different approach for this general case.

The curse of dimensionality is greatly reduced if it is assumed that  $f_j(x_{ij})$  is additive. This is, of course, more restrictive than simply assuming  $f_j(x_{ij})$  is an unknown smooth function, but it is much less restrictive than virtually any parametric model used in this literature. Furthermore, by defining  $x_{ij}$  to include interactions of explanatory variables, some of the restrictions imposed by the additive form can be surmounted. Accordingly, in this section we develop methods for Bayesian inference in the model given in (1.1) with:

$$f_j(x_{ij}) = f_j(x_{ij1}, \dots, x_{ijp}) = f_{j1}(x_{ij1}) + \dots + f_{jp}(x_{ijp}) = \gamma_{ij1} + \dots + \gamma_{ijp}. \quad (2.20)$$

The basic idea underlying our approach to this model is straightforward: define a smoothness prior for each of the  $f_{jr}(x_{ijr})$  for  $j = 1, \dots, m$  and  $r = 1, \dots, p$  and use the methods for Bayesian inference in the semiparametric SUR model with independent Normal-Wishart prior described for Case 1. However, we must further complicate notation to handle this general case. In the following material, the indices run  $i = 1, \dots, N$ ,  $j = 1, \dots, m$  and  $r = 1, \dots, p$ .

For Case 2, we defined matrices,  $I_1^*, \dots, I_m^*$  which were used as explanatory variables for the nonparametric regression lines taking into account the fact that each nonparametric explanatory variable was not necessarily in ascending order. For Case 3, we define analogously  $I_{jr}^*$  which is the re-ordered identity matrix corresponding needed to incorporate  $f_{jr}(x_{ijr})$ , taking into account that the data are not necessarily ordered so that  $x_{1jr} < \dots < x_{Njr}$ . All the other Case 2 definitions can be extended in a similar fashion. Divide the vector of points on each nonparametric regression line,  $\gamma_{jr} = (\gamma_{1jr}, \dots, \gamma_{Njr})'$ , into the  $2 \times 1$  vector of initial conditions,  $\gamma_{jr}^0$ , and the remaining elements,  $\gamma_{jr}^*$ . Similarly, let  $I_{jr}^{**}$  be  $I_{jr}^*$  with the first two columns corresponding deleted and  $I_{jr}^{***}$  be the two deleted columns. Furthermore, let  $\zeta_{ijr}^*$  be  $\zeta_{ijr}$  with the elements corresponding to the initial conditions deleted and  $\zeta_{ijr}^0$  be the two deleted elements, where  $\zeta_{ijr}$  is the  $i^{th}$  row of  $I_{jr}^*$ . Further define  $\zeta_{ij}^0 = (\zeta_{ij1}^0, \dots, \zeta_{ijp}^0)$  and  $\zeta_{im}^* = (\zeta_{ij1}^*, \dots, \zeta_{ijp}^*)$ . Note that these last two definitions differ from Case 2.

With all these definitions, we can write the Case 3 model as a semiparametric SUR as in (2.1) with  $W_i$  as given in (2.15), except that the definition of some of the terms has changed slightly and  $\delta = (\beta_1', \gamma_{11}^0, \dots, \gamma_{1p}^0, \dots, \beta_m', \gamma_{m1}^0, \dots, \gamma_{mp}^0, \gamma_{11}^*, \dots, \gamma_{1p}^*, \dots, \gamma_{m1}^*, \dots, \gamma_{mp}^*)'$  is now a  $K + mpN$  vector of coefficients.

As before, our smoothness prior is expressed in terms of  $R\delta$ , where  $R$  is now an  $mp(N - 2) \times (K + mpN)$  matrix:

$$R = \begin{bmatrix} 0 & D_p^{**} & 0 & \cdot & \cdot & \cdot & \cdot & 0 & D_p^* & 0 & \cdot & \cdot & 0 \\ \cdot & \cdot & 0 & D_p^{**} & \cdot & \cdot & \cdot & \cdot & 0 & D_p^* & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & D_p^{**} & 0 & 0 & \cdot & \cdot & \cdot & D_p^* & 0 & \cdot \\ \cdot & \cdot & \cdot & \cdot & 0 & 0 & D_p^{**} & 0 & \cdot & \cdot & 0 & D_p^* & \cdot \end{bmatrix},$$

where

$$D_p^{**} = \begin{bmatrix} D^{**} & 0 & . & 0 \\ 0 & . & . & . \\ . & . & . & 0 \\ 0 & . & 0 & D^{**} \end{bmatrix}$$

is an  $p(N-2) \times 2p$  matrix and

$$D_p^* = \begin{bmatrix} D^* & 0 & . & 0 \\ 0 & . & . & . \\ . & . & . & 0 \\ 0 & . & 0 & D^* \end{bmatrix}$$

is  $p(N-2) \times p(N-2)$ .

The remainder of the derivations are the same as for Cases 1 and 2. That is, the model can be transformed as in (2.17). An independent Normal-Wishart prior for the transformed parameters is used with prior hyperparameters selected as for Case 2. The Gibbs sampler described in (2.6) through (2.11) can be used for posterior inference. The only difference is that it will usually be desirable to have a different smoothing parameter for every nonparametric regression line in every equation. Thus, we choose the prior covariance matrix for  $\lambda_2$  to be:

$$V(\eta_{11}, \dots, \eta_{1p}, \dots, \eta_{m1}, \dots, \eta_{mp}) = \begin{bmatrix} \eta_{11}I_{N-2} & 0 & . & . & . & . \\ 0 & . & . & . & . & . \\ . & 0 & \eta_{1p}I_{N-2} & . & . & . \\ . & . & . & . & . & . \\ . & . & . & . & . & 0 \\ . & . & . & . & 0 & \eta_{mp}I_{N-2} \end{bmatrix}. \quad (2.21)$$

There is an identification problem with this model in that constants may be added and subtracted to each nonparametric component without changing the likelihood. For instance, the equations  $y_{ij} = z'_{ij}\beta_j + f_{j1}(x_{ij1}) + f_{j2}(x_{ij2}) + \varepsilon_{ij}$  and  $y_{ij} = z'_{ij}\beta_j + g_{j1}(x_{ij1}) + g_{j2}(x_{ij2}) + \varepsilon_{ij}$  are equivalent if  $g_{j1}(x_{ij1}) = f_{j1}(x_{ij1}) + c$  and  $g_{j2}(x_{ij2}) = f_{j2}(x_{ij2}) - c$  where  $c$  is an arbitrary constant. Insofar as interest centers on the shapes of the  $f_{jr}(x_{ijr})$  for  $r = 1, \dots, p$  or the overall fit of the nonparametric regression line, the lack of identification is irrelevant. If desired, identification can be imposed in many ways (e.g. by setting the intercept of the  $r^{th}$  nonparametric component in each equation to be zero for  $r = 2, \dots, p$ ).

#### 4.1 Case 3: Application

In Case 3 we extend Case 2 to also allow for a nonparametric treatment of ABILITY in the wage equation. Thus, ABILITY is treated nonparametrically in both equations and TENURE is treated nonparametrically in the wage equation. The model is:

$$\begin{aligned} s_i &= z_i^{C'}\alpha_1^S + z_i^{S'}\alpha_2^S + f_{11}(x_{i11}) + u_i^S \\ w_i &= \rho s_i + z_i^{C'}\alpha_1^W + z_i^{W'}\alpha_2^W + f_{21}(x_{i21}) + f_{22}(x_{i22}) + u_i^W \end{aligned} \quad (2.22)$$



where definitions are as for Case 2 except that  $x_{i11} = x_{i22}$  is ABILITY and  $x_{i21}$  is TENURE and  $z_i^C$  no longer contains ability in either equation. We identify the model by setting the intercept of one of the nonparametric functions in the wage equation to be zero, i.e.  $f_{22}(x_{122}) = 0$ .

With three nonparametric components, empirical Bayesian methods involve a three-dimensional grid search over the smoothing parameters  $\eta_1, \eta_2$  and  $\eta_3$  for terms relating to ABILITY (in the schooling equation), TENURE and ABILITY (in the wage equation), respectively. We find  $\eta_1 = 10^{-6}, \eta_2 = 10^{-9}$  and  $\eta_3 = 10^{-11}$ . With these values, the log of the Bayes factor in favor of the nonparametric model is 3.837 indicating stronger support for the semiparametric model over the parametric alternative of (2.14) than with previous cases.

Empirical results for the regression coefficients are presented in Table 4 and are found to be similar to those for Case 2. In addition, the posterior mean of the correlation between the errors in the two equations is 0.138 (standard deviation 0.140), values similar to Case 2. Perhaps the most interesting finding is that the posterior mean of the return to schooling parameter is, at 0.042, similar to but smaller than that found for Case 2, and approximately one-half of the size of those reported in Case 1 and the parametric model. Again, upon controlling for nonlinearities in the relationship between ability and log wages, we find even more reduction in the return to schooling coefficient. However, the posterior standard deviation of this parameter is still quite large.

Figures 5, 6 and 7 plot the fitted nonparametric regression lines (after controlling for other explanatory variables in the same manner as for previous cases). Figure 7 indicates the same non-quadratic nonlinearities in the relationship between ABILITY and SCHOOL (after controlling for other explanatory variables) as Figure 4, while Figure 5 is similar to Figures 1 and 3. Figure 6 also appears to exhibit a slightly nonlinear regression relationship between log wages and ABILITY of a non-quadratic form (although the pattern is much weaker than in Figure 7). Specifically, Figure 6 suggests that marginal increments in ability does little to increase the log wages of individuals of low to moderate ability, but does begin to have a reasonable effect on the log wages of those already above the mean of the ability distribution (*i.e.* increasing returns to ability). It is also important to recognize that we are obtaining this result after controlling for the potentially endogenous education variable and also controlling for nonlinearities in the education-ability relationship. The fact that +/- two posterior standard deviation bands in Figure 6 are very tight for the lowest values of ABILITY is due to the identification restriction and the fact that there are very few observations in this region.




Table 4: Posterior Results for the Case 3 Semiparametric Model

Explanatory Variable	Wage Equation		Schooling Equation	
	Mean	St. Dev.	Mean	St. Dev.
SIBED	---	---	0.163	0.049
MOMED	-0.010	0.015	0.202	0.056
DADED	0.022	0.010	0.047	0.039
NON-BROKEN	-0.080	0.079	0.257	0.308
EXPERIENCE	$6.8 \times 10^{-4}$	$2.7 \times 10^{-4}$	---	---
URBAN	0.112	0.063	---	---
UNEMP	-0.010	0.010	---	---
SCHOOL	0.042	0.039	---	---

## 5 Conclusions

In this paper, we have developed methods for carrying out Bayesian inference in the semiparametric seemingly unrelated regressions model and showed how these methods can also be used for semiparametric simultaneous equations models. There are, of course, other methods for carrying out Bayesian inference in semi- or nonparametric extensions of SUR models (e.g. Smith and Kohn, 2000). A distinguishing feature of our approach is that we stay within the simple and familiar framework of the SUR model with independent Normal-Wishart prior. Thus, textbook results for Bayesian inference, model comparison, prediction and

Figure 6: Fitted Nonparametric Regression Line for ABILITY in Wage Equation





Figure 7: Fitted Nonparametric Regression Line in Schooling Equation



posterior computation are immediately available. The focus of this paper is on prior information about the degree of smoothness in the nonparametric regression lines (although, of course, prior information about other parameters can easily be accommodated). We show how empirical Bayesian methods can be used to estimate smoothing parameters, thus minimizing the need for subjective prior elicitation.

The practicality of our approach is demonstrated in a two-equation application involving returns to schooling. In addition to parametric models, we estimate models with a single nonparametric component in one equation and a single nonparametric component in both equations. Our most general model contained an additive specification in the wage equation and a nonparametric ability component in the schooling equation. Although our semiparametric results are, in some cases, similar to those from simpler parametric nonlinear models (e.g. where explanatory variables enter in a quadratic fashion), in other cases our semiparametric approach yields empirical results which could not be easily obtained using standard parametric methods. Using our approach, we found suggestive evidence of nonlinearities in the relationships between ability and the quantity of schooling attained, and that estimates of the return to schooling were sensitive to controlling for these nonlinear relationships. Furthermore, we stress that one clear advantage of a semiparametric approach is that a particular functional form such as the quadratic does not have to be chosen, either in an *ad hoc* fashion or through pre-testing.

Finally, it is worth noting that it is very easy to incorporate the semiparametric SUR model developed here in a more complicated multiple equation model. For instance, Bayesian inference in a multinomial semiparametric probit can be done by adding a data augmentation step in the Gibbs sampler outlined in this paper as in, e.g., McCulloch and Rossi (1994). Bayesian inference in a semiparametric multiple equation model where one (or more) of the dependent variables is censored can be handled in a similar manner. We have assumed Normal errors, but this assumption can easily be relaxed through the use of mixtures of Normals. In short, Bayesian inference in semiparametric variants of a wide range of multiple equation models can be handled in a straightforward manner.

## 6 References

- Altonji, J. and Shakotko, R.A. (1987). "Do wages rise with job seniority?" *Review of Economic Studies* 54(3), 437-459.
- Blackburn, M. and Neumark, D. (1995). "Are OLS estimates of the return to schooling biased downward? Another look," *Review of Economics and Statistics*, 77, 217-230.
- Blundell, R. and A. Duncan (1998). "Kernel Regression in Empirical Microeconometrics" *Journal of Human Resources* 33(1), 62-87.
- Bratsberg, B. and Terrell, D. (1998). "Experience, tenure and wage growth of young black and white men," *Journal of Human Resources* 33(3), 658-677.
- Card, D. (1999). "The causal effect of education on earnings," in *Handbook of Labor Economics Vol 3A*, O. Ashenfelter and D. Card, eds., 1801-1863.

- Cawley, J., Vytlačil, E. and Heckman, J. (1999). "On policies to reward the value added by educators," *Review of Economics and Statistics* 81(4), 720-728.
- Chib, S. and Greenberg, E. (1995). "Hierarchical analysis of SUR models with extensions to correlated serial errors and time-varying parameter models," *Journal of Econometrics*, 68, 339-360.
- Chib, S. and Greenberg, E. (1996). "Markov chain monte carlo simulation methods in econometrics," *Econometric Theory*, 12, 409-431.
- DiNardo, J. and J.L. Tobias (2001). "Nonparametric Density and Regression Estimation" *Journal of Economic Perspectives* 15(4), 11-28.
- Dreze, J. and Richard, J. (1983). "Bayesian analysis of simultaneous equations systems," chapter 9 in *Handbook of Econometrics*, volume 1, edited by Z. Griliches and M. Intriligator, Amsterdam: North Holland.
- Fernandez, C., Osiewalski, J. and Steel, M. (1997). "On the use of panel data in stochastic frontier models with improper priors," *Journal of Econometrics*, 79, 169-193.
- Heckman, J. and Vytlačil, E. (2001). "Identifying the role of cognitive ability in explaining the level of and Change in the return to schooling," *Review of Economics and Statistics* 83(1), 1-12.
- Kleibergen, F. (1997). "Bayesian simultaneous equations analysis using equality restricted random variables," *American Statistical Association, 1997 Proceedings of the Section on Bayesian Statistical Science*, 141-147.
- Kleibergen, F. and van Dijk, H. (1998). "Bayesian simultaneous equations analysis using reduced rank structures," *Econometric Theory*, 14, 699-744.
- Kleibergen, F. and Zivot, E. (2003). "Bayesian and classical approaches to instrumental variable regression" *Journal of Econometrics*, 114, 29-72.
- Koop, G. (2003). *Bayesian Econometrics*, Chichester: John Wiley and Sons.
- Koop, G. and Poirier, D. (2003a). "Bayesian variants of some classical semiparametric regression techniques," *Journal of Econometrics*, forthcoming.
- Koop, G. and Poirier, D. (2003b). "Empirical Bayesian inference in a nonparametric regression model," to appear in a volume from the *Conference in Honour of Professor J. Durbin on State Space Models and Unobserved Components*.
- Light, A. (1998) "Estimating returns to schooling: When does the career begin?" *Economics of Education Review* 17(1), 31-45.
- Light, A. and McGarry, K. (1998). "Job change patterns and the wages of young men," *Review of Economics and Statistics*, 276-286.
- McCulloch, R. and Rossi, P. (1994). "An exact likelihood analysis of the multinomial probit model," *Journal of Econometrics*, 64, 207-240.
- Pagan, A. and Ullah, A. (1999). *Nonparametric Econometrics*, Cambridge: Cambridge University Press.
- Poirier, D. (1995). *Intermediate Statistics and Econometrics*, Cambridge: The MIT Press.
- Rothenberg, T. (1963). "A Bayesian analysis of simultaneous equation systems," *Econometric Institute Report 6315*, Erasmus University, Rotterdam.
- Smith, M. and Kohn, R. (2000). "Nonparametric seemingly unrelated regression," *Journal of Econometrics*, 98, 257-281.
- Tobias, J.L. (2003). "Are returns to schooling concentrated among the most able? A semiparametric analysis of the ability-earnings relationships," *Oxford Bulletin of Economics and Statistics*, 65(1), 1-29.
- Topel, R. (1991). "Specific capital, mobility and wages: Wages rise with job seniority," *Journal of Political Economy* 99(1), 145-176.

van der Linde, A. (2000). "Reference priors for shrinkage and smoothing analysis," *Journal of Statistical Planning and Inference*, 90, 245-274.

Verdinelli, I. and Wasserman, L. (1995). "Computing Bayes Factors using a generalization of the Savage-Dickey density ratio," *Journal of the American Statistical Association*, 90, 614-618.

Wolpin, K. (1992). "The determinants of black-white differences in early employment careers: Search, layoffs, quits and endogenous wage growth" *Journal of Political Economy* 100(3), 525-560.

Yatchew, A. (1998). "Nonparametric regression techniques in economics," *Journal of Economic Literature*, 36, 669-721.