

CIRJE-F-621

Computation in an Asymptotic Expansion Method

Akihiko Takahashi
University of Tokyo

Kohta Takehara
Graduate School of Economics, University of Tokyo

Masashi Toda
Graduate School of Economics, University of Tokyo

May 2009

CIRJE Discussion Papers can be downloaded without charge from:

<http://www.e.u-tokyo.ac.jp/cirje/research/03research02dp.html>

Discussion Papers are a series of manuscripts in their draft form. They are not intended for circulation or distribution except as indicated by the author. For that reason Discussion Papers may not be reproduced or distributed without the written consent of the author.

Computation in an Asymptotic Expansion Method *

Akihiko Takahashi, Kohta Takehara[†] and Masashi Toda

Graduate School of Economics, the University of Tokyo

7-3-1, Hongo, Bunkyo-ku, Tokyo, 113-0033, Japan

May 31, 2009

Abstract

An asymptotic expansion scheme in finance initiated by Kunitomo and Takahashi [15] and Yoshida[68] is a widely applicable methodology for analytic approximation of the expectation of a certain functional of diffusion processes. [46], [47] and [53] provide explicit formulas of conditional expectations necessary for the asymptotic expansion up to the third order. In general, the crucial step in practical applications of the expansion is calculation of conditional expectations for a certain kind of Wiener functionals. This paper presents two methods for computing the conditional expectations that are powerful especially for high order expansions: The first one, an extension of the method introduced by the preceding papers presents a general scheme for computation of the conditional expectations and show the formulas useful for expansions up to the fourth order explicitly. The second one develops a new calculation algorithm for computing the coefficients of the expansion through solving a system of ordinary differential equations that is equivalent to computing the conditional expectations. To demonstrate their effectiveness, the paper gives numerical examples of the approximation for λ -SABR model up to the fifth order and a cross-currency Libor market model with a general stochastic volatility model of the spot foreign exchange rate up to the fourth order.

1 Introduction

This paper presents two alternative schemes for computation in an asymptotic expansion approach based on Watanabe theory(Watanabe [66]) in Malliavin

*This research is supported by the global COE program “The research and training center for new development in mathematics.”

[†]Research Fellow of the Japan Society for the Promotion of Science

calculus by extending the preceding papers and also by developing a new calculation algorithm.

To our best knowledge, the asymptotic expansion is first applied to finance for evaluation of an average option that is a popular derivative in commodity markets. [15] and [46] derive the approximation formulas for an average option by an asymptotic method based on log-normal approximations of an average price distribution when the underlying asset price follows a geometric Brownian motion. [68] applies a formula derived more generally by the asymptotic expansion of small diffusion processes. Thereafter, the asymptotic expansion have been applied to a broad class of problems in finance: See [47], [48], [49], [50], Kunitomo and Takahashi [16], [17], [18], [19], Kawai [11], Matsuoka, Takahashi and Uchida [31], Takahashi and Matsushima [51], Takahashi and Saito [52], Takahashi and Yoshida [57], [58], Kobayashi, Takahashi and Tokioka [13], Muroi [33], Osajima [38], Takahashi and Uchida [56], Kunitomo and Kim [14], Kawai and Jäckel [12], and [53], [54], [55].

For other asymptotic methods in finance which do not depend on Watanabe theory, see also Fouque, Papanicolaou and Sircar [5], [6], Henry-Labordere [24], [25], [26], Kusuoka and Osajima [20], Osajima [39] and Siopacha and Teichmann [45].

In the application of the asymptotic expansion based on Watanabe theory, they calculated certain conditional expectations which appear in their expansions and play a key role in computation, by the formulas up to the third order given explicitly in [46], [47] and [53]. In many applications, these formulas give sufficiently accurate approximation, but in some cases, for example in the cases with long maturities or/and with highly volatile underlying variables, the approximation up to the third order may not provide satisfactory accuracies. Thus, the formulas for the higher order computation are desirable. But to our knowledge, asymptotic expansion formulas higher than the third order have not been given yet. This paper provides the general procedures for the explicit computation of conditional expectations in the asymptotic expansion and show the formulas for the approximation up to the fourth order. Moreover, we develop another calculation algorithm which enables us to derive high order approximation formulas in an automatic manner. As a consequence, our approximation generally shows sufficient accuracy with computation of high order expansions, which is confirmed by numerical experiments.

In the following sections, after a brief explanation of the asymptotic expansion in Section 2, Section 3 will provide a computation procedure explicitly for conditional expectations appearing in the expansion and show the formulas for expansions up to the fourth order. Moreover, Section 4 will introduce our new alternative computation algorithm for the asymptotic expansion and derive the fourth order asymptotic expansion formula. Finally, Section 5 will apply our algorithms described in the previous sections to the concrete financial models, and confirm the effectiveness of the higher order expansions by numerical examples in λ -SABR model and a cross-currency Libor market model with a general stochastic volatility model of the spot foreign exchange rate.

2 Asymptotic Expansion

We consider a d -dimensional diffusion process $X_t^{(\epsilon)} = (X_t^{(\epsilon),1}, \dots, X_t^{(\epsilon),d})$ which is the solution to the following stochastic differential equation:

$$\begin{aligned} dX_t^{(\epsilon),i} &= V_0^i(X_t^{(\epsilon)}, \epsilon)dt + \epsilon V^i(X_t^{(\epsilon)})dW_t \quad (i = 1, \dots, d) \\ X_0^{(\epsilon)} &= x_0 \in \mathbf{R}^d \end{aligned} \quad (1)$$

where $W = (W^1, \dots, W^{d'})$ is a d' -dimensional standard Wiener process, and $\epsilon \in (0, 1]$ is a known parameter. Suppose $V = (V^1, \dots, V^d): \mathbf{R}^d \mapsto \mathbf{R}^d \otimes \mathbf{R}^{d'}$ satisfies some regularity conditions.

Next, suppose that a function $g: \mathbf{R}^d \mapsto \mathbf{R}$ to be smooth and all derivatives have polynomial growth orders. Then, for $\epsilon \downarrow 0$, $g(X_T^{(\epsilon)})$ has its asymptotic expansion;

$$g(X_T^{(\epsilon)}) = \sum_{n=0}^{\infty} \epsilon^n g_{nT}.$$

g_{nT} , the coefficients in the expansion, can be obtained by Taylor's formula and represented based on multiple Wiener-Ito integrals.

Let $A_{kt} = \frac{\partial^k X_t^{(\epsilon)}}{\partial \epsilon^k}|_{\epsilon=0}$ and A_{kt}^i , $i = 1, \dots, d$ denote the i -th elements of A_{kt} . In particular, A_{1t} is represented by

$$A_{1t} = \int_0^t Y_t Y_u^{-1} \left(\partial_\epsilon V_0(X_u^{(0)}, 0) du + V(X_u^{(0)}) dW_u \right) \quad (2)$$

where Y denotes the solution to the differential equation;

$$dY_t = \partial V_0(X_t^{(0)}, 0) Y_t dt; \quad Y_0 = I_d.$$

Here, ∂V_0 denotes the $d \times d$ matrix whose (j, k) -element is $\partial_k V_0^j = \frac{\partial V_0^j(x, \epsilon)}{\partial x_k}$, V_0^j is the j -th element of V_0 , and I_d denotes the $d \times d$ identity matrix.

For $k \geq 2$, A_{kt}^i , $i = 1, \dots, d$ is recursively determined by the following:

$$\begin{aligned} A_{kt}^i &= \int_0^t \partial_\epsilon^k V_0^i(X_s^{(0)}, 0) ds \\ &+ \sum_{l=1}^k \frac{k!}{l!(k-l)!} \sum_{\beta=1}^l \sum_{\vec{l}_\beta \in L_{\beta,l}} \int_0^t \frac{1}{\beta!} \sum_{d_1, \dots, d_\beta=1}^d \partial_{d_1, \dots, d_\beta}^\beta \partial_\epsilon^{k-l} V_0^i(X_s^{(0)}, 0) \prod_{j=1}^\beta A_{l_j, s}^{d_j} ds \\ &+ \sum_{\beta=1}^k \sum_{\vec{l}_\beta \in L_{\beta, k-\beta}} \int_0^t \frac{1}{\beta!} \sum_{\alpha=1}^{d'} \sum_{d_1, \dots, d_\beta=1}^d \partial_{d_1, \dots, d_\beta}^\beta V_\alpha^i(X_s^{(0)}) \prod_{j=1}^\beta A_{l_j, s}^{d_j} dW_s^\alpha, \end{aligned} \quad (3)$$

where $\partial_\epsilon^l = \frac{\partial^l}{\partial \epsilon^l}$, $\partial_{d_1, \dots, d_\beta}^\beta = \frac{\partial^\beta}{\partial x_{d_1} \dots \partial x_{d_\beta}}$ and

$$L_{\beta, k} = \left\{ \vec{l}_\beta = (l_1, \dots, l_\beta); l_j \geq 0 (j = 1, \dots, \beta), \sum_{j=1}^\beta l_j = k \right\}.$$

Then, g_{0T} and g_{1T} can be written as

$$\begin{aligned} g_{0T} &= g(X_T^{(0)}), \\ g_{1T} &= \sum_{i=1}^d \partial_i g(X_T^{(0)}) A_{1T}^i. \end{aligned}$$

For $n \geq 2$, g_{nT} is expressed as follows:

$$g_{nT} = \sum_{\vec{s} \in \mathcal{S}_n} \left(\frac{n!}{s_1! \dots s_n!} \right) \prod_{l=1}^n \left(\frac{1}{l!} \right)^{s_l} \sum_{\vec{p}^s \in P_{s_l}} \left(\frac{s_l!}{p_1^{s_l}! \dots p_d^{s_l}!} \right) \partial_1^{p_1^{s_l}} \dots \partial_d^{p_d^{s_l}} g(X_T^{(0)}) \prod_{i=1}^d (A_{1T}^i)^{p_i^{s_l}} \quad (4)$$

where

$$S_n := \left\{ \vec{s} = (s_1, \dots, s_n); s_l \geq 0 (l = 1, \dots, n), \sum_{l=1}^n l s_l = n \right\},$$

$$P_s := \left\{ \vec{p}^s = (p_1^s, \dots, p_d^s); p_i^s \geq 0 (i = 1, \dots, d), \sum_{i=1}^d p_i^s = s \right\}.$$

Next, normalize $g(X_T^{(\epsilon)})$ to

$$G^{(\epsilon)} = \frac{g(X_T^{(\epsilon)}) - g_{0T}}{\epsilon}$$

for $\epsilon \in (0, 1]$. Then,

$$G^{(\epsilon)} = g_{1T} + \sum_{n=1}^{\infty} \epsilon^n g_{(n+1)T}.$$

Moreover, let

$$a_t = (\partial g(X_T^{(0)}))' [Y_T Y_t^{-1} V(X_t^{(0)})]$$

and make the following assumption:

$$\text{(Assumption 1)} \quad \Sigma_T = \int_0^T a_t a_t' dt > 0.$$

Note that g_{1T} follows a normal distribution with variance Σ_T and hence Assumption 1 means that the distribution of g_{1T} does not degenerate. In application, it is easy to check this condition in most cases.

Next, let Φ be a generalized function. Then, the expectation of $\Phi(G^{(\epsilon)})$ is expanded around $\epsilon = 0$ as follows:

$$\mathbf{E}[\Phi(G^{(\epsilon)})] = \sum_{m=0}^{\infty} \frac{1}{m!} \mathbf{E} \left[\Phi^{(m)}(g_{1T}) \left(\sum_{n=1}^{\infty} \epsilon^n g_{(n+1)T} \right)^m \right] \quad (5)$$

$$= \sum_{m=0}^{\infty} \frac{1}{m!} \mathbf{E} \left[\Phi^{(m)}(g_{1T}) \mathbf{E} \left[\left(\sum_{n=1}^{\infty} \epsilon^n g_{(n+1)T} \right)^m \middle| g_{1T} \right] \right] \quad (6)$$

$$= \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \frac{1}{m!} \mathbf{E} \left[\Phi^{(m)}(g_{1T}) \sum_{k \in K_{j,m}} C^{j,m,k} \mathbf{E} [X^{j,m,k} | g_{1T}] \right] + o(\epsilon^N) \quad (7)$$

where

$$K_{j,m} = \left\{ (k_1, \dots, k_{j-m+1}); k_n \geq 0, \sum_{n=1}^{j-m+1} k_n = m, \sum_{n=1}^{j-m+1} n k_n = j \right\}$$

and

$$X^{j,m,k} = \prod_{n=1}^{j-m+1} g_{(n+1)T}^{k_n},$$

$$C^{j,m,k} = \prod_{n=1}^{j-m+1} \frac{m!}{k_1! \dots k_{j-m+1}!}.$$

3 Computation of Conditional Expectations

3.1 Procedures of Computations

In the previous section, we have

$$\mathbf{E}[\Phi(G^{(\epsilon)})] = \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \frac{1}{m!} \mathbf{E} \left[\Phi^{(m)}(g_{1T}) \sum_{k \in K_{j,m}} C^{j,m,k} \mathbf{E} [X^{j,m,k} | g_{1T}] \right] + o(\epsilon^N). \quad (8)$$

Then, if we obtain conditional expectations appearing in this expression explicitly, it can be easily calculated since g_{1T} follows a normal distribution. In particular, letting Φ be δ_x , the delta function at $x \in \mathbf{R}$, the asymptotic expansion of the density function of $G^{(\epsilon)}$ can be obtained as in (28) in the next section.

Here we describe the procedures of evaluating these conditional expectations.

At the beginning of this subsection, we state the following proposition playing an important role in the evaluation.

Proposition 1 *Let $J_n(f_n)$ denote the n -times iterated Itô integral of $L^2(\mathbf{T}^n)$ -function f_n :*

$$J_n(f_n) := \int_0^T \int_0^{t_1} \cdots \int_0^{t_{n-1}} f_n(t_1, \dots, t_n) dW_{t_n} \cdots dW_{t_2} dW_{t_1}$$

for $n \geq 1$ and $J_0(f_0) := f_0$ (constant).

Then, its expectation conditional on $J_1(q) = x$ is given by

$$\mathbf{E}[J_n(f_n) | J_1(q) = x] = \left(\int_0^T \int_0^{t_1} \cdots \int_0^{t_{n-1}} f_n(t_1, \dots, t_n) q(t_1) \cdots q(t_n) dt_n \cdots dt_2 dt_1 \right) \frac{H_n(x; \|q\|_{L^2(\mathbf{T})}^2)}{(\|q\|_{L^2(\mathbf{T})}^2)^n} \quad (9)$$

where $\mathbf{T} = [0, T]$, $t_i \in \mathbf{T}$ ($i = 1, 2, \dots, n$) and $H_n(x; \Sigma)$ is the Hermite polynomial of degree n which is defined as

$$H_n(x; \Sigma) := (-\Sigma)^n e^{x^2/2\Sigma} \frac{d^n}{dx^n} e^{-x^2/2\Sigma}.$$

(proof) See Section 3.2.□

Next, we show how to compute the conditional expectations in (8). In the rest of this subsection, we assume $\partial_\epsilon V_0(X_t^{(0)}, 0) \equiv (0, \dots, 0)$ with no loss of generality, and set $q(t) = a_t = (\partial g(X_T^{(0)}))' [Y_T Y_t^{-1} V(X_t^{(0)})]$ (then $J_1(q) = g_{1T}$ and $\|q\|_{L^2(\mathbf{T})}^2 = \Sigma_T$). If this assumption is not satisfied, we can obtain almost the same result by taking conditional expectations with respect to

$$\hat{g}_{1T} := g_{1T} - C$$

instead of g_{1T} , where

$$C := \left(\partial g(X_T^{(0)}) \right)' \int_0^T Y_T Y_t^{-1} \partial_\epsilon V_0(X_t^{(0)}, 0) dt.$$

The procedures consist of three steps.

1. The way to derive an expansion of $A_{tt}^i = \frac{\partial^l X_t^{(\epsilon)}}{\partial \epsilon^l} |_{\epsilon=0}$ is explained. In this stage, there are two alternative ways.

- In one way, as in Lemma 2 in Section 3.2, A_{lT}^i can be expanded as a summation of at most l iterated Itô integrals whose integrands are a family of symmetric $L^2(\mathbf{T}^l)$ -functions $\{\hat{f}_{l'}^{i,l}\}_{l'=1}^l$:

$$A_{tt}^i = \sum_{l'=1}^l J_{l'}(\hat{f}_{l'}^{i,l}) \quad (10)$$

The integrand of l' -times iterated Itô integral in this expansion is given by the expectation of the l' -th Malliavin derivative of A_{lT}^i :

$$\hat{f}_{l'}^{i,l}(t_1, \dots, t_{l'}) = \mathbf{E} [D_{t_1, \dots, t_{l'}} A_{lT}^i]. \quad (11)$$

- In fact, as we can see in (3) every A_{lT}^i is given by finite operations of multiplication, (Lebesgue) integration with respect to time parameters and stochastic integrations. Then, the alternative expansion of A_{lT}^i can be directly calculated via iterated use of Itô's formula:

$$A_{lT}^i = \sum_{l'=0}^l J_{l'}(f_{l'}^{i,l}) \quad (12)$$

- We here briefly advert to the relationship between $\hat{f}_{l'}^{i,l}$ and $f_{l'}^{i,l}$. Note that A_{lT}^i has its Wiener-Chaos expansion as in the proof of Lemma 2 which is described in Section 3.2

$$A_{tt}^i = \sum_{l'=1}^l I_{l'}(\tilde{f}_{l'}^{i,l}) \quad (13)$$

and that the integrand of l' -th order multiple Wiener-Itô integral is given by

$$\tilde{f}_{l'}^{i,l}(t_1, \dots, t_{l'}) = \frac{1}{l'!} \mathbf{E} [D_{t_1, \dots, t_{l'}} A_{lT}^i].$$

Then, due to the relationship between an iterated Itô integral and a multiple Wiener-Itô integral of the same order shown in Lemma 1 in Section 3.2, $\hat{f}_{l'}^{i,l} = l'! \tilde{f}_{l'}^{i,l}$ actually coincides with a symmetrization (unnormalized with respect to its norm) of $f_{l'}^{i,l}$;

$$\hat{f}_{l'}^{i,l}(t_1, \dots, t_{l'}) = \sum_{\sigma} 1_{\{t_{\sigma(1)} \geq \dots \geq t_{\sigma(l')}\}} f_{l'}^{i,l}(t_{\sigma(1)}, \dots, t_{\sigma(l')}).$$

2. From the expansion of A_{lT}^i , we derive that of g_{nT} . Recall that for $n \geq 1$

$$g_{nT} = \sum_{\vec{s} \in S_n} \left(\frac{n!}{s_1! \dots s_n!} \right) \prod_{l=1}^n \left(\frac{1}{l!} \right)^{s_l} \sum_{\vec{p}^{s_l} \in P_{s_l}} \left(\frac{s_l!}{p_1^{s_l}! \dots p_d^{s_l}!} \right) \partial_1^{p_1^{s_l}} \dots \partial_d^{p_d^{s_l}} g(X_T^{(0)}) \prod_{i=1}^d (A_{iT}^i)^{p_i^{s_l}}$$

where

$$S_n = \left\{ \vec{s} = (s_1, \dots, s_n); s_l \geq 0, \sum_{l=1}^n l s_l = n \right\} \text{ and } P_s = \left\{ \vec{p}^s = (p_1^s, \dots, p_d^s); p_i^s \geq 0, \sum_{i=1}^d p_i^s = s \right\}.$$

Then, the expansions of g_{nT} are obtained by applying Itô's formula iteratively. Moreover, noting that the highest order of the expansion of g_{nT} is n , g_{nT} is expressed as

$$g_{nT} = \sum_{i=0}^n J_i(f_i^{g_n})$$

with integrands $\{f_i^{g_n}\}_{i=1}^n$ obtained via Itô's formula.

3. Again by iterative applications of Itô's formula to

$$X^{j,m,k} = \prod_{n=1}^{j-m+1} g_{(n+1)T}^{k_n}$$

$$\text{where } K_{j,m} = \left\{ (k_1, \dots, k_{j-m+1}); k_n \geq 0, \sum_{n=1}^{j-m+1} k_n = m, \sum_{n=1}^{j-m+1} nk_n = j \right\},$$

we now have the expansion of $X^{j,m,k}$ in (8) with a finite number of terms as

$$X^{j,m,k} = \sum_{n'=0}^{j+m} J_{n'}(f_{n'}^{j,m,k}) \quad (14)$$

with integrands $\{f_{n'}^{j,m,k}\}_{n'=1}^{j+m}$ which can be deduced from $\{\hat{f}_{l'}^{i,l}\}_{l'=1}^l$ in (10) or $\{f_{l'}^{i,l}\}_{l'=1}^l$ in (12).

4. From Proposition 1, we conclude that the conditional expectations in (8) are given by

$$\mathbf{E} [X^{j,m,k} | g_{1T}] = \sum_{n'=0}^{j+m} \left(\int_0^T \int_0^{t_1} \dots \int_0^{t_{n'-1}} f_{n'}^{j,m,k}(t_1, \dots, t_{n'}) q(t_1) \dots q(t_{n'}) dt_{n'} \dots dt_2 dt_1 \right) \frac{H_{n'}(x; \Sigma_T)}{\Sigma_T^{n'}}. \quad (15)$$

Example 1

At the end of this subsection we show a simple example evaluating $X^{1,1,(1)} = g_{2T}$ in order to make these procedures clear. Let consider the case when $d = d' = 1$ and $V_0(x, \epsilon) \equiv 0$. In this case g_{2T} is given by

$$g_{2T} = \frac{1}{2} \partial^2 g(X_T^{(0)}) (A_{1T})^2 + \frac{1}{2} \partial g(X_T^{(0)}) A_{2T}$$

where

$$\begin{aligned} A_{1T} &= \int_0^T V(X_u^{(0)}) dW_u, \\ A_{2T} &= 2 \int_0^T \partial V(X_u^{(0)}) A_{1u} dW_u. \end{aligned}$$

Then, it can be decomposed as the sum of $J_n(\cdot)$ by Itô's formula;

$$g_{2T} = J_0(f_0^{g_2}) + J_2(f_2^{g_2})$$

where

$$\begin{aligned} f_0^{g_2} &= \frac{1}{2} \partial^2 g(X_T^{(0)}) \int_0^T V(X_u^{(0)})^2 du, \\ f_2^{g_2}(t_1, t_2) &= \partial^2 g(X_T^{(0)}) V(X_{t_2}^{(0)}) V(X_{t_1}^{(0)}) + \partial g(X_T^{(0)}) V(X_{t_2}^{(0)}) \partial V(X_{t_1}^{(0)}). \end{aligned}$$

From Proposition 1, it follows that

$$\begin{aligned} \mathbf{E} \left[X^{1,1,(1)} | g_{1T} \right] &= \left(\partial^2 g(X_T^{(0)}) \partial g(X_T^{(0)})^2 \int_0^T \int_0^{t_1} V(X_{t_2}^{(0)})^2 dt_2 V(X_{t_1}^{(0)})^2 dt_1 \right. \\ &\quad \left. + \partial g(X_T^{(0)})^3 \int_0^T \int_0^{t_1} V(X_{t_2}^{(0)})^2 dt_2 \partial V(X_{t_1}^{(0)}) V(X_{t_1}^{(0)}) dt_1 \right) \frac{H_2(g_{1T}; \Sigma_T)}{\Sigma_T^2} \\ &\quad + \frac{1}{2} \partial^2 g(X_T^{(0)}) \int_0^T V(X_u^{(0)})^2 du. \end{aligned}$$

3.2 Proof of Lemmas and Proposition in Section 3.1

In this subsection we introduce and prove the important proposition and lemmas used in Section 3.1.

Proposition 1 *The expectation of n -times iterated Itô integral $J_n(f_n)$ conditional on $J_1(q) = x$ is given by*

$$\mathbf{E}[J_n(f_n) | J_1(q) = x] = \left(\int_0^T \int_0^{t_1} \cdots \int_0^{t_{n-1}} f_n(t_1, \dots, t_n) q(t_1) \cdots q(t_n) dt_n \cdots dt_2 dt_1 \right) \frac{H_n(x; \|q\|_{L^2(\mathbf{T})}^2)}{(\|q\|_{L^2(\mathbf{T})}^2)^n}.$$

(*proof*) This can be considered as a version of Proposition 3 of Nualart, Üstünel and Zakai [36].

Let $I_n(\hat{f})$ denote the multiple Wiener-Itô integral of n -th order of its integrand $\hat{f} \in L_{sym}^2(\mathbf{T}^n)$, that is \hat{f} is an element of the space of square-integrable symmetric functions from \mathbf{T}^n to \mathbf{R} .

Then, from Proposition 3 of [36], we know

$$\mathbf{E}[I_n(\hat{f}) | I_1(q) = x] = \left(\int \cdots \int_{\mathbf{T}^n} \hat{f}(t_1, \dots, t_n) q(t_1) \cdots q(t_n) dt_n \cdots dt_2 dt_1 \right) \frac{H_n(x; \|q\|_{L^2(\mathbf{T})}^2)}{(\|q\|_{L^2(\mathbf{T})}^2)^n}.$$

Substituting a symmetrization of f_n defined as in (17) in Lemma 1 below, we obtain the result:

$$\begin{aligned} &\mathbf{E}[J_n(f_n) | J_1(q) = x] \\ &= \mathbf{E}[I_n(\hat{f}_n) | I_1(q) = x] \\ &= \left(\int \cdots \int_{\mathbf{T}^n} \hat{f}_n(t_1, \dots, t_n) q(t_1) \cdots q(t_n) dt_n \cdots dt_2 dt_1 \right) \frac{H_n(x; \|q\|_{L^2(\mathbf{T})}^2)}{(\|q\|_{L^2(\mathbf{T})}^2)^n} \\ &= \frac{1}{n!} \sum_{\sigma} \left(\int \cdots \int_{\mathbf{T}^n} \mathbf{1}_{\{t_{\sigma(1)} \geq \cdots \geq t_{\sigma(n)}\}} f_n(t_{\sigma(1)}, \dots, t_{\sigma(n)}) q(t_1) \cdots q(t_n) dt_n \cdots dt_2 dt_1 \right) \frac{H_n(x; \|q\|_{L^2(\mathbf{T})}^2)}{(\|q\|_{L^2(\mathbf{T})}^2)^n} \\ &= \left(\int_0^T \int_0^{t_1} \cdots \int_0^{t_{n-1}} f_n(t_1, \dots, t_n) q(t_1) \cdots q(t_n) dt_n \cdots dt_2 dt_1 \right) \frac{H_n(x; \|q\|_{L^2(\mathbf{T})}^2)}{(\|q\|_{L^2(\mathbf{T})}^2)^n}. \square \end{aligned}$$

The following lemma gives us the relationship between the iterated Itô integral and the multiple Wiener-Itô integral of the same order.

Lemma 1 *For any $L^2(\mathbf{T}^n)$ -function f_n which is not necessarily symmetric it holds that*

$$J_n(f_n) = I_n(\hat{f}_n) \tag{16}$$

where \hat{f}_n is a symmetrization of f_n defined by

$$\hat{f}_n(t_1, \dots, t_n) := \frac{1}{n!} \sum_{\sigma} \mathbf{1}_{\{t_{\sigma(1)} \geq \dots \geq t_{\sigma(n)}\}} f_n(t_{\sigma(1)}, \dots, t_{\sigma(n)}) \quad (17)$$

with taking summation over all permutations σ .

(proof) The assertion can be easily shown:

$$\begin{aligned} I_n(\hat{f}) &= n! J_n(\hat{f}) \\ &= n! J_n \left(\frac{1}{n!} \sum_{\sigma} \mathbf{1}_{\{t_{\sigma(1)} \leq \dots \leq t_{\sigma(n)}\}} f(t_{\sigma(1)}, \dots, t_{\sigma(n)}) \right) \\ &= \sum_{\sigma} \int_0^T \int_0^{t_1} \dots \int_0^{t_{n-1}} \mathbf{1}_{\{t_{\sigma(1)} \leq \dots \leq t_{\sigma(n)}\}} f(t_{\sigma(1)}, \dots, t_{\sigma(n)}) dW_{t_n} \dots dW_{t_2} dW_{t_1} \\ &= \int_0^T \int_0^{t_1} \dots \int_0^{t_{n-1}} f(t_1, \dots, t_n) dW_{t_n} \dots dW_{t_2} dW_{t_1} = J_n(f). \square \end{aligned}$$

Finally we introduce and prove the following lemma.

Lemma 2 Let $A_{t_1}^i := \frac{\partial^l X_t^{(\epsilon)}}{\partial \epsilon^l} |_{\epsilon=0}$ as in Section 2. Then, it has an expansion with a finite number of iterated Itô integrals as

$$A_{t_1}^i = \sum_{l'=1}^l J_{l'}(f_{l'}^{i,l}) \quad (18)$$

with $L^2(\mathbf{T}^{l'})$ -function $\{f_{l'}^{i,l}\}$ whose derivation was explained in Section 3.1.

Before the proof of Lemma 2 we state the following lemma.

Lemma 3 Assume $d' = 1$ for simplicity. Then, the n -th Malliavin derivative of $X_t^{(\epsilon),i}$ are given by

$$\begin{aligned} D_{t_1, \dots, t_n} X_t^{(\epsilon),i} &= \epsilon \sum_{r=1}^n \alpha_{n-1}^i(t_r; t_1, \dots, t_{r-1}, t_{r+1}, \dots, t_n; \epsilon) \\ &\quad + \epsilon \int_{t_1 \vee \dots \vee t_n}^t \alpha_n^i(s, t_1, \dots, t_n; \epsilon) dW_s + \int_{t_1 \vee \dots \vee t_n}^t \beta_n^i(s, t_1, \dots, t_n; \epsilon) ds \end{aligned} \quad (19)$$

for $t \geq t_1 \vee \dots \vee t_n$ and zero for $t < t_1 \vee \dots \vee t_n$, where

$$\alpha_n^i(s, t_1, \dots, t_n; \epsilon) := \sum \partial_{k_1} \dots \partial_{k_\nu} V^i(X_s^{(\epsilon)}) \prod_{p=1}^{\nu} D_{t(M_p)} X_s^{(\epsilon), k_p}, \quad (20)$$

$$\text{and } \beta_n^i(s, t_1, \dots, t_n; \epsilon) := \sum \partial_{k_1} \dots \partial_{k_\nu} V_0^i(X_s^{(\epsilon)}, \epsilon) \prod_{p=1}^{\nu} D_{t(M_p)} X_s^{(\epsilon), k_p} \quad (21)$$

where $t(M_p) = t_{r_1}, \dots, t_{r_p}$ for $M_p = \{r_1, \dots, r_p; r_1 < \dots < r_p\} \subset \{1, \dots, n\}$, and the sums are taken under the set of all partitions $\{M_p\}_{p=1}^{\nu}$ such that $M_1 \cup \dots \cup M_\nu = \{1, \dots, n\}$.

(proof) See pp.123-124 in Nualart [35].□

(proof of Lemma 2) From Lemma 1, it is equivalent to show that A_{lT}^i has a Wiener-Chaos expansion with a finite number of terms as

$$A_{lT}^i = \sum_{l'=1}^l I_{l'}(\tilde{f}_{l'}^{i,l}) \quad (22)$$

with $L_{sym}^2(\mathbf{T}^{l'})$ -functions $\{\tilde{f}_{l'}^{i,l}\}$. Since $\tilde{f}_{l'}^l$ is given by

$$\begin{aligned} \tilde{f}_{l'}^{i,l}(t_1, \dots, t_{l'}) &= \frac{1}{l'!} \mathbf{E} [D_{t_1, \dots, t_{l'}} A_{lT}^i] = \frac{1}{l'!} \mathbf{E} \left[D_{t_1, \dots, t_{l'}} \frac{\partial^l}{\partial \epsilon^l} \Big|_{\epsilon=0} X_T^{(\epsilon), i} \right] \\ &= \frac{1}{l'!} \mathbf{E} \left[\frac{\partial^l}{\partial \epsilon^l} \Big|_{\epsilon=0} D_{t_1, \dots, t_{l'}} X_T^{(\epsilon), i} \right] \end{aligned}$$

where the last equality holds due to uniqueness of the asymptotic expansion of $X_T^{(\epsilon), i}$, in order to prove the expansion (22) it is sufficient to see that for any $l' > l$,

$$Y_{l'T}^{i,l} := \frac{\partial^l}{\partial \epsilon^l} \Big|_{\epsilon=0} D_{t_1, \dots, t_{l'}} X_T^{(\epsilon), i}$$

is equal to zero, which will be proved by induction.

First, it is obvious that this statement holds with $l = 0$, because $X_t^{(\epsilon)}$ becomes deterministic as $\epsilon \downarrow 0$.

Second, from Lemma 3, for $l \geq 1$ we have

$$\begin{aligned} Y_{l't}^{i,l} &= l \sum_{r=1}^{l'} \frac{\partial^{l-1}}{\partial \epsilon^{l-1}} \Big|_{\epsilon=0} (\alpha_{l'-1}^i(t_r; t_1, \dots, t_{r-1}, t_{r+1}, \dots, t_{l'}; \epsilon)) \quad (23) \\ &\quad + l \int_{t_1 \vee \dots \vee t_{l'}}^t \frac{\partial^{l-1}}{\partial \epsilon^{l-1}} \Big|_{\epsilon=0} (\alpha_{l'}^i(s, t_1, \dots, t_{l'}; \epsilon)) dW_s + \int_{t_1 \vee \dots \vee t_{l'}}^t \frac{\partial^l}{\partial \epsilon^l} \Big|_{\epsilon=0} (\beta_{l'}^i(s, t_1, \dots, t_{l'}; \epsilon)) ds. \end{aligned}$$

for $t \geq t_1 \vee \dots \vee t_n$ and $Y_{l't}^{i,l} = 0$ for $t < t_1 \vee \dots \vee t_n$.

First and second terms of the right hand side of (23) is summation of the terms (for the second term whose integrands are)

$$\frac{\partial^{l_0}}{\partial \epsilon^{l_0}} \left(\partial_{k_1} \dots \partial_{k_\nu} V^i(X_s^{(\epsilon)}) \right) \prod_{p=1}^{\nu} \frac{\partial^{l_p}}{\partial \epsilon^{l_p}} \Big|_{\epsilon=0} D_{t(M_p)} X_s^{(\epsilon), k_p}$$

with the conditions $\sum_{p=0}^{\nu} l_p = l - 1$ and $\sum_{p=1}^{\nu} \#(M_p) = k$, $k = l' - 1$ for the first term and $k = l'$ for the second term. Thus, since $k > l - 1$ in both cases, for at least one p we have $l_p < \#(M_p)$. Then, all of these terms will vanish as $\epsilon \downarrow 0$ by the assumption of induction (note that $l_p \leq l - 1$).

For the third terms, almost the same result is obtained except that $\int \partial_j V_0^i(X_s^{(\epsilon)}, \epsilon) Y_{l's}^{j,l} ds$ ($j = 1, \dots, d$) remains.

As a consequence, all terms except for

$$\sum_{j=1}^d \int_{t_1 \vee \dots \vee t_{l'}}^t \partial_j V_0^i(X_s^{(0)}, 0) Y_{l's}^{j,l} ds$$

are equal to zero. Thus we have a trivial linear equation

$$Y_{l't}^l = \int_{t_1 \vee \dots \vee t_{l'}}^t \partial V_0(X_s^{(0)}, 0) Y_{l's}^l ds \quad (24)$$

whose solution is given by $Y_{l't}^l \equiv (0, \dots, 0)'$ where $Y_{l't}^l = (Y_{l't}^{1,l}, \dots, Y_{l't}^{d,l})'$. □

3.3 Useful formulas

Finally, we here list up some formulas of conditional expectations often used in asymptotic expansions. Let $q_i : [0, T] \mapsto \mathbf{R}^m, i = 1, 2, 3, 4, 5, 6, 7$ are non-random functions and we define Σ as

$$\Sigma = \int_0^T q'_{1v} q_{1v} dv,$$

where z' is the transpose of z . We assume that $0 < \Sigma < \infty$ and integrability in the following formulas.

Before the list of formulas, we define a notation of iterated integrations for convenience;

$$F_n(f_1, \dots, f_n) := \int_0^T \int_0^{t_1} \dots \int_0^{t_{n-1}} f_1(t_1) \dots f_n(t_n) dt_n \dots dt_1, \quad n \geq 1. \quad (25)$$

1.

$$\begin{aligned} & \mathbf{E} \left[\int_0^T q'_{2t} dW_t \mid \int_0^T q'_{1v} dW_v = x \right] \\ &= F_1(q'_2 q_1) \frac{H_1(x; \Sigma)}{\Sigma} \end{aligned}$$

2.

$$\begin{aligned} & \mathbf{E} \left[\int_0^T \int_0^t q'_{2u} dW_u q'_{3t} dW_t \mid \int_0^T q'_{1v} dW_v = x \right] \\ &= F_2(q'_2 q_1, q'_3 q_1) \frac{H_2(x; \Sigma)}{\Sigma^2} \end{aligned}$$

3.

$$\begin{aligned} & \mathbf{E} \left[\left(\int_0^T q'_{2u} dW_u \right) \left(\int_0^T q'_{3s} dW_s \right) \mid \int_0^T q'_{1v} dW_v = x \right] \\ &= \left(F_1(q'_2 q_1) \times F_1(q'_3 q_1) \right) \frac{H_2(x; \Sigma)}{\Sigma^2} + F_1(q'_2 q_3) \end{aligned}$$

4.

$$\begin{aligned} & \mathbf{E} \left[\int_0^T \int_0^t \int_0^s q'_{2u} dW_u q'_{3s} dW_s q'_{4t} dW_t \mid \int_0^T q'_{1v} dW_v = x \right] \\ &= F_3(q'_2 q_1, q'_3 q_1, q'_4 q_1) \frac{H_3(x; \Sigma)}{\Sigma^3} \end{aligned}$$

5.

$$\begin{aligned} & \mathbf{E} \left[\int_0^T \left(\int_0^t q'_{2u} dW_u \right) \left(\int_0^t q'_{3s} dW_s \right) q'_{4t} dW_t \mid \int_0^T q'_{1v} dW_v = x \right] \\ &= \left(F_3(q'_2 q_1, q'_3 q_1, q'_4 q_1; T) + F_3(q'_3 q_1, q'_2 q_1, q'_4 q_1) \right) \frac{H_3(x; \Sigma)}{\Sigma^3} + F_2(q'_2 q_3, q'_4 q_1) \frac{H_1(x; \Sigma)}{\Sigma} \end{aligned}$$

6.

$$\begin{aligned} & \mathbf{E} \left[\left(\int_0^T \int_0^t q'_{2s} dW_s q'_{3t} dW_t \right) \left(\int_0^T q'_{4u} dW_u \right) \middle| \int_0^T q'_{1v} dW_v = x \right] \\ &= \left(F_2(q'_2 q_1, q'_3 q_1) \times F_1(q'_4 q_1) \right) \frac{H_3(x; \Sigma)}{\Sigma^3} + \left(F_2(q'_2 q_4, q'_3 q_1) + F_2(q'_2 q_1, q'_3 q_4) \right) \frac{H_1(x; \Sigma)}{\Sigma} \end{aligned}$$

7.

$$\begin{aligned} & \mathbf{E} \left[\left(\int_0^T \int_0^t q'_{2s} dW_s q'_{3t} dW_t \right) \left(\int_0^T \int_0^r q'_{4u} dW_u q'_{5r} dW_r \right) \middle| \int_0^T q'_{1v} dW_v = x \right] \\ &= \left(F_2(q'_2 q_1, q'_3 q_1) \times F_2(q'_4 q_1, q'_5 q_1) \right) \frac{H_4(x; \Sigma)}{\Sigma^4} \\ &+ \left\{ \left(F_3(q'_2 q_4, q'_5 q_1, q'_3 q_1) + F_3(q'_2 q_1, q'_3 q_4, q'_5 q_1) + F_3(q'_2 q_1, q'_4 q_1, q'_3 q_5) \right) \right. \\ &+ \left. \left(F_3(q'_2 q_4, q'_3 q_1, q'_5 q_1) + F_3(q'_4 q_1, q'_2 q_5, q'_3 q_1) + F_3(q'_4 q_1, q'_2 q_1, q'_3 q_5) \right) \right\} \frac{H_2(x; \Sigma)}{\Sigma^2} \\ &+ F_2(q'_2 q_4, q'_3 q_5) \\ &=: \tilde{F}_4^7(q_1, q_2, q_3, q_4, q_5) \frac{H_4(x; \Sigma)}{\Sigma^4} + \tilde{F}_2^7(q_1, q_2, q_3, q_4, q_5) \frac{H_2(x; \Sigma)}{\Sigma^2} + \tilde{F}_0^7(q_1, q_2, q_3, q_4, q_5) \end{aligned}$$

8.

$$\begin{aligned} & \mathbf{E} \left[\left(\int_0^T q'_{2t} dW_t \right) \left(\int_0^T q'_{3s} dW_s \right) \left(\int_0^T \int_0^r q'_{4u} dW_u q'_{5r} dW_r \right) \middle| \int_0^T q'_{1v} dW_v = x \right] \\ &= \left(\tilde{F}_4^7(q_1, q_2, q_3, q_4, q_5) + \tilde{F}_4^7(q_1, q_3, q_2, q_4, q_5) \right) \frac{H_4(x; \Sigma)}{\Sigma^4} \\ &\left(\tilde{F}_2^7(q_1, q_2, q_3, q_4, q_5) + \tilde{F}_2^7(q_1, q_3, q_2, q_4, q_5) + F_1(q'_2 q_3) \times F_2(q'_4 q_1, q'_5 q_1) \right) \frac{H_2(x; \Sigma)}{\Sigma^2} \\ &+ \left(\tilde{F}_2^7(q_1, q_2, q_3, q_4, q_5) + \tilde{F}_2^7(q_1, q_3, q_2, q_4, q_5) \right) \end{aligned}$$

9.

$$\begin{aligned} & \mathbf{E} \left[\int_0^T \int_0^t \int_0^s \int_0^u q'_{2r} dW_r q'_{3u} dW_u q'_{4s} dW_s q'_{5t} dW_t \middle| \int_0^T q'_{1v} dW_v = x \right] \\ &= F_4(q'_2 q_1, q'_3 q_1, q'_4 q_1, q'_5 q_1) \frac{H_4(x; \Sigma)}{\Sigma^4} \end{aligned}$$

10.

$$\begin{aligned} & \mathbf{E} \left[\int_0^T \int_0^t \left(\int_0^u q'_{2s} dW_s \right) \left(\int_0^u q'_{3r} dW_r \right) q'_{4u} dW_u q'_{5t} dW_t \middle| \int_0^T q'_{1v} dW_v = x \right] \\ &= \left(F_4(q'_2 q_1, q'_3 q_1, q'_4 q_1, q'_5 q_1) + F_4(q'_3 q_1, q'_2 q_1, q'_4 q_1, q'_5 q_1) \right) \frac{H_4(x; \Sigma)}{\Sigma^4} \\ &+ F_3(q'_2 q_3, q'_4 q_1, q'_5 q_1) \frac{H_2(x; \Sigma)}{\Sigma^2} \end{aligned}$$

11.

$$\mathbf{E} \left[\int_0^T \left(\int_0^t \int_0^u q'_{2r} dW_r q'_{3u} dW_u \right) \left(\int_0^t q'_{4u} dW_u \right) q'_{5t} dW_t \middle| \int_0^T q'_{1v} dW_v = x \right]$$

$$\begin{aligned}
&= \left(F_4(q'_2q_1, q'_3q_1, q'_4q_1, q'_5q_1) + F_4(q'_2q_1, q'_4q_1, q'_3q_1, q'_5q_1) + F_4(q'_4q_1, q'_2q_1, q'_3q_1, q'_5q_1) \right) \frac{H_4(x; \Sigma)}{\Sigma^4} \\
&\quad + \left(F_3(q'_2q_4, q'_3q_1, q'_5q_1) + F_3(q'_2q_1, q'_3q_4, q'_5q_1) \right) \frac{H_2(x; \Sigma)}{\Sigma^2} \\
&=: \tilde{F}_4^{11}(q_1, q_2, q_3, q_4, q_5) \frac{H_4(x; \Sigma)}{\Sigma^4} + \tilde{F}_2^{11}(q_1, q_2, q_3, q_4, q_5) \frac{H_2(x; \Sigma)}{\Sigma^2}
\end{aligned}$$

12.

$$\begin{aligned}
&\mathbf{E} \left[\int_0^T \left(\int_0^t q'_{2s} dW_s \right) \left(\int_0^t q'_{3r} dW_r \right) \left(\int_0^t q'_{4u} dW_u \right) q'_{5t} dW_t \mid \int_0^T q'_{1v} dW_v = x \right] \\
&= \left(\tilde{F}_4^{11}(q_1, q_2, q_3, q_4, q_5) + \tilde{F}_4^{11}(q_1, q_3, q_2, q_4, q_5) \right) \frac{H_4(x; \Sigma)}{\Sigma^4} \\
&\quad + \left\{ \tilde{F}_2^{11}(q_1, q_2, q_3, q_4, q_5) + \tilde{F}_2^{11}(q_1, q_3, q_2, q_4, q_5) + \left(F_3(q'_2q_3, q'_4q_1, q'_5q_1) + F_3(q'_4q_1, q'_2q_3, q'_5q_1) \right) \right\} \frac{H_2(x; \Sigma)}{\Sigma^2}
\end{aligned}$$

13.

$$\begin{aligned}
&\mathbf{E} \left[\left(\int_0^T \int_0^t q'_{2s} dW_s q'_{3t} dW_t \right) \left(\int_0^T \int_0^t \int_0^u q'_{4s} dW_s q'_{5u} dW_u q'_{6t} dW_t \right) \mid \int_0^T q'_{1v} dW_v = x \right] \\
&= \left(F_2(q'_2q_1, q'_3q_1) \times F_3(q'_4q_1, q'_5q_1, q'_6q_1) \right) \frac{H_5(x; \Sigma)}{\Sigma^5} \\
&\quad + \left\{ \left(F_4(q'_4q_1, q'_5q_1, q'_2q_1, q'_3q_6) + F_4(q'_4q_1, q'_2q_1, q'_5q_1, q'_3q_6) + F_4(q'_2q_1, q'_4q_1, q'_5q_1, q'_3q_6) \right) \right. \\
&\quad + \left(F_4(q'_4q_1, q'_3q_5, q'_2q_1, q'_6q_1) + F_4(q'_4q_1, q'_2q_1, q'_3q_5, q'_3q_1) + F_4(q'_2q_1, q'_4q_1, q'_3q_5, q'_3q_1) \right) \\
&\quad + \left(F_4(q'_2q_1, q'_3q_4, q'_5q_1, q'_6q_1) + F_4(q'_4q_1, q'_2q_5, q'_3q_1, q'_6q_1) \right. \\
&\quad \quad \left. \left. + F_4(q'_2q_4, q'_3q_1, q'_5q_1, q'_6q_1) + F_4(q'_2q_4, q'_5q_1, q'_3q_1, q'_6q_1) \right) \right\} \frac{H_3(x; \Sigma)}{\Sigma^3} \\
&\quad + \left(F_4(q'_4q_1, q'_5q_1, q'_2q_6, q'_3q_1) + F_4(q'_4q_1, q'_2q_5, q'_6q_1, q'_3q_1) + F_4(q'_2q_4, q'_5q_1, q'_6q_1, q'_3q_1) \right) \frac{H_3(x; \Sigma)}{\Sigma^3} \\
&\quad + \left(F_3(q'_4q_1, q'_2q_5, q'_3q_6) + F_3(q'_2q_4, q'_5q_1, q'_3q_6) + F_3(q'_2q_4, q'_3q_5, q'_6q_1) \right) \frac{H_1(x; \Sigma)}{\Sigma} \\
&=: \tilde{F}_5^{13}(q_1, q_2, q_3, q_4, q_5, q_6) \frac{H_5(x; \Sigma)}{\Sigma^5} + \tilde{F}_3^{13}(q_1, q_2, q_3, q_4, q_5, q_6) \frac{H_3(x; \Sigma)}{\Sigma^3} + \tilde{F}_1^{13}(q_1, q_2, q_3, q_4, q_5, q_6) \frac{H_1(x; \Sigma)}{\Sigma}
\end{aligned}$$

14.

$$\begin{aligned}
&\mathbf{E} \left[\left(\int_0^T \int_0^t q'_{2s} dW_s q'_{3t} dW_t \right) \left(\int_0^T \left(\int_0^t q'_{4s} dW_s \right) \left(\int_0^t q'_{5u} dW_u \right) q'_{6t} dW_t \right) \mid \int_0^T q'_{1v} dW_v = x \right] \\
&= \left(\tilde{F}_5^{13}(q_1, q_2, q_3, q_4, q_5, q_6) + \tilde{F}_5^{13}(q_1, q_2, q_3, q_5, q_4, q_6) \right) \frac{H_5(x; \Sigma)}{\Sigma^5} \\
&\quad + \left(\tilde{F}_3^{13}(q_1, q_2, q_3, q_4, q_5, q_6) + \tilde{F}_3^{13}(q_1, q_2, q_3, q_5, q_4, q_6) + F_2(q'_2q_1, q'_3q_1) \times F_2(q'_4q_5, q'_6q_1) \right) \frac{H_3(x; \Sigma)}{\Sigma^3} \\
&\quad + \left(\tilde{F}_1^{13}(q_1, q_2, q_3, q_4, q_5, q_6) + \tilde{F}_1^{13}(q_1, q_2, q_3, q_5, q_4, q_6) \right. \\
&\quad \quad \left. + F_3(q'_4q_5, q'_2q_6, q'_3q_1) + F_3(q'_2q_1, q'_4q_5, q'_3q_6) + F_3(q'_4q_5, q'_2q_1, q'_3q_6) \right) \frac{H_1(x; \Sigma)}{\Sigma}
\end{aligned}$$

15.

$$\begin{aligned}
& \mathbf{E} \left[\left(\int_0^T \int_0^t q'_{2u} dW_s q'_{3t} dW_t \right) \left(\int_0^T \int_0^t q'_{4u} dW_u q'_{5t} dW_r \right) \left(\int_0^T \int_0^t q'_{6u} dW_u q'_{7t} dW_r \right) \middle| \int_0^T q'_{1v} dW_v = x \right] \\
&= \left(F_2(q'_2 q_1, q'_3 q_1) \times F_2(q'_4 q_1, q'_5 q_1) \times F_2(q'_6 q_1, q'_7 q_1) \right) \frac{H_6(x; \Sigma)}{\Sigma^6} \\
&+ \left(\tilde{F}_{4*}^{14}(q_1, q_2, q_3, q_4, q_5, q_6, q_7) + \tilde{F}_{4*}^{14}(q_1, q_4, q_5, q_6, q_7, q_2, q_3) + \tilde{F}_{4*}^{14}(q_1, q_6, q_7, q_2, q_3, q_4, q_5) \right) \frac{H_4(x; \Sigma)}{\Sigma^4} \\
&+ \left(\tilde{F}_{2*}^{14}(q_1, q_2, q_3, q_4, q_5, q_6, q_7) + \tilde{F}_{2*}^{14}(q_1, q_4, q_5, q_6, q_7, q_2, q_3) + \tilde{F}_{2*}^{14}(q_1, q_6, q_7, q_2, q_3, q_4, q_5) \right) \frac{H_2(x; \Sigma)}{\Sigma^2} \\
&+ \left(F_3(q'_2 q_6, q'_4 q_7, q'_3 q_5) + F_3(q'_4 q_6, q'_2 q_7, q'_3 q_5) \right. \\
&\quad \left. + F_3(q'_2 q_4, q'_3 q_6, q'_5 q_7) + F_3(q'_2 q_6, q'_3 q_4, q'_5 q_7) + F_3(q'_4 q_6, q'_2 q_5, q'_3 q_7) + F_3(q'_2 q_4, q'_5 q_6, q'_3 q_7) \right)
\end{aligned}$$

where

$$\begin{aligned}
& \tilde{F}_{4*}^{14}(q_1, q_2, q_3, q_4, q_5, q_6, q_7) \\
&:= \left(F_3(q'_4 q_6, q'_5 q_1, q'_7 q_1) + F_3(q'_4 q_6, q'_7 q_1, q'_5 q_1) + F_3(q'_6 q_1, q'_4 q_7, q'_5 q_1) \right. \\
&\quad \left. + F_3(q'_4 q_1, q'_5 q_6, q'_7 q_1) + F_3(q'_4 q_1, q'_6 q_1, q'_5 q_7) + F_3(q'_6 q_1, q'_4 q_1, q'_5 q_7) \right) \times F_2(q'_2 q_1, q'_3 q_1), \\
& \tilde{F}_{2*}^{14}(q_1, q_2, q_3, q_4, q_5, q_6, q_7) \\
&:= F_2(q'_2 q_1, q'_3 q_1) \times F_2(q'_4 q_6, q'_5 q_7) \\
&\quad + \left(F_4(q'_2 q_4, q'_5 q_1, q'_3 q_6, q'_7 q_1) + F_4(q'_2 q_4, q'_3 q_6, q'_5 q_1, q'_7 q_1) + F_4(q'_2 q_4, q'_3 q_6, q'_7 q_1, q'_5 q_1) \right. \\
&\quad \left. + F_4(q'_2 q_6, q'_7 q_1, q'_3 q_4, q'_5 q_1) + F_4(q'_2 q_6, q'_3 q_4, q'_7 q_1, q'_5 q_1) + F_4(q'_2 q_6, q'_3 q_4, q'_5 q_1, q'_7 q_1) \right) \\
&\quad + \left(F_4(q'_4 q_1, q'_2 q_5, q'_3 q_6, q'_7 q_1) + F_4(q'_4 q_1, q'_2 q_6, q'_3 q_5, q'_7 q_1) + F_4(q'_4 q_1, q'_2 q_6, q'_7 q_1, q'_3 q_5) \right. \\
&\quad \left. + F_4(q'_2 q_6, q'_4 q_1, q'_3 q_5, q'_7 q_1) + F_4(q'_2 q_6, q'_4 q_1, q'_7 q_1, q'_3 q_5) + F_4(q'_2 q_6, q'_7 q_1, q'_4 q_1, q'_3 q_5) \right) \\
&\quad + \left(F_4(q'_6 q_1, q'_2 q_7, q'_3 q_4, q'_5 q_1) + F_4(q'_6 q_1, q'_2 q_4, q'_3 q_7, q'_5 q_1) + F_4(q'_6 q_1, q'_2 q_4, q'_5 q_1, q'_3 q_7) \right. \\
&\quad \left. + F_4(q'_2 q_4, q'_6 q_1, q'_3 q_7, q'_5 q_1) + F_4(q'_2 q_4, q'_6 q_1, q'_5 q_1, q'_3 q_7) + F_4(q'_2 q_4, q'_5 q_1, q'_6 q_1, q'_3 q_7) \right) \\
&\quad + \left(F_4(q'_4 q_1, q'_2 q_5, q'_6 q_1, q'_3 q_7) + F_4(q'_4 q_1, q'_6 q_1, q'_2 q_5, q'_3 q_7) + F_4(q'_4 q_1, q'_6 q_1, q'_2 q_7, q'_3 q_5) \right. \\
&\quad \left. + F_4(q'_6 q_1, q'_4 q_1, q'_2 q_5, q'_3 q_7) + F_4(q'_6 q_1, q'_4 q_1, q'_4 q_1, q'_3 q_5) + F_4(q'_6 q_1, q'_2 q_7, q'_2 q_7, q'_3 q_5) \right)
\end{aligned}$$

4 New Computational Scheme

In this section we propose a new computational scheme in asymptotic expansion which is alternative to the method described in the previous section. To compute conditional expectations in the right hand side of (7), we use the following lemma which can be derived from the property of Hermite polynomials.

Lemma 4 Let $X \in L^2(\Omega)$ and Y be a random variable with Gaussian distribution with mean 0 and variance Σ . Then, the conditional expectation $E[X|Y]$ has following expansion in $L^2(\Omega)$:

$$E[X|Y] = \sum_{n=0}^{\infty} a_n H_n(Y; \Sigma) \quad (26)$$

where $H_n(x; \Sigma)$ is the Hermite polynomial of degree n which is defined as

$$H_n(x; \Sigma) = (-\Sigma)^n e^{x^2/2\Sigma} \frac{d^n}{dx^n} e^{-x^2/2\Sigma}$$

and coefficients a_n are given by

$$a_n = \frac{1}{(i\Sigma)^n} \frac{d^n}{d\xi^n} \Big|_{\xi=0} \left\{ e^{\frac{\xi^2}{2}\Sigma} \mathbf{E}[e^{i\xi Y} X] \right\}. \quad (27)$$

(proof) Since the Hermite polynomials $\{H_n(x; \Sigma)\}$ is the orthogonal basis of $L^2(\mathbf{R}, \mu)$ where μ is the Gaussian measure on \mathbf{R} with mean 0 and variance Σ , and $E[X|Y = y] \in L^2(\mathbf{R}, \mu)$, we have the following unique expansion of $E[X|Y = y]$ in $L^2(\mathbf{R}, \mu)$:

$$E[X|Y = y] = \sum_{n=0}^{\infty} a_n H_n(y; \Sigma)$$

and also we have

$$E[X|Y] = \sum_{n=0}^{\infty} a_n H_n(Y; \Sigma)$$

in $L^2(\Omega)$. And note that

$$e^{i\xi Y} = e^{-\frac{\xi^2}{2}\Sigma} \sum_{n=0}^{\infty} \frac{H_n(Y; \Sigma)}{n!} (i\xi)^n.$$

Then,

$$\begin{aligned} e^{\frac{\xi^2}{2}\Sigma} \mathbf{E}[e^{i\xi Y} X] &= e^{\frac{\xi^2}{2}\Sigma} \mathbf{E}[e^{i\xi Y} \mathbf{E}[X|Y]] \\ &= \mathbf{E}\left[\sum_{m=0}^{\infty} \frac{H_m(Y; \Sigma)}{m!} (i\xi)^m \sum_{n=0}^{\infty} a_n H_n(Y; \Sigma) \right] \\ &= \sum_{n=0}^{\infty} a_n (i\Sigma)^n \xi^n. \end{aligned}$$

Comparing to the coefficients of the Taylor series of $e^{\frac{\xi^2}{2}\Sigma} \mathbf{E}[e^{i\xi Y} X]$ around 0 with respect to ξ , we see that a_n can be written as (27). \square

Recall \hat{g}_{1T} is defined as

$$\hat{g}_{1T} = (\partial g(X_T^{(0)}))' \int_0^T [Y_T Y_t^{-1} V(X_t^{(0)})] dW_t = g_{1T} - C$$

where

$$C = (\partial g(X_T^{(0)}))' \int_0^T Y_T Y_t^{-1} \partial_\epsilon V_0(X_t^{(0)}, 0) dt,$$

and define

$$Z_T^{(\xi)} = \exp\{i\xi\hat{g}_{1T} + \frac{\xi^2}{2}\Sigma_T\}.$$

Then, from Lemma 4, we have the following expression of $\mathbf{E}[\Phi(G^{(\epsilon)})]$:

$$\begin{aligned} \mathbf{E}[\Phi(G^{(\epsilon)})] &= \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \sum_{k \in K_{j,m}} \frac{C^{j,m,k}}{m!} \mathbf{E} \left[\Phi^{(m)}(\hat{g}_{1T} + C) \mathbf{E} [X^{j,m,k} | \hat{g}_{1T}] \right] + o(\epsilon^N) \\ &= \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \sum_{k \in K_{j,m}} \frac{C^{j,m,k}}{m!} \mathbf{E} \left[\Phi^{(m)}(\hat{g}_{1T} + C) \sum_{l=0}^{\infty} a_l^{j,m,k} H_l(\hat{g}_{1T}; \Sigma_T) \right] + o(\epsilon^N) \\ &= \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \sum_{k \in K_{j,m}} \sum_{l=0}^{\infty} \frac{a_l^{j,m,k} C^{j,m,k}}{m!} \mathbf{E} \left[\Phi^{(m)}(\hat{g}_{1T} + C) H_l(\hat{g}_{1T}; \Sigma_T) \right] + o(\epsilon^N) \end{aligned}$$

where

$$a_l^{j,m,k} = \frac{1}{(i\Sigma_T)^l} \left. \frac{d^l}{d\xi^l} \right|_{\xi=0} \left\{ \mathbf{E}[X^{j,m,k} Z_T^{(\xi)}] \right\}.$$

In particular, let Φ be the delta function at $x \in \mathbf{R}$, δ_x , we obtain the asymptotic expansion of density of $G^{(\epsilon)}$:

$$\begin{aligned} f_{G^{(\epsilon)}}(x) &= \mathbf{E}[\delta_x(G^{(\epsilon)})] \\ &= \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \sum_{k \in K_{j,m}} \sum_{l=0}^{\infty} \frac{a_l^{j,m,k} C^{j,m,k}}{m!} \mathbf{E} \left[\delta_x^{(m)}(\hat{g}_{1T} + C) H_l(\hat{g}_{1T}; \Sigma_T) \right] + o(\epsilon^N) \\ &= \sum_{j=0}^N \epsilon^j \sum_{m=0}^j \sum_{k \in K_{j,m}} \sum_{l=0}^{\infty} \frac{a_l^{j,m,k} C^{j,m,k}}{m!} (-1)^m \partial_x^m \{H_l(x - C; \Sigma_T) f_{g_{1T}}(x)\} + o(\epsilon^N) \end{aligned} \tag{28}$$

where

$$f_{g_{1T}}(x) := \frac{1}{\sqrt{2\pi\Sigma_T}} \exp\left(-\frac{x^2}{2\Sigma_T}\right).$$

4.1 Asymptotic Expansion of Density Function

In this subsection, we propose a new computational method for the asymptotic expansion of the density function (28). In particular, we show that coefficients in the expansion is obtained through a system of ordinary differential equations that is solved easily, and derive a concrete expression of the expansion up to ϵ^3 -order.

First, we write down the equation (28) more explicitly up to ϵ^3 -order:

$$\begin{aligned} f_{G^{(\epsilon)}}(x) &= \sum_{l=0}^{\infty} a_l^{0,0,(0)} H_l(x - C; \Sigma_T) f_{g_{1T}}(x) \\ &\quad + \epsilon \left\{ \sum_{l=0}^{\infty} a_l^{1,1,(1)} (-\partial_x) \{H_l(x - C; \Sigma_T) f_{g_{1T}}(x)\} \right\} \\ &\quad + \epsilon^2 \left\{ \sum_{l=0}^{\infty} a_l^{2,1,(0,1)} (-\partial_x) \{H_l(x - C; \Sigma_T) f_{g_{1T}}(x)\} \right. \\ &\quad \left. + \frac{1}{2} \sum_{l=0}^{\infty} a_l^{2,2,(2,0)} \partial_x^2 \{H_l(x - C; \Sigma_T) f_{g_{1T}}(x)\} \right\} \end{aligned}$$

$$\begin{aligned}
& +\epsilon^3 \left\{ \sum_{l=0}^{\infty} a_l^{3,1,(0,0,1)} (-\partial_x) \{H_l(x-C; \Sigma_T) f_{g_{1T}}(x)\} \right. \\
& + \frac{1}{2} \sum_{l=0}^{\infty} a_l^{3,2,(1,1,0)} \partial_x^2 \{H_l(x-C; \Sigma_T) f_{g_{1T}}(x)\} \\
& \left. + \frac{1}{6} \sum_{l=0}^{\infty} a_l^{3,3,(3,0,0)} (-\partial_x^3) \{H_l(x-C; \Sigma_T) f_{g_{1T}}(x)\} \right\} \\
& + o(\epsilon^3),
\end{aligned}$$

where coefficients $a_l^{j,m,k}$ are given by

$$\begin{aligned}
a_l^{0,0,(0)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[Z_T^{(\xi)}] \right\} \\
a_l^{1,1,(1)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[g_{2T} Z_T^{(\xi)}] \right\} \\
a_l^{2,1,(0,1)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[g_{3T} Z_T^{(\xi)}] \right\} \\
a_l^{2,2,(2,0)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[g_{2T}^2 Z_T^{(\xi)}] \right\} \\
a_l^{3,1,(0,0,1)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[g_{4T} Z_T^{(\xi)}] \right\} \\
a_l^{3,2,(1,1,0)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[g_{2T} g_{3T} Z_T^{(\xi)}] \right\} \\
a_l^{3,3,(3,0,0)} &= \frac{1}{(i\Sigma_T)^l} \frac{d^l}{d\xi^l} \Big|_{\xi=0} \left\{ \mathbf{E}[g_{2T}^3 Z_T^{(\xi)}] \right\}. \tag{29}
\end{aligned}$$

Since $E[Z_T^{(\xi)}] = 1$, we have $a_0^{0,0,(0)} = 1$ and $a_l^{0,0,(0)} = 0$ for $l \geq 1$. The other expectations above are expressed in terms of A_{nT} and $Z_T^{(\xi)}$ as follows:

$$\begin{aligned}
\mathbf{E}[g_{2T} Z_T^{(\xi)}] &= \frac{1}{2} \sum_{i,j=1}^d \partial_i \partial_j g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j Z_T^{(\xi)}] + \frac{1}{2} \sum_{i=1}^d \partial_i g(X_T^{(0)}) \mathbf{E}[A_{2T}^i Z_T^{(\xi)}] \\
\mathbf{E}[g_{3T} Z_T^{(\xi)}] &= \frac{1}{6} \sum_{i,j,k=1}^d \partial_i \partial_j \partial_k g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k Z_T^{(\xi)}] + \frac{1}{2} \sum_{i,j=1}^d \partial_i \partial_j g(X_T^{(0)}) \mathbf{E}[A_{2T}^i A_{1T}^j Z_T^{(\xi)}] \\
&+ \frac{1}{6} \sum_{i=1}^d \partial_i g(X_T^{(0)}) \mathbf{E}[A_{3T}^i Z_T^{(\xi)}], \\
\mathbf{E}[g_{2T}^2 Z_T^{(\xi)}] &= \frac{1}{4} \sum_{i,j,k,l=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_k \partial_l g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k A_{1T}^l Z_T^{(\xi)}] \\
&+ \frac{1}{2} \sum_{i,j,k=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_i g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{2T}^k Z_T^{(\xi)}] \\
&+ \frac{1}{4} \sum_{i,j=1}^d \partial_i g(X_T^{(0)}) \partial_j g(X_T^{(0)}) \mathbf{E}[A_{2T}^i A_{2T}^j Z_T^{(\xi)}] \\
\mathbf{E}[g_{4T} Z_T^{(\xi)}] &= \frac{1}{24} \sum_{i,j,k,l=1}^d \partial_i \partial_j \partial_k \partial_l g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k A_{1T}^l Z_T^{(\xi)}]
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{4} \sum_{i,j,k=1}^d \partial_i \partial_j \partial_k g(X_T^{(0)}) \mathbf{E}[A_{2T}^i A_{1T}^j A_{1T}^k Z_T^{(\xi)}] \\
& + \frac{1}{2} \sum_{i,j=1}^d \partial_i \partial_j g(X_T^{(0)}) \mathbf{E}[A_{2T}^i A_{2T}^j Z_T^{(\xi)}] + \frac{1}{2} \sum_{i,j=1}^d \partial_i \partial_j g(X_T^{(0)}) \mathbf{E}[A_{3T}^i A_{1T}^j Z_T^{(\xi)}] \\
& + \frac{1}{6} \sum_{i=1}^d \partial_i g(X_T^{(0)}) \mathbf{E}[A_{4T}^i Z_T^{(\xi)}], \\
\mathbf{E}[g_{2T} g_{3T} Z_T^{(\xi)}] &= \frac{1}{12} \sum_{i,j,k,l,m=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_k \partial_l \partial_m g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k A_{1T}^l A_{1T}^m Z_T^{(\xi)}] \\
& + \frac{1}{12} \sum_{i,j,k,l=1}^d \left\{ \partial_i g(X_T^{(0)}) \partial_j \partial_k \partial_l g(X_T^{(0)}) + 3 \partial_i \partial_j g(X_T^{(0)}) \partial_k \partial_l g(X_T^{(0)}) \right\} \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k A_{2T}^l Z_T^{(\xi)}] \\
& + \frac{1}{4} \sum_{i,j,k=1}^d \partial_k g(X_T^{(0)}) \partial_i \partial_j g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{2T}^j A_{2T}^k Z_T^{(\xi)}] \\
& + \frac{1}{12} \sum_{i,j,k=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_k g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{3T}^k Z_T^{(\xi)}] \\
& + \frac{1}{12} \sum_{i,j=1}^d \partial_i g(X_T^{(0)}) \partial_j g(X_T^{(0)}) \mathbf{E}[A_{2T}^i A_{3T}^j Z_T^{(\xi)}], \\
\mathbf{E}[g_{2T}^3 Z_T^{(\xi)}] &= \frac{1}{8} \sum_{i,j,k,l,m,n=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_k \partial_l g(X_T^{(0)}) \partial_m \partial_n g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k A_{1T}^l A_{1T}^m A_{1T}^n Z_T^{(\xi)}] \\
& + \frac{3}{8} \sum_{i,j,k,l,m=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_k \partial_l g(X_T^{(0)}) \partial_m g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{1T}^k A_{1T}^l A_{2T}^m Z_T^{(\xi)}] \\
& + \frac{3}{8} \sum_{i,j,k,l,m=1}^d \partial_i \partial_j g(X_T^{(0)}) \partial_k g(X_T^{(0)}) \partial_l g(X_T^{(0)}) \mathbf{E}[A_{1T}^i A_{1T}^j A_{2T}^k A_{2T}^l Z_T^{(\xi)}] \\
& + \frac{1}{8} \sum_{i,j,k,l,m=1}^d \partial_i g(X_T^{(0)}) \partial_j g(X_T^{(0)}) \partial_k g(X_T^{(0)}) \mathbf{E}[A_{2T}^i A_{2T}^j A_{2T}^k Z_T^{(\xi)}]
\end{aligned}$$

where A_{1t} is given by (2), and A_{2t} , A_{3t} and A_{4t} are expressed as

$$\begin{aligned}
A_{2t} &= \int_0^t Y_t Y_u^{-1} \left(\sum_{j,k=1}^d \partial_j \partial_k V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k du + 2 \sum_{j=1}^d \partial_\epsilon \partial_j V_0(X_u^{(0)}, 0) A_{1u}^j du \right. \\
&\quad \left. + \partial_\epsilon^2 V_0(X_u^{(0)}, 0) du + 2 \sum_{j=1}^d \partial_j V(X_u^{(0)}) A_{1u}^j dW_u \right), \\
A_{3t} &= \int_0^t Y_t Y_u^{-1} \left(\sum_{j,k,l=1}^d \partial_j \partial_k \partial_l V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k A_{1u}^l du + 3 \sum_{j,k=1}^d \partial_j \partial_k V_0(X_u^{(0)}, 0) A_{1u}^j A_{2u}^k du \right. \\
&\quad + 3 \sum_{j,k=1}^d \partial_j \partial_k \partial_\epsilon V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k du + 3 \sum_{j=1}^d \partial_j \partial_\epsilon V_0(X_u^{(0)}, 0) A_{2u}^j du \\
&\quad \left. + 3 \sum_{j=1}^d \partial_j \partial_\epsilon^2 V_0(X_u^{(0)}, 0) A_{1u}^j du + \partial_\epsilon^3 V_0(X_u^{(0)}, 0) du \right)
\end{aligned}$$

$$\begin{aligned}
& +3 \sum_{j,k=1}^d \partial_j \partial_k V(X_u^{(0)}) A_{1u}^j A_{1u}^k dW_u + 3 \sum_{j=1}^d \partial_j V(X_u^{(0)}) A_{2u}^j dW_u \Big), \\
A_{4t} = & \int_0^t Y_t Y_u^{-1} \left(\sum_{j,k,l,m=1}^d \partial_j \partial_k \partial_l \partial_m V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k A_{1u}^l A_{1u}^m du \right. \\
& +4 \sum_{j,k,l=1}^d \partial_j \partial_k \partial_l V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k A_{2u}^l du + 3 \sum_{j,k=1}^d \partial_j \partial_k V_0(X_u^{(0)}, 0) A_{2u}^j A_{2u}^k du \\
& +4 \sum_{j,k=1}^d \partial_j \partial_k V(X_u^{(0)}) A_{1u}^j A_{3u}^k du + 4 \sum_{j,k,l=1}^d \partial_j \partial_k \partial_l \partial_\epsilon V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k A_{1u}^l du \\
& +6 \sum_{j,k=1}^d \partial_j \partial_k \partial_\epsilon V_0(X_u^{(0)}, 0) A_{1u}^j A_{2u}^k du + 4 \sum_{j=1}^d \partial_j \partial_\epsilon V_0(X_u^{(0)}, 0) A_{3u}^j du \\
& +6 \sum_{j,k=1}^d \partial_j \partial_k \partial_\epsilon^2 V_0(X_u^{(0)}, 0) A_{1u}^j A_{1u}^k du + 6 \sum_{j=1}^d \partial_j \partial_\epsilon^2 V_0(X_u^{(0)}, 0) A_{2u}^j du \\
& +4 \sum_{j=1}^d \partial_j \partial_\epsilon^3 V_0(X_u^{(0)}, 0) A_{1u}^j du + \partial_\epsilon^4 V_0(X_u^{(0)}, 0) du \\
& +4 \sum_{j,k,l=1}^d \partial_j \partial_k \partial_l V(X_u^{(0)}) A_{1u}^j A_{1u}^k A_{1u}^l dW_u + 12 \sum_{j,k=1}^d \partial_j \partial_k V(X_u^{(0)}) A_{1u}^j A_{2u}^k dW_u \\
& \left. +4 \sum_{j=1}^d \partial_j V(X_u^{(0)}) A_{3u}^j dW_u \right).
\end{aligned}$$

Note that each A_{kt}^i ($i = 1, \dots, d, k = 1, 2, 3, 4$) has all finite moments due to a *grading* structure. For the detail of the following definition and theorem, see pp.45-47 in Bichteler, Gravereaux and Jacod [1].

Definition 1 A *grading* of \mathbf{R}^d is a decomposition $\mathbf{R}^d = \mathbf{R}^{d_1} \times \dots \times \mathbf{R}^{d_q}$ with $d = d_1 + \dots + d_q$. The coordinates of a point in \mathbf{R}^d are always arranged in an increasing order along the subspace \mathbf{R}^{d_i} , and we set $M_0 = 0$ and $M_l = d_1 + \dots + d_l$ for $1 \leq l \leq q$. We say that a mapping V on \mathbf{R}^d is *graded* according to the grading $\mathbf{R}^d = \mathbf{R}^{d_1} \times \dots \times \mathbf{R}^{d_q}$ if $V^i(y)$ depends upon only through the coordinates $(y^k)_{1 \leq k \leq M_r}$ when $M_{r-1} \leq i \leq M_r$.

Theorem 1 Consider the stochastic differential equation of the form

$$dY_t = V_0(Y_t, t)dt + V(Y_t, t)dW_t; \quad Y_0 = y_0 \in \mathbf{R}^d \quad (30)$$

where coefficients $V_0 : \mathbf{R}^d \times \mathbf{R}^+ \rightarrow \mathbf{R}^d$ and $V : \mathbf{R}^d \times \mathbf{R}^+ \rightarrow \mathbf{R}^d \otimes \mathbf{R}^d$ have a Lipschitz lower triangular structure, and are graded according to $\mathbf{R}^d = \mathbf{R}^{d_1} \times \dots \times \mathbf{R}^{d_q}$ with respect to Y . Moreover for $F(y, t) = V_0(y, t)$ or $V(y, t)$, we assume F is differentiable in y in \mathbf{R}^d and

1. $|F(0, t)| \leq Z_t$
2. $|D_y F(y, t)| \leq \hat{Z}_t(1 + |y|^\theta)$
3. $|\partial_j F^i(y, t)| \leq \zeta$ if $M_{r-1} \leq i \leq M_r$ for some $r \leq q$

where $\zeta, \theta \geq 0$ are constants, and Z, \hat{Z} are predictable processes such that $\|Z\|_p$ and $\|\hat{Z}\|_p$ are finite for all $p \geq 1$. Then (30) have a unique solution Y , and for

every $p \geq 1$ there are constants c_p and γ_p depending only upon $(\zeta, \theta, \{\|\hat{Z}\|_r\}_{r \geq 1})$, such that

$$\|Y_T\|_p \leq c_p(y_0 + \|Z\|_{\gamma_p}).$$

Applying Theorem 1 to the system of stochastic differential equations consists of $A_{kt}^i (i = 1, \dots, d, k = 1, 2, 3, 4)$ and any products of them, we obtain the following lemma.

Lemma 5 *Each coefficient of the expansion $A_{kt}^i (i = 1, \dots, d, k = 1, 2, 3, 4)$ has all finite moments.*

(proof) Consider the system of stochastic differential equations which $A_1^1, \dots, A_1^d, A_1^1 A_1^1, \dots, A_1^d A_1^d, A_2^1, \dots, A_2^d, \dots$ follow, then it is easily shown that the coefficients of the equation have a grading structure and satisfy the conditions in Theorem 1. Hence the coefficients A_{kt}^i have all finite moments. \square

Here, we redefine $\hat{g}_1 = \{\hat{g}_{1t}; t \in \mathbf{R}^+\}$ and $Z^{(\xi)} = \{Z_t^{(\xi)}; t \in \mathbf{R}^+\}$ as the stochastic processes

$$\hat{g}_{1t} = \int_0^t \hat{V}(X_u^{(0)}, u) dW_u$$

and

$$Z_t^{(\xi)} = \exp\{i\xi \hat{g}_{1t} + \frac{\xi^2}{2} \Sigma_t\},$$

respectively where

$$\hat{V}(x, t) = (\partial g(X_T^{(0)}))' [Y_T Y_t^{-1} V(x)].$$

We define $\eta_{1,1}^i, \eta_{2,1}^i, \eta_{2,2}^{i,k}, \eta_{3,1}^i, \eta_{3,2}^{i,k}, \eta_{3,3}^{i,k}, \eta_{4,1}^i, \eta_{4,2,1}^{i,k}, \eta_{4,2,2}^{i,k}, \eta_{5,2}^{i,k}, \eta_{5,3}^{i,k}$, and $\eta_{6,3}^{i,k,l}$ as

$$\begin{aligned} \eta_{1,1}^i(t) &:= E[A_{1t}^i Z_t], & \eta_{2,1}^i(t) &:= E[A_{2t}^i Z_t], & \eta_{2,2}^{i,k}(t) &:= E[A_{1t}^i A_{1t}^k Z_t], \\ \eta_{3,1}^i(t) &:= E[A_{3t}^i Z_t], & \eta_{3,2}^{i,k}(t) &:= E[A_{1t}^i A_{2t}^k Z_t], & \eta_{3,3}^{i,k,l}(t) &:= E[A_{1t}^i A_{1t}^k A_{1t}^l Z_t], \\ \eta_{4,1}^i(t) &:= E[A_{4t}^i Z_t], & \eta_{4,2,1}^{i,k}(t) &:= E[A_{1t}^i A_{3t}^k Z_t], & \eta_{4,2,2}^{i,k}(t) &:= E[A_{2t}^i A_{2t}^k Z_t], \\ \eta_{4,3}^{i,k,l}(t) &:= E[A_{1t}^i A_{1t}^k A_{2t}^l Z_t], & \eta_{4,4}^{i,k,l,m}(t) &:= E[A_{1t}^i A_{1t}^k A_{1t}^l A_{1t}^m Z_t], \\ \eta_{5,2}^{i,k}(t) &:= E[A_{2t}^i A_{3t}^k Z_t], & \eta_{5,3,1}^{i,k,l}(t) &:= E[A_{1t}^i A_{1t}^k A_{3t}^l Z_t], & \eta_{5,3,2}^{i,k,l}(t) &:= E[A_{1t}^i A_{2t}^k A_{2t}^l Z_t], \\ \eta_{5,4}^{i,k,l,m}(t) &:= E[A_{1t}^i A_{1t}^k A_{1t}^l A_{2t}^m Z_t], & \eta_{5,5}^{i,k,l,m,n}(t) &:= E[A_{1t}^i A_{1t}^k A_{1t}^l A_{1t}^m A_{1t}^n Z_t], \\ \eta_{6,3}^{i,k,l}(t) &:= E[A_{2t}^i A_{2t}^k A_{2t}^l Z_t], & \eta_{6,4}^{i,k,l,m}(t) &:= E[A_{1t}^i A_{1t}^k A_{2t}^l A_{2t}^m Z_t], \\ \eta_{6,5}^{i,k,l,m,n}(t) &:= E[A_{1t}^i A_{1t}^k A_{1t}^l A_{1t}^m A_{2t}^n Z_t], & \eta_{6,5}^{i,k,l,m,n,o}(t) &:= E[A_{1t}^i A_{1t}^k A_{1t}^l A_{1t}^m A_{1t}^n A_{1t}^o Z_t]. \end{aligned} \tag{31}$$

We derive the system of ordinary differential equations of η .

In the followings, for simplicity, we assume that V_0 doesn't depend on ϵ , and write $V_0(x, \epsilon)$ as $V_0(x)$.

Consider the evaluation of $\eta_{2,1}^i(T) = E[A_{2T}^i Z_T^{(\xi)}]$ which appears in the ϵ -order. Applying Ito's formula to $A_{2t}^i Z_t^{(\xi)}$, we have

$$\begin{aligned} d(A_{2t}^i Z_t^{(\xi)}) &= A_{2t}^i dZ_t^{(\xi)} + Z_t^{(\xi)} dA_{2t}^i + dA_{2t}^i dZ_t^{(\xi)} \\ &= \left\{ 2(i\xi) \sum_{i'=1}^d A_{1t}^{i'} Z_t^{(\xi)} \hat{V}(X_t^{(0)}, t) \partial_{i'} V^i(X_t^{(0)})' + \sum_{i'=1}^d A_{2t}^{i'} Z_t^{(\xi)} \partial_{i'} V_0^i(X_t^{(0)}) \right\} \end{aligned}$$

$$\begin{aligned}
& + \left. \sum_{i'=1}^d \sum_{k'=1}^d A_{1t}^{i'} A_{1t}^{k'} Z_t^{(\xi)} \partial_{i'} \partial_{k'} V_0^i(X_t^{(0)}) \right\} dt \\
& + \left\{ (i\xi) A_{2t}^i Z_t^{(\xi)} \hat{V}(X_t^{(0)}, t) + 2 \sum_{i'=1}^d A_{1t}^{i'} Z_t^{(\xi)} \partial_{i'} V^i(X_t^{(0)}) \right\} dW_t
\end{aligned}$$

Since the second and third terms are martingales, taking the expectation on both sides, we have the following ordinary differential equation of $\eta_{2,1}^i$:

$$\begin{aligned}
\frac{d}{dt} \eta_{2,1}^i(t) &= 2(i\xi) \sum_{i'=1}^d \eta_{1,1}^{i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^i(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{2,1}^{i'}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{2,2}^{i',k'}(t) \partial_{i'} \partial_{k'} V_0^i(X_t^{(0)})
\end{aligned}$$

Here, $\eta_{1,1}^i (i = 1, \dots, d)$ appearing in the right hand side of above ODE are evaluated in the similar manner:

$$\begin{aligned}
d(A_{1t}^i Z_t^{(\xi)}) &= A_{1t}^i dZ_t^{(\xi)} + Z_t^{(\xi)} dA_{1t}^i + dA_{1t}^i dZ_t^{(\xi)} \\
&= \left\{ (i\xi) Z_t^{(\xi)} \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \sum_{i'=1}^d A_{1t}^{i'} Z_t^{(\xi)} \partial_{i'} V_0^i(X_t^{(0)}) \right\} dt \\
& + \left\{ (i\xi) A_{1t}^i Z_t^{(\xi)} \hat{V}(X_t^{(0)}, t) + Z_t^{(\xi)} V^i(X_t^{(0)}) \right\} dW_t,
\end{aligned}$$

hence, we have

$$\frac{d}{dt} \eta_{1,1}^i(t) = (i\xi) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \sum_{i'=1}^d \eta_{1,1}^{i'}(t) \partial_{i'} V_0^i(X_t^{(0)}).$$

$\eta_{2,2}^{i,k}$ and other higher order terms can be evaluated in the same way.

The key observation is that each ODE does not involve any higher order terms, and only lower or the same order terms appear in the right hand side of the ODE. So, one can easily solve (analytically or numerically) the system of ODEs and evaluate expectations.

Proposition 2 For $\eta_{j,m,k}$ defined in (31), the following system of ordinary differential equations is hold:

$$\frac{d}{dt} \eta_{1,1}^i(t) = (i\xi) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \sum_{i'=1}^d \eta_{1,1}^{i'}(t) \partial_{i'} V_0^i(X_t^{(0)})$$

$$\begin{aligned}
\frac{d}{dt} \eta_{2,1}^i(t) &= 2(i\xi) \sum_{i'=1}^d \eta_{1,1}^{i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^i(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{2,1}^{i'}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{2,2}^{i',k'}(t) \partial_{i'} \partial_{k'} V_0^i(X_t^{(0)}) \\
\frac{d}{dt} \eta_{2,2}^{i,k}(t) &= (i\xi) \left\{ \eta_{1,1}^k(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' + \eta_{1,1}^i(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' \right\} \\
& + V^i(X_t^{(0)}) V^k(X_t^{(0)})' + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{2,2}^{i',k'}(t) \partial_{i'} V_0^i(X_t^{(0)}) \partial_{k'} V_0^k(X_t^{(0)})
\end{aligned}$$

$$\begin{aligned}
\frac{d}{dt}\eta_{3,1}^i(t) &= (i\xi)\left\{3\sum_{i'=1}^d\eta_{2,1}^{i'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}V^i(X_t^{(0)})'\right. \\
&\quad +3\sum_{i'=1}^d\sum_{k'=1}^d\eta_{2,2}^{i',k'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}\partial_{k'}V^i(X_t^{(0)})'\left.\right\} \\
&\quad +\sum_{i'=1}^d\eta_{3,1}^{i'}(t)\partial_{i'}V_0^i(X_t^{(0)})+3\sum_{i'=1}^d\sum_{k'=1}^d\eta_{3,2}^{i',k'}(t)\partial_{i'}\partial_{k'}V_0^i(X_t^{(0)}) \\
&\quad +\sum_{i'=1}^d\sum_{k'=1}^d\sum_{l'=1}^d\eta_{3,3}^{i',k',l'}(t)\partial_{i'}\partial_{k'}\partial_{l'}V_0^i(X_t^{(0)}) \\
\frac{d}{dt}\eta_{3,2}^{i,k}(t) &= (i\xi)\left\{\eta_{2,1}^k(t)\hat{V}(X_t^{(0)},t)V^i(X_t^{(0)})'+2\sum_{i'=1}^d\eta_{2,2}^{i,i'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}V^k(X_t^{(0)})'\right\} \\
&\quad +2\sum_{i'=1}^d\eta_{1,1}^{i'}(t)V^i(X_t^{(0)})\partial_{i'}V^k(X_t^{(0)})' \\
&\quad +\sum_{i'=1}^d\eta_{3,2}^{i',k}(t)\partial_{i'}V_0^i(X_t^{(0)})+\sum_{k'=1}^d\eta_{3,2}^{i,k'}(t)\partial_{k'}V_0^k(X_t^{(0)}) \\
&\quad +\sum_{i'=1}^d\sum_{k'=1}^d\eta_{3,3}^{i,i',k'}(t)\partial_{i'}\partial_{k'}V_0^k(X_t^{(0)}) \\
\frac{d}{dt}\eta_{3,3}^{i,k,l}(t) &= (i\xi)\left\{\eta_{2,2}^{k,l}(t)\hat{V}(X_t^{(0)},t)V^i(X_t^{(0)})'+\eta_{2,2}^{i,l}(t)\hat{V}(X_t^{(0)},t)V^k(X_t^{(0)})'+\eta_{2,2}^{i,k}(t)\hat{V}(X_t^{(0)},t)V^l(X_t^{(0)})'\right\} \\
&\quad +\eta_{1,1}^i(t)V^k(X_t^{(0)})V^l(X_t^{(0)})'+\eta_{1,1}^k(t)V^i(X_t^{(0)})V^l(X_t^{(0)})'+\eta_{1,1}^l(t)V^i(X_t^{(0)})V^k(X_t^{(0)})' \\
&\quad +\sum_{i'=1}^d\eta_{3,3}^{i',k,l}(t)\partial_{i'}V_0^i(X_t^{(0)})+\sum_{k'=1}^d\eta_{3,3}^{i,k',l}(t)\partial_{k'}V_0^k(X_t^{(0)})+\sum_{l'=1}^d\eta_{3,3}^{i,k,l'}(t)\partial_{l'}V_0^l(X_t^{(0)}) \\
\frac{d}{dt}\eta_{4,1}^i(t) &= (i\xi)\left\{4\sum_{i'=1}^d\eta_{3,1}^{i'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}V^i(X_t^{(0)})'+12\sum_{i'=1}^d\sum_{k'=1}^d\eta_{3,2}^{i',k'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}\partial_{k'}V^i(X_t^{(0)})'\right. \\
&\quad +4\sum_{i'=1}^d\sum_{k'=1}^d\sum_{l'=1}^d\eta_{3,3}^{i',k',l'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}\partial_{k'}\partial_{l'}V^i(X_t^{(0)})'\left.\right\} \\
&\quad +\sum_{i'=1}^d\eta_{4,1}^{i'}(t)\partial_{i'}V_0^i(X_t^{(0)})+4\sum_{i'=1}^d\sum_{k'=1}^d\eta_{4,2,1}^{i',k'}(t)\partial_{i'}\partial_{k'}V_0^i(X_t^{(0)}) \\
&\quad +3\sum_{i'=1}^d\sum_{k'=1}^d\eta_{4,2,2}^{i',k'}(t)\partial_{i'}\partial_{k'}V_0^i(X_t^{(0)})+4\sum_{i'=1}^d\sum_{k'=1}^d\sum_{l'=1}^d\eta_{4,3}^{i',k',l'}(t)\partial_{i'}\partial_{k'}\partial_{l'}V_0^i(X_t^{(0)}) \\
&\quad +\sum_{i'=1}^d\sum_{k'=1}^d\sum_{l'=1}^d\sum_{m'=1}^d\eta_{4,4}^{i',k',l',m'}(t)\partial_{i'}\partial_{k'}\partial_{l'}\partial_{m'}V_0^i(X_t^{(0)}) \\
\frac{d}{dt}\eta_{4,2,1}^{i,k}(t) &= (i\xi)\left\{\eta_{3,1}^k(t)\hat{V}(X_t^{(0)},t)V^i(X_t^{(0)})'+3\sum_{i'=1}^d\eta_{3,2}^{i,i'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}V^k(X_t^{(0)})'\right. \\
&\quad +3\sum_{i'=1}^d\sum_{k'=1}^d\eta_{3,3}^{i,i',k'}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}\partial_{k'}V^k(X_t^{(0)})'\left.\right\} \\
&\quad +3\sum_{i'=1}^d\eta_{2,1}^{i'}(t)V^i(X_t^{(0)})\partial_{i'}V^k(X_t^{(0)})'+3\sum_{i'=1}^d\sum_{k'=1}^d\eta_{2,2}^{i',k'}(t)V^i(X_t^{(0)})\partial_{i'}\partial_{k'}V^k(X_t^{(0)})'
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i'=1}^d \eta_{4,2,1}^{i',k}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{4,2,1}^{i,k'}(t) \partial_{k'} V_0^k(X_t^{(0)}) \\
& + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,3}^{i,i',k'}(t) \partial_{i'} \partial_{k'} V_0^k(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \sum_{l'=1}^d \eta_{4,4}^{i,i',k',l'}(t) \partial_{i'} \partial_{k'} \partial_{l'} V_0^k(X_t^{(0)}) \\
\frac{d}{dt} \eta_{4,2,2}^{i,k}(t) & = (i\xi) \left\{ 2 \sum_{i'=1}^d \eta_{3,2}^{i',k}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^i(X_t^{(0)})' + 2 \sum_{i'=1}^d \eta_{3,2}^{i',i}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^k(X_t^{(0)})' \right\} \\
& + 4 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{2,2}^{i',k'}(t) \partial_{i'} V^i(X_t^{(0)}) \partial_{k'} V^k(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{4,2,2}^{i',k}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{4,2,2}^{i,k'}(t) \partial_{k'} V_0^k(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,3}^{i',k',k}(t) \partial_{i'} \partial_{k'} V_0^i(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,3}^{i',k',i}(t) \partial_{i'} \partial_{k'} V_0^k(X_t^{(0)}) \\
\frac{d}{dt} \eta_{4,3}^{i,k,l}(t) & = (i\xi) \left\{ \eta_{3,2}^{k,l}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{3,2}^{i,l}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right. \\
& + 2 \sum_{i'=1}^d \eta_{3,3}^{i,k,i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^l(X_t^{(0)})' \left. \right\} \\
& + \eta_{2,1}^l(t) V^i(X_t^{(0)}) V^k(X_t^{(0)})' \\
& + 2 \sum_{i'=1}^d \eta_{2,2}^{i,i'}(t) V^k(X_t^{(0)}) \partial_{i'} V^l(X_t^{(0)})' + 2 \sum_{i'=1}^d \eta_{2,2}^{k,i'}(t) V^i(X_t^{(0)}) \partial_{i'} V^l(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{4,3}^{i',k,l}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{4,3}^{i,k',l}(t) \partial_{k'} V_0^k(X_t^{(0)}) + \sum_{l'=1}^d \eta_{4,3}^{i,k,l'}(t) \partial_{l'} V_0^l(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,4}^{i,k,i',k'}(t) \partial_{i'} \partial_{k'} V_0^l(X_t^{(0)}) \\
\frac{d}{dt} \eta_{4,4}^{i,k,l,m}(t) & = (i\xi) \left\{ \eta_{3,3}^{k,l,m}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{3,3}^{i,l,m}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right. \\
& + \eta_{3,3}^{i,k,m}(t) \hat{V}(X_t^{(0)}, t) V^l(X_t^{(0)})' + \eta_{3,3}^{i,k,l}(t) \hat{V}(X_t^{(0)}, t) V^m(X_t^{(0)})' \left. \right\} \\
& + \eta_{2,2}^{i,k}(t) V^l(X_t^{(0)}) V^m(X_t^{(0)})' + \eta_{2,2}^{i,l}(t) V^k(X_t^{(0)}) V^m(X_t^{(0)})' + \eta_{2,2}^{i,m}(t) V^k(X_t^{(0)}) V^l(X_t^{(0)})' \\
& + \eta_{2,2}^{k,l}(t) V^i(X_t^{(0)}) V^m(X_t^{(0)})' + \eta_{2,2}^{k,m}(t) V^i(X_t^{(0)}) V^l(X_t^{(0)})' + \eta_{2,2}^{l,m}(t) V^i(X_t^{(0)}) V^k(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{4,4}^{i',k,l,m}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{4,4}^{i,k',l,m}(t) \partial_{k'} V_0^k(X_t^{(0)}) \\
& + \sum_{l'=1}^d \eta_{4,4}^{i,k,l',m}(t) \partial_{l'} V_0^l(X_t^{(0)}) + \sum_{m'=1}^d \eta_{4,4}^{i,k,l,m'}(t) \partial_{m'} V_0^m(X_t^{(0)}) \\
\frac{d}{dt} \eta_{5,2}^{i,k}(t) & = (i\xi) \left\{ 2 \sum_{i'=1}^d \eta_{4,2,1}^{k,i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^i(X_t^{(0)})' + 3 \sum_{i'=1}^d \eta_{4,2,2}^{i,i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^k(X_t^{(0)})' \right. \\
& + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,3}^{i',k',i}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} \partial_{k'} V^k(X_t^{(0)})' \left. \right\}
\end{aligned}$$

$$\begin{aligned}
& +6 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{3,2}^{i',k'}(t) \partial_{i'} V^i(X_t^{(0)}) \partial_{k'} V^k(X_t^{(0)})' \\
& +6 \sum_{i'=1}^d \sum_{k'=1}^d \sum_{l'=1}^d \eta_{3,3}^{i',k',l'}(t) \partial_{i'} V^i(X_t^{(0)}) \partial_{k'} \partial_{l'} V^k(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{5,2}^{i',k}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{5,2}^{i,k'}(t) \partial_{k'} V_0^k(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{5,3,1}^{i',k',k}(t) \partial_{i'} \partial_{k'} V_0^i(X_t^{(0)}) + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{5,3,2}^{i',k',i}(t) \partial_{i'} \partial_{k'} V_0^k(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \sum_{l'=1}^d \eta_{5,4}^{i',k',l',i}(t) \partial_{i'} \partial_{k'} \partial_{l'} V_0^k(X_t^{(0)}) \\
\frac{d}{dt} \eta_{5,3,1}^{i,k,l}(t) &= (i\xi) \left\{ \eta_{4,2,1}^{k,l}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{4,2,1}^{i,l}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right. \\
& + 3 \sum_{i'=1}^d \eta_{4,3}^{i,k,i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^l(X_t^{(0)})' + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,4}^{i,k,i',k'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} \partial_{k'} V^l(X_t^{(0)})' \left. \right\} \\
& + \eta_{3,1}(t) V^i(X_t^{(0)})' V^k(X_t^{(0)}) \\
& + 3 \sum_{i'=1}^d \eta_{3,2}^{k,i'}(t) V^i(X_t^{(0)}) \partial_{i'} V^l(X_t^{(0)})' + 3 \sum_{i'=1}^d \eta_{3,2}^{i,i'}(t) V^k(X_t^{(0)}) \partial_{i'} V^l(X_t^{(0)})' \\
& + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{3,3}^{k,i',k'}(t) V^i(X_t^{(0)}) \partial_{i'} \partial_{k'} V^l(X_t^{(0)})' + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{3,3}^{i,i',k'}(t) V^k(X_t^{(0)}) \partial_{i'} \partial_{k'} V^l(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{5,3,1}^{i',k,l}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{5,3,1}^{i,k',l}(t) \partial_{k'} V_0^k(X_t^{(0)}) + \sum_{l'=1}^d \eta_{5,3,1}^{i,k,l'}(t) \partial_{l'} V_0^l(X_t^{(0)}) \\
& + 3 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{5,4}^{i,k,i',k'}(t) \partial_{i'} \partial_{k'} V_0^l(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \sum_{l'=1}^d \eta_{5,5}^{i,k,i',k',l'}(t) \partial_{i'} \partial_{k'} \partial_{l'} V_0^l(X_t^{(0)}) \\
\frac{d}{dt} \eta_{5,3,2}^{i,k,l}(t) &= (i\xi) \left\{ \eta_{4,2}^{k,l}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' \right. \\
& + 2 \sum_{i'=1}^d \eta_{4,3}^{i',i,l}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^k(X_t^{(0)})' + 2 \sum_{i'=1}^d \eta_{4,3}^{i',i,k}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^l(X_t^{(0)})' \left. \right\} \\
& + 2 \sum_{i'=1}^d \eta_{3,2}^{i',k}(t) V^i(X_t^{(0)}) \partial_{i'} V^l(X_t^{(0)})' + 2 \sum_{i'=1}^d \eta_{3,2}^{i',l}(t) V^i(X_t^{(0)}) \partial_{i'} V^k(X_t^{(0)})' \\
& + 4 \sum_{i'=1}^d \sum_{k'=1}^d \eta_{3,3}^{i,i',k'}(t) \partial_{i'} V^k(X_t^{(0)}) \partial_{k'} V^l(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{5,3,2}^{i',k,l}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{5,3,2}^{i,k',l}(t) \partial_{k'} V_0^k(X_t^{(0)}) + \sum_{l'=1}^d \eta_{5,3,2}^{i,k,l'}(t) \partial_{l'} V_0^l(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{5,4}^{i,i',k',l}(t) \partial_{i'} \partial_{k'} V_0^k(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{5,4}^{i,i',k',k}(t) \partial_{i'} \partial_{k'} V_0^l(X_t^{(0)}) \\
\frac{d}{dt} \eta_{5,4}^{i,k,l,m}(t) &= (i\xi) \left\{ \eta_{4,3}^{k,l,m}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{4,3}^{l,i,m}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right. \\
& \left. + \eta_{4,3}^{i,k,m}(t) \hat{V}(X_t^{(0)}, t) V^l(X_t^{(0)})' + 2 \sum_{i'=1}^d \eta_{4,4}^{i,k,l,i'}(t) \hat{V}(X_t^{(0)}, t) \partial_{i'} V^m(X_t^{(0)})' \right\}
\end{aligned}$$

$$\begin{aligned}
& +\eta_{3,2}^{l,m}(t)V^i(X_t^{(0)})V^k(X_t^{(0)})' + \eta_{3,2}^{k,m}(t)V^i(X_t^{(0)})V^l(X_t^{(0)})' + \eta_{3,2}^{i,m}(t)V^k(X_t^{(0)})V^l(X_t^{(0)})' \\
& + 2\sum_{l'=1}^d \eta_{3,3}^{k,l,i'}(t)V^i(X_t^{(0)})\partial_{i'}V^m(X_t^{(0)})' + 2\sum_{l'=1}^d \eta_{3,3}^{l,i,i'}(t)V^k(X_t^{(0)})\partial_{i'}V^m(X_t^{(0)})' \\
& + 2\sum_{l'=1}^d \eta_{3,3}^{i,k,i'}(t)V^l(X_t^{(0)})\partial_{i'}V^m(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{5,4}^{i',k,l,m}(t)\partial_{i'}V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{5,4}^{i,k',l,m}(t)\partial_{k'}V_0^k(X_t^{(0)}) + \sum_{l'=1}^d \eta_{5,4}^{i,k,l',m}(t)\partial_{l'}V_0^l(X_t^{(0)}) \\
& + \sum_{m'=1}^d \eta_{5,4}^{i,k,l,m'}(t)\partial_{m'}V_0^m(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{5,5}^{i,k,l,i',k'}(t)\partial_{i'}\partial_{k'}V_0^m(X_t^{(0)}) \\
\frac{d}{dt}\eta_{5,5}^{i,k,l,m,n}(t) = & (i\xi)\left\{\eta_{4,4}^{k,l,m,n}(t)\hat{V}(X_t^{(0)},t)V^i(X_t^{(0)})' + \eta_{4,4}^{i,l,m,n}(t)\hat{V}(X_t^{(0)},t)V^k(X_t^{(0)})' \right. \\
& + \eta_{4,4}^{i,k,m,n}(t)\hat{V}(X_t^{(0)},t)V^l(X_t^{(0)})' + \eta_{4,4}^{i,k,l,n}(t)\hat{V}(X_t^{(0)},t)V^m(X_t^{(0)})' \\
& \left. + \eta_{4,4}^{i,k,l,m}(t)\hat{V}(X_t^{(0)},t)V^n(X_t^{(0)})'\right\} \\
& + \eta_{3,3}^{l,m,n}(t)V^i(X_t^{(0)})V^k(X_t^{(0)})' + \eta_{3,3}^{k,m,n}(t)V^i(X_t^{(0)})V^l(X_t^{(0)})' \\
& + \eta_{3,3}^{k,l,n}(t)V^i(X_t^{(0)})V^m(X_t^{(0)})' + \eta_{3,3}^{k,l,m}(t)V^i(X_t^{(0)})V^n(X_t^{(0)})' \\
& + \eta_{3,3}^{i,m,n}(t)V^k(X_t^{(0)})V^l(X_t^{(0)})' + \eta_{3,3}^{i,l,n}(t)V^k(X_t^{(0)})V^m(X_t^{(0)})' \\
& + \eta_{3,3}^{i,l,m}(t)V^k(X_t^{(0)})V^n(X_t^{(0)})' + \eta_{3,3}^{i,k,n}(t)V^l(X_t^{(0)})V^m(X_t^{(0)})' \\
& + \eta_{3,3}^{i,k,m}(t)V^l(X_t^{(0)})V^n(X_t^{(0)})' + \eta_{3,3}^{i,k,l}(t)V^m(X_t^{(0)})V^n(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{5,5}^{i',k,l,m,n}(t)\partial_{i'}V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{5,5}^{i,k',l,m,n}(t)\partial_{k'}V_0^k(X_t^{(0)}) \\
& + \sum_{l'=1}^d \eta_{5,5}^{i,k,l',m,n}(t)\partial_{l'}V_0^l(X_t^{(0)}) + \sum_{m'=1}^d \eta_{5,5}^{i,k,l,m',n}(t)\partial_{m'}V_0^m(X_t^{(0)}) \\
& + \sum_{n'=1}^d \eta_{5,5}^{i,k,l,m,n'}(t)\partial_{n'}V_0^n(X_t^{(0)})
\end{aligned}$$

$$\begin{aligned}
\frac{d}{dt}\eta_{6,3}^{i,k,l}(t) = & (i\xi)\left\{2\sum_{i'=1}^d \eta_{5,3,2}^{i',k,l}(t)\hat{V}(X_t^{(0)},t)\partial_{i'}V^i(X_t^{(0)})' + 2\sum_{k'=1}^d \eta_{5,3,2}^{i,k',l}(t)\hat{V}(X_t^{(0)},t)\partial_{k'}V^k(X_t^{(0)})' \right. \\
& \left. + 2\sum_{l'=1}^d \eta_{5,3,2}^{i,k,l'}(t)\hat{V}(X_t^{(0)},t)\partial_{l'}V^l(X_t^{(0)})'\right\} \\
& + 4\sum_{i'=1}^d \sum_{k'=1}^d \eta_{4,3}^{i',k',l}(t)\partial_{i'}V^i(X_t^{(0)})\partial_{k'}V^k(X_t^{(0)})' + 4\sum_{i'=1}^d \sum_{l'=1}^d \eta_{4,3}^{i',k,l'}(t)\partial_{i'}V^i(X_t^{(0)})\partial_{l'}V^l(X_t^{(0)})' \\
& + 4\sum_{k'=1}^d \sum_{l'=1}^d \eta_{4,3}^{i,k',l'}(t)\partial_{k'}V^k(X_t^{(0)})\partial_{l'}V^l(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{6,3}^{i',k,l}(t)\partial_{i'}V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{6,3}^{i,k',l}(t)\partial_{k'}V_0^k(X_t^{(0)}) + \sum_{l'=1}^d \eta_{6,3}^{i,k,l'}(t)\partial_{l'}V_0^l(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{6,4}^{i',k',k,l}(t)\partial_{i'}\partial_{k'}V_0^i(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{6,4}^{i',k',i,l}(t)\partial_{i'}\partial_{k'}V_0^k(X_t^{(0)})
\end{aligned}$$

$$\begin{aligned}
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{6,4}^{i',k',i,k}(t) \partial_{i'} \partial_{k'} V_0^l(X_t^{(0)}) \\
\frac{d}{dt} \eta_{6,4}^{i,k,l,m}(t) & = (i\xi) \left\{ \eta_{5,3,2}^{k,l,m}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{5,3,2}^{i,l,m}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right. \\
& + 2 \sum_{l'=1}^d \eta_{5,4}^{i,k,l',m}(t) \hat{V}(X_t^{(0)}, t) \partial_{l'} V^l(X_t^{(0)})' + 2 \sum_{m'=1}^d \eta_{5,4}^{i,k,l,m'}(t) \hat{V}(X_t^{(0)}, t) \partial_{m'} V^m(X_t^{(0)})' \left. \right\} \\
& + \eta_{4,2,2}^{l,m}(t) V^i(X_t^{(0)}) V^k(X_t^{(0)})' \\
& + 2 \sum_{l'=1}^d \eta_{4,3}^{k,l',m}(t) V_j^i(X_t^{(0)}) \partial_{l'} V^l(X_t^{(0)})' + 2 \sum_{l'=1}^d \eta_{4,3}^{i,l',m}(t) V_j^k(X_t^{(0)}) \partial_{l'} V^l(X_t^{(0)})' \\
& + 2 \sum_{m'=1}^d \eta_{4,3}^{k,m',l}(t) V_j^i(X_t^{(0)}) \partial_{m'} V^m(X_t^{(0)})' + 2 \sum_{m'=1}^d \eta_{4,3}^{i,m',l}(t) V_j^k(X_t^{(0)}) \partial_{m'} V^m(X_t^{(0)})' \\
& + 4 \sum_{l'=1}^d \sum_{m'=1}^d \eta_{4,4}^{i,k,l',m'}(t) \partial_{l'} V^l(X_t^{(0)}) \partial_{m'} V^m(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{6,4}^{i',k,l,m}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{6,4}^{i,k',l,m}(t) \partial_{k'} V_0^k(X_t^{(0)}) \\
& + \sum_{l'=1}^d \eta_{6,4}^{i,k,l',m}(t) \partial_{l'} V_0^l(X_t^{(0)}) + \sum_{m'=1}^d \eta_{6,4}^{i,k,l,m'}(t) \partial_{m'} V_0^m(X_t^{(0)}) \\
& + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{6,5}^{i,k,i',k',m}(t) \partial_{i'} \partial_{k'} V_0^l(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{6,5}^{i,k,l',k',l}(t) \partial_{i'} \partial_{k'} V_0^m(X_t^{(0)}) \\
\frac{d}{dt} \eta_{6,5}^{i,k,l,m,n}(t) & = (i\xi) \left\{ \eta_{5,4}^{k,l,m,n}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{5,4}^{i,l,m,n}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right. \\
& + \eta_{5,4}^{i,k,m,n}(t) \hat{V}(X_t^{(0)}, t) V^l(X_t^{(0)})' + \eta_{5,4}^{i,k,l,n}(t) \hat{V}(X_t^{(0)}, t) V^m(X_t^{(0)})' \\
& + 2 \sum_{n'=1}^d \eta_{5,5}^{i,k,l,m,n'}(t) \hat{V}(X_t^{(0)}, t) \partial_{n'} V^n(X_t^{(0)})' \left. \right\} \\
& + \eta_{4,3}^{l,m,n}(t) V^i(X_t^{(0)}) V^k(X_t^{(0)})' + \eta_{4,3}^{k,m,n}(t) V^i(X_t^{(0)}) V^l(X_t^{(0)})' + \eta_{4,3}^{k,l,n}(t) V^i(X_t^{(0)}) V^m(X_t^{(0)})' \\
& + \eta_{4,3}^{i,m,n}(t) V^k(X_t^{(0)}) V^l(X_t^{(0)})' + \eta_{4,3}^{i,l,n}(t) V^k(X_t^{(0)}) V^m(X_t^{(0)})' + \eta_{4,3}^{i,k,n}(t) V^l(X_t^{(0)}) V^m(X_t^{(0)})' \\
& + 2 \sum_{n'=1}^d \eta_{4,4}^{k,l,m,n'}(t) V^i(X_t^{(0)}) \partial_{n'} V^n(X_t^{(0)})' + 2 \sum_{n'=1}^d \eta_{4,4}^{i,l,m,n'}(t) V^k(X_t^{(0)}) \partial_{n'} V^n(X_t^{(0)})' \\
& + 2 \sum_{n'=1}^d \eta_{4,4}^{i,k,m,n'}(t) V^l(X_t^{(0)}) \partial_{n'} V^n(X_t^{(0)})' + 2 \sum_{n'=1}^d \eta_{4,4}^{i,k,l,n'}(t) V^m(X_t^{(0)}) \partial_{n'} V^n(X_t^{(0)})' \\
& + \sum_{i'=1}^d \eta_{6,5}^{i',k,l,m,n}(t) \partial_{i'} V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{6,5}^{i,k',l,m,n}(t) \partial_{k'} V_0^k(X_t^{(0)}) \\
& + \sum_{l'=1}^d \eta_{6,5}^{i,k,l',m,n}(t) \partial_{l'} V_0^l(X_t^{(0)}) + \sum_{m'=1}^d \eta_{6,5}^{i,k,l,m',n}(t) \partial_{m'} V_0^m(X_t^{(0)}) \\
& + \sum_{n'=1}^d \eta_{6,5}^{i,k,l,m,n'}(t) \partial_{n'} V_0^n(X_t^{(0)}) + \sum_{i'=1}^d \sum_{k'=1}^d \eta_{6,6}^{i,k,l,m,i',k'}(t) \partial_{i'} \partial_{k'} V_0^n(X_t^{(0)}) \\
\frac{d}{dt} \eta_{6,6}^{i,k,l,m,n,o}(t) & = (i\xi) \left\{ \eta_{5,5}^{k,l,m,n,o}(t) \hat{V}(X_t^{(0)}, t) V^i(X_t^{(0)})' + \eta_{5,5}^{i,l,m,n,o}(t) \hat{V}(X_t^{(0)}, t) V^k(X_t^{(0)})' \right.
\end{aligned}$$

$$\begin{aligned}
& +\eta_{5,5}^{i,k,m,n,o}(t)\hat{V}(X_t^{(0)},t)V^l(X_t^{(0)})' + \eta_{5,5}^{i,k,l,n,o}(t)\hat{V}(X_t^{(0)},t)V^m(X_t^{(0)})' \\
& +\eta_{5,5}^{i,k,m,l,o}(t)\hat{V}(X_t^{(0)},t)V^n(X_t^{(0)})' + \eta_{5,5}^{i,k,m,l,n}(t)\hat{V}(X_t^{(0)},t)V^o(X_t^{(0)})' \} \\
& +\eta_{4,4}^{l,m,n,o}(t)V^i(X_t^{(0)})V^k(X_t^{(0)})' + \eta_{4,4}^{k,m,n,o}(t)V^i(X_t^{(0)})V^l(X_t^{(0)})' \\
& +\eta_{4,4}^{k,l,n,o}(t)V^i(X_t^{(0)})V^m(X_t^{(0)})' + \eta_{4,4}^{k,l,m,o}(t)V^i(X_t^{(0)})V^n(X_t^{(0)})' \\
& +\eta_{4,4}^{k,l,m,n}(t)V^i(X_t^{(0)})V^o(X_t^{(0)})' + \eta_{4,4}^{i,m,n,o}(t)V^k(X_t^{(0)})V^l(X_t^{(0)})' \\
& +\eta_{4,4}^{i,l,n,o}(t)V^k(X_t^{(0)})V^m(X_t^{(0)})' + \eta_{4,4}^{i,l,m,o}(t)V^k(X_t^{(0)})V^n(X_t^{(0)})' \\
& +\eta_{4,4}^{i,l,m,n}(t)V^k(X_t^{(0)})V^o(X_t^{(0)})' + \eta_{4,4}^{i,k,n,o}(t)V^l(X_t^{(0)})V^m(X_t^{(0)})' \\
& +\eta_{4,4}^{i,k,m,o}(t)V^l(X_t^{(0)})V^n(X_t^{(0)})' + \eta_{4,4}^{i,k,m,n}(t)V^l(X_t^{(0)})V^o(X_t^{(0)})' \\
& +\eta_{4,4}^{i,k,l,o}(t)V^m(X_t^{(0)})V^n(X_t^{(0)})' + \eta_{4,4}^{i,k,l,n}(t)V^m(X_t^{(0)})V^o(X_t^{(0)})' \\
& +\eta_{4,4}^{i,k,l,m}(t)V^n(X_t^{(0)})V^o(X_t^{(0)})' \\
& +\sum_{i'=1}^d \eta_{6,6}^{i',k,l,m,n,o}(t)\partial_{i'}V_0^i(X_t^{(0)}) + \sum_{k'=1}^d \eta_{6,6}^{i,k',l,m,n,o}(t)\partial_{k'}V_0^k(X_t^{(0)}) \\
& +\sum_{l'=1}^d \eta_{6,6}^{i,k,l',m,n,o}(t)\partial_{l'}V_0^l(X_t^{(0)}) + \sum_{m'=1}^d \eta_{6,6}^{i,k,l,m',n,o}(t)\partial_{m'}V_0^m(X_t^{(0)}) \\
& +\sum_{n'=1}^d \eta_{6,6}^{i,k,l,m,n',o}(t)\partial_{n'}V_0^n(X_t^{(0)}) + \sum_{o'=1}^d \eta_{6,6}^{i,k,l,m,n,o'}(t)\partial_{o'}V_0^o(X_t^{(0)})
\end{aligned}$$

From the derivation of differential equations, it is easily shown that each $\eta_{j,m,k}(T)$ is expressed as a polynomial of degree j with respect to $(i\xi)$, and also it is shown that $\mathbf{E}[X^{j,m,k}Z_T^{(\xi)}]$ is a polynomial of $(i\xi)$ of degree less than or equal to $2j$. Thus, in the asymptotic expansion scheme, the infinite sum in the expansion (26) is replaced by a finite sum.

We summarize the discussion above as the following theorem:

Theorem 2 *The asymptotic expansion of density of $G^{(\epsilon)}$ up to ϵ^3 -order is given by*

$$\begin{aligned}
f_{G^{(\epsilon)}}(x) &= f_{g_{1T}}(x) \\
&+ \epsilon \left\{ \sum_{l=1}^3 C_{1l} H_l(x; \Sigma_T) \right\} f_{g_{1T}}(x) \\
&+ \epsilon^2 \left\{ \sum_{l=1}^6 C_{2l} H_l(x; \Sigma_T) \right\} f_{g_{1T}}(x) \\
&+ \epsilon^3 \left\{ \sum_{l=1}^9 C_{3l} H_l(x; \Sigma_T) \right\} f_{g_{1T}}(x) \\
&+ o(\epsilon^3).
\end{aligned}$$

where

$$\begin{aligned}
C_{1l} &= \Sigma_T a_{l-1}^{1,1,(1)}, \\
C_{2l} &= \Sigma_T a_{l-1}^{2,1,(0,1)} + \frac{1}{2} \Sigma_T^2 a_{l-2}^{2,2,(2,0)}, \\
C_{3l} &= \Sigma_T a_{l-1}^{3,1,(0,0,1)} + \frac{1}{2} \Sigma_T^2 a_{l-2}^{3,2,(1,2,0)} + \frac{1}{6} \Sigma_T^3 a_{l-2}^{3,3,(3,0,0)}.
\end{aligned}$$

$a_l^{j,m,k}$ are given by (29), and expectations in (29) are obtained as the solutions to the system of ordinary differential equations given in Proposition 2.

4.2 Asymptotic Expansion of Option Prices

We apply the asymptotic expansion to option pricing. We consider the plain vanilla option on the underlying asset $g(X_T^{(\epsilon)})$ whose dynamics is given by (1). For example, the call option price with strike K and maturity T is given by

$$C(K, T) = \epsilon P(0, T) \int_{-k^{(\epsilon)}}^{\infty} (x + k^{(\epsilon)}) f_{G^{(\epsilon)}}(x) dx$$

where $k^{(\epsilon)} = \frac{G^{(0)} - K}{\epsilon}$, $P(0, T)$ denotes the price at time 0 of a zero coupon bond with maturity T , and $f_{G^{(\epsilon)}}$ is the normal asymptotic expansion of density of $G^{(\epsilon)}$ given by (28). In particular, using the result of the previous subsection, the approximated price to the option up to the fourth order can be expressed as

$$\begin{aligned} C(K, T) &= \epsilon P(0, T) \int_{-k^{(\epsilon)}}^{\infty} (x + k^{(\epsilon)}) f_{g_{1T}}(x) dx \\ &+ \epsilon^2 P(0, T) \int_{-k^{(\epsilon)}}^{\infty} (x + k^{(\epsilon)}) \left\{ \sum_{l=1}^3 C_{1l} H_l(x; \Sigma_T) \right\} f_{g_{1T}}(x) dx \\ &+ \epsilon^3 P(0, T) \int_{-k^{(\epsilon)}}^{\infty} (x + k^{(\epsilon)}) \left\{ \sum_{l=1}^6 C_{2l} H_l(x; \Sigma_T) \right\} f_{g_{1T}}(x) dx \\ &+ \epsilon^4 P(0, T) \int_{-k^{(\epsilon)}}^{\infty} (x + k^{(\epsilon)}) \left\{ \sum_{l=1}^9 C_{3l} H_l(x; \Sigma_T) \right\} f_{g_{1T}}(x) dx \\ &+ o(\epsilon^4). \end{aligned}$$

Integrals appeared in the right hand side can be calculated using following formulas related to the Hermite polynomial

$$\begin{aligned} \int_{-y}^{\infty} H_k(x; \Sigma) f_{g_{1T}}(x) dx &= \Sigma H_{k-1}(-y; \Sigma) f_{g_{1T}}(y) \quad (k \geq 1), \\ \int_{-y}^{\infty} x H_k(x; \Sigma) f_{g_{1T}}(x) dx &= -\Sigma y H_{k-1}(-y; \Sigma) f_{g_{1T}}(y) \\ &+ \Sigma^2 H_{k-2}(-y; \Sigma) f_{g_{1T}}(y) \quad (k \geq 2). \end{aligned}$$

4.3 Log-Normal Asymptotic Expansion

Suppose that the underlying asset process $S^{(\epsilon)}$ follows

$$\begin{aligned} dS_t^{(\epsilon)} &= g(X_t^{(\epsilon)}) S_t^{(\epsilon)} \bar{\sigma} dW_t; \quad S_0^{(\epsilon)} = s_0 \\ dX_t^{(\epsilon)} &= V_0(X_t^{(\epsilon)}, \epsilon) dt + \epsilon V(X_t^{(\epsilon)}) dW_t; \quad X_0^{(\epsilon)} = x_0 \in \mathbf{R}^d. \end{aligned}$$

Define $\hat{X}^{(\epsilon)}$ as

$$\hat{X}_t^{(\epsilon)} = \log \left(\frac{S_t^{(\epsilon)}}{s_0} \right).$$

Then, we have

$$\hat{X}_t^{(\epsilon)} = -\frac{|\bar{\sigma}|^2}{2} \int_0^t g(X_u^{(\epsilon)})^2 du + \int_0^t g(X_u^{(\epsilon)}) \bar{\sigma} dW_u,$$

and note that

$$\hat{X}_T^{(0)} \sim N(\hat{\mu}_T, \hat{\Sigma}_T),$$

where

$$\begin{aligned}\hat{\mu}_T &= -\frac{|\bar{\sigma}|^2}{2} \int_0^T g(X_u^{(0)})^2 du = -\frac{1}{2} \hat{\Sigma}_T, \\ \hat{\Sigma}_T &= |\bar{\sigma}|^2 \int_0^T g(X_u^{(0)})^2 du.\end{aligned}$$

Moreover, an asymptotic expansion of $\hat{X}_T^{(\epsilon)}$ up to ϵ^N -order is expressed as

$$\hat{X}_T^{(\epsilon)} = \hat{X}_T^{(0)} + \sum_{n=1}^N \frac{\epsilon^n}{n!} \hat{A}_{nT} + o(\epsilon^N),$$

where $\hat{A}_{nt} = \frac{\partial^n \hat{X}_t^{(\epsilon)}}{\partial \epsilon^n} |_{\epsilon=0}$. Note that $S^{(\epsilon)}$ is expanded around a log-normal distribution since $\hat{X}_T^{(0)}$ has a Gaussian distribution.

Next, define $Z_t^{(\xi)}$ as

$$Z_t^{(\xi)} = \exp\left(i\xi \int_0^t g(X_u^{(0)}) \bar{\sigma} dW_u\right).$$

Then, the result in the previous subsection is applied to deriving the density function of $\hat{X}_T^{(\epsilon)}$ if $G^{(\epsilon)}$ is replaced by $\hat{X}_T^{(\epsilon)}$.

Similar to the normal case, the log-normal asymptotic expansion of the price of the call option on $\hat{X}_T^{(\epsilon)}$ is given by

$$C(K, T) = P(0, T) \int_{\log \frac{K}{s_0}}^{\infty} (s_0 e^x - K) f_{\hat{X}_T^{(\epsilon)}}(x) dx$$

5 Numerical Examples

5.1 λ -SABR model

To test the validity of the expansion, we first consider the European plain-vanilla call and put prices under the following λ -SABR model [24] (interest rate=0%) :

$$\begin{aligned}dS^{(\epsilon)}(t) &= \epsilon \sigma^{(\epsilon)}(t) (S^{(\epsilon)}(t))^{\beta} dW_t^1, \\ d\sigma^{(\epsilon)}(t) &= \lambda(\theta - \sigma^{(\epsilon)}(t)) dt + \epsilon \nu_1 \sigma^{(\epsilon)}(t) dW_t^1 + \epsilon \nu_2 \sigma^{(\epsilon)}(t) dW_t^2,\end{aligned}$$

where $\nu_1 = \rho \nu$, $\nu_2 = (\sqrt{1 - \rho^2}) \nu$. (The correlation between S and σ is $\rho \in [-1, 1]$.)

Approximated prices by the asymptotic expansion method are calculated up to the fifth order. Note that all the solutions to differential equations are obtained analytically. Benchmark values are computed by Monte Carlo simulations. ϵ is set to be one and other parameters used in the test are given in Table 1:

Table 1:

Parameter	λ	$\sigma(0)$	β	ρ	θ	ν	T
i	0.1	3.0	0.5	-0.7	3.0	0.3	10
ii	0.1	3.0	0.5	-0.7	3.0	0.1	10
iii	0.1	3.0	0.5	-0.7	3.0	0.3	1

In Monte Carlo simulations for benchmark values, we use Euler-Maruyama scheme as a discretization scheme with 1024, 1024, and 512 time steps for case i, ii, and iii respectively, and generate 10^8 paths in each simulation.

For the case of $\beta = 1$ in the λ -SABR model, we can apply the log-normal asymptotic expansion method given in the previous section. To test the efficiency of the high order log-normal asymptotic expansion method, we consider the European plain-vanilla call and put prices under the following parameters (and $\epsilon = 1$ as well as in the previous examples) with different maturities:

Table 2:

Parameter	λ	$\sigma(0)$	β	ρ	θ	ν	T
iv	0.1	0.3	1.0	-0.7	0.3	0.3	10
v	0.1	0.3	1.0	-0.7	0.3	0.3	20
vi	0.1	0.3	1.0	-0.7	0.3	0.3	30

We calculate approximated prices by the log-normal asymptotic expansion method up to the fourth order. Benchmark prices are computed by Monte Carlo simulations. In the simulations, we adapt the second order discretization scheme given by Ninomiya-Victoir [34] with 128, 256, 256 time steps respectively.

Results are in Table 3 and Table 4.

From the results, in each case, the higher order asymptotic expansion or log-normal asymptotic expansion almost always improve the accuracy of approximation by the lower expansions. Improvement is significant especially in long-term cases in which the lower order asymptotic expansions cannot approximate the price well.

Table 3:

Case	Strike(C/P)	MC	A.E.(Difference)					A.E.(Relative Difference)					
			1st	2nd	3rd	4th	5th	1st	2nd	3rd	4th	5th	
i	50 Put	13.109	4.876	5.000	2.313	1.067	0.260	37.20 %	38.14 %	17.64 %	8.14 %	1.98 %	
	60 Put	16.618	4.544	4.648	1.931	0.938	0.195	27.34 %	27.97 %	11.62 %	5.65 %	1.17 %	
	70 Put	20.482	4.241	4.322	1.585	0.844	0.149	20.71 %	21.10 %	7.74 %	4.12 %	0.73 %	
	80 Put	24.720	3.965	4.020	1.269	0.778	0.117	16.04 %	16.26 %	5.14 %	3.15 %	0.47 %	
	90 Put	29.347	3.710	3.738	0.980	0.735	0.094	12.64 %	12.74 %	3.34 %	2.51 %	0.32 %	
	100 Call	34.375	3.472	3.472	0.712	0.712	0.077	10.10 %	10.10 %	2.07 %	2.07 %	0.22 %	
	110 Call	29.811	3.246	3.217	0.459	0.704	0.063	10.89 %	10.79 %	1.54 %	2.36 %	0.21 %	
	120 Call	25.659	3.026	2.971	0.220	0.711	0.050	11.79 %	11.58 %	0.86 %	2.77 %	0.19 %	
	130 Call	21.914	2.809	2.728	-0.010	0.731	0.035	12.82 %	12.45 %	-0.04 %	3.33 %	0.16 %	
	140 Call	18.571	2.591	2.487	-0.230	0.762	0.018	13.95 %	13.39 %	-1.24 %	4.10 %	0.10 %	
	150 Call	15.615	2.370	2.246	-0.441	0.804	-0.002	15.18 %	14.38 %	-2.83 %	5.15 %	-0.02 %	
	ii	50 Put	1.682	-0.914	0.030	0.475	0.182	-0.016	-54.33 %	1.81 %	28.25 %	10.84 %	-0.92 %
		60 Put	2.607	-1.056	0.129	0.445	0.103	-0.003	-40.52 %	4.94 %	17.06 %	3.95 %	-0.13 %
		70 Put	3.950	-1.047	0.214	0.364	0.061	0.008	-26.51 %	5.41 %	9.22 %	1.55 %	0.20 %
		80 Put	5.883	-0.825	0.254	0.258	0.048	0.013	-14.03 %	4.32 %	4.39 %	0.82 %	0.23 %
90 Put		8.631	-0.390	0.237	0.150	0.047	0.016	-4.52 %	2.75 %	1.74 %	0.54 %	0.18 %	
100 Call		12.450	0.166	0.166	0.048	0.048	0.018	1.33 %	1.33 %	0.39 %	0.39 %	0.14 %	
110 Call		7.577	0.664	0.037	-0.050	0.053	0.022	8.76 %	0.49 %	-0.67 %	0.70 %	0.29 %	
120 Call		4.131	0.927	-0.153	-0.149	0.062	0.027	22.43 %	-3.70 %	-3.60 %	1.49 %	0.65 %	
130 Call		2.008	0.894	-0.367	-0.217	0.086	0.033	44.52 %	-18.27 %	-10.79 %	4.30 %	1.64 %	
140 Call		0.887	0.663	-0.522	-0.205	0.136	0.030	74.77 %	-58.78 %	-23.16 %	15.35 %	3.36 %	
150 Call		0.372	0.396	-0.548	-0.104	0.189	-0.009	106.35 %	-147.29 %	-27.82 %	50.82 %	-2.34 %	
iii		50 Put	0.633	-0.038	0.094	0.061	0.015	0.005	-6.05 %	14.84 %	9.64 %	2.33 %	0.85 %
		60 Put	1.335	-0.063	0.111	0.058	0.013	0.006	-4.74 %	8.32 %	4.34 %	0.97 %	0.42 %
		70 Put	2.571	-0.072	0.121	0.048	0.011	0.006	-2.79 %	4.72 %	1.87 %	0.45 %	0.22 %
		80 Put	4.579	-0.046	0.124	0.034	0.010	0.005	-1.00 %	2.71 %	0.75 %	0.22 %	0.12 %
	90 Put	7.608	0.019	0.119	0.019	0.008	0.004	0.25 %	1.57 %	0.26 %	0.11 %	0.05 %	
	100 Call	11.857	0.111	0.111	0.008	0.008	0.004	0.94 %	0.94 %	0.07 %	0.07 %	0.03 %	
	110 Call	7.430	0.197	0.096	-0.004	0.008	0.003	2.65 %	1.29 %	-0.05 %	0.10 %	0.05 %	
	120 Call	4.289	0.244	0.074	-0.015	0.009	0.004	5.70 %	1.74 %	-0.36 %	0.20 %	0.09 %	
	130 Call	2.260	0.239	0.046	-0.027	0.009	0.003	10.57 %	2.03 %	-1.21 %	0.40 %	0.14 %	
	140 Call	1.080	0.192	0.017	-0.036	0.009	0.002	17.77 %	1.62 %	-3.30 %	0.88 %	0.19 %	
	150 Call	0.466	0.129	-0.004	-0.036	0.010	0.001	27.62 %	-0.75 %	-7.81 %	2.13 %	0.13 %	

Table 4:

Case	Strike(C/P)	MC	Log Normal A.E.(Difference)				Log Normal A.E.(Relative Difference)						
			Log-Norm	1st	2nd	3rd	4th	Log-Norm	1st	2nd	3rd	4th	
iv	50 Put	9.429	-0.896	0.250	0.470	-0.223	0.021	-9.51 %	2.65 %	4.99 %	-2.36 %	0.22 %	
	60 Put	13.095	-0.187	0.168	0.449	-0.215	0.028	-1.43 %	1.29 %	3.43 %	-1.64 %	0.21 %	
	70 Put	17.307	0.678	0.045	0.431	-0.203	0.034	3.92 %	0.26 %	2.49 %	-1.17 %	0.19 %	
	80 Put	22.041	1.620	-0.099	0.414	-0.190	0.039	7.35 %	-0.45 %	1.88 %	-0.86 %	0.18 %	
	90 Put	27.272	2.577	-0.253	0.397	-0.177	0.045	9.45 %	-0.93 %	1.45 %	-0.65 %	0.17 %	
	100 Call	32.971	3.503	-0.416	0.379	-0.163	0.051	10.62 %	-1.26 %	1.15 %	-0.49 %	0.15 %	
	110 Call	29.110	4.367	-0.589	0.360	-0.149	0.057	15.00 %	-2.02 %	1.24 %	-0.51 %	0.20 %	
	120 Call	25.655	5.149	-0.773	0.338	-0.135	0.063	20.07 %	-3.01 %	1.32 %	-0.53 %	0.25 %	
	130 Call	22.576	5.837	-0.972	0.315	-0.120	0.069	25.85 %	-4.30 %	1.39 %	-0.53 %	0.31 %	
	140 Call	19.842	6.424	-1.186	0.289	-0.104	0.076	32.38 %	-5.98 %	1.46 %	-0.53 %	0.38 %	
	150 Call	17.420	6.912	-1.416	0.261	-0.088	0.083	39.68 %	-8.13 %	1.50 %	-0.50 %	0.47 %	
	v	50 Put	15.350	0.961	-0.125	0.782	-0.523	0.148	6.26 %	-0.82 %	5.10 %	-3.41 %	0.96 %
		60 Put	20.207	1.990	-0.391	0.823	-0.513	0.153	9.85 %	-1.93 %	4.07 %	-2.54 %	0.76 %
		70 Put	25.499	3.062	-0.664	0.857	-0.495	0.153	12.01 %	-2.60 %	3.36 %	-1.94 %	0.60 %
		80 Put	31.184	4.134	-0.937	0.884	-0.472	0.150	13.26 %	-3.00 %	2.84 %	-1.51 %	0.48 %
90 Put		37.228	5.175	-1.207	0.908	-0.446	0.145	13.90 %	-3.24 %	2.44 %	-1.20 %	0.39 %	
100 Call		43.598	6.168	-1.474	0.928	-0.417	0.137	14.15 %	-3.38 %	2.13 %	-0.96 %	0.31 %	
110 Call		40.267	7.101	-1.741	0.946	-0.387	0.129	17.63 %	-4.32 %	2.35 %	-0.96 %	0.32 %	
120 Call		37.208	7.967	-2.009	0.962	-0.356	0.119	21.41 %	-5.40 %	2.59 %	-0.96 %	0.32 %	
130 Call		34.399	8.763	-2.278	0.977	-0.323	0.107	25.47 %	-6.62 %	2.84 %	-0.94 %	0.31 %	
140 Call		31.818	9.487	-2.551	0.990	-0.289	0.095	29.82 %	-8.02 %	3.11 %	-0.91 %	0.30 %	
150 Call		29.447	10.142	-2.829	1.003	-0.255	0.082	34.44 %	-9.61 %	3.41 %	-0.87 %	0.28 %	
vi		50 Put	19.801	2.280	-0.889	1.143	-0.592	0.182	11.51 %	-4.49 %	5.77 %	-2.99 %	0.92 %
		60 Put	25.471	3.371	-1.248	1.254	-0.581	0.154	13.23 %	-4.90 %	4.93 %	-2.28 %	0.60 %
		70 Put	31.500	4.459	-1.594	1.351	-0.560	0.120	14.15 %	-5.06 %	4.29 %	-1.78 %	0.38 %
		80 Put	37.847	5.520	-1.927	1.437	-0.535	0.081	14.59 %	-5.09 %	3.80 %	-1.41 %	0.21 %
	90 Put	44.476	6.541	-2.246	1.515	-0.505	0.039	14.71 %	-5.05 %	3.41 %	-1.14 %	0.09 %	
	100 Call	51.357	7.512	-2.555	1.586	-0.474	-0.005	14.63 %	-4.98 %	3.09 %	-0.92 %	-0.01 %	
	110 Call	48.465	8.430	-2.856	1.652	-0.442	-0.051	17.39 %	-5.89 %	3.41 %	-0.91 %	-0.10 %	
	120 Call	45.780	9.291	-3.150	1.715	-0.409	-0.098	20.30 %	-6.88 %	3.75 %	-0.89 %	-0.21 %	
	130 Call	43.281	10.097	-3.439	1.774	-0.376	-0.147	23.33 %	-7.94 %	4.10 %	-0.87 %	-0.34 %	
	140 Call	40.954	10.848	-3.724	1.831	-0.342	-0.197	26.49 %	-9.09 %	4.47 %	-0.84 %	-0.48 %	
	150 Call	38.782	11.545	-4.007	1.886	-0.309	-0.248	29.77 %	-10.33 %	4.86 %	-0.80 %	-0.64 %	

5.2 Currency Option under a Libor Market Model of Interest Rates and a Stochastic Volatility of a Spot Exchange Rate

In this subsection, we apply our methods to pricing options on currencies under Libor Market Models(LMMs) of interest rates and a stochastic volatility of the spot foreign exchange rate(Forex). Due to limitation of space, only the structure of the stochastic differential equations of our model is described here. For details of the underlying model, see Takahashi and Takehara [53].

5.2.1 Cross-Currency Libor Market Models

Let $(\Omega, \mathcal{F}, \tilde{P}, \{\mathcal{F}_t\}_{0 \leq t \leq T^* < \infty})$ be a complete probability space with a filtration satisfying the usual conditions. We consider the following pricing problem for the call option with maturity $T \in (0, T^*]$ and strike rate $K > 0$;

$$V^C(0; T, K) = P_d(0, T) \times \mathbf{E}^P [(S(T) - K)^+] = P_d(0, T) \times \mathbf{E}^P [(F_T(T) - K)^+] \quad (32)$$

where $V^C(0; T, K)$ denotes the value of an European call option at time 0 with maturity T and strike rate K , $S(t)$ denotes the spot exchange rate at time $t \geq 0$ and $F_T(t)$ denotes the time t value of the forex forward rate with maturity T . Similarly, for the put option we consider

$$V^P(0; T, K) = P_d(0, T) \times \mathbf{E}^P [(K - S(T))^+] = P_d(0, T) \times \mathbf{E}^P [(K - F_T(T))^+] \quad (33)$$

It is well known that the arbitrage-free relation between the forex spot rate and the forex forward rate are given by $F_T(t) = S(t) \frac{P_f(t, T)}{P_d(t, T)}$ where $P_d(t, T)$ and $P_f(t, T)$ denote the time t values of domestic and foreign zero coupon bonds with maturity T respectively. $\mathbf{E}^P[\cdot]$ denotes an expectation operator under EMM(Equivalent Martingale Measure) P whose associated numeraire is the domestic zero coupon bond maturing at T .

For these pricing problems, a market model and a stochastic volatility model are applied to modeling interest rates' and the spot exchange rate's dynamics respectively.

We first define domestic and foreign forward interest rates as $f_{dj}(t) = \left(\frac{P_d(t, T_j)}{P_d(t, T_{j+1})} - 1 \right) \frac{1}{\tau_j}$ and $f_{fj}(t) = \left(\frac{P_f(t, T_j)}{P_f(t, T_{j+1})} - 1 \right) \frac{1}{\tau_j}$ respectively, where $j = n(t), n(t) + 1, \dots, N$, $\tau_j = T_{j+1} - T_j$, and $P_d(t, T_j)$ and $P_f(t, T_j)$ denote the prices of domestic/foreign zero coupon bonds with maturity T_j at time $t (\leq T_j)$ respectively; $n(t) = \min\{i : t \leq T_i\}$. We also define spot interest rates to the nearest fixing date denoted by $f_{d, n(t)-1}(t)$ and $f_{f, n(t)-1}(t)$ as $f_{d, n(t)-1}(t) = \left(\frac{1}{P_d(t, T_{n(t)})} - 1 \right) \frac{1}{(T_{n(t)} - t)}$ and $f_{f, n(t)-1}(t) = \left(\frac{1}{P_f(t, T_{n(t)})} - 1 \right) \frac{1}{(T_{n(t)} - t)}$. Finally, we set $T = T_{N+1}$ and will abbreviate $F_{T_{N+1}}(t)$ to $F_{N+1}(t)$ in what follows.

Under the framework of the asymptotic expansion in the standard cross-currency libor market model, we have to consider the following system of stochastic differential equations(henceforth called S.D.E.s) under the domestic terminal measure P to price options. For detailed arguments on the framework of these S.D.E.s see [53].

As for the domestic and foreign interest rates we assume forward market

models; for $j = n(t) - 1, n(t), n(t) + 1, \dots, N$,

$$f_{dj}^{(\epsilon)}(t) = f_{dj}(0) + \epsilon^2 \sum_{i=j+1}^N \int_0^t g_{di}^{0,(\epsilon)}(u)' \gamma_{dj}(u) f_{dj}^{(\epsilon)}(u) du + \epsilon \int_0^t f_{dj}^{(\epsilon)}(u) \gamma'_{dj}(u) dW_u, \quad (34)$$

$$\begin{aligned} f_{fj}^{(\epsilon)}(t) &= f_{fj}(0) - \epsilon^2 \sum_{i=0}^j \int_0^t g_{fi}^{0,(\epsilon)}(u)' \gamma_{fj}(u) f_{fj}^{(\epsilon)}(u) du + \epsilon^2 \sum_{i=0}^N \int_0^t g_{di}^{0,(\epsilon)}(u)' \gamma_{fj}(u) f_{fj}^{(\epsilon)}(u) du \\ &\quad - \epsilon^2 \int_0^t \sigma^{(\epsilon)}(u) \bar{\sigma}' \gamma_{fj}(u) f_{fj}^{(\epsilon)}(u) du + \epsilon \int_0^t f_{fj}^{(\epsilon)}(u) \gamma'_{fj}(u) dW_u, \end{aligned} \quad (35)$$

where

$$g_{dj}^{0,(\epsilon)}(t) := \frac{-\tau_j f_{dj}^{(\epsilon)}(t)}{1 + \tau_j f_{dj}^{(\epsilon)}(t)} \gamma_{dj}(t), \quad g_{fj}^{0,(\epsilon)}(t) := \frac{-\tau_j f_{fj}^{(\epsilon)}(t)}{1 + \tau_j f_{fj}^{(\epsilon)}(t)} \gamma_{fj}(t);$$

x' denotes the transpose of x , $\hat{J}_{j+1} := \{0, 1, \dots, j\}$, and W is a d' -dimensional standard Wiener process under the domestic terminal measure P ; $\gamma_{dj}(s)$, $\gamma_{fj}(s)$ are d' -dimensional vector-valued functions of time-parameter s ; $\bar{\sigma}$ denotes a d' -dimensional constant vector satisfying $\|\bar{\sigma}\| = 1$ and $\sigma(t)$, the volatility of the spot exchange rate, is specified to follow a \mathbf{R}_{++} -valued general time-inhomogeneous Markovian process as follows:

$$\sigma(t) = \sigma(0) + \int_0^t \mu(u, \sigma^{(\epsilon)}(u)) du + \epsilon^2 \sum_{j=1}^N \int_0^t g_{dj}^{0,(\epsilon)}(u)' \omega(u, \sigma^{(\epsilon)}(u)) du + \epsilon \int_0^t \omega'(u, \sigma^{(\epsilon)}(u)) dW_u, \quad (36)$$

where $\mu(s, x)$ and $\omega(s, x)$ are functions of s and x .

Finally, we consider the process of the forex forward $F_{N+1}(t)$. Since $F_{N+1}(t) \equiv F_{T_{N+1}}(t)$ can be expressed as $F_{N+1}(t) = S(t) \frac{P_f(t, T_{N+1})}{P_d(t, T_{N+1})}$, we easily notice that it is a martingale under the domestic terminal measure. In particular, it satisfies the following stochastic differential equation

$$F_T^{(\epsilon)}(t) = F_T(0) + \epsilon \int_0^t \sigma_F^{(\epsilon)}(u)' F^{(\epsilon)}(u) dW_u \quad (37)$$

where

$$\sigma_F^{(\epsilon)}(t) := \sum_{j=0}^N \left(g_{fj}^{0,(\epsilon)}(t) - g_{dj}^{0,(\epsilon)}(t) \right) + \sigma^{(\epsilon)}(t).$$

5.2.2 Numerical Examples

We here specify our model and parameters, and confirm the effectiveness of our method in this cross-currency framework.

First of all, the processes of domestic and foreign forward interest rates and of the volatility of the spot exchange rate are specified. We suppose $d' = 4$, that is the dimension of a Brownian motion is set to be four; it represents the uncertainty of domestic and foreign interest rates, the spot exchange rate, and its volatility. Note that in this framework correlations among all factors are allowed.

Next, we specify a volatility process of the spot exchange rate in (36) with

$$\begin{cases} \mu(s, x) = \kappa(\theta - x), \\ \omega(s, x) = \omega x, \end{cases} \quad (38)$$

Table 5: Initial domestic/foreign forward interest rates and their volatilities

	f_d	γ_d^*	f_f	γ_f^*
case (i)	0.05	0.12	0.05	0.12
case (ii)	0.02	0.3	0.05	0.12
case (iii)	0.05	0.12	0.02	0.3
case (iv)	0.02	0.3	0.02	0.3

where θ and κ represent the level and speed of its mean-reversion respectively, and ω denotes a volatility vector on the volatility. In this section the parameters are set as follows; $\epsilon = 1$, $\sigma(0) = \theta = 0.1$, and $\kappa = 0.1$; $\omega = \omega^* \bar{v}$ where $\omega^* = 0.3$ and \bar{v} denotes a four dimensional constant vector given below.

We further suppose that initial term structures of domestic and foreign forward interest rates are flat, and their volatilities also have flat structures and are constant over time: that is, for all j , $f_{dj}(0) = f_d$, $f_{fj}(0) = f_f$, $\gamma_{dj}(t) = \gamma_d^* \bar{\gamma}_d 1_{\{t < T_j\}}(t)$ and $\gamma_{fj}(t) = \gamma_f^* \bar{\gamma}_f 1_{\{t < T_j\}}(t)$. Here, γ_d^* and γ_f^* are constant scalars, and $\bar{\gamma}_d$ and $\bar{\gamma}_f$ denote four dimensional constant vectors. Moreover, given a correlation matrix \underline{C} among all four factors, the constant vectors $\bar{\gamma}_d$, $\bar{\gamma}_f$, $\bar{\sigma}$ and \bar{v} can be determined to satisfy $\|\bar{\gamma}_d\| = \|\bar{\gamma}_f\| = \|\bar{\sigma}\| = \|\bar{v}\| = 1$ and $V'V = \underline{C}$ where $V := (\bar{\gamma}_d, \bar{\gamma}_f, \bar{\sigma}, \bar{v})$.

In this subsection, we consider four different cases for f_d , γ_d^* , f_f and γ_f^* as in Table 5. For correlations, four sets of parameters are considered: In the case ‘‘Corr.1’’, all the factors are independent: In ‘‘Corr.2’’, there exists only the correlation of -0.5 between the spot exchange rate and its volatility (i.e. $\bar{\sigma}'\bar{v} = -0.5$) while there are no correlations among the others: In ‘‘Corr.3’’, the correlation between interest rates and the spot exchange rate are allowed while there are no correlations among the others; the correlation between domestic ones and the spot forex is 0.5 ($\bar{\gamma}_d'\bar{\sigma} = 0.5$) and the correlation between foreign ones and the spot forex is -0.5 ($\bar{\gamma}_f'\bar{\sigma} = -0.5$): Finally in ‘‘Corr.4’’, more intricately correlated structure is considered; $\bar{\gamma}_d'\bar{\sigma} = 0.5$, $\bar{\gamma}_f'\bar{\sigma} = -0.5$ between interest rates and the spot forex; and $\bar{\sigma}'\bar{v} = -0.5$ between the spot forex and its volatility. It is well known that (both of exact and approximate) evaluation of the long-term options is a hard task in the case with complex structures of correlations such as in ‘‘Corr.3’’ or ‘‘Corr.4’’.

Lastly, we make an assumption that $\gamma_{dn(t)-1}(t)$ and $\gamma_{fn(t)-1}(t)$, volatilities of the domestic and foreign interest rates applied to the period from t to the next fixing date $T_{n(t)}$, are equal to be zero for arbitrary $t \in [t, T_{n(t)}]$.

In Table 6-9 and Figure 1, we compare our estimations of the values of call and put options by an asymptotic expansion up to the fourth order to the benchmarks estimated by 10^6 trials of Monte Carlo simulation which is discretized by Euler-Maruyama scheme with time step 0.05 and applied the Antithetic Variable Method. For the moneynesses (defined by $K/F_{N+1}(0)$) less than one, the prices of put options are shown; otherwise, the prices of call options are displayed.

As seen in these tables and figure, in general the estimators show more accuracy as the order of the expansion increases. Especially, for the deep OTM options the fourth order approximation performs much better and is stabler than the approximation with lower orders.

Table 6

Case (i)		Corr.1																			
Moneyness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0021	0.0136	0.051	0.1504	0.3865	0.8867	1.8303	3.4047	5.7408	8.861	6.5943	4.9199	3.7043	2.8275	2.1939	1.7316	1.3904	1.1351	0.9451	0.7922
A.E.	1st	0.0325	0.0781	0.1749	0.3663	0.7181	1.3209	2.2859	3.7325	5.7694	8.4733	5.7694	3.7324	2.2859	1.3209	0.7181	0.3663	0.1749	0.078	0.0325	0.0126
	2nd	-0.102	-0.163	-0.217	-0.209	-0.037	0.4454	1.4093	3.0141	5.3628	8.4733	6.1759	4.4508	3.1625	2.1964	1.4728	0.9414	0.5673	0.3194	0.1671	0.0809
	3rd	0.2001	0.2571	0.3089	0.386	0.5755	1.0293	1.9409	3.4945	5.8091	8.9081	6.6222	4.9312	3.6941	2.7802	2.085	1.5363	1.0936	0.7398	0.4693	0.2763
	4th	-0.181	-0.129	-0.031	0.1267	0.4012	0.9201	1.8706	3.4461	5.7786	8.8977	6.633	4.9629	3.7517	2.8805	2.2531	1.7907	1.4292	1.1223	0.847	0.6014
	Diff.																				
A.E.	1st	0.0304	0.0645	0.1239	0.2159	0.3316	0.4342	0.4556	0.3278	0.0286	-0.388	-0.825	-1.188	-1.418	-1.507	-1.476	-1.365	-1.216	-1.057	-0.913	-0.78
	2nd	-0.104	-0.177	-0.268	-0.359	-0.423	-0.441	-0.421	-0.391	-0.378	-0.388	-0.418	-0.469	-0.542	-0.631	-0.721	-0.79	-0.823	-0.816	-0.778	-0.711
	3rd	0.198	0.2435	0.2579	0.2356	0.189	0.1426	0.1106	0.0898	0.0683	0.0471	0.0279	0.0113	-0.01	-0.047	-0.109	-0.195	-0.297	-0.395	-0.476	-0.516
	4th	-0.183	-0.143	-0.082	-0.024	0.0147	0.0334	0.0403	0.0414	0.0378	0.0367	0.0387	0.043	0.0474	0.053	0.0592	0.0591	0.0388	-0.013	-0.098	-0.191

Case (ii)		Corr.1																			
Moneyness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0021	0.0142	0.0526	0.1534	0.3897	0.8898	1.8359	3.4182	5.7667	8.9083	6.6647	5.0085	3.8038	2.9298	2.2932	1.8239	1.4742	1.2101	1.0074	0.8502
A.E.	1st	0.0329	0.0789	0.1764	0.3688	0.7221	1.3266	2.2935	3.7418	5.78	8.4844	5.78	3.7418	2.2935	1.3266	0.722	0.3688	0.1764	0.0788	0.0329	0.0128
	2nd	-0.107	-0.172	-0.231	-0.227	-0.059	0.4214	1.3878	3	5.3603	8.4844	6.1996	4.4836	3.1992	2.2318	1.5032	0.9648	0.5835	0.3297	0.173	0.084
	3rd	0.2151	0.2758	0.3296	0.4068	0.5957	1.0506	1.9676	3.5318	5.8608	8.9744	6.7001	5.0153	3.7789	2.861	2.1579	1.5987	1.1439	0.7775	0.4954	0.2929
	4th	-0.197	-0.142	-0.04	0.1192	0.3925	0.9108	1.8666	3.4573	5.8147	8.9631	6.7246	5.0714	3.8655	2.99	2.3531	1.8798	1.5075	1.1897	0.9026	0.6444
	Diff.																				
A.E.	1st	0.0308	0.0647	0.1238	0.2154	0.3324	0.4368	0.4576	0.3236	0.0133	-0.424	-0.885	-1.267	-1.51	-1.603	-1.571	-1.455	-1.298	-1.131	-0.975	-0.837
	2nd	-0.109	-0.186	-0.283	-0.38	-0.449	-0.468	-0.448	-0.418	-0.406	-0.424	-0.465	-0.525	-0.605	-0.698	-0.79	-0.859	-0.891	-0.88	-0.834	-0.766
	3rd	0.213	0.2616	0.277	0.2534	0.206	0.1608	0.1317	0.1136	0.0941	0.0661	0.0354	0.0068	-0.025	-0.069	-0.135	-0.225	-0.33	-0.433	-0.512	-0.557
	4th	-0.199	-0.156	-0.093	-0.034	0.0028	0.021	0.0307	0.0391	0.048	0.0548	0.0599	0.0629	0.0617	0.0602	0.0599	0.0559	0.0333	-0.02	-0.105	-0.206

Case (iii)		Corr.1																			
Moneyness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0031	0.0205	0.0776	0.2271	0.5707	1.2768	2.5725	4.6977	7.8173	11.978	8.9033	6.6303	4.9801	3.7907	2.9308	2.3056	1.8445	1.5008	1.2392	1.0365
A.E.	1st	0.0442	0.1059	0.2369	0.4953	0.9697	1.7815	3.08	5.0249	7.7621	11.394	7.762	5.0249	3.08	1.7815	0.9696	0.4952	0.2369	0.1058	0.0441	0.0172
	2nd	-0.133	-0.211	-0.277	-0.257	-0.017	0.6386	1.9365	4.0883	7.2322	11.394	8.2919	5.9615	4.2235	2.9244	1.9559	1.2477	0.7509	0.4226	0.2211	0.1071
	3rd	0.2663	0.348	0.4292	0.5517	0.8304	1.4644	2.707	4.8005	7.9041	12.052	8.9638	6.6737	4.994	3.7502	2.8029	2.0566	1.4573	0.9815	0.6202	0.364
	4th	-0.249	-0.182	-0.044	0.1866	0.5865	1.3198	2.6254	4.7519	7.8739	12.037	8.9651	6.6975	5.0565	3.8811	3.0373	2.4145	1.9246	1.5062	1.1312	0.7989
	Diff.																				
A.E.	1st	0.0411	0.0854	0.1593	0.2682	0.399	0.5047	0.5075	0.3272	-0.055	-0.584	-1.141	-1.605	-1.9	-2.009	-1.961	-1.81	-1.608	-1.395	-1.195	-1.019
	2nd	-0.136	-0.231	-0.355	-0.484	-0.587	-0.638	-0.636	-0.609	-0.585	-0.584	-0.611	-0.669	-0.757	-0.866	-0.975	-1.058	-1.094	-1.078	-1.018	-0.929
	3rd	0.2632	0.3275	0.3516	0.3246	0.2597	0.1876	0.1345	0.1028	0.0868	0.0738	0.0605	0.0434	0.0139	-0.041	-0.128	-0.249	-0.387	-0.519	-0.619	-0.673
	4th	-0.252	-0.203	-0.122	-0.041	0.0158	0.043	0.0529	0.0542	0.0566	0.0586	0.0618	0.0672	0.0764	0.0904	0.1065	0.1089	0.0801	0.0054	-0.108	-0.238

Case (iv)		Corr.1																			
Moneyness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0032	0.0202	0.0771	0.2279	0.5732	1.2829	2.5829	4.7165	7.8575	12.047	9.0009	6.7503	5.1155	3.9308	3.069	2.4361	1.9648	1.6092	1.3368	1.1245
A.E.	1st	0.0447	0.107	0.2389	0.4987	0.975	1.7892	3.0902	5.0375	7.7763	11.409	7.7763	5.0375	3.0902	1.7892	0.9749	0.4986	0.2389	0.1069	0.0447	0.0174
	2nd	-0.14	-0.223	-0.295	-0.282	-0.047	0.6064	1.9077	4.0695	7.2288	11.409	8.3237	6.0056	4.2727	2.972	1.9966	1.279	0.7727	0.4364	0.2291	0.1113
	3rd	0.2864	0.3732	0.4574	0.5802	0.8582	1.4935	2.743	4.8501	7.9726	12.14	9.0675	6.7862	5.108	3.8592	2.9016	2.1409	1.5251	1.0322	0.6552	0.3862
	4th	-0.271	-0.199	-0.056	0.1764	0.5752	1.3079	2.6203	4.7667	7.9215	12.124	9.0872	6.8426	5.2092	4.0286	3.1724	2.5351	2.0306	1.5973	1.2061	0.8566
	Diff.																				
A.E.	1st	0.0415	0.0868	0.1618	0.2708	0.4018	0.5063	0.5073	0.321	-0.081	-0.638	-1.225	-1.713	-2.025	-2.142	-2.094	-1.938	-1.726	-1.502	-1.292	-1.107
	2nd	-0.143	-0.243	-0.372	-0.51	-0.62	-0.677	-0.675	-0.647	-0.629	-0.638	-0.677	-0.745	-0.843	-0.959	-1.072	-1.157	-1.192	-1.173	-1.108	-1.013
	3rd	0.2832	0.353	0.3803	0.3523	0.285	0.2106	0.1601	0.1386	0.1151	0.093	0.0666	0.0359	-0.008	-0.072	-0.167	-0.295	-0.44	-0.577	-0.682	-0.738
	4th	-0.274	-0.219	-0.133	-0.051	0.002	0.025	0.0374	0.0502	0.064	0.0766	0.0863	0.0923	0.0937	0.0978	0.1034	0.099	0.0658	-0.012	-0.131	-0.268

Table 7

Case (i)		Corr.2																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0065	0.0366	0.1137	0.2784	0.5955	1.1579	2.0956	3.5506	5.6542	8.4874	5.9496	4.0731	2.7427	1.8319	1.2232	0.8223	0.5598	0.3875	0.2736	0.1975
A.E.	1st	0.0325	0.0781	0.1749	0.3663	0.7181	1.3209	2.2859	3.7325	5.7694	8.4733	5.7694	3.7324	2.2859	1.3209	0.7181	0.3663	0.1749	0.0781	0.0325	0.0126
	2nd	-8E-04	0.0182	0.0777	0.2238	0.5311	1.104	2.0687	3.5545	5.6686	8.4733	5.8701	3.9104	2.5031	1.5378	0.9051	0.5088	0.2722	0.1378	0.0658	0.0295
	3rd	0.0853	0.1526	0.266	0.4596	0.7925	1.3576	2.2812	3.7078	5.7701	8.5543	5.9716	4.0638	2.7156	1.7914	1.1665	0.7446	0.4605	0.2721	0.152	0.0794
	4th	0.0149	0.0627	0.1598	0.3408	0.665	1.2264	2.1559	3.6041	5.7064	8.5437	6.0151	4.1499	2.8269	1.912	1.2862	0.8576	0.5625	0.3588	0.2201	0.1282
	Diff.																				
A.E.	1st	0.026	0.0415	0.0612	0.0879	0.1226	0.163	0.1903	0.1819	0.1152	-0.014	-0.18	-0.341	-0.457	-0.511	-0.505	-0.456	-0.385	-0.31	-0.241	-0.185
	2nd	-0.007	-0.018	-0.036	-0.055	-0.064	-0.054	-0.027	0.0039	0.0144	-0.014	-0.08	-0.163	-0.24	-0.294	-0.318	-0.314	-0.288	-0.25	-0.208	-0.168
	3rd	0.0788	0.116	0.1523	0.1812	0.197	0.1997	0.1856	0.1572	0.1159	0.0669	0.022	-0.009	-0.027	-0.041	-0.057	-0.078	-0.099	-0.115	-0.122	-0.118
	4th	0.0084	0.0261	0.0461	0.0624	0.0695	0.0685	0.0603	0.0535	0.0522	0.0563	0.0655	0.0768	0.0842	0.0801	0.063	0.0353	0.0027	-0.029	-0.054	-0.069
Case (ii)		Corr.2																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0063	0.0361	0.114	0.2801	0.5966	1.1597	2.0998	3.5616	5.6763	8.5259	6.0095	4.1544	2.8405	1.9379	1.3285	0.9215	0.6489	0.4641	0.3376	0.2497
A.E.	1st	0.0329	0.0789	0.1764	0.3688	0.7221	1.3266	2.2935	3.7418	5.78	8.4844	5.78	3.7418	2.2935	1.3266	0.722	0.3688	0.1764	0.0788	0.0329	0.0128
	2nd	-0.005	0.0101	0.0648	0.2055	0.508	1.0786	2.0453	3.5385	5.665	8.4844	5.895	3.9451	2.5417	1.5747	0.9361	0.5321	0.288	0.1475	0.0713	0.0323
	3rd	0.0983	0.1716	0.291	0.489	0.8238	1.3882	2.3105	3.7381	5.8067	8.6031	6.0367	4.1447	2.8069	1.8844	1.2519	0.8157	0.5142	0.309	0.1751	0.0926
	4th	-0.002	0.0439	0.1412	0.3246	0.6523	1.2179	2.1532	3.6116	5.7307	8.5918	6.091	4.2521	2.9482	2.042	1.4134	0.9722	0.6579	0.4322	0.2721	0.1621
	Diff.																				
A.E.	1st	0.0266	0.0428	0.0624	0.0887	0.1255	0.1669	0.1937	0.1802	0.1037	-0.041	-0.23	-0.413	-0.547	-0.611	-0.607	-0.553	-0.473	-0.385	-0.305	-0.237
	2nd	-0.012	-0.026	-0.049	-0.075	-0.089	-0.081	-0.055	-0.023	-0.011	-0.041	-0.115	-0.209	-0.299	-0.363	-0.392	-0.389	-0.361	-0.317	-0.266	-0.217
	3rd	0.092	0.1355	0.177	0.2089	0.2272	0.2285	0.2107	0.1765	0.1304	0.0772	0.0272	-0.01	-0.034	-0.053	-0.077	-0.106	-0.135	-0.155	-0.163	-0.157
	4th	-0.008	0.0078	0.0272	0.0445	0.0557	0.0582	0.0534	0.05	0.0544	0.0659	0.0815	0.0977	0.1077	0.1041	0.0849	0.0507	0.009	-0.032	-0.066	-0.088
Case (iii)		Corr.2																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0092	0.0513	0.1611	0.3953	0.8444	1.6351	2.93	4.9079	7.7376	11.521	8.0786	5.5251	3.7142	2.476	1.6485	1.1038	0.7472	0.5144	0.361	0.2583
A.E.	1st	0.0442	0.1059	0.2369	0.4953	0.9697	1.7815	3.08	5.0249	7.7621	11.394	7.762	5.0249	3.08	1.7815	0.9696	0.4952	0.2369	0.1058	0.0441	0.0172
	2nd	0.0039	0.0337	0.1198	0.3239	0.7449	1.521	2.8194	4.8115	7.6683	11.394	7.8828	5.2384	3.3406	2.042	1.1944	0.6667	0.354	0.178	0.0845	0.0377
	3rd	0.1137	0.2071	0.3669	0.6403	1.1075	1.8916	3.1576	5.0929	7.87	11.601	8.1114	5.5198	3.6788	2.4125	1.557	0.9832	0.6011	0.3514	0.1943	0.1007
	4th	0.0182	0.0824	0.2172	0.473	0.9317	1.7175	2.9984	4.9654	7.792	11.585	8.159	5.6209	3.817	2.571	1.7216	1.1425	0.7452	0.4722	0.2872	0.1656
	Diff.																				
A.E.	1st	0.035	0.0546	0.0758	0.1	0.1253	0.1464	0.15	0.117	0.0245	-0.127	-0.317	-0.5	-0.634	-0.695	-0.679	-0.609	-0.51	-0.409	-0.317	-0.241
	2nd	-0.005	-0.018	-0.041	-0.071	-0.099	-0.114	-0.111	-0.096	-0.069	-0.127	-0.196	-0.287	-0.374	-0.434	-0.454	-0.437	-0.393	-0.336	-0.277	-0.221
	3rd	0.1045	0.1558	0.2058	0.245	0.2631	0.2565	0.2276	0.185	0.1324	0.0801	0.0328	-0.005	-0.035	-0.063	-0.092	-0.121	-0.146	-0.163	-0.167	-0.158
	4th	0.009	0.0311	0.0561	0.0777	0.0873	0.0824	0.0684	0.0575	0.0544	0.0641	0.0804	0.0958	0.1028	0.095	0.0731	0.0387	-0.002	-0.042	-0.074	-0.093
Case (iv)		Corr.2																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0087	0.05	0.1589	0.3929	0.8411	1.6312	2.9244	4.9046	7.7428	11.548	8.136	5.6174	3.8329	2.6083	1.7833	1.233	0.8668	0.6217	0.4539	0.3375
A.E.	1st	0.0447	0.107	0.2389	0.4987	0.975	1.7892	3.0902	5.0375	7.7763	11.409	7.7763	5.0375	3.0902	1.7892	0.9749	0.4986	0.2389	0.1069	0.0447	0.0174
	2nd	-0.002	0.0227	0.1025	0.2993	0.7138	1.4869	2.788	4.7901	7.6364	11.409	7.9162	5.2849	3.3924	2.0915	1.236	0.6981	0.3753	0.1911	0.0918	0.0414
	3rd	0.1312	0.2328	0.4007	0.6803	1.1502	1.9332	3.197	5.1332	7.9181	11.666	8.1979	5.628	3.8015	2.5378	1.6724	1.0791	0.6736	0.4011	0.2254	0.1184
	4th	-0.005	0.0566	0.1915	0.4506	0.9145	1.7067	2.996	4.9766	7.825	11.649	8.259	5.7561	3.9789	2.7457	1.8937	1.2981	0.8747	0.5714	0.3572	0.2109
	Diff.																				
A.E.	1st	0.036	0.057	0.08	0.1058	0.1339	0.158	0.1658	0.1329	0.0335	-0.14	-0.36	-0.58	-0.743	-0.819	-0.808	-0.734	-0.628	-0.515	-0.409	-0.32
	2nd	-0.011	-0.027	-0.056	-0.094	-0.127	-0.144	-0.136	-0.115	-0.106	-0.14	-0.22	-0.333	-0.441	-0.517	-0.547	-0.535	-0.492	-0.431	-0.362	-0.296
	3rd	0.1225	0.1828	0.2418	0.2874	0.3091	0.302	0.2726	0.2286	0.1753	0.1174	0.0619	0.0106	-0.031	-0.071	-0.111	-0.154	-0.193	-0.221	-0.229	-0.219
	4th	-0.013	0.0066	0.0326	0.0577	0.0734	0.0755	0.0716	0.072	0.0822	0.1007	0.123	0.1387	0.146	0.1374	0.1104	0.0651	0.0079	-0.05	-0.097	-0.127

Table 8

Case (i)		Corr.3																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0027	0.0209	0.0845	0.2576	0.6535	1.4218	2.7132	4.6397	7.2473	10.515	8.2679	6.5157	5.1621	4.12	3.3181	2.6992	2.2193	1.8439	1.5475	1.3115
A.E.	1st	0.148	0.2812	0.5103	0.8852	1.4704	2.3415	3.5808	5.268	7.4705	10.234	7.4705	5.268	3.5807	2.3415	1.4703	0.8852	0.5102	0.2812	0.1478	0.0742
	2nd	-0.317	-0.388	-0.385	-0.223	0.2094	1.0401	2.3909	4.3541	6.9731	10.234	7.9679	6.1818	4.7706	3.6429	2.7313	1.9936	1.4053	0.9503	0.6131	0.3762
	3rd	0.3084	0.294	0.2991	0.4108	0.7602	1.5013	2.7796	4.6979	7.2963	10.552	8.2911	6.5256	5.1594	4.1041	3.2821	2.6276	2.0892	1.6323	1.2388	0.9042
	4th	-0.073	0.0023	0.097	0.2787	0.6712	1.4339	2.7229	4.6503	7.2603	10.53	8.2853	6.5353	5.1839	4.1458	3.3509	2.7436	2.2774	1.9111	1.6074	1.3374
	Diff.																				
A.E.	1st	0.1453	0.2603	0.4258	0.6276	0.8169	0.9197	0.8676	0.6283	0.2232	-0.281	-0.797	-1.248	-1.581	-1.779	-1.848	-1.814	-1.709	-1.563	-1.4	-1.237
	2nd	-0.32	-0.409	-0.469	-0.481	-0.444	-0.382	-0.322	-0.286	-0.274	-0.281	-0.3	-0.334	-0.392	-0.477	-0.587	-0.706	-0.814	-0.894	-0.934	-0.935
	3rd	0.3057	0.2731	0.2146	0.1532	0.1067	0.0795	0.0664	0.0582	0.049	0.037	0.0232	0.0099	-0.003	-0.016	-0.036	-0.072	-0.13	-0.212	-0.309	-0.407
	4th	-0.076	-0.019	0.0125	0.0211	0.0177	0.0121	0.0097	0.0106	0.013	0.0154	0.0174	0.0196	0.0218	0.0258	0.0328	0.0444	0.0581	0.0672	0.0599	0.0259

Case (ii)		Corr.3																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0027	0.0201	0.0821	0.2509	0.6367	1.3919	2.6766	4.6112	7.2423	10.541	8.3293	6.6101	5.2844	4.2629	3.4733	2.8596	2.3782	1.9985	1.6961	1.4529
A.E.	1st	0.1501	0.2846	0.5155	0.8928	1.4808	2.3549	3.5971	5.2868	7.491	10.255	7.491	5.2868	3.5971	2.3549	1.4807	0.8928	0.5155	0.2846	0.15	0.0755
	2nd	-0.358	-0.445	-0.458	-0.311	0.1128	0.9446	2.3087	4.2978	6.9529	10.255	8.0291	6.2757	4.8856	3.7653	2.8487	2.0968	1.4893	1.0138	0.658	0.4059
	3rd	0.3816	0.3594	0.3452	0.4292	0.7493	1.4682	2.7385	4.6669	7.292	10.585	8.3682	6.6448	5.3154	4.2889	3.4853	2.8372	2.2927	1.8178	1.3975	1.0313
	4th	-0.085	0.0123	0.1162	0.2903	0.6627	1.4037	2.6816	4.6166	7.2535	10.564	8.3645	6.6566	5.339	4.3257	3.5492	2.9569	2.504	2.148	1.8477	1.5703
	Diff.																				
A.E.	1st	0.1474	0.2645	0.4334	0.6419	0.8441	0.963	0.9205	0.6756	0.2487	-0.287	-0.838	-1.323	-1.687	-1.908	-1.993	-1.967	-1.863	-1.714	-1.546	-1.377
	2nd	-0.361	-0.465	-0.54	-0.562	-0.524	-0.447	-0.368	-0.313	-0.289	-0.287	-0.3	-0.334	-0.399	-0.498	-0.625	-0.763	-0.889	-0.985	-1.038	-1.047
	3rd	0.3789	0.3393	0.2631	0.1783	0.1126	0.0763	0.0619	0.0557	0.0497	0.0442	0.0389	0.0347	0.031	0.026	0.012	-0.022	-0.086	-0.181	-0.299	-0.422
	4th	-0.088	-0.008	0.0341	0.0394	0.026	0.0118	0.005	0.0054	0.0112	0.0224	0.0352	0.0465	0.0546	0.0628	0.0759	0.0973	0.1258	0.1495	0.1516	0.1174

Case (iii)		Corr.3																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0049	0.0374	0.1471	0.424	1.0083	2.0787	3.8183	6.3764	9.8256	14.154	11.085	8.6932	6.8513	5.4403	4.3571	3.5245	2.8796	2.3778	1.9844	1.6731
A.E.	1st	0.2016	0.3822	0.6924	1.199	1.9886	3.1624	4.8306	7.0997	10.06	13.771	10.06	7.0997	4.831	3.1624	1.9885	1.199	0.6923	0.3822	0.2014	0.1014
	2nd	-0.38	-0.453	-0.423	-0.18	0.422	1.5473	3.3552	5.9672	9.4436	13.771	10.676	8.2322	6.3061	4.7775	3.5551	2.5778	1.8047	1.2173	0.7832	0.4798
	3rd	0.4065	0.4188	0.47	0.6696	1.1815	2.2002	3.9131	6.4584	9.8987	14.215	11.131	8.7235	6.864	5.4304	4.3146	3.4271	2.7001	2.0889	1.5699	1.1353
	4th	-0.137	-0.023	0.1426	0.4431	1.0283	2.0924	3.8318	6.3952	9.8518	14.186	11.121	8.7349	6.9007	5.5019	4.4392	3.6309	3.0086	2.5136	2.0971	1.7249
	Diff.																				
A.E.	1st	0.1967	0.3448	0.5453	0.775	0.9803	1.0837	1.0123	0.7233	0.2343	-0.383	-1.025	-1.594	-2.02	-2.278	-2.369	-2.326	-2.187	-1.996	-1.783	-1.572
	2nd	-0.385	-0.49	-0.57	-0.604	-0.586	-0.531	-0.463	-0.409	-0.382	-0.383	-0.409	-0.461	-0.545	-0.663	-0.802	-0.947	-1.075	-1.161	-1.201	-1.193
	3rd	0.4016	0.3814	0.3229	0.2456	0.1732	0.1215	0.0948	0.082	0.0731	0.0612	0.0466	0.0303	0.0127	-0.01	-0.042	-0.097	-0.18	-0.289	-0.415	-0.538
	4th	-0.142	-0.06	-0.005	0.0191	0.02	0.0137	0.0135	0.0188	0.0262	0.0319	0.0367	0.0417	0.0494	0.0616	0.0821	0.1064	0.129	0.1358	0.1127	0.0518

Case (iv)		Corr.3																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0048	0.0361	0.1429	0.4129	0.9836	2.0364	3.7688	6.3345	9.8133	14.189	11.174	8.8387	7.0468	5.6752	4.6197	3.8027	3.1649	2.6627	2.2626	1.9413
A.E.	1st	0.2045	0.3869	0.6995	1.2093	2.0025	3.1804	4.8526	7.125	10.087	13.8	10.087	7.125	4.8525	3.1804	2.0025	1.2093	0.6994	0.3869	0.2043	0.103
	2nd	-0.435	-0.529	-0.521	-0.298	0.2923	1.419	3.2447	5.8916	9.4164	13.8	10.758	8.3584	6.4604	4.9418	3.7127	2.7164	1.9202	1.3026	0.8436	0.5198
	3rd	0.5061	0.51	0.5373	0.7009	1.173	2.1592	3.8572	6.4112	9.8835	14.25	11.225	8.878	7.0728	5.682	4.5934	3.7151	2.9787	2.3415	1.7844	1.306
	4th	-0.16	-0.014	0.1678	0.4623	1.0231	2.0574	3.7778	6.3458	9.8339	14.221	11.218	8.8909	7.1061	5.745	4.7111	3.9264	3.324	2.8427	2.4288	2.0431
	Diff.																				
A.E.	1st	0.1997	0.3508	0.5566	0.7964	1.0189	1.144	1.0838	0.7905	0.274	-0.389	-1.087	-1.714	-2.194	-2.495	-2.617	-2.593	-2.466	-2.276	-2.058	-1.838
	2nd	-0.44	-0.565	-0.664	-0.711	-0.691	-0.617	-0.524	-0.443	-0.397	-0.389	-0.416	-0.48	-0.586	-0.733	-0.907	-1.086	-1.245	-1.36	-1.419	-1.422
	3rd	0.5013	0.4739	0.3944	0.288	0.1894	0.1228	0.0884	0.0767	0.0702	0.0613	0.0513	0.0393	0.026	0.0068	-0.026	-0.088	-0.186	-0.321	-0.478	-0.635
	4th	-0.165	-0.05	0.0249	0.0494	0.0395	0.021	0.009	0.0113	0.0206	0.032	0.0439	0.0522	0.0593	0.0698	0.0914	0.1237	0.1591	0.18	0.1662	0.1018

Table 9


Case (i)		Corr.4																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.008	0.0478	0.1569	0.3962	0.8563	1.6506	2.8984	4.7016	7.1253	10.182	7.7353	5.8173	4.3474	3.2373	2.4089	1.7966	1.3448	1.0114	0.7649	0.5809
A.E.	1st	0.0008	0.0038	0.0148	0.0504	0.1506	0.3971	0.9292	1.9422	3.6525	6.2344	3.6525	1.9422	0.9292	0.397	0.1505	0.0503	0.0148	0.0038	0.0009	0.0001
	2nd	-0.005	-0.015	-0.037	-0.07	-0.081	0.0289	0.4584	1.4826	3.3637	6.2344	3.9414	2.4017	1.4	0.7651	0.3822	0.1705	0.0668	0.0228	0.0067	0.0017
	3rd	0.1517	0.2005	0.3075	0.5388	0.9896	1.7746	3.0122	4.802	7.2072	10.244	7.7815	5.8571	4.386	3.2773	2.4455	1.8187	1.341	0.9731	0.6888	0.4719
	4th	0.0085	0.0794	0.2113	0.4622	0.923	1.7101	2.9479	4.7434	7.1633	10.222	7.783	5.8772	4.4173	3.3149	2.4906	1.878	1.423	1.0821	0.8215	0.6177
	Diff.																				
A.E.	1st	-0.007	-0.044	-0.142	-0.346	-0.706	-1.254	-1.969	-2.759	-3.473	-3.948	-4.083	-3.875	-3.418	-2.84	-2.258	-1.746	-1.33	-1.008	-0.764	-0.581
	2nd	-0.013	-0.063	-0.194	-0.466	-0.937	-1.622	-2.44	-3.219	-3.762	-3.948	-3.794	-3.416	-2.947	-2.472	-2.027	-1.626	-1.278	-0.989	-0.758	-0.579
	3rd	0.1437	0.1527	0.1506	0.1426	0.1333	0.124	0.1138	0.1004	0.0819	0.0622	0.0462	0.0398	0.0386	0.04	0.0366	0.0221	-0.004	-0.038	-0.076	-0.109
	4th	0.0005	0.0316	0.0544	0.066	0.0667	0.0595	0.0495	0.0418	0.038	0.0403	0.0477	0.0599	0.0699	0.0776	0.0817	0.0814	0.0782	0.0707	0.0566	0.0368

Case (ii)		Corr.4																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0081	0.0469	0.1531	0.387	0.8384	1.6206	2.8589	4.6608	7.0958	10.181	7.7743	5.9033	4.4778	3.4072	2.6084	2.0139	1.5708	1.2384	0.9868	0.7954
A.E.	1st	0.0008	0.0038	0.0147	0.0501	0.1499	0.3957	0.9269	1.9388	3.6483	6.2299	3.6483	1.9388	0.9269	0.3956	0.1498	0.005	0.0147	0.0038	0.0008	0.0001
	2nd	-0.005	-0.015	-0.038	-0.071	-0.085	0.0225	0.4491	1.4722	3.3549	6.2299	3.9416	2.4054	1.4046	0.7689	0.3845	0.1716	0.0673	0.0229	0.0067	0.0017
	3rd	0.2233	0.2721	0.3683	0.5781	1.0005	1.7572	2.9748	4.7605	7.1813	10.253	7.8391	5.9693	4.5497	3.4811	2.6726	2.0499	1.5586	1.1635	0.8442	0.5905
	4th	-0.029	0.0612	0.2091	0.4663	0.9216	1.6949	2.919	4.7102	7.1422	10.231	7.8355	5.9805	4.5715	3.5145	2.7272	2.1409	1.6994	1.3576	1.0816	0.849
	Diff.																				
A.E.	1st	-0.007	-0.043	-0.138	-0.337	-0.689	-1.225	-1.932	-2.722	-3.448	-3.951	-4.126	-3.965	-3.551	-3.012	-2.459	-2.009	-1.556	-1.235	-0.986	-0.795
	2nd	-0.013	-0.062	-0.191	-0.459	-0.923	-1.598	-2.41	-3.189	-3.741	-3.951	-3.833	-3.498	-3.073	-2.638	-2.224	-1.842	-1.504	-1.216	-0.98	-0.794
	3rd	0.2152	0.2252	0.2152	0.1911	0.1621	0.1366	0.1159	0.0997	0.0855	0.0723	0.0648	0.066	0.0719	0.0739	0.0642	0.036	-0.012	-0.075	-0.143	-0.205
	4th	-0.037	0.0143	0.056	0.0793	0.0832	0.0743	0.0601	0.0494	0.0464	0.0503	0.0612	0.0772	0.0937	0.1073	0.1188	0.127	0.1286	0.1192	0.0948	0.0536

Case (iii)		Corr.4																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0127	0.076	0.2452	0.6053	1.2708	2.3779	4.0731	6.484	9.7001	13.751	10.409	7.789	5.7858	4.2827	3.1695	2.3523	1.7541	1.3161	0.9951	0.7588
A.E.	1st	0.0011	0.0051	0.0198	0.0673	0.2013	0.5313	1.2447	2.6036	4.8994	8.3662	4.8993	2.6036	1.2447	0.5313	0.2012	0.0672	0.0197	0.0051	0.0011	0.0002
	2nd	-0.007	-0.02	-0.049	-0.091	-0.104	0.0464	0.624	1.9974	4.5182	8.3662	5.2805	3.2099	1.8654	1.0163	0.5061	0.2252	0.088	0.0299	0.0088	0.0022
	3rd	0.1987	0.287	0.4639	0.8148	1.4597	2.5435	4.2166	6.6089	9.8067	13.838	10.477	7.8412	5.8221	4.3008	3.1642	2.315	1.6771	1.1953	0.8315	0.5605
	4th	-0.015	0.089	0.2934	0.6755	1.3486	2.4543	4.1435	6.5492	9.7608	13.808	10.465	7.848	5.8496	4.3525	3.2453	2.4306	1.8286	1.3775	1.0322	0.7629
	Diff.																				
A.E.	1st	-0.012	-0.071	-0.225	-0.538	-1.07	-1.847	-2.828	-3.88	-4.801	-5.385	-5.509	-5.185	-4.541	-3.751	-2.968	-2.285	-1.734	-1.311	-0.994	-0.759
	2nd	-0.019	-0.096	-0.294	-0.696	-1.374	-2.332	-3.449	-4.487	-5.182	-5.385	-5.128	-4.579	-3.92	-3.266	-2.663	-2.127	-1.666	-1.286	-0.986	-0.757
	3rd	0.186	0.211	0.2187	0.2095	0.1889	0.1656	0.1435	0.1249	0.1066	0.0873	0.0686	0.0522	0.0363	0.0181	-0.005	-0.037	-0.077	-0.121	-0.164	-0.198
	4th	-0.027	0.013	0.0482	0.0702	0.0778	0.0764	0.0704	0.0652	0.0607	0.0573	0.0564	0.059	0.0638	0.0698	0.0758	0.0783	0.0745	0.0614	0.0371	0.0041

Case (iv)		Corr.4																			
Moneyiness		0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9	1	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2
	MC	0.0123	0.0741	0.2413	0.596	1.2523	2.3474	4.0298	6.4348	9.66	13.74	10.447	7.8881	5.9458	4.4945	3.4194	2.6239	2.0349	1.596	1.266	1.0148
A.E.	1st	0.0011	0.005	0.0196	0.0669	0.2004	0.5295	1.2416	2.5991	4.8937	8.36	4.8936	2.5991	1.2416	0.5295	0.2003	0.0668	0.0196	0.005	0.0011	0.0002
	2nd	-0.007	-0.02	-0.049	-0.093	-0.109	0.0377	0.6116	1.9834	4.5064	8.36	5.2808	3.2148	1.8717	1.0213	0.5092	0.2266	0.0886	0.0301	0.0088	0.0022
	3rd	0.2962	0.3862	0.5509	0.8741	1.4804	2.5236	4.1657	6.5476	9.7624	13.839	10.545	7.9863	6.0411	4.5781	3.4752	2.632	1.9747	1.4544	1.0416	0.7198
	4th	-0.072	0.0576	0.2856	0.6804	1.3509	2.441	4.1112	6.5068	9.7281	13.809	10.521	7.9736	6.0486	4.6204	3.5707	2.7969	2.215	1.7612	1.3908	1.0778
	Diff.																				
A.E.	1st	-0.011	-0.069	-0.222	-0.529	-1.052	-1.818	-2.788	-3.836	-4.766	-5.38	-5.553	-5.289	-4.704	-3.965	-3.219	-2.557	-2.015	-1.591	-1.265	-1.015
	2nd	-0.019	-0.094	-0.291	-0.689	-1.361	-2.31	-3.418	-4.451	-5.154	-5.38	-5.166	-4.673	-4.074	-3.473	-2.91	-2.397	-1.946	-1.566	-1.257	-1.013
	3rd	0.2839	0.3121	0.3096	0.2781	0.2281	0.1762	0.1359	0.1128	0.1024	0.0988	0.0981	0.0982	0.0953	0.0836	0.0558	0.0081	-0.06	-0.142	-0.224	-0.295
	4th	-0.084	-0.016	0.0443	0.0844	0.0986	0.0936	0.0814	0.072	0.0681	0.069	0.0746	0.0855	0.1028	0.1259	0.1513	0.173	0.1801	0.1652	0.1248	0.063

Figure 1



References

- [1] Bichteler, K., Gravereaux, J.-B. and Jacod, J. [1987], Malliavin Calculus for Processes with Jumps, volume 2 of *Stochastic Monographs*, Gordon and Breach Science Publishers.
- [2] Björk, T. [2004] “Arbitrage Theory in Continuous Time(second ed.)”, Oxford University Press, 2004.
- [3] Carr, P., Jarrow, R. and Myneni, R. [1992], “Alternative characterizations of American put options,” *Mathematical Finance*, Vol. 2, 87-106.
- [4] Dermoune A. and Kutoyants, Y. [1995], “Expansion of distribution of Maximum Likelihood Estimate for Misspecified Diffusion Type Observation,” *Stochastics and Stochastic Reports*, Vol. 52, no. 1-2, 121-145.
- [5] Fouque, J.-P., Papanicolaou, G. and Sircar, K. R.[1999], “Financial Modeling in a Fast Mean-reverting Stochastic Volatility Environment, ” *Asia-Pacific Financial Markets*, Vol. 6(1), pp.37-48.
- [6] Fouque, J.-P., Papanicolaou, G. and Sircar, K. R.[2000], *Derivatives in financial Markets with Stochastic Volatility*, Cambridge University Press.
- [7] Heath, D. Jarrow, R. and Morton, A. [1992], “Bond Pricing and the Term Structure of Interest Rates : A New Methodology for Contingent Claims Valuation,” *Econometrica*, Vol. 60, 77-105.
- [8] Ikeda, N. and Watanabe, S. [1989], *Stochastic Differential Equations and Diffusion Processes*, Second Edition, North-Holland/Kodansha, Tokyo.
- [9] Karatzas, I. and Shreve, S. [1998], “Methods of Mathematical Finance,” Springer.
- [10] Kashiwakura, K. and Yoshida, N. [2001], “Numerical Experiment for Hybrid Expansion,” Preprint, Graduate School of Mathematical Sciences, the University of Tokyo.
- [11] Kawai, A. [2003], “A New Approximate Swaption Formula in the LIBOR Market Model: An Asymptotic Expansion Approach,” *Applied Mathematical Finance*, Vol.10, 49-74.
- [12] Kawai, A. and Jäckel, P. [2007], ” An Asymptotic FX Option Formula in the Cross Currency Libor Market Model,” *Willmot Magazine*, March, 74-84.
- [13] Kobayashi, T., Takahashi, A. and Tokioka, N. [2005], “Dynamic Optimality of Yield Curve Strategies”, *International Review of Finance*, Vol.4, 49-78.
- [14] Kunitomo, N. and Kim, Y-J.[2007], “Effects of Stochastic Interest Rates and Volatility on Contingent Claims,” *Japanese Economic Review*, Vol.58, 71-106.
- [15] Kunitomo, N. and Takahashi, A. [1992], “Pricing Average Options,” *Japan Financial Review*, Vol. 14, 1-20. (in Japanese).
- [16] Kunitomo, N. and Takahashi, A. [2001], “The Asymptotic Expansion Approach to the Valuation of Interest Rate Contingent Claims,” *Mathematical Finance*, Vol. 11, 117-151.
- [17] Kunitomo, N. and Takahashi, A. [2003a], “On Validity of the Asymptotic Expansion Approach in Contingent Claim Analysis,” *Annals of Applied Probability* Vol. 13-3, 914-952.

- [18] Kunitomo, N. and Takahashi, A. [2003b], "Foundation of Mathematical Finance -Application of Malliavin Calculus and Asymptotic Expansion Method-," Toyo-Keizai (in Japanese).
- [19] Kunitomo, N. and Takahashi, A. [2004], "Applications of the Asymptotic Expansion Approach based on Malliavin-Watanabe Calculus in Financial Problems," *Stochastic Processes and Applications to Mathematical Finance*, World Scientific, 195-232.
- [20] Kusuoka S. and Osajima, Y. [2007], "A Remark on the Asymptotic Expansion of Density Function of Wiener Functionals," Preprint, Graduate School of Mathematical Sciences, the University of Tokyo.
- [21] Kusuoka S. and Strook, D. [1991], "Precise Asymptotics of Certain Wiener Functionals," *Journal of Functional Analysis*, Vol.99, 1-74.
- [22] Kutoyants, Y. [1994], "Identification of Dynamical Systems with Small Noise," *Mathematics and its Applications*, 300, Kluwer Academic Publishers Group.
- [23] Kutoyants, Y. [1998], "Semiparametric Estimation for Dynamic Systems with Small Noise," *Mathematical Methods of Statistics*, 7, no. 4, 457-465.
- [24] Labordere, P.H. [2005a], A General Asymptotic Implied Volatility for Stochastic Volatility Models, cond-mat/0504317.
- [25] Labordere, P.H. [2005b], "Solvable Local and Stochastic Volatility Models: Supersymmetric Methods in Option Pricing," Working Paper.
- [26] Labordere, P.H. [2006], "Unifying the BGM and SABR Models: A short Ride in Hyperbolic Geometry," Working Paper.
- [27] Lütkebohmert, K. [2004], "An Asymptotic Expansion for a Black-Scholes Type Model," *Bull. Sci. math.*, 128, 661-685.
- [28] Malliavin, P. [1997], "Stochastic Analysis," Springer.
- [29] Malliavin, P. and Thalmaier, A. [2006], *Stochastic Calculus of Variations in Mathematical Finance*, Springer.
- [30] Masuda, H. and Yoshida, N. [2004], "An Application of the Double Edgeworth Expansion to a Filtering Model with Gaussian Limit," *Statistics and Probability Letters*, 70, no. 1, 37-48.
- [31] Matsuoka, R. Takahashi, A. and Uchida, Y. [2004], "A New Computational Scheme for Computing Greeks by the Asymptotic Expansion Approach," *Asia-Pacific Financial Markets*, Vol.11, 393-430.
- [32] Merton, R.C. [1976], "Option Pricing When Underlying Stock Returns are Discontinuous," *Journal of Financial Economics*, Vol. 3, 125-144.
- [33] Muroi, Y. [2005], "Pricing Contingent Claims with Credit Risk: Asymptotic Expansion Approach," *Finance and Stochastics*, Vol 9(3), 415-427.
- [34] Ninomiya, S. and Victoir, N. [2006], Weak Approximation of Stochastic Differential Equations and Application to Derivative Pricing, preprint.
- [35] Nualart, D. [1995], "The Malliavin Calculus and Related Topics," Springer.

- [36] Nualart, D., Üstünel A. S. and Zalai M. [1988], “On the moments of a multiple Wiener-Itô integral and the space induced by the polynomials of the integral,” *Stochastics*, Vol 25, 233-340.
- [37] Ocone, D. and Karatzas, I. [1991], “A Generalized Clark Representation Formula, with Application to Optimal Portfolios,” *Stochastics and Stochastics Reports*, Vol. 34, 187-220.
- [38] Osajima, Y. [2006], ”The Asymptotic Expansion Formula of Implied Volatility for Dynamic SABR Model and FX Hybrid Model,” Preprint, Graduate School of Mathematical Sciences, the University of Tokyo.
- [39] Osajima, Y. [2007], ”General Asymptotics of Wiener Functionals and Application to Mathematical Finance,” Preprint, Graduate School of Mathematical Sciences, the University of Tokyo.
- [40] Sakamoto, Y., Takada, Y. and Yoshida, N. [2004], “Expansions of the Coverage Probabilities of Prediction Region Based on a Shrinkage Estimator,” *Statistics*, Vol. 38, no. 5, 381-390.
- [41] Sakamoto, Y. and Yoshida, N. [1996], “Expansion of Perturbed Random Variables Based on Generalized Wiener Functionals,” *Journal of Multivariate Analysis*, Vol. 59, no. 1, 34-59.
- [42] Sakamoto, Y. and Yoshida, N. [1998], “Asymptotic Expansion of M -estimator over Wiener Space,” *Statistical Inference for Stochastic Processes*, Vol. 1-1, 85-103.
- [43] Schoutens, W. [2000], *Stochastic Processes and Orthogonal Polynomials*, volume 146 of *Lecture Notes in Statistics*, Springer-Verlag.
- [44] Shigekawa, I. [1998], “Stochastic Analysis,” Iwanami(in Japanese).
- [45] Siopacha, M. and Teichmann, J.[2007], “Weak and Strong Taylor Methods for Numerical Solutions of Stochastic Differential Equations,” Working paper.
- [46] Takahashi, A. [1995], “Essays on the Valuation Problems of Contingent Claims,” Unpublished Ph.D. Dissertation, Haas School of Business, University of California, Berkeley.
- [47] Takahashi, A. [1999], “An Asymptotic Expansion Approach to Pricing Contingent Claims,” *Asia-Pacific Financial Markets*, Vol. 6, 115-151.
- [48] Takahashi, A. [2005], “A Note on Computing Greeks by an Asymptotic Expansion Scheme,” Preprint.
- [49] Takahashi, A. [2007a], “On an Asymptotic Expansion Approach to Numerical Problems in Finance,” Sugaku, Mathematical Society of Japan, Vol.59-1, 75-91. (in Japanese).
- [50] Takahashi, A. [2007b], An Asymptotic Expansion Approach in Finance, CIRJE Discussion Papers.
- [51] Takahashi, A. and Matsushima, S. [2004], “Monte Carlo Simulation with an Asymptotic Expansion in HJM Framework,” *FSA Research Review 2004*, 82-103, Financial Services Agency (in Japanese).

- [52] Takahashi, A. and Saito, T. [2003], “An Asymptotic Expansion Approach to Pricing American Options,” *Monetary and Economic Studies*, Vol. 22, 35-87. (in Japanese).
- [53] Takahashi, A. and Takehara, K.[2007], “Pricing Currency Options with a Market Model of Interest Rates under Jump-Diffusion Stochastic Volatility Processes of Spot Exchange Rates,” *Asia-Pacific Financial Markets*, Vol.14 , pp. 69-121.
- [54] Takahashi, A. and Takehara, K.[2008a], “Fourier Transform Method with an Asymptotic Expansion Approach: an Applications to Currency Options,” *International Journal of Theoretical and Applied Finance*, Vol. 11(4), pp. 381-401.
- [55] Takahashi, A. and Takehara, K. [2008b], “A Hybrid Asymptotic Expansion Scheme: an Application to Currency Options,” Working paper, CARF-F-116, the University of Tokyo, <http://www.carf.e.u-tokyo.ac.jp/workingpaper/>
- [56] Takahashi, A. and Uchida, Y. [2006], “New Acceleration Schemes with the Asymptotic Expansion in Monte Carlo Simulation,” *Advances in Mathematical Economics*, Vol. 8, 411-431.
- [57] Takahashi, A. and Yoshida, N. [2004], “An Asymptotic Expansion Scheme for Optimal Investment Problems,” *Statistical Inference for Stochastic Processes*, Vol. 7-2, 153-188.
- [58] Takahashi, A. and Yoshida, N. [2005], “Monte Carlo Simulation with Asymptotic Method,” *The Journal of Japan Statistical Society*, Vol. 35-2, 171-203.
- [59] Takanobu, S. [1988], “Diagonal Short Time Asymptotics of Heat Kernels for Certain Degenerate Second Order Differential Operators of Hormander Type,” *Publications of Research Institute for Mathematical Sciences, Kyoto University*, Vol. 24, 169-203.
- [60] Takanobu, S. and Watanabe, S. [1990], “Asymptotic Expansion formula of the Schilder Type for a Class of Conditional Wiener Functional Integrations,” In *Asymptotic Expansion Problems in Probability Theory: Proceedings of the Taniguchi International Symposium*, Sanda and Kyoto, 1990, Longman, 194-241.
- [61] Taniguchi, M. and Kakizawa, Y. [2000], “Asymptotic Theory of Statistical Inference for Time Series,” Springer Series in Statistics, Springer-Verlag, New York.
- [62] Uchida, M. and Yoshida, N. [2004a], “Information Criteria for Small Diffusions via the Theory of Malliavin-Watanabe,” *Statistical Inference for Stochastic Processes*, Vol. 7-1, 35-67.
- [63] Uchida, M. and Yoshida, N. [2004b], “Asymptotic Expansion for Small Diffusions Applied to Option Pricing,” *Statistical Inference for Stochastic Processes*, Vol. 7-3, 189-223.
- [64] Uemura, H. [1987], “On a Short Time Expansion of the Fundamental Solution of the Heat Equations by the Method of Wiener Functionals,” *Journal of Mathematics of Kyoto University*, Vol. 27-3, 417-431.

- [65] Watanabe, S. [1984], *Lectures on Stochastic Differential Equations and Malliavin Calculus*, Tata Institute of Fundamental Research, Springer-Verlag.
- [66] Watanabe, S. [1987], “Analysis of Wiener Functionals (Malliavin Calculus) and its Applications to Heat Kernels,” *The Annals of Probability*, Vol. 15, 1-39.
- [67] Yoshida, N. [1992a], “Asymptotic Expansion for Small Diffusions via the Theory of Malliavin-Watanabe,” *Probability Theory and Related Fields*, Vol. 92, 275-311.
- [68] Yoshida, N. [1992b], “Asymptotic Expansions for Statistics Related to Small Diffusions,” *The Journal of Japan Statistical Society*, Vol. 22, 139-159.
- [69] Yoshida, N. [1993], “Asymptotic Expansion of Bayes Estimators for Small Diffusions,” *Probability Theory and Related Fields*, Vol. Vol. 95, no. 4, 429-450.
- [70] Yoshida, N. [1996], “Asymptotic Expansion for Perturbed Systems on Wiener Space: Maximum Likelihood Estimators,” *Journal of Multivariate Analysis*, Vol. 57, no. 1, 1-36.
- [71] Yoshida, N. [2003], “Conditional Expansions and their Applications,” *Stochastic Processes and their Applications*, Vol. 107, no. 1, 53-81.
- [72] Yoshida, N. [2003], “Malliavin Calculus and Mathematical Statistics,” *Sugaku*, Vol. 55, no. 3, 225-244, (in Japanese).