

CIRJE-F-837

Combinatorial Bootstrap Inference IN in Prtially Identified Incomplete Structural Models

Marc Henry
Université de Montréal

Romuald Méango
Université de Montréal

Maurice Queyranne
Sauder School of Business

January 2012

CIRJE Discussion Papers can be downloaded without charge from:

<http://www.cirje.e.u-tokyo.ac.jp/research/03research02dp.html>

Discussion Papers are a series of manuscripts in their draft form. They are not intended for circulation or distribution except as indicated by the author. For that reason Discussion Papers may not be reproduced or distributed without the written consent of the author.

COMBINATORIAL BOOTSTRAP INFERENCE IN PARTIALLY IDENTIFIED INCOMPLETE STRUCTURAL MODELS

MARC HENRY, ROMUALD MÉANGO AND MAURICE QUEYRANNE

ABSTRACT. We propose a computationally feasible inference method in finite games of complete information. Galichon and Henry (2011) and Beresteanu, Molchanov, and Molinari (2011) show that such models are equivalent to a collection of moment inequalities that increases exponentially with the number of discrete outcomes. We propose an equivalent characterization based on classical combinatorial optimization methods that alleviates this computational burden and allows the construction of confidence regions with an efficient combinatorial bootstrap procedure that runs in linear computing time. The method can also be applied to the empirical analysis of cooperative and noncooperative games, instrumental variable models of discrete choice and revealed preference analysis. We propose an application to the determinants of long term elderly care choices.

Keywords: Incomplete structural models, multiple equilibria, partial identification, sharp bounds, confidence regions, max-flow min-cut, combinatorial bootstrap, elderly care.

JEL subject classification: C13, C72

Date: December 3, 2011. Parts of this paper were written while Henry was visiting the Graduate School of Economics at the University of Tokyo; he gratefully acknowledges the CIRJE for its hospitality and support. Financial support from SSHRC Grant 410-2010-242 is gratefully acknowledged. The authors thank Victor Chernozhukov, Russell Davidson, Alfred Galichon, Silvia Gonçalves, Hidehiko Ichimura and Ismael Mourifié for helpful discussions, Jean-Marc Robin and three anonymous referees for helpful suggestions, David Straley for providing the data and Daniel Stubbs for computing assistance. Comments from seminar audiences in Cambridge, the Bank of Japan, Harvard-MIT, Hitotsubashi, UBC Sauder School, UCL, U-Kyoto, U-Tokyo, Rochester, Vanderbilt, and participants at the Inference in Incomplete Models conference in Montreal, the First Workshop on New Challenges in Distributed Systems in Valparaíso and the Second Workshop on Optimal Transportation and Applications to Economics in Vancouver are also gratefully acknowledged. Correspondence address: Département de sciences économiques, Université de Montréal, C.P. 6128, succursale Centre-ville, Montréal QC H3C 3J7, Canada, marc.henry@umontreal.ca.

CIRJE-F-837

Combinatorial Bootstrap Inference IN in Prtially Identified Incomplete Structural Models

Marc Henry
Université de Montréal

Romuald Méango
Université de Montréal

Maurice Queyranne
Sauder School of Business

January 2012

CIRJE Discussion Papers can be downloaded without charge from:

<http://www.cirje.e.u-tokyo.ac.jp/research/03research02dp.html>

Discussion Papers are a series of manuscripts in their draft form. They are not intended for circulation or distribution except as indicated by the author. For that reason Discussion Papers may not be reproduced or distributed without the written consent of the author.

INTRODUCTION

With the conjoined advent of powerful computing capabilities and rich data sets, the empirical evaluation of complex structural models with equilibrium data is becoming prevalent, particularly in the analysis of social networks and industrial organization. However, in such models, multiple equilibria are the norm rather than the exception. Though multiplicity of equilibria and identifiability of the model's structural parameters are conceptually distinct, the former often leads to a failure of the latter, thereby invalidating traditional inference methods. This is generally remedied by imposing additional assumptions to achieve identification, such as imposing an equilibrium selection mechanism or a refinement of the equilibrium concept. Manski (1993) and Jovanovic (1989) were among the first to advocate a new inference approach that dispenses with identification assumptions and delivers confidence regions for partially identified structural parameters. A large literature has developed on the general problem of inference on partially identified parameters defined as minimizers of objective functions or more specifically as solutions to moment inequality restrictions, following the seminal work of Chernozhukov, Hong, and Tamer (2007).

In structural estimation using equilibrium conditions, the partial identification approach was initially applied, as in Haile and Tamer (2003), to achieve simple and robust inference from implications of the model in the form of a small number of moment inequalities. This partial identification approach was applied to inference in games by Andrews, Berry, and Jia (2003), Pakes, Porter, Ho, and Ishii (2004), Ciliberto and Tamer (2009), Jia (2008) among others. However, this approach brings only part of the empirical content of the model to bear on the estimation, resulting in unnecessary loss of informativeness. In models with multiple equilibria and no additional prior information, nothing is known of the equilibrium selection mechanism. If a particular equilibrium selection mechanism is posited, the model likelihood can be derived and inference based on it. Jovanovic (1989) characterizes compatibility of an economic structure with the true data generating process as the existence of some (unknown) equilibrium selection mechanism, for which the likelihood is equal to the true data generating mechanism. Berry and Tamer (2006) define the identified set as the collection of structural parameter values for which the structure is compatible with the data generating mechanism in the sense of Jovanovic (1989). This definition of the identified set is not directly conducive to inference, as it involves an infinite dimensional (nuisance) parameter (the equilibrium selection mechanism). However, in the case of finite non cooperative games of complete information, Galichon and Henry (2011) and Beresteanu, Molchanov, and Molinari (2011) show equivalence of the Jovanovic (1989) definition with a system of inequalities. Hence, they show that the empirical content of such models is characterized by a finite collection of moment inequalities.

A large literature has developed on inference in moment inequality models since the seminal contribution of Chernozhukov, Hong, and Tamer (2007). We discuss and review it in Section 3. However, a major challenge in the framework of this paper is that the number of inequalities characterizing the empirical content of the model grows exponentially with the number of equilibrium strategy profiles. Hence the combinatorial optimization approach that we propose in this paper is to the best of our knowledge the only computationally feasible inference procedure for empirically relevant incomplete economic structures. The growing literature on “inference with many moment inequalities” addresses theoretical issues relating to the case, where the number of inequalities grows with sample size and does not alleviate the computational burden mentioned here. This problem of exponential complexity goes a long way towards explaining the dearth of empirical studies using partial identification in such models. However, abandoning this partial identification approach would mean abandoning robust inference not only in non cooperative games of perfect information but also in large classes of models that share exactly the same feature, and fall into the framework of this paper. They include cooperative games, such as matching games and network formation games, revealed preference analysis of spacial preferences and matching markets and instrumental variable models of discrete choice.

The objective of this paper is to propose a combinatorial solution to this problem, where the number of inequality restrictions grows exponentially with the number of strategy profiles or discrete outcomes. Ekeland, Galichon, and Henry (2010) have shown that generic partial identification problems can be formulated as optimal transportation problems. Developing ideas in Galichon and Henry (2011), we exploit the special structure of discrete choice problems and show that correct specification can be formulated as a problem of maximizing flow through a network, and that the identified set can be obtained from the Max-Flow Min-Cut Theorem. The dual problems of maximizing flow through a network and finding a minimum capacity cut are classics in combinatorial optimization and operations research, with applications in many areas such as traffic, communications, routing and scheduling; see, for example Schrijver (2004) for the theory and history, and Ahuja, Magnanti, and Orlin (1993) for numerous applications. To our knowledge this is the first application of the Max-Flow Min-Cut Theorem to statistical inference for equilibrium models. We apply this powerful combinatorial method to the problem of constructing confidence regions for structural parameters. We construct a functional quantile for the bootstrap process using a linear computing time algorithm and replace the unknown empirical process by this quantile in the system of moment inequalities to obtain the least relaxation of the moment inequalities, hence maximum informativeness, while controlling the confidence level of the covering region. Since the procedure involves bootstrapping the empirical process only, it does not suffer from the problems of bootstrap validity in partially identified models described in Chernozhukov, Hong, and Tamer (2007)

and Bugni (2010). We illustrate and assess our procedure on a very simple full information game with 2 players and 3 strategies, easily derived equilibria and yet a large number of inequalities to characterize its empirical content (namely 127). We simulate the game under a variety of parameter values and assumptions on the data generating process and with explanatory variables. Finally, we illustrate the approach, the procedure and the interpretation of results on an application to the determinants of long term elderly care choices of American families.

In summary, the main contributions of this paper are as follows:

1. We present a unified approach to inference in incomplete structural models.
2. We provide a simplified and insightful new proof for a characterization of the identified set.
3. We present a computationally efficient, combinatorial procedure that allows feasible inference in empirically relevant incomplete structural models. We demonstrate its practical efficiency in extensive simulations of a simple game.
4. We apply this methodology to an empirical example and demonstrate the type of econometric analysis and insights that it allows.

The paper is organized as follows. The next section introduces the general framework and the object of study. Section 2 derives the characterization of the identified set with the Min-Cut Max-Flow Theorem. Section 3 describes the combinatorial procedure to efficiently construct the confidence region. Section 4 contains the simulation evidence and Section 5 the empirical application. The last section concludes. Proofs are collected in an appendix.

1. ANALYTICAL FRAMEWORK

1.1. Model specification. We consider the following model specification.

$$Y \in G(X, \varepsilon; \theta), \tag{1.1}$$

where Y is an observable outcome variable, which takes values in a finite set $\mathcal{Y} = \{y_1, \dots, y_K\}$, X is a vector of exogenous explanatory variables with domain \mathcal{X} , ε is a vector of unobservable heterogeneity variables with domain $\Xi \subset \mathbb{R}^l$ and $\theta \in \Theta \subset \mathbb{R}^d$ is a vector of unknown parameters. Finally, $G : (X, \varepsilon) \rightrightarrows G(X, \varepsilon; \theta)$ is a multi-valued mapping. The random elements X , Y and ε are defined on a common probability space $(\Omega, \mathcal{F}, \mathbb{P})$. The sample consists in n observational units $i = 1, \dots, n$, which are independent and identical in distribution. To each unit i is attached a vector $(Y_i, X_i, \varepsilon_i)$, only the first two elements of which shall be observed. For each potential outcome $y \in \mathcal{Y}$, we denote by $P(y|X)$ the conditional probability $\mathbb{P}(Y = y|X)$. If Z is a subset of \mathcal{Y} , $P(Z|X)$ will denote $\sum_{y \in Z} P(y|X)$. It is important to emphasize here the fact that $P(\cdot|X)$ denotes the true outcome data generating process, which is unknown, but can be estimated from the data. It is not

a function of the structural parameter vector and cannot be construed as the likelihood from the model. The vector of unobservable variables ε in the economic structure has conditional cumulative distribution function $F(\varepsilon|X; \theta)$ for some known function F parameterized by θ (the same notation is used for the parameters of the model correspondence and for the parameters of the error distribution to indicate that they may have common components). The economic structure is summarized by the multi-valued mapping G . A special case of specification (1.1) arises when G is a function, in which case model (1.1) is a nonlinear non separable single equation discrete choice model as in Chesher (2010). Here, however, we entertain the possibility of G having multiple values arising from multiple equilibria, data censoring or endogeneity. G is entirely given by the economic structural model, up to an unknown parameter vector θ .

The analytical framework, concepts and procedures proposed throughout the paper will be illustrated and discussed with the following simple example.

Example 1 (Partnership game). *Our example is a simple non cooperative full information game of complementarities.*

- **STRATEGIES:** *There are two players, who simultaneously decide, whether to invest strongly (strategy H), weakly (strategy L) or not at all (strategy O) in a partnership.*
- **PAYOFFS:** *Players pay a cost $c \geq 0$ (respectively $2c$) for a weak (respectively strong) investment. Benefits that accrue to players depend on the overall level of investment in the partnership and explanatory variables J_i , $i = 1, 2$, where $J_i = 1$ if player i is female, and zero otherwise. The benefits for player i are $3c(1 + \beta J_i)$ in case both players invest strongly, $2c(1 + \beta J_i)$ in case one player invests weakly and the other strongly and $c(1 + \beta J_i)$ in case both players invest weakly. Finally player i also experiences an idiosyncratic random participation payoff ε_i , $i = 1, 2$ with a density with respect to Lebesgue measure. The payoff matrix for the game is given in Table 1.*
- **EQUILIBRIUM CONCEPT:** *We assume that outcomes are Nash equilibria in pure strategies. Other equilibrium concepts could be entertained, in particular with mixed strategies, as will be discussed in Section 3.1 and illustrated in the empirical application.*

The strategies, payoffs and equilibrium concept together define the economic structure. Y is an observed equilibrium strategy profile. $J = (J_1, J_2)$ is also observed by the analyst. The idiosyncratic participation benefit $\varepsilon = (\varepsilon_1, \varepsilon_2)$ is not, but it is common knowledge to the players. The structural parameter vector is $\theta = (c, \beta)$. The equilibrium correspondence, i.e., the set of equilibria for each value of ε , J and θ , can be easily derived, and defines the multi-valued mapping G in model specification (1.1), which is represented in the $(\varepsilon_1, \varepsilon_2)$ space in Figure 1 for the case $\beta = 0$. Since we

TABLE 1. Payoff matrix for the partnership game. In each cell, the top expression is player 1's payoff and the bottom term is player 2's payoff.

		<i>Player 2 :</i>		
		<i>H</i>	<i>L</i>	<i>O</i>
<i>Player 1 :</i>	<i>H</i>	$3c(1 + \beta J_i) - 2c + \varepsilon_1$	$2c(1 + \beta J_i) - 2c + \varepsilon_1$	$-2c + \varepsilon_1$
		$3c(1 + \beta J_i) - 2c + \varepsilon_2$	$2c(1 + \beta J_i) - c + \varepsilon_2$	0
	<i>L</i>	$2c(1 + \beta J_i) - c + \varepsilon_1$	$c(1 + \beta J_i) - c + \varepsilon_1$	$-c + \varepsilon_1$
		$2c(1 + \beta J_i) - 2c + \varepsilon_2$	$c(1 + \beta J_i) - c + \varepsilon_2$	0
	<i>O</i>	0	0	0
		$-2c + \varepsilon_2$	$-c + \varepsilon_2$	0

assume that ε has absolutely continuous distribution with respect to Lebesgue measure, we do not include zero probability predictions, such as $\{OO, OL\}$ when $\varepsilon_2 = c$ and $\varepsilon_1 < -c$ for instance.

1.2. Object of inference. Model (1.1) has the fundamental feature that G is multi-valued (because of multiple equilibria in the example above for instance). For a given value of (X, ε, θ) , the model predicts a set of possible outcomes $G(X, \varepsilon; \theta)$. Only one of them, namely Y , is actually realized, but the economic structure is silent about how that particular Y was selected among $G(X, \varepsilon; \theta)$. In other words, the economic structure holds no information about the equilibrium selection mechanism. If the true (unknown) equilibrium selection mechanism is denoted $\pi_0(y|\varepsilon, X)$, which is a probability on $G(X, \varepsilon; \theta)$, then the likelihood of observation y can be written

$$L(\theta|y, X) = \int_{\Xi} \pi_0(y|\varepsilon, X) dF(\varepsilon|X; \theta),$$

and the true parameter θ_0 satisfies

$$P(y|X) = \int_{\Xi} \pi_0(y|\varepsilon, X) dF(\varepsilon|X; \theta_0), \quad X\text{-a.s., for all } y. \quad (1.2)$$

Jovanovic (1989) points out that the incomplete model (incomplete because the equilibrium selection is not modeled) is compatible with the true data generating process $P(\cdot|X)$ if and only if there exists a (generally non unique) equilibrium selection mechanism π_0 such that (1.2) holds. The identified set is then defined as the set Θ_I of parameter values θ such that model (1.1) is compatible in the sense of Jovanovic (1989).

Definition 1 (Identified set). *The identified set Θ_I is the set of parameter values $\theta \in \Theta$ such that there exists a probability kernel $\pi(\cdot|\varepsilon, X)$ with support $G(X, \varepsilon; \theta)$ for which (1.2) holds.*




FIGURE 1. Representation of the equilibrium correspondence $G(J, \varepsilon; \theta)$ in the $(\varepsilon_1, \varepsilon_2)$ space, when $\beta = 0$.

The identified set is empty if no value of the parameter can rationalize the data generating process, in which case the structural model is misspecified. The identified set is a singleton in case of point identification, which occurs if G happens to be single valued under the true parameter values (in case $c = \beta = 0$ in Example 1) or in very special cases under large support assumptions on X , as in Tamer (2003). The identified set is totally uninformative, i.e., $\Theta_I = \Theta$, in case the model has no empirical content (if for instance $G(X, \varepsilon; \theta_0)$ contains all selected outcome values for almost all ε at the true value θ_0).

1.3. Applications of the framework. Specification (1.1), hence the inference procedure presented in this paper, has a wide range of applications. Some of the most compelling ones are the empirical analysis of games, instrumental variable models of discrete choice with endogeneity and revealed preference analysis.

- **EMPIRICAL ANALYSIS OF GAMES:** As illustrated in Example 1, Model (1.1) applies to the empirical analysis of noncooperative games of perfect information (normal form games). They include the classic entry game of Bresnahan and Reiss (1990) and Berry (1992) as well as the social interaction game of Soetevent and Kooreman (2007). Noncooperative games of private information make for a less compelling application of this framework as point identification conditions are more easily derived and justified than in their perfect information counterparts (see for instance Aradillas-Lopez (2010) and Bajari, Hahn, Hong, and Ridder (2011) for a discussion). Finally, some cooperative games can be analyzed and estimated within the present framework, in particular matching and social network formation games, where the equilibrium correspondence is characterized by pairwise stability. Uetake and Watanabe (2011) present an empirical analysis of entry by merger, where the present inference procedure can be applied.
- **DISCRETE CHOICE MODELS WITH ENDOGENEITY:** Chesher, Rosen, and Smolinski (2011) show that instrumental variable models of discrete choice fall under model (1.1) and they use Theorem 1 of Galichon and Henry (2011) or equivalently Theorem 3.2 of Beresteanu, Molchanov, and Molinari (2011) to characterize the identified set. The present work complements Chesher, Rosen, and Smolinski (2011) in proposing the first feasible inference procedure for such models.
- **REVEALED PREFERENCE ANALYSIS:** Henry and Mourifié (2011) apply the inference procedure proposed here to analyze voting behaviour from a revealed preference standpoint. The same approach can be applied to revealed preference testing in matching markets as in Echenique, Lee, Shum, and Yenmez (2011) or the revealed preference approach to games taken in Pakes, Porter, Ho, and Ishii (2004).

2. OPERATIONAL CHARACTERIZATION OF THE IDENTIFIED SET

As noted in Berry and Tamer (2006), Definition 1 is not an operational definition of the identified set, as it includes the equilibrium selection mechanism as an infinite dimensional parameter. Galichon and Henry (2011) and Beresteanu, Molchanov, and Molinari (2011) show a characterization of the identified set with a finite collection of moment inequalities. In this section, we give an equivalent characterization of the identified set, whose proof is much simpler and relies on the Min-Cut Max-Flow Theorem, which brings classical efficient combinatorial optimization methods to bear on the problem. This will prove crucial for the feasibility of the inference procedure in realistic and relevant empirical examples.

First, we set out the main heuristic for the operational characterization of the identified set. Model specification (1.1) is a discrete choice model, hence the set \mathcal{Y} of outcomes is finite and the correspondence G takes only a finite number of values, which we label $\mathcal{U} = \{u_1, \dots, u_J\}$. Each u is a set (possibly singleton) of outcomes in \mathcal{Y} . Because the model is incomplete, it does not predict the probabilities of individual outcomes in \mathcal{Y} , but it predicts the probability of each combination of equilibria listed in \mathcal{U} . We denote these probabilities $Q(u|X; \theta)$ as they depend on the structural parameter value.

Definition 2 (Predicted probabilities). *For each $u \in \mathcal{U}$, we define $Q(u|X; \theta) := \mathbb{P}(G(X, \varepsilon; \theta) = u|X, \theta)$. If V is a subset of \mathcal{U} , we write $Q(V|X; \theta) = \sum_{u \in V} Q(u|X; \theta)$.*

In most applications, it will be difficult to obtain closed forms for $Q(u|X; \theta)$. However, ε can be randomly generated. Given a sample $(\varepsilon^r)_{r=1, \dots, R}$ of simulated values, $Q(u|X; \theta)$ can be approximated by $\sum_{r=1}^R 1\{u = G(X, \varepsilon^r; \theta)\}/R$. Bajari, Hong, and Ryan (2010) propose an importance sampling procedure that greatly reduces the computational burden of this stage of the inference. The simulation procedure is now standard and cannot be avoided if one wishes, as we do here, to exhaust the empirical content of the structural model.

Example 1 continued: *In the partnership example with $\beta = 0$, the model predicts the following values for the equilibrium correspondence: $\mathcal{U} = \{ \{OL\}, \{LH, OL, HH\}, \{HH, LH, OO\}, \{OO\}, \{HH, OO\}, \{HH, LL, HL, LH\}, \{HH, LL, OO, HL, LH\}, \{HH, OO, HL\}, \{HH, HL, LO\}, \{LO\} \}$. The set \mathcal{Y} of equilibrium strategy profiles (that may be observed) is $\{HH, HL, LH, LL, LO, OL, OO\}$ with 7 elements, while the set of predicted collections of equilibria (possible values of the equilibrium correspondence) \mathcal{U} has 10 elements. The predicted probabilities can be computed in the following way. For instance, $Q(\{OL\}|c) = \mathbb{P}(\varepsilon_1 \leq -c \text{ and } \varepsilon_2 \leq c)$ and $Q(\{HH, LH, OL\}|c) = \mathbb{P}(-c \leq \varepsilon_1 \leq 0 \text{ and } \varepsilon_2 \leq c)$ and the remaining 8 probabilities are determined similarly from Figure 1.*

The model structure imposes a set of restrictions on the relation between the predicted probabilities of equilibrium combinations and the true probabilities of outcomes. For instance, the predicted probability $Q(\{HH, LH, OL\}|X; \theta)$ in the above example cannot be larger than the sum $P(HH) + P(LH) + P(OL)$ of probabilities of occurrence of each individual equilibrium in u , since Y is either HH , LH or OL , when $u = \{HH, LH, OL\}$ is predicted. More generally, since P and Q are the marginals of the joint distribution of (Y, U) given X , we must have for all $u \in \mathcal{U}$:

$$Q(u|X; \theta) = \sum_{y \in u} \mathbb{P}(Y = y \text{ and } U = u|X; \theta) \leq \sum_{y \in u} \mathbb{P}(Y = y|X; \theta) = \sum_{y \in u} P(y|X). \quad (2.1)$$

Note that $Q(u|X; \theta)$ may be strictly smaller than $\sum_{y \in u} P(y|X)$ when some outcome $y \in u$ also belongs to other combinations u' that may arise under different values of ε , as its (marginal) probability $P(y|X)$ must then be split between $Q(u|X; \theta)$ and the probabilities $Q(u'|X; \theta)$ of such other combinations $u' \in \mathcal{U}$ containing y . However, inequalities (2.1) do not exhaust the information in the structure. They may all be satisfied and yet the structure may be incompatible with the data generating process as the following example shows. Hence more inequalities will be needed as derived below.

Example 1 continued: *In the partnership example with $\beta = 0$, suppose that the true equilibrium selection mechanism is such that $Q(\{OL\}|\theta) = P(OL) > 0$ and $Q(\{HH, LH, OL\}|\theta) = P(HH) + P(LH) + P(OL)$. Then $Q(\{OL\} \cup \{HH, LH, OL\}|\theta) = Q(\{OL\}|\theta) + Q(\{HH, LH, OL\}|\theta) > P(HH) + P(LH) + P(OL)$ so that $\theta \notin \Theta_I$.*

Extending this observation, consider a subset $V \subseteq \mathcal{U}$ and define

$$V^\cup := \{y \in Y : y \in u \text{ for some } u \in V\} = \bigcup_{u \in V} u.$$

Then we must have

$$\begin{aligned} Q(V|X; \theta) &= \sum_{u \in V} \sum_{y \in u} \mathbb{P}(Y = y \text{ and } U = u|X; \theta) \\ &= \sum_{y \in V^\cup} \sum_{u \in V: y \in u} \mathbb{P}(Y = y \text{ and } U = u|X; \theta) \\ &\leq \sum_{y \in V^\cup} \sum_{u \in \mathcal{U}} \mathbb{P}(Y = y \text{ and } U = u|X; \theta) \\ &= \sum_{y \in V^\cup} P(y|X) \end{aligned}$$

where the inequality is again due to the fact that some $y \in V^\cup$ may also belong to some $u' \notin V$. Since this inequality holds for every $V \subseteq \mathcal{U}$, we must have

$$\max_{V \subseteq \mathcal{U}} \left(\sum_{u \in V} Q(u|X; \theta) - \sum_{y \in V^\cup} P(y|X) \right) \leq 0.$$

This inequality must also hold for every realization x of X in the domain \mathcal{X} of the explanatory variables, implying that every θ in the identified set $\Theta_I(P)$ must satisfy

$$\sup_{x \in \mathcal{X}} \max_{V \subseteq \mathcal{U}} \left(\sum_{u \in V} Q(u|x; \theta) - \sum_{y \in V^\cup} P(y|x) \right) \leq 0.$$

So far, we have shown implications of the model. It is far more difficult to show that these implication actually exhaust all the empirical content of the model, i.e., that they involve no loss of information and constitute sharp bounds. In Theorem 1 below, we will show this with an appeal to the classical

Max-Flow Min-Cut Theorem of combinatorial optimization, providing our characterization (2.2) of the identified set. We thereby provide, for the case of a finite set of possible outcomes, a new and simpler proof of the characterization of the identified set with a finite collection of inequalities, without the complicated apparatus of the theory of random sets. This allows us to emphasize the combinatorial optimization formulation of our inference problem, which is key to its tractable solution in empirically relevant instances. Theorem 1 below also provides an alternative characterization (2.3) of the identified set from the “dual” perspective of outcome subsets $Z \subseteq \mathcal{Y}$, in addition to the preceding characterization (2.2) based on combination subsets $V \subseteq \mathcal{U}$, with the notation

$$Z^\cap := \{u \in \mathcal{U} : u \subseteq Z\} \text{ and } Z^{-1} := \{u \in \mathcal{U} : u \cap Z \neq \emptyset\}.$$

This alternative characterization may be useful in situations where the number of possible outcomes is much smaller than the number of possible combinations (as is the case in Example 1, where the number of equilibrium outcomes (cardinality of \mathcal{Y}) is 7, so the corresponding number of inequalities to be checked is $2^7 - 1 = 127$, whereas the number of predicted equilibrium combinations (cardinality of \mathcal{U}) is 10, so the corresponding number of inequalities to check would be $2^{10} - 1 = 1023$). Finally, it is also equivalent to the characterization of the identified set first derived in Galichon and Henry (2011) and Beresteanu, Molchanov, and Molinari (2011), which we give in (2.4) in our notation.

Theorem 1. *The identified set is*

$$\Theta_I(P) = \left\{ \theta \in \Theta : \sup_{x \in \mathcal{X}} \max_{V \subseteq \mathcal{U}} \left(Q(V|x; \theta) - P(V^\cup|x) \right) \leq 0 \right\} \quad (2.2)$$

$$= \left\{ \theta \in \Theta : \sup_{x \in \mathcal{X}} \max_{Z \subseteq \mathcal{Y}} \left(Q(Z^\cap|x; \theta) - P(Z|x; \theta) \right) \leq 0 \right\} \quad (2.3)$$

$$= \left\{ \theta \in \Theta : \sup_{x \in \mathcal{X}} \max_{Z \subseteq \mathcal{Y}} \left(P(Z|x; \theta) - Q(Z^{-1}|x; \theta) \right) \leq 0 \right\}. \quad (2.4)$$

Theorem 1 gives three characterizations of the identified set $\Theta_I(P)$, sometimes called *sharp identified region* in the literature. $\Theta_I(P)$ contains all the values of the parameter such that (1.1) holds and only such values. Moreover, all elements of $\Theta_I(P)$ are observationally equivalent. Hence no value of the parameter vector θ contained in $\Theta_I(P)$ can be rejected on the basis of the information available to the analyst. Thus, $\Theta_I(P)$ completely characterizes the empirical content of the model.

Example 1 continued: *To illustrate the computation of the identified set, consider the case, where it is known that $\beta = 0$. Assume that the true parameter value is $c_0 = 1/4$ and the idiosyncratic shocks are independent and uniformly distributed over $[-1/2, 1/2]$. Suppose further that the true data generating process is equal to the distribution implied by a uniform equilibrium selection rule, whereby all equilibrium strategy profiles within the equilibrium correspondence are selected with*

equal probability. For example, when $\varepsilon_1 \geq c_0 = 1/4$ and $-1/4 = -c_0 \leq \varepsilon_2 \leq 0$, each strategy profile within the equilibrium correspondence $\{HH, HL, LO\}$ is equally likely. The probability distribution of the true data generating process in this case is defined by $P(HH) = 167/960$, $P(OO) = 191/480$, $P(OL) = P(LO) = 1/12$, $P(LL) = 19/320$ and $P(HL) = P(LH) = 97/960$. The identified set is derived as the set of values of c such that the $2^7 - 1 = 127$ inequalities of the form $P(Z) \geq Q(Z^\cap | c)$, all $Z \subseteq \{HH, HL, LH, LL, LO, OL, OO\}$, are satisfied. For instance, one of those inequalities is $59/320 = P(LO \text{ or } HL) \geq Q(\{LO\} | c) = (1/2 - c)^2$ if $c \leq 1/2$ and zero otherwise. The identified set can be computed using a Min-Cut Max-Flow algorithm, which yields $[1/2 - 1/\sqrt{12}, 1/3] \simeq [0.2113, 0.3333]$ where the lower bound of the interval happens to be the smallest value of $c > 0$ for which the inequality in (2.3) with $Z = \{LO, OL\}$ is satisfied, and the upper bound happens to be the largest value for which that with $Z = \{HH, HL, LH, LL, OO\}$ is satisfied.

As illustrated in Example 1, even in simple examples, where the equilibria are very easy to compute, the exponential size of the characterization of the identified set is a severe computational burden that is best approached with combinatorial optimization techniques, as developed in the next section.

3. CONFIDENCE REGION

3.1. Objective. We now turn to the problem of inference on $\Theta_I(P)$ based on a sample of observations $((Y_1, X_1), \dots, (Y_n, X_n))$. We seek coverage of the identified set with prescribed probability $1 - \alpha$, for some $\alpha \in (0, 1)$. It would be tempting to appeal to the large literature on inference in moment inequality models. This includes several proposals for the construction of confidence regions covering each point in the identified set, which are generally preferred on account of the fact that they may be more informative (although this may sometimes be misleading as pointed out in Henry and Onatski (2011)). Such proposals include Section 5 of Chernozhukov, Hong, and Tamer (2007), Romano and Shaikh (2008), Rosen (2008), Galichon and Henry (2009) and Andrews and Soares (2010) among others. All of the above propose to construct confidence regions by inverting specification tests. Hence, the confidence region is constructed through a search in the parameter space, with a computationally demanding testing procedure at each parameter value visited in the search. This becomes computationally infeasible for realistic parameter vector dimensions. With a reasonably precise grid search and 5 parameters (for example), the number of points to be visited is in the tens of billions. If the identified set is known to be convex, the search can be conducted from a central point with a dichotomy in polar coordinates, yet it remains computationally impractical to conduct a statistical procedure for each point in the search.

Hence, each parameter value in the search must be accepted or rejected based on a deterministic criterion. This means the significance of the confidence region must be controlled independently of

the parameter value. This will automatically produce a confidence region that covers the identified set. Proposals for the construction of confidence regions covering the identified set include Chernozhukov, Hong, and Tamer (2007), Romano and Shaikh (2010), Galichon and Henry (2006) and Bugni (2010) among others. These can be applied to realistic models defined by a small number of moment inequality restrictions. However, a major challenge in the framework of this paper is that the number of inequalities characterizing the empirical content of the model in Theorem 1 grows exponentially with the cardinality of \mathcal{Y} , which in the case of games is the number of equilibrium strategy profiles (in the very simple partnership game of Example 1, the number of inequalities is 127). Hence the combinatorial optimization approach that we propose in this paper is to the best of our knowledge the only computationally feasible inference procedure for empirically relevant economic structures defined by finite games and other models of discrete choice with endogeneity.

Definition 3 (Confidence region). *A confidence region of asymptotic level $1 - \alpha$ for the identified set Θ_I is defined as a sequence of regions Θ_n , $n \in \mathbb{N}$, satisfying $\liminf_n \mathbb{P}(\Theta_I \subseteq \Theta_n) \geq 1 - \alpha$.*

We seek coverage of the set of values of the parameter θ such that $Q(V|x, \theta) \leq P(V^\cup|x)$ for all values of x and all subset V of \mathcal{U} . Q is determined from the model, but P is unknown. However, if we can construct random functions $\bar{P}_n(A|x)$ that dominate the probabilities $P(A|x)$ for all values of x and all subsets A of \mathcal{Y} with high probability, then in particular, $\bar{P}_n(V^\cup|x) \geq P(V^\cup|x)$ for each x and each subset V of \mathcal{U} . Hence any θ satisfying $Q(V|x, \theta) \leq P(V^\cup|x)$ for all values of x and all subsets V of \mathcal{U} also satisfies $Q(V|x, \theta) \leq \bar{P}_n(V^\cup|x)$ for all values of x and all subsets V of \mathcal{U} . There remains to control the level of confidence of the covering region, which is achieved by requiring that \bar{P}_n dominate P with probability asymptotically no less than the desired confidence level. Equivalently, when working from characterization (2.4), we impose the same requirement for dominated functions \underline{P}_n . Hence the following assumption.

Assumption 1. *Let the random functions $A \mapsto \bar{P}_n(A|x)$, $A \subseteq \mathcal{Y}$, satisfy*

$$\liminf_n \mathbb{P} \left(\sup_{x \in \mathcal{X}} \max_{A \subseteq \mathcal{Y}} [P(A|x) - \bar{P}_n(A|x)] \leq 0 \right) \geq 1 - \alpha. \quad (3.1)$$

Suppose now a value θ_0 of the parameter vector belongs to the identified set Θ_I . Then, by Theorem 1, for all x and $V \subseteq \mathcal{U}$, $Q(V|x; \theta_0) \leq P(V^\cup|x)$, so that with probability tending to no less than $1 - \alpha$, $Q(V|x; \theta_0) \leq \bar{P}_n(V^\cup|x)$, hence Theorem 2.

Theorem 2 (Confidence region). *Under Assumption 1, the sets*

$$\Theta_I(\bar{P}_n) = \left\{ \theta \in \Theta : \sup_{x \in \mathcal{X}} \max_{V \subseteq \mathcal{U}} (Q(V|x; \theta) - \bar{P}_n(V^\cup|x)) \leq 0 \right\} \quad (3.2)$$

define a confidence region of asymptotic level $1 - \alpha$ for Θ_I (according to Definition 3).

Theorem 2 has the fundamental feature that it dissociates search in the parameter space (or even possibly search over a class of models) from the statistical procedure necessary to control the confidence level. The upper probabilities \bar{P}_n can be determined independently of θ in a procedure that is performed once and for all using only sample information, i.e. fully nonparametrically. Once the upper probabilities are determined, probabilities Q over predicted sets of outcomes are computed for particular chosen specifications of the structure and values of the parameter, and such specifications and values are tested with inequalities defining $\Theta_n(\bar{P}_n)$. This dissociation of the statistical procedure to control confidence level from the search in the parameter space is crucial to the computational feasibility of the proposed inference procedure in realistic examples (i.e. sample sizes in the thousands, two-digit dimension of the parameter space and two-digit cardinality of the set of observed outcomes, as in the application to teen behavior in Soetevent and Kooreman (2007), or to entry in the airline market in Ciliberto and Tamer (2009)). The latter consider only equilibria in pure strategies, as we have until now. If equilibria in mixed strategies are also considered, as in Bajari, Hong, and Ryan (2010) and in the family bargaining application below, we can appeal to results in Beresteanu, Molchanov, and Molinari (2011) and Galichon and Henry (2011). In particular, Galichon and Henry (2011) show that if the game has a *Shapley regular core* (which is the case in the family bargaining application, by Lemma 2 of Galichon and Henry (2011)), then the identified set is characterized by (2.3) of Theorem 1 with the caveat that the set function $Z \mapsto Q(Z^\cap|x)$ is replaced by

$$\mathcal{L}(Z|x) = \int \min_{\sigma \in G(\varepsilon|X;\theta)} \sigma(Z) d\nu(\varepsilon), \quad (3.3)$$

where $G(\varepsilon|X;\theta)$ is now a set of mixed strategies, i.e. a set of probabilities on the set of outcomes, as opposed to a subset of the set of outcomes. Hence the methodology is easily adapted, as in the application of Section 5.

3.2. Control of confidence level. We now turn to the determination of random functions satisfying Assumption 1. First, for each $y \in \mathcal{Y}$, let $\hat{P}_n(y|x)$ be the empirical analog (or more generally a nonparametric estimator) of $P(y|x)$ and $\hat{P}_n(A|x) = \sum_{y \in A} \hat{P}_n(y|x)$ for each $A \subseteq \mathcal{Y}$. A simple way of achieving (3.1) is by considering the random variable

$$M_n := \sup_{x \in \mathcal{X}} \max_{A \subseteq \mathcal{Y}} [P(A|x) - \hat{P}_n(A|x)].$$

Denoting by c_n^α the $(1 - \alpha)$ -quantile of the distribution of M_n , we have $\mathbb{P}(M_n \leq c_n^\alpha) = 1 - \alpha$ by construction, hence

$$\mathbb{P} \left(\sup_{x \in \mathcal{X}} \max_{A \subseteq \mathcal{Y}} [P(A|x) - \hat{P}_n(A|x) - c_n^\alpha] \leq 0 \right) \geq 1 - \alpha, \quad (3.4)$$

and the desired result with $\bar{P}(A|x) = \hat{P}_n(A|x) + c_n^\alpha$. However, by construction, c_n^α is independent of A and x , so that the region obtained by plugging $\bar{P}(A|x) = \hat{P}_n(A|x) + c_n^\alpha$ into (3.2) of Theorem 2 will be unnecessarily conservative. We propose, instead, to replace c_n^α by a function $\beta_n(A|x)$ of A and x , which we interpret as a *functional quantile* of the distribution of the random function $P(A|x) - \hat{P}_n(A|x)$. Analogously to (3.4), we require it to satisfy

$$\mathbb{P} \left(\sup_{x \in \mathcal{X}} \max_{A \subseteq \mathcal{Y}} [P(A|x) - \hat{P}_n(A|x) - \beta_n(A|x)] \leq 0 \right) \geq 1 - \alpha. \quad (3.5)$$

We first give a heuristic description of our proposed functional quantile before precisely spelling out the bootstrap procedure involved in approximating it. If \mathcal{X} is finite, the random matrix $P(A|x) - \hat{P}_n(A|x)$, with $A \subseteq \mathcal{Y}$ and $x \in \mathcal{X}$ has a finite population of possible realizations, at most one for each possible sample draw. These realizations can be ordered according to the maximum entry in the matrix $\max_{x \in \mathcal{X}} \max_{A \subseteq \mathcal{Y}} [P(A|x) - \hat{P}_n(A|x)]$. Now take all realizations that never exceed the $(1 - \alpha)$ -quantile c_n^α of $\max_{x \in \mathcal{X}} \max_{A \subseteq \mathcal{Y}} [P(A|x) - \hat{P}_n(A|x)]$ and define $\bar{P}_n(A|x) = \hat{P}_n(A|x) + \beta_n(A|x)$, where $\beta_n(A|x)$ is the pointwise maximum over all realizations that never exceed c_n^α . This guarantees that the resulting confidence region obtained in (3.2) of Theorem 2 with $\bar{P}_n(A|x) = \hat{P}_n(A|x) + \beta_n(A|x)$ will be valid and will be contained in the region obtained with $\bar{P}_n(A|x) = \hat{P}_n(A|x) + c_n^\alpha$ (hence more informative than the latter). In case the conditioning variables are finitely supported, it is well known (see Singh (1981) and Bickel and Freedman (1981)) that the nonparametric bootstrap version of c_n^α is a valid approximation, which in turns guarantees the validity of the bootstrap procedure described below. In case X has continuous components, Chernozhukov, Lee, and Rosen (2009) derive the asymptotic distribution of the supremum (over \mathcal{X}) of the conditional empirical process, but nothing is known of its nonparametric bootstrap approximation.

Definition 4 (Nonparametric Bootstrap). *Let \mathbb{P}_n^* denote probability statements relative to the bootstrap distribution and conditional on the original sample $((Y_1, X_1), \dots, (Y_n, X_n))$. A bootstrap sample takes the form $((Y_1^*, X_1), \dots, (Y_n^*, X_n))$, where the explanatory variable is not resampled and for each i , Y_i^* is drawn from distribution $\hat{P}_n(\cdot|X_i)$. Let $((Y_1^b, X_1), \dots, (Y_n^b, X_n))$, $b = 1, \dots, B$ be a sequence of B bootstrapped samples. Denote by $\hat{P}_n^*(\cdot)$ the bootstrap version (i.e., constructed identically from a bootstrap sample) of $\hat{P}_n(\cdot)$ and \hat{P}_n^b , $b = 1, \dots, B$ its values taken on the B realized bootstrap samples. Finally, for each $A \subseteq \mathcal{Y}$ and $1 \leq j \leq n$, denote $\zeta_n^*(A|X_j) = \sum_{y \in A} [\hat{P}_n(y|X_j) - \hat{P}_n^*(y|X_j)]$ and define $\zeta_n^b(A|X_j)$ analogously.*

In the bootstrap version of the problem, we are seeking functions β_n satisfying

$$\mathbb{P}_n^* \left(\max_{1 \leq j \leq n} \max_{A \subseteq \mathcal{Y}} [\hat{P}_n(A|X_j) - \hat{P}_n^*(A|X_j) - \beta_n(A|X_j)] \leq 0 \right) \geq 1 - \alpha \text{ * -a.s.}$$

If there was a total order on the space of realizations of ζ_n^* , we could choose β_n as the quantile of level $1 - \alpha$ of the distribution of ζ_n^* . However, the $\zeta_n^*(\cdot, X_j)$'s are random functions defined on $2^{\mathcal{Y}} \times \{X_1, \dots, X_n\}$, hence there is no such total order. We propose to determine β_n from a subset of $\lceil B(1 - \alpha) \rceil$ bootstrap realizations determined as follows (where $\lfloor x \rfloor$ is the largest integer below x).

Step 1: Draw bootstrap samples $((Y_1^b, X_1), \dots, (Y_n^b, X_n))$, for $b = 1, \dots, B$.

Step 2: For each $b \leq B$, $j \leq n$ and $A \subseteq \mathcal{Y}$, compute $\zeta_n^b(A|X_j) = \hat{P}_n(A|X_j) - \hat{P}_n^b(A|X_j)$.

Step 3: Discard at most a proportion α of the bootstrap indices, and compute $\beta_n(A|X_j)$ as the maximum over the remaining bootstrap realizations $\zeta_n^b(A|X_j)$.

Discarding at most $B\alpha$ among the bootstrap realizations guarantees the control of the level of confidence, and we wish to choose the set $D \subseteq \{1, \dots, B\}$ of discarded indices so as to make β_n as small as possible, to maximize informativeness of the resulting confidence region. Again, if there was a total order, we would be similarly discarding the $B\alpha$ largest realizations of ζ_n^b , effectively choosing β_n as the quantile of the distribution of ζ_n^b , $b = 1 \dots, B$. Instead, we discard all realizations of the matrix $\zeta_n^b(A|X_j)$ that have at least one entry that strictly exceeds the $(1 - \alpha)$ -quantile of $w_b = \max_{1 \leq j \leq n} \max_{A \subseteq \mathcal{Y}} \zeta_n^b(A|X_j)$. Hence, we choose D solving the optimization problem

$$\min \left\{ \max_{b \notin D} w_b : D \subseteq \{1, \dots, B\}, |D| \leq B\alpha \right\}. \quad (3.6)$$

The procedure is explained graphically in Figure 2.




FIGURE 2. Stylized representation of the determination of the functional quantile β_n in a case without explanatory variables. The subsets A of \mathcal{Y} are represented on the horizontal axis, ranging from \emptyset to \mathcal{Y} . ζ_n^d is one of two discarded realization of the empirical process (dotted lines), whereas ζ_n^k is one of three realizations that are not discarded (solid lines). β_n is the pointwise maximum over the realizations that were not discarded (thick line).

Problem (3.6) can be solved by the following *Bootstrap Realization Selection (BRS) algorithm*:

BRS Step 1: For each $b \leq B$, set $w'_b = \max_{1 \leq j \leq n} \sum_{y \in \mathcal{Y}} \max\{0, \hat{P}_n(y|X_j) - \hat{P}_n^b(y|X_j)\}$.

BRS Step 2: Let D be the set of indices b of the $\lfloor B\alpha \rfloor$ largest w'_b .

Proposition 1. *The BRS algorithm determines an optimal solution to problem (3.6) in $O(nB|\mathcal{Y}|)$ time.*

Remark 1. *Problem (3.6) may have alternate optimum solutions. As observed by a referee, this may arise when the sample size n is small, since $\hat{P}_n(y|X_j)$ and $\hat{P}_n^b(y|X_j)$ are multiples of $1/n$ and thus distinct w_b 's are more likely to have the same value when the sample size n is small. In case of ties, any optimum solution D to Problem (3.6) may be used to discard bootstrap realizations and determine functions β_n . If one desires a specific tie-breaking rule, e.g., for robustness or reproducibility, then we suggest the following lexicographic selection rule as a refinement to BRS Step 2: let w^b be the vector with components $w_j^b = \sum_{y \in \mathcal{Y}} \max\{0, \hat{P}_n(y|X_j) - \hat{P}_n^b(y|X_j)\}$ for $j = 1, \dots, n$; and let $[w]^b$ be the vector w^b with its components sorted in nonincreasing order, i.e., with $[w]_1^b = w_b \geq [w]_2^b \geq \dots \geq [w]_n^b = \min_j w_j^b$; then discard the $\lfloor B\alpha \rfloor$ bootstrap realizations b with the lexicographically largest vector $[w]^b$. In other words, we refine problem (3.6) as $\text{lexmin}\{\text{lexmax}_{b \notin D} [w]^b : D \subseteq \{1, \dots, B\}, |D| \leq B\alpha\}$ where lexmin and lexmax denote the minimum and maximum relative to the lexicographic total order of vectors with n components. This rule aims at simultaneously minimizing all the values $\beta(A|X_j)$ without going through extensive additional computations.*

In problem (3.6), we chose to minimize the maximum, over all $j \in \{1, \dots, n\}$ and $A \subseteq \mathcal{Y}$, of the non-discarded bootstrap realizations $\zeta_n^b(A|X_j)$. Other objectives are possible, for example the \mathbb{L}^1 objective $\sum_{b \notin D} w_b$. The main justification for the \mathbb{L}^∞ norm objective $\max_{b \notin D} w_b$ in (3.6) is that it leads to a problem solvable in linear time. In contrast, the problem with an \mathbb{L}^1 objective is computationally difficult, namely NP-hard in the strong sense, as shown in the next result.

Proposition 2. *Minimization of $\{\sum_{b \notin D} w_b : |D| \leq \lfloor B\alpha \rfloor, D \subseteq \{1, \dots, B\}\}$ is NP-hard in the strong sense.*

This result implies that unless $P = NP$, there exists no algorithm for this problem that runs in polynomial time. This is a severe computational drawback relative to the linear-time algorithm achieved with BRS.

3.3. Search in the parameter space. Once the functional quantile has been computed, there remains to search in the parameter space for the values of θ that satisfy (3.2). As shown in the Lemma 1, the function to be optimized in characterization (2.2) of the identified set is supermodular.

Definition 5 (Supermodular function). *A set function $\rho : A \mapsto \rho(A) \in \mathbb{R}$ is called supermodular (resp. submodular) if for all pairs of sets (A, B) , $\rho(A \cup B) + \rho(A \cap B) \geq$ (resp. \leq) $\rho(A) + \rho(B)$.*

Lemma 1. *The function $V \mapsto P(V^\cup | x)$ is submodular for all $x \in \mathcal{X}$.*

In the computation of $\Theta_n(\bar{P}_n)$, it may be desirable to require $\bar{P}_n(V^\cup | x)$ to also be submodular as a function of $V \subseteq \mathcal{U}$, so that the function to be maximized in (3.2) can be maximized using submodular optimization techniques. This can be achieved by adding the following additional linear constraints (see Schrijver (2004)): $\forall u \neq v \in \mathcal{U}, \forall V \subseteq \mathcal{U} \setminus \{u, v\}, j = 1, \dots, n$,

$$\bar{P}_n([V \cup \{u\} \cup \{v\}]^\cup | X_j) - \bar{P}_n([V \cup \{u\}]^\cup | X_j) - \bar{P}_n([V \cup \{v\}]^\cup | X_j) + \bar{P}_n([V]^\cup | X_j) \leq 0. \quad (3.7)$$

The problem of checking whether θ is in the confidence regions can then be solved in polynomial time. Moreover, since submodular optimization has far ranging applications in all areas of operations research, many extremely efficient algorithms and implementations are readily available.

4. SIMULATION BASED ON EXAMPLE 1

We now illustrate and assess the performance of our procedure on the game described in Example 1. Throughout the experiment, we assume that $(\varepsilon_1, \varepsilon_2)$ is uniformly distributed on $[-1/2, 1/2]^2$ and $J = (J_1, J_2)$ is a vector of independent Bernoulli(1/2) random variables. True values for the parameters are indicated with a 0 subscript. We consider the following true parameter specifications: $(\beta_0, c_0) = (0, 0)$ (point identified case) and $(\beta_0, c_0) = (0, 1/4)$ (which corresponds in some sense to the greatest possible indeterminacy). For the true data generating process, we consider two distinct equilibrium selection rules (which, like the true parameter values, are of course supposed unknown in the inference procedure). The first rule specifies that in case of multiplicity, all equilibrium strategy profiles in the equilibrium correspondence are selected with equal probability: we call this case “uniform selection”. The second selection rule specifies that in case of multiplicity, the equilibrium with largest aggregate investment is selected; suppose for instance that the equilibrium correspondence takes the value $\{HH, HL, LO\}$, then equilibrium strategy profile HH is realized: we call this case “maximal selection”. In the case of maximal selection with $c_0 = 0.25$, $\beta_0 = 0$ is assumed known a priori by the analyst performing inference (to avoid an unbounded identified set in the simulations). In the remaining 3 cases, β_0 is unknown a priori. The experiment is run as follows. We calculate in each of the 4 cases above the distribution of the true data generating process. With the latter, we compute the identified set. In the point identified case, the identified set is equal to the true value. In the case $c_0 = 0.25$, with $\beta = 0$ known a priori and maximal selection, the identified set is $[0.2113, 0.3333]$ as explained in the example at the end of Section 2. In case $(c_0 = 0.25, \beta_0 = 0)$ with uniform selection, the identified set projects to $[0, 0.375]$ on the c coordinate and to $[0, 0.320]$

on the β coordinate. We then simulate 5000 samples of sizes $n = 100$, $n = 500$ and $n = 1000$ from this distribution and construct confidence regions for the identified set using lower probabilities \underline{P}_n (based on characterization 2.4), which turned out to have better coverage properties. We use 999 bootstrap replications for the first two sample sizes, and 399 bootstrap replications for $n = 1000$. We consider confidence levels 90%, 95% and 99%. Coverage probabilities of the true value and of the identified set by the confidence region, as computed from the 5000 samples, are displayed in Table 2 for the data generating process obtained with maximal selection and Table 3 for the data generating process obtained with uniform selection. Alongside coverage of the identified set and of the true value, we report the effective level at which Condition (3.5) is satisfied to directly assess the bootstrap functional quantile approximation. Monte Carlo coverage of the identified set is close to the theoretical level in the case of maximal selection and tends to be very high in case of uniform selection. In cases of maximal and uniform selection alike, coverage of Condition (3.5) is almost identical to point coverage in the point identified case ($c_0 = 0$), but lower in the set identified case ($c_0 = 0.25$). Overall the procedure over rejects in all but 13 out of a possible 90 cases. Improvements with sample size occur only in 21 cases (out of a possible 60). These improvements tend to occur when going from $n = 500$ to $n = 1000$ and given the nonparametric procedure, there are doubt as to the accuracy of the procedure for $n = 100$. Finally, the coverage of the true value (as opposed to the whole identified set) is only marginally greater than the coverage of the whole identified set.

5. APPLICATION TO LONG TERM ELDERLY CARE DECISIONS

We estimate the determinants of long term care option choices for elderly parents in American families. The model we use closely follows the one proposed by Engers and Stern (2002) who present these choices as the result of a non family participation game. The family members decide simultaneously whether to participate in a family reunion where the care option maximizing the participants' utility is chosen. Profits are then split among these participants according to some benefit-sharing rule. The data consists of a sample of 1,212 elderly Americans with two children drawn from the National Long Term Care Survey, sponsored by the National Institute of Aging and conducted by the Duke University Center for Demographic Studies under Grant number U01-AG007198, Duke (1999). Elderly people were interviewed in 1984 about their living and care arrangements. The survey questions include gender and age of the children, the distance between homes of the elderly parent and each of the children, the disability status of the elderly parent (where disability is referred to as problems with "Activities of Daily Living or Instrumental Activities of Daily Living (ADL)") and the number of days per week each of the children devotes to the care of the elderly parent. The dependent variable is the care provision for the parent. The parent is asked to list children

TABLE 2. Coverage probabilities of (α_0, β_0) and of the identified set by the confidence region, as computed from 5000 samples. The last column shows the level at which Condition (3.5) is satisfied. Case, where the data generating process obtained with maximal selection.

(α_0, β_0)	n	Level	Point Coverage	Set Coverage	Condition (3.5)
(0, 0)	100	0.99	-	0.9826	0.9792
		0.95	-	0.9574	0.9564
		0.90	-	0.9324	0.9392
	500	0.99	-	0.9894	0.9894
		0.95	-	0.9770	0.9760
		0.90	-	0.9592	0.9584
	1000	0.99	-	0.9714	0.9712
		0.95	-	0.9564	0.9554
		0.90	-	0.9362	0.9352
(0.5, 0)	100	0.99	0.9364	0.9364	0.9286
		0.95	0.9356	0.9354	0.9122
		0.90	0.9232	0.9220	0.8830
	500	0.99	0.9906	0.9902	0.9656
		0.95	0.9810	0.9804	0.9518
		0.90	0.9640	0.9632	0.9330
	1000	0.99	0.9878	0.9870	0.9772
		0.95	0.9746	0.9730	0.9532
		0.90	0.9594	0.9570	0.9210

(either at home or away from home) and how much each provides help. If only one child is listed as providing significant help, that child is designated the primary care giver. If more than one child is listed, the one providing the most time is designated the primary care giver. If the elderly parent lives in a nursing home, then the nursing home is the primary care giver. If no child is listed and the parent does not live in a nursing home, then the parent is designated as “living alone”. Table 4 presents the list of variables used in the analysis. They include parent characteristics, characteristics

TABLE 3. Coverage probabilities of (α_0, β_0) and of the identified set by the confidence region, as computed from 5000 samples. The last column shows the level at which condition (3.5) is satisfied. Case where the data generating process is obtained with uniform selection.

(α_0, β_0)	n	Level	Point Coverage	Set Coverage	Condition (3.5)
(0, 0)	100	0.99	-	0.9872	0.9846
		0.95	-	0.9784	0.9746
		0.90	-	0.9680	0.9652
	500	0.99	-	0.9950	0.9944
		0.95	-	0.9886	0.9872
		0.90	-	0.9814	0.9794
	1000	0.99	-	0.9790	0.9738
		0.95	-	0.9706	0.9640
		0.90	-	0.9628	0.9548
(0.5, 0)	100	0.99	0.9998	0.9986	0.9850
		0.95	0.9998	0.9984	0.9792
		0.90	0.9998	0.9978	0.9704
	500	0.99	1.0000	0.9996	0.9850
		0.95	1.0000	0.9974	0.9664
		0.90	1.0000	0.9980	0.9382
	1000	0.99	1.0000	0.9964	0.9792
		0.95	1.0000	0.9956	0.9694
		0.90	1.0000	0.9938	0.9578

of the children and the care option chosen. A more detailed discussion and summary statistics and additional results can be found in the supplementary material.

5.1. The game. The observable choice of care option is modeled as in Engers and Stern (2002) as the outcome of a family bargaining game. We index family members as follows. Parent: 0, Firstborn child: 1 and Second born child: 2. The payoff to family member i , $i = 0, 1, 2$, is the sum of three terms. The first term V_{ij} is the value to parent 0 and to child i of care option j , where $j \in 1, 2$ means child j becomes the primary care giver, $j = 0$ means the parent remains self-reliant and $j = 3$, the parent is moved to a nursing home. The matrix $V = (V_{ij})_{ij}$ is known to both children and the

Variables	Equal to 1 if:	Percentage of sample
Care Option		
	Living with child 1	26.81
	Living with child 2	6.75
	Living in nursing home	19.92
	Living home alone	46.54
Parent Variables		
DA	Highly disabled	33.81
DM	Living with the spouse	40.36
Children Variables		
DD	Living with parent	11.55
$DD1$	Distance from parent: 31 min and more	49.45
DS	Female	49.26

TABLE 4. List of variables.

parent. We suppose it takes the form

$$V_{ij} = \gamma_{ij} + W\beta_{ij} + Z_j\psi_{ij}$$

where W indicates the characteristics of the parents (DA and DM), and Z_j indicates the characteristics of care option j (DS , $DD1$ and $DD2$) and $X = (W, Z)$. $\theta = (\gamma_{ij}, \beta_{ij}, \psi_{ij})'$ is unknown to the analyst and the object of inference.

Example 2. Consider the following family, in which the matrix where given value of X and θ result in V that takes the form:

$$V = \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 4 & -1 & 1 \\ 0 & -1 & 4 & 1 \end{bmatrix}$$

Rows indicate family member $i = 0, 1, 2$, and columns represents care giving options $j = 0, 1, 2, 3$, in that order. In this example, the parent is indifferent between all the care options, except the one where she has to move to the Nursing home. Each child prefers to be the primary care giver to any other care option, followed by the parent living in a nursing home, living at home and being taken care of by the other child, in that order.

The second term in the payoff results from the family bargaining process as follows. We assume that it is always in the interest of the parent to attend the family reunion. However, child i ($i = 1, 2$) can refrain from participating in the meeting. By choosing not to participate, a member of the family agrees on whatever is decided but can neither assume the role of primary care giver, nor can he be involved in any side payment. Both children simultaneously decide whether or not to participate in the long term care decision. Suppose M is the set of children who participate. The option chosen is option $j \in M \cup \{0, 3\}$ which maximizes the participants's total utility $\sum_{i \in M} V_{ij}$. It is assumed that participants abide by the decision and that benefits are then shared equally among parent and children participating in the decision through a monetary transfer s_i , which is the second term in the children's payoff. The third term ϵ_i in the payoff is a random benefit from participation, which is 0 for children who decide not to participate and distributed according to absolutely continuous distribution $\nu(\cdot|\theta)$ for each child who participates. All children observe the realizations of ϵ , whereas the analyst only knows its distribution. The Payoff matrix is given in Table 5, where overall benefit shares w_i^{IJ} , $i = 1, 2$, $I, J = N, P$ are defined and derived in the supplementary appendix. Multiple Nash equilibria in pure and mixed strategies are also derived in the appendix. Each equilibrium action profile results in a (almost surely) unique care option choice, hence for each participation shock ϵ , we can derive $G(\epsilon|X; \theta)$ as the set of probability measures on the set of care options $\{0, 1, 2, 3\}$ induced by mixed strategy profiles, which are probabilities on the set of participation profiles $\{NN, NP, PN, PP\}$.

Child 1	Child 2	
	N	P
N	w_1^{NN}, w_2^{NN}	$w_1^{NP}, \epsilon_2 + w_2^{NP}$
P	$\epsilon_1 + w_1^{PN}, w_2^{PN}$	$\epsilon_1 + w_1^{PP}, \epsilon_2 + w_2^{PP}$

TABLE 5. Payoffs for the family participation game.

5.2. Specification. We provide estimates for the following utility specification (an alternative with altruistic utility specification was estimated and results are reported in the supplementary material).

$$V(X; \theta) = \begin{bmatrix} \begin{pmatrix} \beta_{00} \\ +\beta_m DM \\ +\beta_{ah} DA \end{pmatrix} & \begin{pmatrix} \alpha \\ +\psi_s DS_1 \end{pmatrix} & \psi_s DS_2 & 0 \\ \begin{pmatrix} \beta_m DM \\ +\beta_{ah} DA \end{pmatrix} & \begin{pmatrix} \beta_{11} \\ +\psi_1 DD_1 \\ +\beta_{ac} DA \end{pmatrix} & 0 & 0 \\ \begin{pmatrix} \beta_m DM \\ +\beta_{ah} DA \end{pmatrix} & 0 & \begin{pmatrix} \beta_{11} \\ +\psi_1 DD_2 \\ +\beta_{ac} DA \end{pmatrix} & 0 \end{bmatrix}$$

Recall that the columns indicate the options, in the following order $\{0, 1, 2, 3\}$, and the rows represent each member of the family, in the following order Parent, Child 1, Child 2. For example, the value the first born child (family member 1) living less than 30 minutes away from the parent's home attaches to the fact that she takes care of a non disabled, non-married parent is measured by β_{11} , whereas for a disabled parent, it is $\beta_{11} + \beta_{ac}$.

5.3. Estimation methodology. The methodology proposed in the paper allows the construction of the identified set based on the hypothetical knowledge of the true distribution of the data. As described in Section 3, we account for sampling uncertainty and control the level of confidence by constructing set functions $A \mapsto \overline{P}(A|X)$, which dominate $P(A|X)$ (uniformly over $A \subseteq \{0, 1, 2, 3\}$ and X) with probability $1 - \alpha$ (the chosen level of confidence, here 0.95). We implement the method detailed in Section 3 (except that the *pairs* or *cases* bootstrap was used instead of the nonparametric bootstrap advocated above) with a number of bootstrap replications $B = 2500$. Second, we obtain the model likelihood by simulating the valuation matrix and computing the Equilibrium correspondence from the payoff matrix, for given values of X and θ . The procedure is as follow. For a given X and θ ,

- We generate and store R draws of ε from the distribution ν_θ . Here, $R = 5000$ and ν_θ is normally distributed with mean μ and variance σ_ε^2 , where $(\mu, \sigma_\varepsilon^2)$ belong to the parameter θ .
- For each value ε^r , we compute the valuation matrix $V(X, \varepsilon^r, \theta)$ and the corresponding payoff matrix.
- Then, we determine the equilibrium correspondence $G(X, \varepsilon; \theta)$ from the analytical results derived in the preceding section. The Gambit software provides an alternative for computing numerically the set NE for more complex games.

- The last step of the simulation is to compute an estimator of the model likelihood \mathcal{L} defined in (3.3) as follows: $\hat{\mathcal{L}}(A|X;\theta) = \frac{1}{R} \sum_{r=1}^R \min\{\sigma(A) : \sigma \in G(X, \varepsilon^r; \theta)\}$.

Having constructed those two elements, the identified set comprises all values of θ such that for all observed values of the explanatory variables, the minimum over $A \subseteq \{0, 1, 2, 3\}$ of the function $\bar{P}(A|X;\theta) - \hat{\mathcal{L}}(A|X;\theta)$ is non negative, as explained in Section 3. We construct an n-dimensional grid to conduct the search over the parameter space. Each value of the parameter can be tested in a fraction of a second on a standard laptop, and a region of small dimensionality (1 to 4) can be constructed in a few hours, again on a standard laptop without parallel processing. However, estimation time grows exponentially with the number of parameters induced by the model. In our case, each specification involves a 12-dimensional parameter space. Parallel processing becomes therefore necessary. We use an Open-MP procedure for parallel processing, which is perfectly suited to the method we propose. The computation resources have been provided by the Réseau Québécois de Calcul de Haute Performance (RQCHP). All computation where made under the system ‘‘Cottos’’ which provides up to 128 computation nodes (1024 CPU cores) equipped with two Intel Xeon E5462 quad-core processors at 3 GHz. Under 1 node, approximately 10^7 parameters points can be tested in 24 hours.

5.4. Results. We perform the estimation under different values of the mean and variance of the error term. To alleviate the computational burden, we first test the significance of some of the individual parameters by checking whether the hyper planes defined by $\theta_i = 0$ - where θ_i is a component of θ - intersect the 95% confidence region. We fail to reject the Null Hypothesis if the estimation procedure returns a non-empty set. We then obtain a constrained confidence region for the remaining parameters. For each value of mean and variance of the error term, we find a non empty intersection between the confidence region and the hyperplane defined by $\beta_{11} = 0$. This means we fail to reject (at the 5% level) the null hypothesis that there is no additional constant disutility for a child to take care of an elderly parent. Since, this hypothesis is not rejected, we obtain a constrained confidence region for the remaining parameters. We then obtain confidence regions for different values of β_{11} and discuss the latter’s effect on the regions. We note that the Null hypothesis $H_0 : \beta_{00} = 0$ is always rejected. Hence, when we control for all other effects, parents are not indifferent between the first two options. They show a clear preference in favor of living in their own home (option called ‘‘living alone’’) instead of living in a nursing home (β_{00} is always positive). The results we present are then for given values of β_{00} . We provide an insight of how different values of this parameter change the results. We report the range for each parameters in Table 6. Note that the identified set is not a compact set. In particular, β_{ac} , β_{ah} , β_m and ψ are allowed to diverge to $-\infty$. Results are generally consistent with expectations and previous results

Parameters	Min	Max	Min	Max	Min	Max	Min	Max
β_{00}	2	2	3	3	1	1	1	1
β_{11}	0	0	0	0	0	0	0	0
β_{ah}	$-\infty$	-3.57	$-\infty$	-3.57	$-\infty$	-2.86	$-\infty$	-2.14
$\beta_{ac} = -\beta_m$	$-\infty$	-2.86	$-\infty$	-2.86	$-\infty$	-3.57	$-\infty$	-3.57
α	0.00	8.00	0.00	8.00	1.00	5.00	0.00	4.00
ψ_s	0.00	5.00	0.00	4.00	1.00	4.00	0.00	2.00
ψ_d	$-\infty$	-2.86	$-\infty$	-2.14	$-\infty$	-1.43	$-\infty$	-3.57
μ	-1	-1	0	0	-1	-1	0	0
σ_ε	1	1	1	1	1	1	1	1
σ_u^2	1	1	1	1	0.25	0.25	0.25	0.25
$p\xi$	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1

TABLE 6. Parameters Range for estimation of Specification 1 at $\beta_{11} = 0$, $\beta_{ac} = -\beta_m$ and for different values of the error terms and of β_{00} .

on the subject. Namely:

- (1) The existence of several problems with the parent's functional ability is a key determinant of the decision to enter a nursing home. β_{ah} and β_{ac} are both negative and can both be (very) large. The negative sign of β_{ah} captures the fact that a parent's disability increases the value of care provided by the family or a specialized institution. In addition, $\beta_{ac} < 0$ means that the disability entails a utility cost for the child if he is chosen as primary care giver.
- (2) Parameter β_m associated with the parent living with a spouse is positive and large. This implies that married parents are more likely to remain self-reliant. In families where the parent is disabled, the effect of living with the spouse compensates the disutility of disability and preserves the incentive for parents to live at home.
- (3) While we cannot rule out parents being indifferent to the gender or birth order of their primary care giver, estimation shows a tilt of the confidence interval toward positive values for both parameters, with a possible positive and large magnitude of the parameter α . In case $\mu = -1$ and $\sigma_u^2 = 0.25$, the data reveal that parents exhibit a preference for an older and for a female care giver.

- (4) Children living more than 30 minutes from the parents are less likely to provide care than those living closer to the parents. Distance has a (possibly strong) disutility effect on children's incentives to participate in the care decision.

The shape of the confidence region also conveys a considerable amount of information. Figure 3 shows two dimensional dimensional projections and cuts of the confidence region for column 2 of Table 6, i.e $\mu_\varepsilon = 0$, $\sigma_\varepsilon^2 = 1$, $\sigma_u^2 = 1$. Of great interest is the projection of the identified set in the plan β_{ah}, β_m . Figure 3(a) reveals an almost linear relation between the two parameters of the type $\beta_{ah} = -\beta_m$. The estimation rejects models for which the absolute value of the two parameters are significantly different. The data suggest therefore that the disutility induced by the disability of the parent can be entirely compensated by the presence of a spouse in the same household. Notice the triangular shape of the region plotted in Figure 3(b) which entails that simultaneous large values of ψ_s and α are rejected. This finding means that only one of the effects (gender or birth order) can be large, not both. In other words, firstborn daughter are not the only possible care givers. Note also that both effects can be very small, though not jointly insignificant. We observe similar types of constraints for the pairs (α, β_{ah}) , (α, β_{ac}) , (α, ψ_d) , (ψ_s, β_{ac}) , (ψ_s, ψ_d) as large values of parameters α or ψ_s are only permitted when the other parameters are jointly large (see Figure 3(c) to 3(f)). For example, we obtain a constrained confidence region at $\beta_{ac} = -3.5$. The ranges for the two parameters, α and ψ_s , are tighter, as $\alpha \in [1, 2]$ and $\psi \in [0, 1]$. Figure 4 shows the effect of the variation of parameter β_{11} on ψ_s and α . Recall that β_{11} represents a fixed cost or benefit for the child chosen as care giver. We observe negative relations between β_{11} and ψ_s , and β_{11} and α . Negative values of ψ_s and α are only admissible for positive values of β_{11} . Hence a model where parents exhibit no favoritism for a daughter and/or a firstborn, or favoritism for a son and/or a second born, will be consistent with our data if and only if there exist a strictly positive constant benefit for a child to be caregiver.

CONCLUSION

We have considered the problem of statistical inference in incomplete partially identified structural models, such as models of discrete choice with interactions and other forms of endogeneity. A characterization of the identified set for structural parameters was given, with an appeal to a classical theorem in combinatorial optimization, the Max-Flow Min-Cut Theorem, thereby emphasizing the optimization formulation of the problem of inference in such models. Finally, we have shown how to apply combinatorial optimization methods within a bootstrap procedure in order to compute informative confidence regions very efficiently, hence feasibly in empirically relevant applications. An application of the methodology was carried out on a family bargaining example and

it was shown that most findings in the literature on the determinants of long term elderly care by American families were supported in this more robust framework, where the effects of interaction are accounted for. This procedure applies to very general classes of models and its efficiency and coverage properties could no doubt be improved, when tailored to more specific applications. In particular, the application to matching games and revealed preference testing of stability in matching still poses considerable challenges. Other perspectives for further work include the application of Max-Flow Min-Cut algorithms to the detection of redundant inequalities at the identification stage, to improve the performance at the inference stage, possibly by appealing to other existing procedures if the number of non redundant inequalities is small enough.

APPENDIX A. PROOFS OF RESULTS IN THE MAIN TEXT

Proof of Theorem 1. By Proposition 1 of Galichon and Henry (2011), a value θ of the parameter vector belongs to $\Theta_I(P)$ if and only if $\mathbb{P}(Y \in G(X, \varepsilon; \theta)) = 1$, X -a.s. (which we drop from the notation from this point on). Hence if there exists a pair (Y, U) of random vectors on $\mathcal{Y} \times \mathcal{U}$ such that Y has probability mass $P(y|X)$, $y \in \mathcal{Y}$, U has probability mass $Q(u|X; \theta)$, $u \in \mathcal{U}$, and $\mathbb{P}(Y \in U|X) = 1$. This is equivalent to the existence of non negative weights π_y^u , $(y, u) \in \mathcal{Y} \times \mathcal{U}$, such that $\sum_{u \in \mathcal{U}} \pi_y^u = P(y|X)$, $\sum_{y \in \mathcal{Y}} \pi_y^u = Q(u|X)$, and $\pi_y^u = 0$ when $y \notin u$. The latter is equivalent to the following programming problem with auxiliary variables a_y , $y \in \mathcal{Y}$ and a^u , $u \in \mathcal{U}$ having zero as a solution. The programming problem is the following: $\min(\sum_{y \in \mathcal{Y}} a_y + \sum_{u \in \mathcal{U}} a^u)$ subject to the constraints $\sum_{u \in \mathcal{U}} \pi_y^u + a_y \leq P(y|X)$, $\sum_{y \in \mathcal{Y}} \pi_y^u + a^u \leq Q(u|X; \theta)$, $a_y, a^u, \pi_y^u \geq 0$, and $\pi_y^u = 0$ when $y \notin u$. Since $\sum_{y \in \mathcal{Y}} a_y + \sum_{u \in \mathcal{U}} a^u \leq \sum_{y \in \mathcal{Y}} P(y|X) + \sum_{u \in \mathcal{U}} Q(u|X; \theta) - 2 \sum_{y \in \mathcal{Y}} \sum_{u \in \mathcal{U}} \pi_y^u = 2 - 2 \sum_{y \in \mathcal{Y}} \sum_{u \in \mathcal{U}} \pi_y^u$, the latter is also equivalent to $\max \sum_{y \in \mathcal{Y}} \sum_{u \in \mathcal{U}} \pi_y^u \geq 1$ subject to the constraints $\sum_{u \in \mathcal{U}} \pi_y^u \leq P(y|X)$, $\sum_{y \in \mathcal{Y}} \pi_y^u \leq Q(u|X)$, $\pi_y^u \geq 0$ and $\pi_y^u = 0$ when $y \notin u$. This is called a maximum flow problem, i.e. the problem of maximizing quantity flowing through a network under capacity constraints. A network is a collection of nodes, including a *source* S and a *sink* T , and directed edges between the nodes. For instance, (N_1, N_2) is an edge leading from node N_1 to node N_2 . Here the network involved in the maximum flow problem is comprised of a source S , K nodes corresponding to the K elements of \mathcal{Y} , J nodes corresponding to the J elements of \mathcal{U} and a sink T . The source S is connected to each of the nodes y_1, \dots, y_K in \mathcal{Y} . A node $y \in \mathcal{Y}$ is connected to a node $u \in \mathcal{U}$ if and only if $y \in u$. All nodes u_1, \dots, u_J in \mathcal{U} are connected to the sink T . To each edge is attached a capacity, which is the maximum amount that can flow through it. Capacity is constrained to $P(y|X)$ between S and node y . Capacity is unconstrained (i.e. infinite) between node y and node u such that $y \in u$. The capacity of edges between a node u and the sink T is constrained to $Q(u|X; \theta)$.

We have shown that $\theta \in \Theta_I$ if and only if the maximum flow in the network described above is equal to 1. We now appeal to a classical result in combinatorial optimization called the *Max-Flow Min-Cut Theorem*, see for instance Theorem 10.3 page 150 of Schrijver (2004). A *cut* through a network is partition of the nodes into two sets separating the source from the sink. The *capacity* of a cut is defined as the sum of the capacities of edges in the network that cross the cut from the source side to the sink side. Let a cut be defined by the set V of elements of \mathcal{U} and the set Z of elements of \mathcal{Y} on the sink side of the cut. Since the capacity of an edge from y to u such that $y \in u$ is infinite, the cut defined by V and Z has finite capacity if and only if $y \in u$ and $u \in V$ jointly imply $y \in Z$. Such a cut has capacity $C(Z, V) = \sum_{y \in Z} P(y|X) + \sum_{u \in \mathcal{U} \setminus V} Q(u|X; \theta) = \sum_{y \in Z} P(y|X) + 1 - \sum_{u \in V} Q(u|X; \theta)$. A cut has minimum capacity if no node can be moved between the source side of the cut and the sink side of the cut without increasing capacity, hence if $y \notin u$ and $u \in V$ jointly imply $y \notin Z$, hence if $Z = V^\cup = \bigcup\{u : u \in V\}$. Therefore, the capacity of a minimum cut is $C(V^\cup, V) = \sum_{y \in V^\cup} P(y|X) + 1 - \sum_{u \in V} Q(u|X; \theta) = P(V^\cup|X) + 1 - Q(V|X; \theta)$. By the Max-Flow Min-Cut Theorem, the capacity of any minimum cut is equal to the maximum flow through the network, hence $\theta \in \Theta_I(P)$ if and only if for all subset V of \mathcal{U} , $P(V^\cup|X) + 1 - Q(V|X; \theta) \geq 1$, i.e. $Q(V|X; \theta) \leq P(V^\cup|X)$, and the result follows. \square

Proof of Lemma 1. Take an $x \in \mathcal{X}$. Take any $u \in \mathcal{U}$ and $V \subseteq \mathcal{U} \setminus \{u\}$. We have $P([V \cup \{u\}]^\cup|x) - P(V^\cup|x) = \sum_{y \in \bigcup_{v \in V \cup \{u\}} v} P(y|x) - \sum_{y \in \bigcup_{v \in V} v} P(y|x) = \sum_{y \in u \setminus V^\cup} P(y|x) = P(u \setminus V^\cup|x)$, which is non-increasing in V , hence the result. \square

Proof of Theorem 2. Given a value $\theta \in \Theta_I$, by Theorem 1, we have $\sup_{x \in \mathcal{X}} \max_{V \subseteq \mathcal{U}} (Q(V|x; \theta) - P(V^\cup|x)) \leq 0$. Under Assumption 1, $\sup_{x \in \mathcal{X}} \max_{V \subseteq \mathcal{U}} (P(V^\cup|x) - \bar{P}_n(V^\cup|x)) \leq 0$, with limiting probability larger than $1 - \alpha$. Hence, with probability at least $1 - \alpha$, $\sup_{x \in \mathcal{X}} \max_{V \subseteq \mathcal{U}} (Q(V|x; \theta) - \bar{P}_n(V^\cup|x)) \leq 0$, and thus $\theta \in \Theta_I(\bar{P}_n)$. \square

Proof of Proposition 1. We first justify the BRS Step 1 by showing that $w_b = w'_b$ for all b . Indeed observe that for any $j \in \{1, \dots, n\}$ and $A \subseteq \mathcal{Y}$, we have

$$\zeta_n^b(A|X_j) = \sum_{y \in A} \hat{P}_n(y|X_j) - \sum_{y \in A} \hat{P}_n^b(y|X_j) = \sum_{y \in A} [\hat{P}_n(y|X_j) - \hat{P}_n^b(y|X_j)]$$

and thus $\max_{A \subseteq \mathcal{Y}} \zeta_n^b(A|X_j)$ is attained by selecting all the elements $y \in \mathcal{Y}$ with $\hat{P}_n(y|X_j) - \hat{P}_n^b(y|X_j) > 0$. It follows that $w'_b = \max_{1 \leq j \leq n} \max_{A \subseteq \mathcal{Y}} [\sum_{y \in \mathcal{Y}} \hat{P}_n(y|X_j) - \sum_{y \in A} \hat{P}_n^b(y|X_j)]$ and therefore $w_b = w'_b$. To justify BRS Step 2, let w^{opt} denote the optimum objective value of problem (3.6). If D fails to include any b such that $w_b > w^{opt}$ then $\max_{b \notin D} w_b > w^{opt}$, therefore an optimal D must include all b such that $w_b > w^{opt}$. On the other hand, if D is any optimal subset and some $b' \in D$ satisfies $w_{b'} \leq w^{opt}$ then discarding b' from D yields a feasible subset $D \setminus \{b'\}$

(since $|D \setminus \{b'\}| < |D| \leq \bar{d}$) such that $\max_{b \in D \setminus \{b'\}} w_b \leq \max_{b \in D} w_b$ hence $D \setminus \{b'\}$ is an alternate optimal solution. Therefore an optimal D consists of all indices b such that $w_b > w^{opt}$. Concerning the running time, BRS Step 1 requires $O(nB|\mathcal{Y}|)$ time, and BRS Step 2 requires $O(B)$ time using a linear time selection (or median-finding) algorithm (see Blum, Floyd, Pratt, Rivest, and Tarjan (1973)). \square

Proof of Proposition 2. The problem corresponds to the following decision problem: given an $n \times m$ matrix H , an integer k and a target value t , can one find a subset $S \subseteq \{1, \dots, n\}$ such that $|S| \geq k$ and $\sum_{i=1}^m \max_{j \in S} H_{ij} \leq t$? Denote (H, k, t) an instance of the latter problem. Consider the well-known NP-hard decision problem CLIQUE (see for instance section 4.8 page 43 of Schrijver (2004)): given a graph $G = (V, E)$ and an integer q satisfying $2 \leq q \leq |V|$, does there exist a subset $Q \subseteq V$ such that $|Q| \geq q$ and for all $i, j \in V$, $ij \in E$ (i.e. Q is a *clique*). To any instance (G, q) of the problem CLIQUE, we associate an instance (H, k, t) of our decision problem, where lines of H corresponds to vertices of G (elements of V), columns of H corresponds to edges in G (elements of E) and $H_{ij} = 1$ if vertex i belongs to edge j , and 0 otherwise. For any subset $S \subseteq E$ of edges in G , we have for all $i \in V$, $\max_{j \in S} H_{ij} = 1$ if i belongs to at least one element of S , and 0 otherwise. Hence, $\sum_{i \in V} \max_{j \in S} H_{ij}$ is the number of vertices that belong to at least one edge in S . Define $k = q(q-1)/2$ and $t = q$. Then, a set S of k edges involves at least (hence exactly) q vertices if and only if S is the set of edges of a CLIQUE. Hence the answer to the decision problem (H, k, t) thus defined is YES if and only if G contains a CLIQUE with q vertices. Since CLIQUE is NP-complete, it follows that our decision problem is NP-hard. Since $k = O(|V|^2)$ and $t = O(|V|)$, the input size (in unitary notation) of such instances of our problem is polynomially bounded by the input size (in unitary or binary notation) $\Omega(|V|)$ of the corresponding instance of CLIQUE. Hence our decision problem is NP-hard in the strong sense. \square

REFERENCES

- AHUJA, R. K., T. L. MAGNANTI, and J. B. ORLIN (1993): *Network Flows : Theory, Algorithms, and Applications*. Prentice Hall.
- ANDREWS, D., S. BERRY, and P. JIA (2003): “Placing bounds on parameters of entry games in the presence of multiple equilibria,” unpublished manuscript.
- ANDREWS, D., and G. SOARES (2010): “Inference for Parameters Defined by Moment Inequalities Using Generalized Moment Selection,” *Econometrica*, 78, 119–157.
- ARADILLAS-LOPEZ, A. (2010): “Semiparametric estimation of a simultaneous game with incomplete information,” *Journal of Econometrics*, 157, 409–431.

- BAJARI, P., J. HAHN, H. HONG, and G. RIDDER (2011): “A note on semiparametric estimation of finite mixtures of discrete choice models with applications to game theoretic models,” *International Economic Review*, 52, 807–824.
- BAJARI, P., H. HONG, and S. RYAN (2010): “Identification and estimation of a discrete game of complete information,” *Econometrica*, 78, 1529–1568.
- BERESTEANU, A., I. MOLCHANOV, and F. MOLINARI (2011): “Sharp identification regions in models with convex predictions,” *Econometrica*, 79, 1785–1821.
- BERRY, S. (1992): “Estimation of a model of entry in the airline industry,” *Econometrica*, 60, 889–917.
- BERRY, S., and E. TAMER (2006): “Identification in models of oligopoly entry,” in *Advances in Economics and Econometrics*, pp. 46–85. Cambridge University Press.
- BICKEL, P., and D. FREEDMAN (1981): “Some asymptotic theory for the bootstrap,” *Annals of Statistics*, 9, 1196–1217.
- BLUM, M., R. FLOYD, V. PRATT, R. RIVEST, and R. TARJAN (1973): “Time bounds for selection,” *Journal of Computational Systems Science*, 7, 448–461.
- BRESNAHAN, T., and P. REISS (1990): “Entry in monopoly markets,” *Review of Economic Studies*, 57, 531–553.
- BUGNI, F. (2010): “Bootstrap inference in partially identified models defined by moment inequalities: coverage of the identified set,” *Econometrica*, 78, 735–753.
- CHERNOZHUKOV, V., H. HONG, and E. TAMER (2007): “Estimation and Confidence Regions for Parameter Sets in Econometric Models,” *Econometrica*, 75, 1243–1285.
- CHERNOZHUKOV, V., S. LEE, and A. ROSEN (2009): “Inference on intersection bounds,” CEMMAP Working Paper CWP19/09.
- CHESHER, A. (2010): “Instrumental variable models for discrete outcomes,” *Econometrica*, 78, 575–601.
- CHESHER, A., A. ROSEN, and K. SMOLINSKI (2011): “An instrumental variable model of multiple discrete choice,” CEMMAP Working Paper CWP06/11.
- CILIBERTO, F., and E. TAMER (2009): “Market structure and multiple equilibria in airline markets,” *Econometrica*, 70, 1791–1828.
- DUKE (1999): “National Long Term Care Survey,” Public use data set produced and distributed by the Duke University Center for Demographic Studies with funding from the National Institute on Aging under Grant No. U01-AG007198.
- ECHENIQUE, F., S. LEE, M. SHUM, and B. YENMEZ (2011): “The revealed preference theory of stable and extremal stable matchings,” unpublished manuscript.

- EKELAND, I., A. GALICHON, and M. HENRY (2010): “Optimal transportation and the falsifiability of incompletely specified economic models,” *Economic Theory*, 42, 355–374.
- ENGERS, M., and S. STERN (2002): “Family bargaining and long term care,” *International Economic Review*, 43, 73–114.
- GALICHON, A., and M. HENRY (2006): “Dilation Bootstrap,” unpublished manuscript.
- GALICHON, A., and M. HENRY (2009): “A test of non-identifying restrictions and confidence regions for partially identified parameters,” *Journal of Econometrics*, 152, 186–196.
- GALICHON, A., and M. HENRY (2011): “Set identification in models with multiple equilibria,” *Review of Economic Studies*, 78, 1264–1298.
- HAILE, P., and E. TAMER (2003): “Inference with an Incomplete Model of English Auctions,” *Journal of Political Economy*, 111, 1–51.
- HENRY, M., and I. MOURIFIÉ (2011): “Euclidean revealed preferences: testing the spatial voting model,” *Journal of Applied Econometrics*, forthcoming.
- HENRY, M., and A. ONATSKI (2011): “Set coverage and robust policy,” *Economics Letters*, forthcoming.
- JIA, P. (2008): “What Happens When Wal-Mart Comes to Town: An Empirical Analysis of the Discount Retailing Industry,” *Econometrica*, 76, 1263–1316.
- JOVANOVIĆ, B. (1989): “Observable implications of models with multiple equilibria,” *Econometrica*, 57, 1431–1437.
- MANSKI, C. (1993): “Identification of endogenous social effects: the reflection problem,” *Review of Economic Studies*, 60, 531–542.
- PAKES, A., J. PORTER, K. HO, and J. ISHII (2004): “Moment inequalities and their application,” unpublished manuscript.
- ROMANO, J., and A. SHAIKH (2008): “Inference for identifiable parameters in partially identified econometric models,” *Journal of Statistical Planning and Inference*, 139, 2786–2807.
- ROMANO, J., and A. SHAIKH (2010): “Inference for the identified set in partially identified econometric models,” *Econometrica*, 78, 169–211.
- ROSEN, A. (2008): “Confidence sets for partially identified parameters that satisfy a finite number of moment inequalities,” *Journal of Econometrics*, 146, 107–117.
- SCHRIJVER, A. (2004): *Combinatorial optimization: polyhedra and efficiency*. Springer: Berlin.
- SINGH, K. (1981): “On the asymptotic accuracy of Efron’s bootstrap,” *Annals of Statistics*, 9, 1187–1195.
- SOETEVEENT, A., and KOOREMAN (2007): “A discrete choice model with social interactions: with an application to high school teen behaviour,” *Journal of Applied Econometrics*, 22, 599–624.

- TAMER, E. (2003): “Incomplete simultaneous discrete response model with multiple equilibria,” *Review of Economic Studies*, 70, 147–165.
- UETAKE, K., and Y. WATANABE (2011): “Entry by acquisition: estimates from a two-sided matching model with externality,” unpublished manuscript.







(a) (β_{ah}, β_m) region(b) (α, ψ_s) region(c) (α, β_{ac}) region(d) (α, ψ_d) region(e) (ψ_s, β_{ac}) region(f) (ψ_s, ψ_d) region

FIGURE 3. Two dimensional representations of the confidence region at $\beta_{00} = 3$, $\beta_{11} = 0$, $\mu = 0$, $\sigma_\varepsilon = 1$, $\sigma_u = 1$, $p_\xi = 0.1$


 (a) (α, β_{11}) region


 (b) (ψ_s, β_{11}) region

 (c) (α, ψ_s) regions for different values of β_{11}

FIGURE 4. Parameter β_{11} in relation with other parameters: $\beta_{00} = 3$, $\mu = 0$, $\sigma_\varepsilon = 1$, $\sigma_u = 1$, $p_\xi = 0.1$