

Der Open-Access-Publikationsserver der ZBW – Leibniz-Informationzentrum Wirtschaft
The Open Access Publication Server of the ZBW – Leibniz Information Centre for Economics

Hermeling, Claudia; Mennel, Tim

Working Paper

Sensitivity Analysis in Economic Simulations: A Systematic Approach

ZEW Discussion Papers, No. 08-068

Provided in cooperation with:

Zentrum für Europäische Wirtschaftsforschung (ZEW)

Suggested citation: Hermeling, Claudia; Mennel, Tim (2008) : Sensitivity Analysis in Economic Simulations: A Systematic Approach, ZEW Discussion Papers, No. 08-068, <http://hdl.handle.net/10419/24764>

Nutzungsbedingungen:

Die ZBW räumt Ihnen als Nutzerin/Nutzer das unentgeltliche, räumlich unbeschränkte und zeitlich auf die Dauer des Schutzrechts beschränkte einfache Recht ein, das ausgewählte Werk im Rahmen der unter

→ <http://www.econstor.eu/dspace/Nutzungsbedingungen> nachzulesenden vollständigen Nutzungsbedingungen zu vervielfältigen, mit denen die Nutzerin/der Nutzer sich durch die erste Nutzung einverstanden erklärt.

Terms of use:

The ZBW grants you, the user, the non-exclusive right to use the selected work free of charge, territorially unrestricted and within the time limit of the term of the property rights according to the terms specified at

→ <http://www.econstor.eu/dspace/Nutzungsbedingungen>
By the first use of the selected work the user agrees and declares to comply with these terms of use.

Discussion Paper No. 08-068

**Sensitivity Analysis
in Economic Simulations –
A Systematic Approach**

Claudia Hermeling and Tim Mennel

ZEW

Zentrum für Europäische
Wirtschaftsforschung GmbH

Centre for European
Economic Research

Discussion Paper No. 08-068

**Sensitivity Analysis
in Economic Simulations –
A Systematic Approach**

Claudia Hermeling and Tim Mennel

Download this ZEW Discussion Paper from our ftp server:

<ftp://ftp.zew.de/pub/zew-docs/dp/dp08068.pdf>

Die Discussion Papers dienen einer möglichst schnellen Verbreitung von neueren Forschungsarbeiten des ZEW. Die Beiträge liegen in alleiniger Verantwortung der Autoren und stellen nicht notwendigerweise die Meinung des ZEW dar.

Discussion Papers are intended to make results of ZEW research promptly available to other economists in order to encourage discussion and suggestions for revisions. The authors are solely responsible for the contents which do not necessarily represent the opinion of the ZEW.

Executive Summary

This paper gives a systematic introduction to sensitivity analysis in economic simulations. Sensitivity analysis studies how the variation in the numerical output of a model can be quantitatively apportioned to different sources of variation in basic input parameters. It thus provides a check of robustness of the numerical results of a model. We present a formalisation of the deterministic and the stochastic approach to sensitivity analysis. The former assumes that the basic economic parameter stems from a known interval (in higher dimensions: a compact set) and quantifies the spread of the corresponding equilibrium output variables. The latter treats the parameter as a stochastic variable with known distribution and calculates mean and variance of output variables accordingly. Deterministic sensitivity analysis can be numerically implemented by peacemeal formulae. Stochastic sensitivity analysis is implemented by a Monte-Carlo or a Gauss-Quadrature algorithm. We develop an improved version of Gauss-Quadrature based on orthogonal polynomials. In an application of both algorithms to a CGE model we demonstrate its advantages: It significantly reduces the computational burden of sensitivity analysis and tends to be more stable numerically.

Das Wichtigste in Kürze

Dieses Discussion Paper ist eine systematische Anleitung zur Sensitivitätsanalyse bei ökonomischen Simulationen. Eine Sensitivitätsanalyse untersucht, wie die Variation in den numerischen Ergebnissen eines Modells der Variation verschiedener Grundparameter zugeordnet werden kann. Auf diese Weise ermöglicht sie eine Überprüfung der Robustheit numerischer Modellergebnisse. Wir stellen eine Formalisierung des deterministischen und des stochastischen Ansatzes zur Sensitivitätsanalyse vor. Bei ersterem geht man davon aus, dass der ökonomische Grundparameter aus einem bekannten Intervall (in höheren Dimensionen: aus einer kompakten Menge) stammt und man quantifiziert die Ausdehnung der zugehörigen Modellvariablen im Gleichgewicht. Bei letzterem behandelt man den Grundparameter als stochastische Variable mit bekannter Verteilung und berechnet entsprechend Erwartungswert und Varianz der Modellvariablen. Deterministische Sensitivitätsanalyse kann numerische durch Stückformeln (engl. "piece-meal formulae") implementiert werden. Stochastische Sensitivitätsanalyse wird durch das Monte-Carlo Verfahren oder Gauss-Quadratur implementiert. Wir entwickeln eine verbesserte Version des Gauss-Quadratur-Verfahrens, die auf orthogonalen Polynomen beruht. In einer Anwendung der beiden Verfahren auf ein allgemeines Gleichgewichtsmodell zeigen wir seine Vorteile: Es reduziert den numerischen Aufwand der Sensitivitätsanalyse erheblich und ist im allgemeinen numerisch stabiler.

Sensitivity analysis in economic simulations - a systematic approach

Claudia Hermeling
Tim Mennel*

Center for European Economic Research (ZEW)
P.O. Box 103443
D-68034 Mannheim, Germany

September 2008

Abstract

Sensitivity analysis studies how the variation in the numerical output of a model can be quantitatively apportioned to different sources of variation in basic input parameters. Thus, it serves to examine the robustness of numerical results with respect to input parameters, which is a prerequisite for deriving economic conclusions from them. In practice, modellers apply different methods, often chosen ad hoc, to do sensitivity analysis. This paper pursues a systematic approach. It formalizes deterministic and stochastic methods used for sensitivity analysis. Moreover, it presents the numerical algorithms to apply the methods, in particular, an improved version of a Gauss-Quadrature algorithm, applicable to one as well as multidimensional sensitivity analysis. The advantages and disadvantages of different methods and algorithms are discussed as well as their applicability.

Keywords: Sensitivity Analysis, Computational Methods
JEL Classification: C15, C63, D50

*Corresponding Author, e-mail: mennel@zew.de. We would like to thank Kerstin Hesse for helpful comments. Financial Support by the EU project "TranSust.Scan" is gratefully acknowledged.

1 Introduction

In economics as well as other model based sciences, a modeller has to do a sensitivity analysis to show the validity of results of his numerical model simulations. A sensitivity analysis is the study of how the variation in the output of a model (numerical or otherwise) can be apportioned, qualitatively or quantitatively, to different sources of variation in input parameters. It thus allows for an assessment of the robustness of numerical results, as it translates the range (confidence intervals) of fundamental (input) parameters into the model into ranges (confidence intervals) of economic (output) variables. The econometrician Edward Leamer makes it quite clear: "A fragile inference is not worth taking seriously. All scientific disciplines routinely subject their inferences to studies of fragility. Why should economics be different? ... What we need are organized sensitivity analyses." (Leamer, 1985)

In the context of CGE models, we ask whether the choice of basic parameters of the model, e.g. elasticities or time preference parameters, lead to stable equilibrium values of economic variables, e.g. GDP or labor participation. Usually, we refer to the equilibrium of the benchmark scenario. Quite importantly, a sensitivity analysis depends on the existence of equilibria for a sufficient range of parameters: If the model is not solvable for parameter values close to the ones we have chosen as benchmark values, model results are instable and thus worthless.

Basically, there are two methodological approaches to sensitivity analysis: a deterministic and a stochastic approach. Deterministic sensitivity analysis assumes that the tuple of basic parameters is an element of a given subset of all possible parameter choices. It seeks to determine upper and lower bounds on the corresponding subset of economic outcomes of the model. Stochastic sensitivity analysis treats the vector of parameters as a stochastic variable with a given distribution, rendering economic equilibria of the model into stochastic variables. It aims at calculating the first moments of these variables, with the variance indicating the robustness of the results. Both approaches are presented and discussed in section 2.

The choice a modeller has to make in a sensitivity analysis is, however, not only a methodological, but also a numerical one. Sensitivity analysis can involve more or less calculations of equilibria, so that usually there is a trade-off between accuracy and calculation time. This holds already true for a comparison of the deterministic and the stochastic approach, and is particularly relevant for the case of a multidimensional sensitivity analysis. In section 3, we present and discuss different algorithms. In particular, an improved version of an algorithm based on Gauss-Quadrature is developed.

Section 4 presents a simple CGE model in Markusen's (2002) spirit. We conduct a stochastic sensitivity analysis with respect to demand elasticities with

a Monte-Carlo algorithm and Gauss-Quadrature. From the results we see that Monte Carlo is more burdensome computationally. Also, it appears less stable numerically than Gauss Quadrature.

2 Theory of Sensitivity Analysis

2.1 Mathematical Preliminaries

Before conducting a sensitivity analysis, indeed even before implementing a numerical model, the modeller has to understand whether his model is in fact solvable. Proving existence (and sometimes uniqueness) of economic equilibria is a challenge of its own. There exist, however, a set of mathematical theorems that - if applicable - guarantee the existence of solutions. For example, the existence of an equilibrium in a general equilibrium framework is proved by recurring on Kakutani's fix-point theorem (cf. Mas-Collel et al. 1995, chapter 17). In the sequel, we assume that models are (uniquely) solvable for at least some parameter values and discuss under which circumstances a sensitivity analysis is possible.

An equilibrium of a computable general equilibrium (CGE) model takes the mathematical form of a solution to a system of (non-linear) equations

$$F(x^*, a) = 0,$$

where $x^* \in \mathbf{R}^n$ is a vector of (equilibrium) state variables of the economy (such as capital or wage) and $a \in \mathbf{R}^d$ a vector of parameters of the economy (such as demand elasticity or time preference). G is a continuously differentiable function

$$\mathbf{R}^n \times \mathbf{R}^d \rightarrow \mathbf{R},$$

that consists of first order conditions and (budget) constraints. This CGE model will be our standard example. Note, however, that the methods for sensitivity analysis presented can be generally be applied to economic models: The decisive distinction is the one between economic state variables x and basic parameters a .

Sensitivity analysis is concerned with the effect that (minor) changes of basic parameters Δa have on equilibrium state variables Δx^* . This notion will be formalized subsequently. At this point we ask under what circumstances an equilibrium x^{**} exists for a parameter value a' in a neighborhood of a , e.i. $a' \in B_\epsilon(a) = \{\tilde{a} \text{ s.t. } |\tilde{a} - a| < \epsilon\}$. This is an important question: The existence of economic equilibria in a neighborhood of a is the theoretical prerequisite for sensitivity analysis. The implicit function theorem gives a definite answer.

Theorem 1 (Implicit function theorem) *If $\det|\nabla_{x^*} F(x^*, a)| \neq 0$, then there exists an open neighbourhood $\mathcal{U}(a) \subset \mathbf{R}^d$ of a and a continuously differentiable function $h : \mathcal{U}(a) \rightarrow \mathbf{R}^n$ that maps any vector of parameters on the corresponding equilibrium vector.*

Proof. see Rudin (1976), ch. 9., p. 223. ■

Thus we learn that there is some $\epsilon > 0$ so that the existence of equilibria in a ϵ -neighborhood of a is guaranteed whenever F is a regular function at a . Generally speaking, we would expect F to be regular as long as first order conditions and constraints are independent. While it may be difficult to prove the assertion in some cases, it can be checked without problem numerically.

We have formulated the implicit function theorem for the case of CGE models. Similar formulations can be given for partial equilibrium models that are characterized by first and second order conditions and, possibly, additional constraints. In the case of economic optimization problems, the role of the implicit function theorem is taken by the theorem of the maximum (cf. MasCollé et al. (1995), Mathematical Appendix M.K).

So far we have neglected the notion of uniqueness of equilibria. While in principle sensitivity analysis can be conducted in the presence of multiple equilibria as well, uniqueness facilitates the analysis considerably. It is usually ensured by adequate convexity assumptions (cf. MasCollé et al. (1995), ch. 17). In the more general case of multiple equilibria, caution is warranted. In this case,

$$h : \mathcal{U}(a) \rightarrow \mathcal{P}(\mathbf{R}^n)$$

is a correspondence, mapping the vector of parameters into a set of solutions

$$h(a) = \{x_1^*, x_2^*, \dots, x_m^*\}.$$

Ignoring multiplicity can seriously blur a sensitivity analysis whenever a numerical solver 'jumps' from a solution x_j^* to some other solution x_i^* along changes of underlying parameters a . In that case, sensitivity of an equilibrium with respect to the basic parameters can be seriously exaggerated: Instead of following the initial equilibrium x_j^* along a continuous path for changes of a (cf. Judd (1998), p. 179), a numerical discontinuity occurs and the new equilibrium \tilde{x}_i^* is more distant to x_j^* than the correct equilibrium \tilde{x}_j^* . The best provision against such fallacies is the calculation of all equilibria along the path of change; however, this can entail a considerable computational effort. At this point, we will not discuss the issue any further.

2.2 Deterministic Sensitivity Analysis

Sensitivity analysis is sometimes called robustness analysis. This term highlights its motivation: Assuming that we do not know the basic set of parameters exactly, how robust are the economic state variables in an equilibrium with respect to changes in the parameters? The **deterministic approach** to sensitivity analysis states is that there exists one true vector of economic parameters $a^* \in \mathbf{R}^d$, but that -instead of a^* - we only know its neighbourhood \mathcal{A} . Usually, we choose one vector of parameters $\hat{a} \in \mathcal{A}$ and call it the *benchmark scenario*. The point of sensitivity analysis then is to investigate whether equilibria vary considerably across $h(\mathcal{A})$ in comparison to the benchmark equilibrium $h(\hat{a})$.

Mathematically speaking, deterministic sensitivity analysis amounts to a geometric problem: Determine the relation of the volume of the image of \mathcal{A} under h and the size of $h(\hat{a})$, weighted with a scaling factor w_k in each dimension¹

$$\frac{\text{vol}(\text{im}(w * h))}{\|w * h(\hat{a})\|} = \frac{\int_{\mathcal{A}} \sqrt{\det(\partial_{ij}(w * h(a)))} da_1 \dots da_d}{(w * h(\hat{a}), w * h(\hat{a}))},$$

where the vector w specifies the relative weight we want to attach to the different economic variables in equilibrium². The findings of the model are robust whenever the relation is sufficiently small, where the assessment of sufficiency is left to the reader. The formal definition we have just given is a generalization of the more familiar notion of sensitivity analysis in one dimension: E.g. we might ask how big is maximal interval of values of GDP engendered by a model for a given interval of demand elasticities. We will discuss the issue in more detail in the next section where we present the *piecemeal approach* to sensitivity analysis.

2.3 Stochastic Sensitivity Analysis

The **stochastic approach** to sensitivity analysis takes a different view of the basic problem: It treats the vector of basic parameters as a stochastic variable a with values $a \in \mathcal{A} \subset \mathbf{R}^d$. The distribution G of a is given. While somewhat counterintuitive in the first place, the approach is in line with econometric estimations. These do not only produce mean values for parameters such as demand elasticities, but confidence intervals and higher moments for them. Under stochastic sensitivity analysis, h becomes a mapping onto a stochastic variable $x^* = h(a)$ of equilibria.

¹By $'*$ ' we denote component-wise multiplication of vectors.

²We do not have to attach a weight to an economic variable explicitly - instead we assess sensitivity by comparing the effect of changes in a on each dimension of the equilibrium state variable x . Implicitly, however, we will attach some weight to each dimension by accepting a certain outcome of the sensitivity analysis.

We then calculate the mean and the variation of the equilibrium vector x^* :

$$m = E[h(a)] = \int_{\mathcal{A}} h(a)dG, \quad (1)$$

$$v = Var[h(a)] = E[(h(a) - m)^2] = \int_{\mathcal{A}} (h(a) - m)^2 dG. \quad (2)$$

Attaching different weights to different economic variables, the stochastic sensitivity analysis assesses the size of

$$\sum_{k=1}^n w_k \frac{v_k}{m_k},$$

where index k is running over the dimension of x . In words: Given a distribution of basic parameters, we investigate the most likely equilibrium (the mean). We assess its robustness by assessing the relative size of the variance of equilibria with respect to the mean, possibly attaching different weights to different economic variables.

3 Practical Sensitivity Analysis

Having formalized the notion of sensitivity analysis in the preceding section we now present the practical implementation of the (somewhat abstract) concepts. For the sake of clarity in the sequel we assume that all dimensions are weighted equally, thus dropping the vector w from all formulae.

3.1 The Piecemeal Approach

In a **piecemeal approach** to sensitivity analysis, we calculate

$$\Delta = \max_{a_i, a_j \in \{a_1, \dots, a_M\}} |h(a_i) - h(a_j)|$$

for a set of representative parameters $a_i \in \mathcal{A}$. The relation of Δ to the weighted benchmark equilibrium $h(\hat{a})$ is used to assess the sensitivity of the model at equilibrium $h(\hat{a})$. The piecemeal approach is kindred to a deterministic sensitivity analysis. But instead of calculating the volume of \mathcal{A} under h , it focusses on the maximal intervals of economic variables engendered by the set of parameters $a \in \mathcal{A}$.

If the set contains $\underline{a} = \mathbf{argmin}_{a \in \mathcal{A}} \mathbf{h}(a)$ and $\bar{a} = \mathbf{argmax}_{a \in \mathcal{A}} \mathbf{h}(a)$, then the following inequality holds (to facilitate the presentation and without loss of generality, we set $n = 1$):

$$\frac{vol(im(h))}{|h(\hat{a})|} \leq vol(\mathcal{A}) \frac{h(\bar{a}) - h(\underline{a})}{|h(\hat{a})|} = vol(\mathcal{A}) \frac{\Delta}{|h(\hat{a})|}.$$

A piecemeal approach can give a good idea of the sensitivity of the model if the set of parameters $a_i \in \mathcal{A}$ is sufficiently representative.

As an example, let both $n = 1$ and $d = 1$. Then

$$h : [\underline{a}, \bar{a}] \rightarrow [\underline{x}, \bar{x}]$$

for some scalar parameters \underline{a} , \bar{a} , \underline{x} and \bar{x} . If h is monotonously increasing, then

$$\frac{vol(im(h))}{h(\hat{a})} \leq (\bar{a} - \underline{a}) \frac{h(\bar{a}) - h(\underline{a})}{h(\hat{a})}.$$

In the one-dimensional case, the piecemeal approach and the more formal definition of deterministic sensitivity analysis given in the first section virtually coincide. Generally speaking, while our formal definition captures the intention of sensitivity analysis more accurately, the piecemeal approach is more practicable.

3.2 The Monte-Carlo Approach

The **Monte-Carlo** approach is the first of two practical implementations of stochastic sensitivity analysis. Both mean and variance of equilibrium x^* , as defined by equation 1 and 2, are approximated in the following way: Draw a (large) set of realisations $\{a_1, \dots, a_M\}$ from the distribution $G(\mathcal{A})$ and calculate

$$m = E[h(a)] \approx \frac{1}{M} \sum_{i=1}^M h(a_i) = \tilde{m}, \quad (3)$$

$$v = Var[h(a)] \approx \frac{1}{M} \sum_{i=1}^M (h(a_i) - \tilde{m})^2 = \tilde{v}. \quad (4)$$

The sums of the right-hand side converge stochastically to the true values of m and v . Beyond mean and variance of the stochastic variable x^* , we can easily approximate its distribution $h \circ G$. A great disadvantage of the Monte-Carlo approach is that in order to assure convergence, the number of draws M has to be high and thus the approximation is numerically costly. This is a problem in particular when the space of parameters a is high dimensional - the curse of dimensionality drives up the number of necessary draws exponentially (cf. Judd).

3.3 The Gauss-Quadrature Approach

The second way of practically implementing stochastic sensitivity analysis is by **Gauss quadrature** - in fact a numerical method to approximate integrals (cf. Stoer). Remember that we intent to approximate mean and variance, that are defined by integrals of the distribution of basic parameters a . We want to do so using a rather small number L of function evaluations $h(\cdot)$.

Essentially, the Gauss quadrature gives us nodes x_i and weights ω_i to approximate the (one dimensional) integral

$$\int_a^b f(x)\omega(x)dx \approx \sum_{i=1}^L \omega_i f(x_i). \quad (5)$$

In our specific case, we look for nodes a_i and weights g_i to approximate mean, equation 1, and variance, equation 2, of equilibria. Here and in the following, we assume that probability distribution G can be represented by a (continuous) probability density function $g(\cdot)$. While somewhat limiting the applicability of the procedure, we can safely say that all economically sensible distributions should fulfill this assumption.

$$m = \int_{\mathcal{A}} h(a)dG = \int_{\mathcal{A}} h(a)g(a)da \approx \sum_{i=1}^L g_i h(a_i) = \tilde{m}, \quad (6)$$

$$v = \int_{\mathcal{A}} (h(a) - m)^2 dG = \int_{\mathcal{A}} (h(a) - m)^2 g(a) da \approx \sum_{i=1}^L g_i (h(a_i) - \tilde{m})^2 = \tilde{v}, \quad (7)$$

where again, the dimension of economic variables is set to $n = 1$ (we present the generalisation to several variables below).

In the following, we develop a version of Gauss quadrature new to computational economics, in that it builds on *orthogonal polynomials*. While somewhat complicating the straightforward Gauss quadrature algorithm commonly used in economics (cf. Arndt 1996 and DeVuyst and Preckel 1997) conceptually, our approach simplifies the computation of a sensitivity analysis in cases of standard probability distributions, increasing the approximation quality at the same time. This is possible because the optimal nodes x_i turn out to be zeros of orthogonal polynomials. They have to be linearly transformed to fit the respective interval but can otherwise be taken from an existing table. In contrast, in Arndt's (1996) algorithm, the nodes are the solution of a system of non-linear equations (similarly in DeVuyst and Preckel 1997).

We define orthogonality in this context by the scalar product

$$(f_1, f_2)_g = \int_{\mathcal{A}} f_1(a)f_2(a)dG = \int_{\mathcal{A}} f_1(a)f_2(a)g(a)da \quad (8)$$

defines a scalar product $(\cdot, \cdot)_g$. We refer to orthogonality with respect to this scalar product. The following lemma holds:

Lemma 2 (Gram-Schmidt, Weierstrass) *For any scalar product (\cdot, \cdot) on the space of continuous functions $\mathcal{C}([\underline{a}, \bar{a}])$, there is a complete system of orthogonal polynomials $\{p_0, p_1, \dots | (p_i, p_j) = 0, i \neq j\}$.*

Proof.

For any given scalar product, orthogonal polynomials can be constructed from monomials $1, x, x^2, \dots$ by the Gram-Schmidt procedure

$$p_0 \equiv 1 \quad p_i(x) = x^i - \sum_{j=1}^{i-1} \frac{(p_j, x^i)}{(p_j, p_j)} p_j.$$

We thus obtain an infinite sequence of orthogonal polynomials. As for completeness, we know that the polynomials $(p_0(x), p_1(x), \dots, p_n(x))$ span the same linear subspace of the space of continuous functions as the monomials $(1, x, x^2, \dots, x^n)$. Consequently, we can apply Weierstrass' approximation theorem which states that the space of polynomials is dense in the space of continuous functions (cf. Rudin 1976, ch. 7, p. 159) and the completeness of the family of orthogonal polynomials ensues.

■

There are well known examples of orthogonal polynomials, the best known being Legendre, Tchebychev, Laguerre and Hermite polynomials.

Examples of families of orthogonal polynomials

Name	$g(x)$	$[a, b]$	Definition
Legendre	1	$[-1, 1]$	$P_k(x) = \frac{(-1)^k}{2^k k!} \frac{d^k}{dx^k} [(1-x^2)^k]$
Tschebyscheff	$(1-x^2)^{-\frac{1}{2}}$	$[-1, 1]$	$T_k(x) = \cos(k \cos^{-1}(x))$
Laguerre	$\exp(-x)$	$[0, \infty)$	$L_k(x) = \frac{\exp(x)}{k!} \frac{d^k}{dx^k} (x^k \exp(-x))$
Hermite	$\exp(-x^2)$	$(-\infty, \infty)$	$H_k(x) = (-1)^k \exp(x^2) \frac{d^k}{dx^k} (\exp(-x^2))$

The proof of lemma is constructive, so that for any density function $g(a)$ orthogonal polynomials can be constructed from monomials $1, x, x^2, \dots$. For a general distribution probability distribution $G(a)$, their calculation can entail considerable

To proceed, we need one property of orthogonal polynomials.

Lemma 3 *The zeros $\{a_1, a_2, \dots, a_l\}$ of $p_l(a)$ are real and distinct.*

Proof. Stoer and Bulirsch (1990), ch 3.6, p. 173. ■

It is because that they are real and distinct that the zeros of a orthogonal polynomial are a possible choice of nodes for the evaluation of the approximation formula 5. The following theorem shows that they are indeed a good choice.

Theorem 4 *Let $\{a_1, a_2, \dots, a_l\}$ be the zeros of $p_l(a)$ and g_1, \dots, g_l be the solution of the system of linear equations*

$$\sum_{i=1}^n g_i p_k(a_i) = \begin{cases} (p_0, p_0) & : k = 0 \\ 0 & : k = 1, 2, \dots, l - 1 \end{cases}$$

Then $g_i > 0$ for $i = 1, 2, \dots, l$ and

$$\int_{\underline{a}}^{\bar{a}} p(a)g(a)da = \sum_{i=1}^l g_i p(a_i)$$

for all $p \in \Pi_{2l-1} = \ll p_0, \dots, p_{2l-1} \gg$.

Proof. Stoer and Bulirsch (1990), ch 3.6, p. 176. ■

In words: For a given density function $g(a)$ (i.e. probability distribution G), we calculate the zeros a_1, \dots, a_l of the corresponding orthogonal polynomial of degree l . Calculating the weights g_1, \dots, g_l from a suitable system of linear equations, we obtain an integration formula of type 5 that integrates polynomials up to degree $2l-1$ exactly.

Thus, for our purpose of numerical integration, we have to calculate the zeros of orthogonal polynomials and weights corresponding to the probability distribution G with weight function $g(a)$. *However, we have to do so only once for a given G .* While in general the numerical determination of the zeros of orthogonal polynomials for a given distribution may be tricky, in the case of standard probability distributions we have no problem. A look at the table of orthogonal polynomials confirms that for uniform distributions, we can use Legendre polynomials, and Hermite polynomials for normal distributions. This facilitates our task considerably: We can either (easily) calculate the zeros numerically from the

defining formulae for Legendre or Hermite polynomials, or take these from published tables. The next section shows an application of the method presented here to a simple example.

In higher dimensions $n > 1$, integrals can be approximated by product rules, combining one-dimensional nodes and weights

$$\int_{\underline{a}^1}^{\bar{a}^1} \dots \int_{\underline{a}^d}^{\bar{a}^d} f(a^1, \dots, a^d) g^1(a^1) \dots g^d(a^d) da^d \dots da^1 \\ \approx \sum_{i_1=1}^n \dots \sum_{i_d=1}^n g_{i_1}^1 \dots g_{i_d}^d f(a_{i_1}^1, \dots, a_{i_d}^d).$$

We can thus approximate joint distributions of economic parameters instead of analysis the robustness of results with respect to single parameters separately. Note that by specifying probability density functions $g^i(a^i)$ we implicitly assume that the probability distributions are independent. In this case, higher-dimensional Gauss-Quadrature is straightforward - we only have to combine the sums for each dimension. As we will see in the example in the next section, it is in this case that Gauss-Quadrature integration has a great advantage over Monte-Carlo simulations, as the evaluation of nodes increases exponentially with each dimension, making MC simulations simply too expensive.

4 A Simple Example

In this section we present a simple numerical example to demonstrate the use of our approach to sensitivity analysis. We have chosen a CGE framework, the example is implemented in GAMS (Brooke, Kendrick and Meeraus, 1988).

We consider a two-by-two closed economy in the spirit of Markusen (2002). It represents an economy with two commodity goods, X and Y, two factors (capital K and labor L), and one single representative agent. The goods are produced through constant returns to scale production activities which combine primary factor inputs. As usual, we use a balanced equilibrium data set given by the square accounting matrix below. The accounts labeled X and Y in this matrix refer to markets for final commodities; account W correspond to final consumption. The RA account corresponds to the representative agent. It defines both the endowment and expenditures for the model's single representative agent.

Accounting matrix with benchmark flows:

Markets	X	Y	W	RA	Row sum
PX	100		-100		0
PY		100	-100		0
PW			200	-200	0
PL	-40	-60		100	0
PK	-60	-40		100	0
Column sum	0	0	0	0	

The accounts of the matrix do not by themselves completely characterize a general equilibrium framework because they provide a variety of benchmark value shares. Hence, a model formulation additionally relies on assumptions about elasticities of substitution in the various sectors. In our simple model there are three elasticities of substitution: e_{subx} denotes the elasticity of substitution between inputs to X production, e_{suby} the elasticity of substitution between inputs to Y production and e_{subw} the one between inputs to final consumption. In this section we present a sensitivity analysis with respect to these three elasticities which are examples of economic fundamentals and typical parameters for a sensitivity analysis.

Numerically, we formulate the model as a mixed complementarity problem (MCP) using the share form for the specification of functions forms (cost and expenditure functions). The algebraic formulation is implemented in GAMS. For our example, we choose $e_{subx} = e_{suby} = e_{subw} = 0.5$ as starting point and introduce an exogenous labor tax on good X by setting $TAX_L X = 1$.

In the following we compare the two different implementation methods of stochastic sensitivity analysis that were presented in Section 3.2 and in Section 3.3: the Monte-Carlo (MC) analysis and the Gauss-Quadrature (GQ). We calculate both the case of a uniform and a normal distribution in one dimension, as well as the multidimensional case of a uniform distribution.

4.1 One-Dimensional Sensitivity Analysis

First, we assume that the elasticity of substitution e_{subx} is **uniformly distributed** on the interval 0.25 and 0.75 and that the other two elasticities in question are constant and equal to 0.5. Due to the assumption of uniform distribution we choose Legendre polynomials for the Gauss-Quadrature.

For the sensitivity analysis according to the **Monte-Carlo** approach we randomly draw values for e_{subx} from the interval $[0.25, 0.75]$: $e_{subx} = \text{UNIFORM}[0.25, 0.75]$. We calculate the mean and variance of normalized equilibrium variables X, Y and

W according to the following formulas:

$$\begin{aligned}\text{mean}(\text{output}) &= \text{SUM}(i, \text{results}(i, \text{output}))/\text{card}(i); \\ \text{variance}(\text{output}) &= \text{SUM}(i, (\text{sqr}(\text{results}(i, \text{output}) - \text{mean}(\text{output}))))/\text{card}(i);\end{aligned}$$

$\text{card}(i)$ is equal to the number of drawings of esubx from the interval $[0.25, 0.75]$. $\text{results}(i, \text{output})$ denotes the (normalized) equilibrium values for the set $\text{output} = \{X, Y, Z\}$ given the i -th drawing of the elasticity. sqr denotes the square root.

For the sensitivity analysis using **Gauss-Quadrature with Legendre polynomials** we use a MATLAB routine to calculate the zeros of Legendre polynomials, `legendrenodes`, and the associated weights, `legendreweights`. Hereby, we use the recursive formula for polynomials given by the Gram-Schmidt procedure. We calculate the zeros for these polynomials and solve the equation system given in Theorem 4 with respect to the corresponding weights. Zeros and weights are saved in a `gdx` file. Alternatively, schedules with zeros and weights for Legendre polynomials can also be taken from existing tables which are published in the internet or in math books (see for instance Belousov 1962). Once this `gdx` file is created, it can be used to run sensitivity analysis for every CGE model where the parameters are assumed to be uniformly distributed.

In our GAMS routine, the variables `legendrenodes` and `legendreweights` are loaded from the `gdx` file³ and transformed into the variables `grid` and `weights`. We choose `maxdegree` which is the degree of the maximal Legendre polynomial that we want to consider. It also refers to the rows with zeros and weights that are loaded from the `gdx` file. A higher value of `maxdegree` raises the quality of the Legendre approach but is computationally more expensive. Additionally we have to transform `legendrenodes` linearly from the interval $[-1, 1]$ into a new vector `grid` on interval $[a, b] = [0.25, 0.75]$:

$$\text{grid}(i) = (b - a) * (\text{zerosval}('maxdegree', i) + 1)/2 + a.$$

This is due to the fact that standard Legendre polynomials are defined on the interval $[-1, 1]$. $\text{zerosval}('maxdegree', i)$ denotes the i -th zero point for a Legendre polynomial of degree `maxdegree` which is loaded from `legendrenodes`. We calibrate the equilibrium values for X , Y and W by a loop of the model and vary the elasticity esubx across `grid`. We calculate the means and variances of these economic variables by summing up `results`, weighted by `weights` which can be drawn from the `gdx` file directly:

$$\begin{aligned}\text{mean}(\text{output}) &= \text{SUM}(i, \text{weights}(i) * \text{results}(i, \text{output})), \\ \text{variance}(\text{output}) &= \text{SUM}(i, \text{weights}(i) * (\text{sqr}(\text{results}(i, \text{output}) - \text{mean}(\text{output}))))).\end{aligned}$$

³We make the files available on our website.

This implements the formulas (2) and (3) from Section 3. Table 1 reports the results for the Monte-Carlo approach (MC) and the Gauss-Quadrature with Legendre polynomials (GQ). For Monte-Carlo we choose 100, 1,000 and 6,000 drawings of the random variable e_{subx} from the interval $[0.25, 0.75]$. For each drawing the equilibrium values of our model have to be calculated. Hence, computing time is proportional to the number of drawings. For the Gauss-Quadrature we choose the degree of the maximal Legendre polynomial to be equal to 10, 20 and 40. These values correspond to the numbers of zeros of the polynomial function (i.e. to the length of grid) at which the equilibrium problem has to be solved. Thus, they are also equal to the number of model runs. Table 1 shows the means and variances of the variables X, Y and W depending on the chosen sensitivity approach and the number of runs. We can see directly, that the results for the two different approaches converge with increasing number of model evaluations. Gauss-Quadrature converges more quickly to both mean and variance than Monte Carlo. At the same time, it is numerically much more efficient. It reduces calculation time by a factor of at least 50 in comparison to Monte Carlo.

Table 1: Uniformly distributed variable e_{subx}

Means of X,Y and W

Name (Runs)	MC (100)	MC (1000)	MC (6000)	GQ (10)	GQ (20)	GQ (40)
x	-8.990	-9.022	-9.030	-9.036	-9.026	-9.030
Y	7.582	7.575	7.572	7.576	7.568	7.570
w	-1.396	-1.418	-1.421	-1.426	-1.427	-1.421

Variances of X,Y, and W

Name (Runs)	MC (100)	MC (1000)	MC (6000)	GQ (10)	GQ (20)	GQ (40)
x	0.027	0.031	0.031	0.031	0.031	0.031
Y	0.004	0.005	0.004	0.004	0.004	0.005
w	0.014	0.016	0.016	0.018	0.015	0.016

4.2 Multidimensional Sensitivity Analysis

The sensitivity analysis can be conducted in an equivalent way for the elasticities e_{subx} and e_{subw} separately or for all elasticities simultaneously. In this section we assume that they are independently, uniform distributed. For the case of Monte Carlo, the three elasticities of substitution are drawn randomly but independently from given intervals $[a_x, b_x]$, $[a_y, b_y]$ and $[a_w, b_w]$. For the **Gauss-Quadrature** the Legendre nodes are transformed three times from the `gdx` file. In both cases, the equilibrium values are calculated by three loops over the three grids. Mean and variance can be derived by weighting the results with the associated weights:

$$\begin{aligned} \text{mean}(\text{output}) &= \\ & \text{SUM}(i, \text{SUM}(ii, (\text{SUM}(iii, \text{weights}_x(i) * \text{weights}_y(ii) * \\ & \quad \text{weights}_w(iii) * \text{results}(i, ii, iii, \text{output})))))) \\ \\ \text{variance}(\text{output}) &= \\ & \text{SUM}(i, \text{SUM}(ii, \text{SUM}(iii, \text{weights}_x(i) * \text{weights}_y(ii) * \\ & \text{weights}_w(iii) * \text{sqr}(\text{results}(i, ii, iii, \text{output}) - \text{mean}(\text{output})))))) \quad . \end{aligned}$$

$\text{results}(i, ii, iii, \text{output})$ denotes the equilibrium values for the set $\text{output} = \{X, Y, Z\}$ given the (i)-th zero of the chosen Legendre polynomial for e_{subx} , the (ii)-th zero for the Legendre polynomial of e_{suby} and the (iii)-th zero of the one for e_{subw} . weights_x , weights_y and weights_w refer to the Legendre weights belonging to the elasticities e_{subx} , e_{suby} , and e_{subw} . Please note that different values for maxdegree_x , maxdegree_y and maxdegree_w can be chosen corresponding to different degrees of polynomials.

In our example run, we set $[a_x, b_x] = [a_y, b_y] = [a_w, b_w] = [0.25, 0.75]$ and $\text{maxdegree} := \text{maxdegree}_x = \text{maxdegree}_y = \text{maxdegree}_w$. To test the Monte-Carlo (MC) approach we choose $10 \times 10 \times 10$, $20 \times 20 \times 20$, $30 \times 30 \times 30$ and $40 \times 40 \times 40$ independent drawings from $[0.25, 0.75] \times [0.25, 0.75] \times [0.25, 0.75]$ which correspond to 1,000, 8,000, 27,000 and 64,000 model runs. To apply the Gauss-Quadrature (GQ) approach we study the cases $\text{maxdegree} = 10, = 20, = 30$ and $= 40$ with 1,000, 8,000, 27,000 and 64,000 model evaluations. The results are reported in Table 2.

We notice the rapid convergence for the Gauss-Quadrature. Using this approach, the approximated mean does not change anymore after 8,000 runs. In contrast, there is no trend observable from the results of the Monte Carlo approach, even after 64,000 runs. Besides, the values for the means and variances differ considerably from the values derived by Gauss-Quadrature. It can be suggested that the number of model evaluations for Monte-Carlo is still far too low to

achieve acceptable results, despite the high computational burden.

Table 2: Uniformly distributed variable esubx, esuby and esubw

Means of X,Y and W

Name (Runs)	MC (1,000)	MC (8,000)	MC (27,000)	MC (64,000)	GQ (1,000)	GQ (8,000)	GQ (27,000)	GQ (64,000)
x	-9.852	-11.189	-10.901	-11.165	-9.002	-9.001	-9.001	-9.001
y	8.435	9.574	9.323	9.680	7.571	7.573	7.573	7.573
w	-1.475	-1.702	-1.652	-1.628	-1.404	-1.406	-1.406	-1.406

Variances of X,Y, and W

Name (Runs)	MC (1,000)	MC (8,000)	MC (27,000)	MC (64,000)	GQ (1,000)	GQ (8,000)	GQ (27,000)	GQ (64,000)
x	5.719	4.570	4.570	3.462	5.435	5.438	5.441	5.440
y	5.605	3.758	4.358	3.324	5.331	5.318	5.318	5.317
w	0.068	0.095	0.083	0.071	0.062	0.062	0.062	0.062

4.3 Sensitivity Analysis with Normal Distribution

Similarly, sensitivity analysis can be conducted for normally distributed parameters. In our model we consider the case of a log-normal distributed elasticity esubx. Note that we cannot assume a normal distribution for an the elasticity because it is not defined for negative numbers. Instead, we assume that $\ln(\text{esubx})$ is normal distributed with mean 0.5 and variance 0.4. For the **Monte-Carlo** simulation, random draws are generated by

$$\text{esubx} = \exp(\text{NORMAL}[0.5, 0.4]).$$

The **Gauss-Quadrature** method used here is based on Hermite polynomials corresponding to the normal distribution. We calculate the zeros and weights of the Hermite polynomials again in a separate MATLAB program and store them in a gdx-files as hermitenodes and hermiteweights. It is also possible to take these

values from published tables like e.g. from Greenwood and Miller (1948). Then, we proceed in the same way as in the case of uniform distribution. The results are reported in Table 3. Again, the values imply convergence of the Gauss-Quadrature results. But also an additional disadvantage of the Monte Carlo approach becomes visible. Our Monte-Carlo simulation was only feasible for small numbers of runs up to 628. After that, the random generator of GAMS started to produce repeatedly values of $exubx$ for which our program failed to find equilibrium solutions. Dropping these values from our sample produces an unknown bias to the MC estimations. Unfortunately, this procedure is very common in practice. In contrast, the Gauss-Quadrature is numerically much more stable.

Table 3: Log-normal distributed variable $esubx$ with $\sigma^2 = 0.4$, and $\mu = 0.5$

Means of X,Y and W

Name (Runs)	MC (100)	MC (500)	MC (628)	GQ (10)	GQ (20)	GQ (40)
X	-9.804	-9.844	-9.842	-9.835	-9.842	-9.853
Y	7.559	7.572	7.572	7.571	7.578	7.592
W	-1.878	-1.901	-1.900	-1.883	-1.891	-1.910

Variances of X,Y and W

Name (Runs)	MC (100)	MC (500)	MC (628)	GQ (10)	GQ (20)	GQ (40)
X	0.069	0.075	0.070	0.178	0.168	0.168
Y	0.004	0.004	0.003	0.006	0.007	0.005
W	0.019	0.022	0.020	0.051	0.042	0.052

5 Conclusion

Due to its general importance in economic modelling, sensitivity analysis merits a systematic understanding by economic modellers. This paper hopes to contribute to such an understanding and to serve as a guide in applying the appropriate algorithm. Dimensionality of sensitivity analysis (the number of parameters that are varied simultaneously) makes an important difference. Moreover, we argue that generally stochastic sensitivity analysis gives more and better insights than deterministic sensitivity analysis. However, it is also more burdensome computationally. Comparing stochastic methods, the paper shows that Monte-Carlo methods are easily applicable, but computationally expensive. Gauss-Quadrature methods reduce the computational burden and thus are suitable for higher dimensional problems. In an application of both methods Gauss-Quadrature turns out to be more stable than Monte-Carlo, which can be explained by the fact that it avoids the extreme parts of the distribution.

References

- [1] Arndt, C. (1996), "An Introduction to Systematic Sensitivity Analysis via Gaussian Quadrature", GTAP Technical Paper No. 2, Purdue University
- [2] Belousov, S. L. (1962), "Tables of normalized associated Legendre polynomials", Mathematical tables series Vol. 18, Pergamon Press
- [3] Brooke, A., Kendrick, D. and A. Meeraus (1988), "GAMS: A User's Guide", The Scientific Press
- [4] DeVuyst, E.A. and P.V. Preckel (1997), "Sensitivity Analysis Revisited: A Quadrature-Based Approach", Journal of Policy Modeling 19, 175-185.
- [5] Greenwood, R. E. and J. J. Miller (1948), "Zeros of the Hermite polynomials and weights for Gauss' mechanical quadrature formula", Bull. Amer. Math. Soc. 54 (8), 765-769
- [6] Judd, K. (1998), "Numerical Methods in Economics", MIT Press
- [7] Leamer, E. (1985), "Sensitivity Analyses would Help", American Economic Review 75 (3), 308-313
- [8] Mas-Colell, A., Whinston, M. D. and J. R. Green (1995), "Microeconomic Theory", Oxford University Press
- [9] Markusen, J. A. (2002), "General-Equilibrium Modeling using GAMS and MPS/GE", mimeo, University of Colorado
- [10] Rudin, W. (1976), "Principles of Mathematical Analysis", McGraw-Hill
- [11] Stoer, J. and R. Bulirsch (1990) "Introduction to Numerical Analysis", Springer