

ECO

CBM
R




UNIVERSITEIT
BRABANT

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS

7626
1990
452



* C I N O 1 1 5 3 *



DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM



STADT
D'Y. O'HEERK
TILBURG

STRONGLY REGULAR GRAPHS INDUCED BY
POLARITIES OF SYMMETRIC DESIGNS

W.H. Haemers
D.G. Higman, S.A. Hobart

FEW 452

198
2009

STRONGLY REGULAR GRAPHS INDUCED BY POLARITIES OF SYMMETRIC DESIGNS

by

W.H. HAEMERS,

Tilburg University, Tilburg, The Netherlands,

D.G. HIGMAN,

University of Michigan, Ann Arbor, Michigan, USA,

and

S.A. HOBART,

University of Wyoming, Laramie, Wyoming, USA.

ABSTRACT. We study symmetric designs with a polarity for which the graph induced by the absolute points and the graph induced by the non-absolute points both are strongly regular. The main result is that all parameters can be expressed in terms of only one parameter.

INTRODUCTION.

Throughout, we shall assume that A is the $(0,1)$ incidence matrix of a symmetric $2-(v,k,\lambda)$ design D with a polarity with $v_1 (\neq 0)$ absolute points and $v_2 = v - v_1 (\neq 0)$ nonabsolute points. So A takes the form:

$$A = \begin{bmatrix} A_1 & C \\ C^T & A_2 \end{bmatrix},$$

where A_1 and A_2 are symmetric matrices, A_1 has ones on the diagonal and A_2 has zeros on the diagonal. Thus $A_1 - I$ and A_2 are adjacency matrices of graphs Γ_1 and Γ_2 , say. We call Γ_1 and Γ_2 the graphs *induced* by the polarity of D . The above decomposition of A is called the *polar decomposition*. A polar decomposition is *regular* if Γ_1 and Γ_2 are regular, and *strongly regular* if Γ_1 and Γ_2 are strongly regular, complete or empty. The incidence structure of absolute points and nonabsolute blocks, given by the matrix C , is called the *polar structure* of the design D .

We assume that D is nontrivial, i.e. $0 < \lambda < k-1 < v-2$. Polar decompositions of trivial designs are obvious and not very interesting.

In the present paper we study strongly regular polar decompositions. It is, in a certain sense, an extension to a paper on strongly regular graphs with strongly regular decompositions by the first two authors [1]. The subject is motivated by the following example. Let D be the design of points and hyperplanes of $PG(n,4)$ ($n \geq 2$). Then the graphs induced by the unitary polarity are strongly regular (see for instance [3]). The parameters for this example are:

$$v = \frac{4^{n+1}-1}{3}, \quad k = \frac{4^n-1}{3}, \quad \lambda = \frac{4^{n-1}-1}{3}, \quad v_1 = \frac{2^{2n+1}-(-2)^{n-1}}{3}.$$

We shall prove that all parameters of a strongly regular polar decomposition can be expressed in terms of only one parameter a . If a is a power of -2 we find the values above, but unfortunately, other values of a for which no examples are known, remain feasible.

For strongly regular Γ_i ($i = 1, 2$) the parameters are denoted by v_i, k'_i, λ_i and μ_i , and the eigenvalues of A_i by k_i, ρ_i and σ_i , such that $|\rho_i| \geq |\sigma_i|$. Note that $k_1 = k'_1 + 1$ and $k_2 = k'_2$. The multiplicity of σ_i is denoted by φ_i . If Γ_i is complete or empty we put $\varphi_i = v_i - 1$ (ρ_i disappears).

PRELIMINARIES.

The following three Lemmas are the analogues of 2.2, 2.4 and 2.5 of [1]. Since the proofs are the same, we omit them here.

Lemma 1. Suppose Γ_1 is regular of degree $k'_1 = k_1 - 1$. Then

$$\left| \frac{k_1 v - v_1 k}{v - v_1} \right| \leq \sqrt{k - \lambda}.$$

Equality holds if and only if the polar decomposition is regular.

Note that for the polar decomposition to be regular $k - \lambda$ (the order of D) must be a square.

For a regular polar decomposition we define k_2 to be the degree of Γ_2 and

$$(1) \quad a = \frac{k_1 v - v_1 k}{v - v_1}.$$

Then we easily have

$$(2) \quad a = k_1 + k_2 - k = \pm \sqrt{k - \lambda}, \quad |a| \geq 2.$$

So $a, -a$ and k are the eigenvalues of A with multiplicities φ (say), $v - 1 - \varphi$ and 1, respectively.

Lemma 2. Suppose A_1 has eigenvalues k_1, σ_1 and ρ_1 ($|\sigma_1| \leq |\rho_1|$) with multiplicities 1, φ_1 and $v_1 - 1 - \varphi_1$ ($1 \leq \varphi_1 \leq v_1 - 1$), respectively. Let the

polar decomposition be regular. Then A_2 has eigenvalues $k_2, -\sigma_1, -\rho_1, a, -a$ with multiplicities $1, \varphi_1, v_1-1-\varphi_1, \varphi-v_1$ and $v_2-\varphi$, respectively. Also $|\sigma_1| \leq |\rho_1| \leq |a|$, and $|\sigma_1| \neq |a|$.

Lemma 3. With the hypotheses of Lemma 2, the polar decomposition is strongly regular if and only if one of the following occurs.

- i. $v_1 = v_2 = \varphi$,
- ii. $\rho_1 = a, v_1 = \varphi$,
- iii. $\rho_1 = -a, v_2 = \varphi$,

It is easy to see that if ii occurs for a matrix A , then iii occurs for the complementary case, that is for the design with incidence matrix (J denotes the all-one matrix).

$$J = \begin{bmatrix} A_2 & C^T \\ C & A_1 \end{bmatrix},$$

Note that Lemma 3 also applies if Γ_1 is complete or empty (the conditions $\rho_1 = \pm a$ are meaningless in this case). Next we show that case i does not occur.

Lemma 4. A strongly regular polar decomposition with $v_1 = v_2 = \varphi$ does not exist.

Proof. Assume $v \geq 2k$ (otherwise consider the complement). Trace $A = v_1 = \varphi = k + \varphi a - (v-\varphi-1)a$ implies $k = v_1 - a$. Hence, by (1) and (2), $k_1 = k_2 = v_1/2 = v_2/2$. Therefore, since Γ_2 is strongly regular, $k_2(k_2 - \lambda_2 - 1) = \mu_2(v_2 - k_2 - 1) = \mu_2(k_2 - 1)$. This implies $\mu_2 = 0 \pmod{k_2}$, hence Γ_2 is complete bipartite, so $\rho_2 = -k_2$. This is impossible, since then, by Lemma 2, $|a| \geq |\rho_2| = k_2 = v_2/2 = v/4$, which is a contradiction to $a^2 < k \leq v/2$ and $|a| \geq 2$. \square

MAIN THEOREM.

Theorem 5. If D has a strongly regular polar decomposition then $a \equiv 4 \pmod{6}$, and, up to taking complements, all other parameters are expressible in terms of a as follows:

$$v = \frac{1}{3}(16a^2 - 1), \quad k = \frac{1}{3}(4a^2 - 1), \quad \lambda = \frac{1}{3}(a^2 - 1),$$

$$v_1 = \varphi = \frac{1}{3}(2a+1)(4a-1), \quad v_2 = \frac{2}{3}a(4a-1),$$

$$k'_1 = k_1 - 1 = \frac{2}{3}(a-1)(a+2), \quad k'_2 = k_2 = \frac{1}{3}a(2a-1),$$

$$\rho_1 = -\rho_2 = a, \quad \sigma_2 = -\sigma_1 = \frac{a}{2},$$

$$\varphi_1 = \varphi_2 = \frac{8}{9}(a-1)(2a+1),$$

$$\mu_1 = \frac{1}{6}(a-1)(a+2), \quad \lambda_1 = \frac{1}{6}(a^2 + 4a - 2),$$

$$\mu_2 = \frac{1}{6}a(a+2), \quad \lambda_2 = \frac{1}{6}a(a-1).$$

Proof. We may restrict ourselves to case ii of Lemma 3. By this result we can express all parameters in terms of k and a only. We have $\lambda = k - a^2$, and hence, using $\lambda(v-1) = k(k-1)$:

$$v = \frac{k^2 - a^2}{k - a^2}.$$

Trace $A = v_1 = \varphi = k + \varphi a - (v-1-\varphi)a$ yields

$$v_1 = \frac{k(k+a)(a-1)}{(k-a^2)(2a-1)}, \quad v_2 = \frac{a(k+a)(k+1-2a)}{(k-a^2)(2a-1)}.$$

Next, using (1) and (2), we find

$$k_2 = \frac{ak}{2a-1}, \quad v_2 - k_2 - 1 = \frac{k(a+1)(a-1)^2}{(k-a^2)(2a-1)}.$$

Since $k-a^2 = \lambda > 0$ and $|a| \geq 2$ it follows that $0 < k_2 < v_2 - 1$, so Γ_2 is not the empty graph or the complete graph. Moreover, $2a-1$ divides k . Define $x = k/(2a-1)$, then $k_2 = ax$ and a and x have the same sign. By use of $\rho_2 = -a$ (by Lemma 2 and 3) and $(k_2 - \rho_2)(k_2 - \sigma_2) = (k_2 + \rho_2 \sigma_2)v_2$ (because Γ_2 is strongly regular) we find after some computation:

$$\sigma_2 = \frac{a(ax+a-2x)}{2ax-x-1}.$$

Define

$$d = 4\sigma_2 - 2a + 3 = \frac{4a^2 + 2a - 3x - 3}{2ax - x - 1},$$

then $d \geq 0$, since $d \leq -1$ would imply $(2a-1)(a+1) \leq x(1-a) < 0$, which contradicts $|a| \geq 2$. Moreover, using $2ax - x - 1 = k - 1 \geq a^2$, we have

$$d \leq 4 + \frac{2}{a} - \frac{3(x+1)}{2ax-x-1} < \begin{cases} 4 + \frac{2}{a} \leq 5 & \text{if } a \text{ and } x \text{ are positive,} \\ 4 - \frac{3}{2a-1} < 5 & \text{if } a \text{ and } x \text{ are negative.} \end{cases}$$

Since d is odd we can conclude that $d = 1$ or $d = 3$. If $d = 1$, then $x = 2a-1$ and $v_1 = (2a-1)(4a^2 - 3a + 1)/(3a-1)$, which is no integer. Hence $d = 3$, so

$$x = \frac{2a+1}{3}, \quad \sigma_2 = \frac{a}{2}, \quad a = 4 \pmod{6}.$$

All other parameter now follow in a straightforward manner. □

If $a = (-2)^{n-1}$ we find the parameters of the example given in the introduction. No other examples are known. This leaves $a = 10$ (D is a $2-(533, 133, 33)$ design) as the smallest unsolved case. By Theorem 5, only for $a = -2$ one of the graphs induced by the polarity is complete or empty. For $a = -2$, however, the structure is unique. So $PG(2, 4)$ with its unitary polarity and the complement provide the only strongly regular polar decomposition for which one of the induced graphs is empty or complete. In this case the polar structure is $AG(2, 3)$, the affine plane of order 3. If $|a| > 2$, it can be shown (by similar, but less complicated, arguments as for Theorem 2.8 of [1]) that the polar structure is a strongly regular design (see [2] for definition and notation) with parameters

$$(3) \quad (n_1, S_1, a_1, b_1, n_2, S_2, a_2, b_2) = (v_1, k-k_1, \lambda-2-\lambda_1, \lambda-\mu_1, v_2, k-k_2, \lambda-\lambda_2, \lambda-\mu_2),$$

and that, conversely, a strongly regular design whose parameters satisfy (3) with v_1, k, k_1 etc. as given in Theorem 5, is the polar structure of some symmetric design with a strongly regular polar decomposition.

A symmetric design with a polarity can be seen as a strongly regular graph for which loops are admitted. More precisely, if we allow loops in graphs, a strongly regular graph is a simple strongly regular graph or a symmetric design with a polarity. In this more general setting, a strongly regular graph with strongly regular decomposition either belongs to the case without loops treated in [1], or is a design with a strongly regular polar decomposition. This can be proved as follows. For a strongly regular decomposition of a symmetric design with a polarity it follows by eigenvalue arguments as in Lemma 3, that if Γ_1 is a symmetric design with a polarity, then so is Γ_2 . This, however is impossible by Rahilly [4]. Thus both Γ_1 and Γ_2 must be strongly regular graphs without loops, so we have a strongly regular polar decomposition.

REFERENCES

- [1] W.H. Haemers and D.G. Higman, Strongly regular graphs with strongly regular decomposition, *Linear Algebra Appl.* 114/115: 379-398 (1989).
- [2] D.G. Higman, Strongly regular designs and coherent configurations of type $\begin{pmatrix} 3 & 2 \\ & 3 \end{pmatrix}$, *European J. Combin.* 9: 411-422 (1988).
- [3] X.L. Hubaut, Strongly regular graphs, *Discrete Math.* 13: 357-381 (1975).
- [4] A. Rahilly, Divisions of symmetric designs into two parts, *Graphs and Combinatorics* 4: 67-73 (1988).

IN 1989 REEDS VERSCHENEN

- 368 Ed Nijssen, Will Reijnders
"Macht als strategisch en tactisch marketinginstrument binnen de distributieketen"
- 369 Raymond Gradus
Optimal dynamic taxation with respect to firms
- 370 Theo Nijman
The optimal choice of controls and pre-experimental observations
- 371 Robert P. Gilles, Pieter H.M. Ruys
Relational constraints in coalition formation
- 372 F.A. van der Duyn Schouten, S.G. Vanneste
Analysis and computation of (n,N) -strategies for maintenance of a two-component system
- 373 Drs. R. Hamers, Drs. P. Verstappen
Het company ranking model: a means for evaluating the competition
- 374 Rommert J. Casimir
Infogame Final Report
- 375 Christian B. Mulder
Efficient and inefficient institutional arrangements between governments and trade unions; an explanation of high unemployment, corporatism and union bashing
- 376 Marno Verbeek
On the estimation of a fixed effects model with selective non-response
- 377 J. Engwerda
Admissible target paths in economic models
- 378 Jack P.C. Kleijnen and Nabil Adams
Pseudorandom number generation on supercomputers
- 379 J.P.C. Blanc
The power-series algorithm applied to the shortest-queue model
- 380 Prof. Dr. Robert Bannink
Management's information needs and the definition of costs, with special regard to the cost of interest
- 381 Bert Bettonvil
Sequential bifurcation: the design of a factor screening method
- 382 Bert Bettonvil
Sequential bifurcation for observations with random errors

- 383 Harold Houba and Hans Kremers
Correction of the material balance equation in dynamic input-output models
- 384 T.M. Doup, A.H. van den Elzen, A.J.J. Talman
Homotopy interpretation of price adjustment processes
- 385 Drs. R.T. Frambach, Prof. Dr. W.H.J. de Freytas
Technologische ontwikkeling en marketing. Een oriënterende beschouwing
- 386 A.L.P.M. Hendriks, R.M.J. Heuts, L.G. Hoving
Comparison of automatic monitoring systems in automatic forecasting
- 387 Drs. J.G.L.M. Willems
Enkele opmerkingen over het inversificerend gedrag van multinationale ondernemingen
- 388 Jack P.C. Kleijnen and Ben Annink
Pseudorandom number generators revisited
- 389 Dr. G.W.J. Hendrikse
Speltheorie en strategisch management
- 390 Dr. A.W.A. Boot en Dr. M.F.C.M. Wijn
Liquiditeit, insolventie en vermogensstructuur
- 391 Antoon van den Elzen, Gerard van der Laan
Price adjustment in a two-country model
- 392 Martin F.C.M. Wijn, Emanuel J. Bijnen
Prediction of failure in industry
An analysis of income statements
- 393 Dr. S.C.W. Eijffinger and Drs. A.P.D. Gruijters
On the short term objectives of daily intervention by the Deutsche Bundesbank and the Federal Reserve System in the U.S. Dollar - Deutsche Mark exchange market
- 394 Dr. S.C.W. Eijffinger and Drs. A.P.D. Gruijters
On the effectiveness of daily interventions by the Deutsche Bundesbank and the Federal Reserve System in the U.S. Dollar - Deutsche Mark exchange market
- 395 A.E.M. Meijer and J.W.A. Vingerhoets
Structural adjustment and diversification in mineral exporting developing countries
- 396 R. Gradus
About Tobin's marginal and average q
A Note
- 397 Jacob C. Engwerda
On the existence of a positive definite solution of the matrix equation $X + A^T X^{-1} A = I$

- 398 Paul C. van Batenburg and J. Kriens
Bayesian discovery sampling: a simple model of Bayesian inference in auditing
- 399 Hans Kremers and Dolf Talman
Solving the nonlinear complementarity problem
- 400 Raymond Gradus
Optimal dynamic taxation, savings and investment
- 401 W.H. Haemers
Regular two-graphs and extensions of partial geometries
- 402 Jack P.C. Kleijnen, Ben Annink
Supercomputers, Monte Carlo simulation and regression analysis
- 403 Ruud T. Frambach, Ed J. Nijssen, William H.J. Freytas
Technologie, Strategisch management en marketing
- 404 Theo Nijman
A natural approach to optimal forecasting in case of preliminary observations
- 405 Harry Barkema
An empirical test of Holmström's principal-agent model that tax and signally hypotheses explicitly into account
- 406 Drs. W.J. van Braband
De begrotingsvoorbereiding bij het Rijk
- 407 Marco Wilke
Societal bargaining and stability
- 408 Willem van Groenendaal and Aart de Zeeuw
Control, coordination and conflict on international commodity markets
- 409 Prof. Dr. W. de Freytas, Drs. L. Arts
Tourism to Curacao: a new deal based on visitors' experiences
- 410 Drs. C.H. Veld
The use of the implied standard deviation as a predictor of future stock price variability: a review of empirical tests
- 411 Drs. J.C. Caanen en Dr. E.N. Kertzman
Inflatieneutrale belastingheffing van ondernemingen
- 412 Prof. Dr. B.B. van der Genugten
A weak law of large numbers for m -dependent random variables with unbounded m
- 413 R.M.J. Heuts, H.P. Seidel, W.J. Selen
A comparison of two lot sizing-sequencing heuristics for the process industry

- 414 C.B. Mulder en A.B.T.M. van Schaik
Een nieuwe kijk op structuurwerkloosheid
- 415 Drs. Ch. Caanen
De hefboomwerking en de vermogens- en voorraadaf trek
- 416 Guido W. Imbens
Duration models with time-varying coefficients
- 417 Guido W. Imbens
Efficient estimation of choice-based sample models with the method of moments
- 418 Harry H. Tigelaar
On monotone linear operators on linear spaces of square matrices

IN 1990 REEDS VERSCHENEN

- 419 Bertrand Melenberg, Rob Alessie
A method to construct moments in the multi-good life cycle consumption model
- 420 J. Kriens
On the differentiability of the set of efficient (μ, σ^2) combinations in the Markowitz portfolio selection method
- 421 Steffen Jørgensen, Peter M. Kort
Optimal dynamic investment policies under concave-convex adjustment costs
- 422 J.P.C. Blanc
Cyclic polling systems: limited service versus Bernoulli schedules
- 423 M.H.C. Paardekooper
Parallel normreducing transformations for the algebraic eigenvalue problem
- 424 Hans Gremmen
On the political (ir)relevance of classical customs union theory
- 425 Ed Nijssen
Marketingstrategie in Machtspectief
- 426 Jack P.C. Kleijnen
Regression Metamodels for Simulation with Common Random Numbers: Comparison of Techniques
- 427 Harry H. Tigelaar
The correlation structure of stationary bilinear processes
- 428 Drs. C.H. Veld en Drs. A.H.F. Verboven
De waardering van aandelenwarrants en langlopende call-opties
- 429 Theo van de Klundert en Anton B. van Schaik
Liquidity Constraints and the Keynesian Corridor
- 430 Gert Nieuwenhuis
Central limit theorems for sequences with $m(n)$ -dependent main part.
- 431 Hans J. Gremmen
Macro-Economic Implications of Profit Optimizing Investment Behaviour
- 432 J.M. Schumacher
System-Theoretic Trends in Econometrics
- 433 Peter M. Kort, Paul M.J.J. van Loon, Mikuláš Luptacik
Optimal Dynamic Environmental Policies of a Profit Maximizing Firm
- 434 Raymond Gradus
Optimal Dynamic Profit Taxation: The Derivation of Feedback Stackelberg Equilibria

- 435 Jack P.C. Kleijnen
Statistics and Deterministic Simulation Models: Why Not?
- 436 M.J.G. van Eijs, R.J.M. Heuts, J.P.C. Kleijnen
Analysis and comparison of two strategies for multi-item inventory systems with joint replenishment costs
- 437 Jan A. Weststrate
Waiting times in a two-queue model with exhaustive and Bernoulli service
- 438 Alfons Daems
Typologie van non-profit organisaties
- 439 Drs. C.H. Veld en Drs. J. Grazell
Motieven voor de uitgifte van converteerbare obligatieleningen en warrantobligatieleningen
- 440 Jack P.C. Kleijnen
Sensitivity analysis of simulation experiments: regression analysis and statistical design
- 441 C.H. Veld en A.H.F. Verboven
De waardering van conversierechten van Nederlandse converteerbare obligaties
- 442 Drs. C.H. Veld en Drs. P.J.W. Duffhues
Verslaggevingsaspecten van aandelenwarrants
- 443 Jack P.C. Kleijnen and Ben Annink
Vector computers, Monte Carlo simulation, and regression analysis: an introduction
- 444 Alfons Daems
"Non-market failures": Imperfecties in de budgetsector
- 445 J.P.C. Blanc
The power-series algorithm applied to cyclic polling systems
- 446 L.W.G. Strijbosch and R.M.J. Heuts
Modelling (s,Q) inventory systems: parametric versus non-parametric approximations for the lead time demand distribution
- 447 Jack P.C. Kleijnen
Supercomputers for Monte Carlo simulation: cross-validation versus Rao's test in multivariate regression
- 448 Jack P.C. Kleijnen, Greet van Ham and Jan Rotmans
Techniques for sensitivity analysis of simulation models: a case study of the CO₂ greenhouse effect
- 449 Harrie A.A. Verbon and Marijn J.M. Verhoeven
Decision-making on pension schemes: expectation-formation under demographic change

- 450 Drs. W. Reijnders en Drs. P. Verstappen
Logistiek management marketinginstrument van de jaren negentig
- 451 Alfons J. Daems
Budgeting the non-profit organization
An agency theoretic approach

Bibliotheek K. U. Brabant



17 000 01066413 5